

Crystal structure of catena-poly[[diaquacobalt(II)]-bis[μ -5-(4-carboxylatophenyl)picolinate]- $\kappa^3 N, O^2: O^5; \kappa^3 O^5: N, O^2$ -[diaquacobalt(II)]- μ -1-[4-(1H-imidazol-1-yl)phenyl]-1H-imidazole- $\kappa^2 N^3: N^3'$]

Guo-Wang Xu,^a Ye-Nan Wang,^b Hai-Bing Wang^b and Zhong-Long Wang^{a*}

^aCollege of Science, China Three Gorges University, Yichang 443002, People's Republic of China, and ^bCollege of Mechanical and Power Engineering, China Three Gorges University, Yichang 443002, People's Republic of China. *Correspondence e-mail: wzlsanxia@163.com

Received 26 May 2015; accepted 25 June 2015

Edited by D.-J. Xu, Zhejiang University (Yuquan Campus), China


The asymmetric unit of the title polymeric Co^{II} complex, [Co₂(C₁₃H₇NO₄)₂(C₁₂H₁₀N₄)(H₂O)₄]_n, contains a Co^{II} cation, a 5-(4-carboxylatophenyl)picolinate dianion, two coordination water molecules and half of 1-[4-(1H-imidazol-1-yl)phenyl]-1H-imidazole ligand. The Co^{II} cation is coordinated by two picolinate dianions, two water molecules and one 1-[4-(1H-imidazol-1-yl)phenyl]-1H-imidazole molecule in a distorted N₂O₄ octahedral coordination geometry. The two picolinate dianions are related by an inversion centre and link two Co^{II} cations, forming a binuclear unit, which is further bridged by the imidazole molecules, located about an inversion centre, into the polymeric chain propagating along the [111] direction. In the crystal, the three-dimensional supramolecular architecture is constructed by O—H...O hydrogen bonds between the coordinating water molecules and the non-coordinating carboxylate O atoms of adjacent polymeric chains.

Keywords: crystal structure; 5-(4-carboxylphenyl)picolinate; 1-[4-(1H-imidazol-1-yl)phenyl]-1H-imidazole; one-dimensional coordination polymer; cobalt(II) complex.

CCDC reference: 1062328

1. Related literature

For the structure of a related 5-(4-carboxylphenyl)picolinate complex, see: Meng *et al.* (2012). For a related 1,4-bis(1-imidazolyl)benzene compound, see: Li *et al.* (2009).



2. Experimental

2.1. Crystal data

[Co ₂ (C ₁₃ H ₇ NO ₄) ₂ (C ₁₂ H ₁₀ N ₄)(H ₂ O) ₄] _n	$\beta = 79.90 (2)^\circ$
$M_r = 441.28$	$\gamma = 64.080 (15)^\circ$
Triclinic, $P\bar{1}$	$V = 895.1 (7) \text{ \AA}^3$
$a = 7.055 (3) \text{ \AA}$	$Z = 2$
$b = 7.190 (3) \text{ \AA}$	Mo $K\alpha$ radiation
$c = 20.038 (10) \text{ \AA}$	$\mu = 1.00 \text{ mm}^{-1}$
$\alpha = 80.25 (2)^\circ$	$T = 293 \text{ K}$
	$0.20 \times 0.20 \times 0.17 \text{ mm}$

2.2. Data collection

Bruker SMART 1000 CCD diffractometer	9552 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2001)	4104 independent reflections
$T_{\min} = 0.825$, $T_{\max} = 0.848$	3636 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.045$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.090$	$\Delta\rho_{\text{max}} = 0.34 \text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\text{min}} = -0.41 \text{ e \AA}^{-3}$
4104 reflections	
274 parameters	
4 restraints	

Table 1
Selected bond lengths (Å).

Co1—N1 ⁱ	2.1403 (18)	Co1—O4 ⁱ	2.1028 (16)
Co1—N2	2.0815 (18)	Co1—O5	2.0773 (18)
Co1—O2	2.1575 (16)	Co1—O6	2.0889 (19)

Symmetry code: (i) $-x, -y + 1, -z + 1$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O5—H5A...O3 ⁱⁱⁱ	0.81 (2)	1.89 (2)	2.697 (3)	178 (4)
O5—H5B...O1 ⁱⁱⁱ	0.82 (2)	2.01 (2)	2.825 (4)	175 (3)
O6—H6A...O3 ^{iv}	0.81 (3)	1.98 (3)	2.769 (3)	165 (3)
O6—H6B...O1 ^v	0.81 (2)	1.99 (2)	2.795 (3)	171 (3)

Symmetry codes: (ii) $-x, -y + 2, -z + 1$; (iii) $x - 1, y, z$; (iv) $-x + 1, -y + 1, -z + 1$; (v) $x, y - 1, z$.

Data collection: SMART (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine

structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *publCIF* (Westrip, 2010).

Acknowledgements

This work was supported financially by the Graduate Student Research Innovation Fund of CTGU (CX2014094) and Training Excellent Master's Thesis Fund of CTGU (Nos. PY2015074, PY2015030 and PY2015024), China.

Supporting information for this paper is available from the IUCr electronic archives (Reference: XU5853).

References

- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Li, Z.-X., Xu, Y., Zuo, Y., Li, L., Pan, Q.-H., Hu, T.-L. & Bu, X.-H. (2009). *Cryst. Growth Des.* **9**, 3904–3909.
Meng, F.-J., Jia, H.-Q., Hu, N.-H. & Zhou, H. (2012). *Acta Cryst.* **E68**, m1364.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supporting information

Acta Cryst. (2015). E71, m145–m146 [doi:10.1107/S2056989015012190]

Crystal structure of *catena*-poly[[diaquacobalt(II)]-bis[μ -5-(4-carboxyylato-phenyl)picolinato]- $\kappa^3 N, O^2: O^5; \kappa^3 O^5: N, O^2$ -[diaquacobalt(II)]- μ -1-[4-(1*H*-imidazol-1-yl)phenyl]-1*H*-imidazole- $\kappa^2 N^3: N^3'$]

Guo-Wang Xu, Ye-Nan Wang, Hai-Bing Wang and Zhong-Long Wang

S1. Comment

Recent years, many successful examples have been made based on mixed ligands. Pyridine carboxylic acids and N-donor ligand as good kinds of organic linkers were always selected to construct various metal-organic frameworks (MOFs) with specially properties. In this work, we choose 5-(4-carboxyphenyl)picolinic acid and 1,4-bis(1-imidazolyl)benzene to construct a new kind of Co-based MOFs. By now, a related 5-(4-carboxyphenyl)picolinate complex (Meng et al., 2012) and a related 1,4-bis(1-imidazolyl)benzene compound (Li et al., 2009) have been reported.

S2. Experimental

An aqueous mixture of cobalt(II) nitrate hexahydrate (58.2 mg, 0.2 mmol), 5-(4-carboxyphenyl)picolinic acid (48.6 mg, 0.2 mmol) and 1,4-bis(1-imidazolyl)benzene (42 mg, 0.2 mmol) was placed in a Teflon-lined, stainless-steel reactor. The reactor was heated to 413 K for 72 hours. It was then cooled to room temperature at the rate of 15 K per hour. Red crystals were isolated in 73% yield (based on Co). C, H, N elemental analysis. calcd. for C₁₉H₁₆O₆N₃Co: C 51.71, H 3.65, N 9.52%; found C 51.88, H 3.71, N 9.92%.

S3. Refinement

Water H atoms were located in a difference Fourier map and refined with a distance constraint of O—H = 0.82 (1) Å, $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. Other H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.




Figure 1

A part of the crystal structure of the title compound with labelling and displacement ellipsoids drawn at the 30% probability level. Symmetry codes: (i) $-x, 1 - y, 1 - z$; (ii) $1 - x, 2 - y, -z$.

catena-Poly[[diaquacobalt(II)]-bis[μ -5-(4-carboxylatophenyl)picolinate]- $\kappa^3N,O^2:O^5;\kappa^3O^5:N,O^2$ -[diaquacobalt(II)]- μ -1-[4-(1H-imidazol-1-yl)phenyl]-1H-imidazole- $\kappa^2N^3:N^3$]

Crystal data

$[\text{Co}_2(\text{C}_{13}\text{H}_7\text{NO}_4)_2(\text{C}_{12}\text{H}_{10}\text{N}_4)(\text{H}_2\text{O})_4]$

$M_r = 441.28$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.055\ (3)\ \text{\AA}$

$b = 7.190\ (3)\ \text{\AA}$

$c = 20.038\ (10)\ \text{\AA}$

$\alpha = 80.25\ (2)^\circ$

$\beta = 79.90\ (2)^\circ$

$\gamma = 64.080\ (15)^\circ$

$V = 895.1\ (7)\ \text{\AA}^3$

$Z = 2$

$F(000) = 452$

$D_x = 1.637\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2498 reflections

$\theta = 2.1\text{--}27.6^\circ$

$\mu = 1.00\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, red

$0.20 \times 0.20 \times 0.17\ \text{mm}$

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

phi and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.825$, $T_{\max} = 0.848$

9552 measured reflections

4104 independent reflections

3636 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.045$

$\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 3.1^\circ$

$h = -9 \rightarrow 9$

$k = -9 \rightarrow 9$

$l = -26 \rightarrow 25$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.090$

$S = 1.05$

4104 reflections

274 parameters

4 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.035P)^2 + 0.341P]$
where $P = (F_o^2 + 2F_c^2)/3$

$$\begin{aligned}(\Delta/\sigma)_{\max} &= 0.001 \\ \Delta\rho_{\max} &= 0.34 \text{ e } \text{\AA}^{-3} \\ \Delta\rho_{\min} &= -0.41 \text{ e } \text{\AA}^{-3}\end{aligned}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Co1	−0.00343 (4)	0.48246 (4)	0.218107 (13)	0.02287 (10)
N1	0.1553 (3)	0.6510 (3)	0.69865 (8)	0.0240 (4)
N2	0.1229 (3)	0.5882 (3)	0.12579 (9)	0.0286 (4)
N3	0.2772 (3)	0.7531 (3)	0.04950 (8)	0.0291 (4)
O1	0.2958 (3)	0.7975 (3)	0.25034 (8)	0.0359 (4)
O2	0.1274 (3)	0.5901 (3)	0.28498 (8)	0.0368 (4)
O3	0.3160 (3)	0.8566 (2)	0.81611 (8)	0.0325 (3)
O4	0.1426 (3)	0.6548 (3)	0.83101 (7)	0.0341 (4)
O5	−0.2808 (3)	0.7538 (3)	0.22533 (10)	0.0404 (4)
O6	0.2678 (3)	0.2026 (3)	0.22547 (11)	0.0443 (4)
C1	0.2182 (3)	0.6971 (3)	0.29511 (10)	0.0247 (4)
C2	0.2347 (3)	0.7058 (3)	0.36903 (10)	0.0242 (4)
C3	0.2170 (3)	0.5541 (3)	0.41956 (11)	0.0279 (4)
H3	0.1991	0.4436	0.4078	0.033*
C4	0.2260 (3)	0.5653 (3)	0.48755 (11)	0.0286 (4)
H4	0.2190	0.4596	0.5204	0.034*
C5	0.2452 (3)	0.7331 (3)	0.50710 (10)	0.0248 (4)
C6	0.2625 (4)	0.8848 (3)	0.45621 (11)	0.0294 (5)
H6	0.2748	0.9983	0.4681	0.035*
C7	0.2618 (3)	0.8700 (3)	0.38789 (11)	0.0287 (4)
H7	0.2794	0.9701	0.3546	0.034*
C8	0.2451 (3)	0.7502 (3)	0.58043 (10)	0.0253 (4)
C9	0.1620 (3)	0.6403 (3)	0.63224 (10)	0.0270 (4)
H9	0.1085	0.5552	0.6198	0.032*
C10	0.2369 (3)	0.7693 (3)	0.71770 (10)	0.0246 (4)
C11	0.3202 (4)	0.8855 (4)	0.67012 (11)	0.0330 (5)
H11	0.3740	0.9681	0.6840	0.040*
C12	0.3220 (4)	0.8769 (4)	0.60140 (11)	0.0329 (5)
H12	0.3752	0.9563	0.5690	0.039*
C13	0.2319 (3)	0.7604 (3)	0.79457 (10)	0.0246 (4)
C14	0.2106 (4)	0.7192 (4)	0.11613 (11)	0.0341 (5)

H14	0.2254	0.7813	0.1508	0.041*
C15	0.2261 (4)	0.6350 (4)	0.01419 (11)	0.0346 (5)
H15	0.2507	0.6269	-0.0326	0.042*
C16	0.1328 (4)	0.5334 (4)	0.06182 (11)	0.0332 (5)
H16	0.0830	0.4409	0.0529	0.040*
C17	0.5370 (4)	0.8784 (4)	0.06193 (11)	0.0373 (6)
H17	0.5615	0.7960	0.1034	0.045*
C18	0.3908 (4)	0.8791 (3)	0.02369 (10)	0.0286 (5)
C19	0.3533 (4)	0.9994 (4)	-0.03836 (11)	0.0346 (5)
H19	0.2557	0.9982	-0.0639	0.042*
H5A	-0.288 (5)	0.869 (2)	0.2128 (14)	0.052*
H5B	-0.403 (2)	0.767 (5)	0.2352 (15)	0.052*
H6A	0.383 (3)	0.191 (5)	0.2064 (14)	0.052*
H6B	0.263 (5)	0.090 (2)	0.2309 (15)	0.052*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Co1	0.02701 (16)	0.02627 (15)	0.02110 (15)	-0.01781 (12)	-0.00122 (10)	-0.00029 (10)
N1	0.0276 (9)	0.0289 (9)	0.0216 (8)	-0.0184 (8)	-0.0011 (7)	-0.0018 (7)
N2	0.0365 (10)	0.0367 (10)	0.0220 (9)	-0.0261 (9)	-0.0009 (7)	0.0002 (7)
N3	0.0420 (11)	0.0399 (10)	0.0188 (8)	-0.0316 (9)	0.0007 (7)	-0.0013 (7)
O1	0.0512 (10)	0.0455 (9)	0.0246 (8)	-0.0345 (9)	-0.0010 (7)	-0.0021 (7)
O2	0.0505 (10)	0.0487 (10)	0.0291 (8)	-0.0366 (9)	-0.0077 (7)	-0.0026 (7)
O3	0.0429 (9)	0.0393 (9)	0.0288 (8)	-0.0294 (8)	-0.0038 (7)	-0.0044 (7)
O4	0.0496 (10)	0.0474 (9)	0.0220 (7)	-0.0378 (9)	-0.0002 (6)	-0.0017 (7)
O5	0.0300 (9)	0.0252 (8)	0.0647 (12)	-0.0139 (8)	-0.0014 (8)	0.0008 (8)
O6	0.0285 (9)	0.0289 (8)	0.0732 (13)	-0.0152 (8)	0.0035 (8)	0.0000 (9)
C1	0.0246 (10)	0.0273 (10)	0.0256 (10)	-0.0140 (9)	-0.0030 (8)	-0.0031 (8)
C2	0.0218 (10)	0.0308 (10)	0.0231 (10)	-0.0139 (9)	-0.0022 (7)	-0.0036 (8)
C3	0.0294 (11)	0.0306 (11)	0.0298 (11)	-0.0180 (9)	-0.0027 (8)	-0.0042 (9)
C4	0.0338 (12)	0.0310 (11)	0.0267 (11)	-0.0205 (10)	-0.0026 (8)	0.0007 (8)
C5	0.0228 (10)	0.0323 (10)	0.0235 (10)	-0.0158 (9)	-0.0010 (8)	-0.0031 (8)
C6	0.0390 (12)	0.0331 (11)	0.0270 (10)	-0.0250 (10)	-0.0037 (9)	-0.0035 (9)
C7	0.0374 (12)	0.0313 (11)	0.0240 (10)	-0.0217 (10)	-0.0047 (8)	0.0016 (8)
C8	0.0256 (10)	0.0323 (10)	0.0216 (10)	-0.0163 (9)	-0.0009 (8)	-0.0022 (8)
C9	0.0314 (11)	0.0331 (11)	0.0251 (10)	-0.0213 (9)	-0.0039 (8)	-0.0026 (8)
C10	0.0253 (10)	0.0283 (10)	0.0256 (10)	-0.0167 (9)	-0.0009 (8)	-0.0035 (8)
C11	0.0445 (13)	0.0420 (12)	0.0279 (11)	-0.0331 (11)	-0.0004 (9)	-0.0050 (9)
C12	0.0442 (14)	0.0445 (13)	0.0229 (10)	-0.0332 (12)	-0.0005 (9)	0.0007 (9)
C13	0.0271 (11)	0.0272 (10)	0.0236 (10)	-0.0156 (9)	-0.0018 (8)	-0.0028 (8)
C14	0.0519 (14)	0.0485 (13)	0.0187 (10)	-0.0381 (12)	0.0015 (9)	-0.0040 (9)
C15	0.0461 (14)	0.0486 (13)	0.0215 (10)	-0.0322 (12)	0.0015 (9)	-0.0066 (9)
C16	0.0468 (14)	0.0405 (12)	0.0271 (11)	-0.0323 (11)	-0.0012 (9)	-0.0055 (9)
C17	0.0531 (15)	0.0522 (14)	0.0211 (10)	-0.0385 (13)	-0.0090 (10)	0.0098 (10)
C18	0.0404 (13)	0.0369 (12)	0.0189 (9)	-0.0283 (10)	0.0020 (8)	-0.0014 (8)
C19	0.0433 (13)	0.0516 (14)	0.0233 (10)	-0.0339 (12)	-0.0081 (9)	0.0032 (9)

Geometric parameters (Å, °)

Co1—N1 ⁱ	2.1403 (18)	C3—H3	0.9300
Co1—N2	2.0815 (18)	C4—C5	1.396 (3)
Co1—O2	2.1575 (16)	C4—H4	0.9300
Co1—O4 ⁱ	2.1028 (16)	C5—C6	1.393 (3)
Co1—O5	2.0773 (18)	C5—C8	1.495 (3)
Co1—O6	2.0889 (19)	C6—C7	1.392 (3)
N1—C9	1.338 (3)	C6—H6	0.9300
N1—C10	1.347 (2)	C7—H7	0.9300
N1—Co1 ⁱ	2.1403 (18)	C8—C12	1.394 (3)
N2—C14	1.309 (3)	C8—C9	1.400 (3)
N2—C16	1.385 (3)	C9—H9	0.9300
N3—C14	1.354 (3)	C10—C11	1.384 (3)
N3—C15	1.383 (3)	C10—C13	1.526 (3)
N3—C18	1.435 (3)	C11—C12	1.386 (3)
O1—C1	1.257 (2)	C11—H11	0.9300
O2—C1	1.254 (2)	C12—H12	0.9300
O3—C13	1.251 (2)	C14—H14	0.9300
O4—C13	1.255 (2)	C15—C16	1.357 (3)
O4—Co1 ⁱ	2.1028 (16)	C15—H15	0.9300
O5—H5A	0.810 (10)	C16—H16	0.9300
O5—H5B	0.813 (10)	C17—C19 ⁱⁱ	1.386 (3)
O6—H6A	0.811 (10)	C17—C18	1.387 (3)
O6—H6B	0.814 (10)	C17—H17	0.9300
C1—C2	1.520 (3)	C18—C19	1.384 (3)
C2—C3	1.390 (3)	C19—C17 ⁱⁱ	1.386 (3)
C2—C7	1.395 (3)	C19—H19	0.9300
C3—C4	1.392 (3)		
O5—Co1—N2	94.70 (8)	C6—C5—C8	121.13 (18)
O5—Co1—O6	172.08 (8)	C4—C5—C8	120.96 (18)
N2—Co1—O6	92.55 (8)	C7—C6—C5	121.32 (19)
O5—Co1—O4 ⁱ	92.29 (8)	C7—C6—H6	119.3
N2—Co1—O4 ⁱ	92.21 (7)	C5—C6—H6	119.3
O6—Co1—O4 ⁱ	90.60 (8)	C6—C7—C2	120.39 (19)
O5—Co1—N1 ⁱ	86.11 (8)	C6—C7—H7	119.8
N2—Co1—N1 ⁱ	169.23 (7)	C2—C7—H7	119.8
O6—Co1—N1 ⁱ	87.36 (8)	C12—C8—C9	116.25 (18)
O4 ⁱ —Co1—N1 ⁱ	77.02 (7)	C12—C8—C5	122.94 (18)
O5—Co1—O2	89.39 (8)	C9—C8—C5	120.80 (18)
N2—Co1—O2	97.93 (7)	N1—C9—C8	123.55 (18)
O6—Co1—O2	86.46 (8)	N1—C9—H9	118.2
O4 ⁱ —Co1—O2	169.55 (6)	C8—C9—H9	118.2
N1 ⁱ —Co1—O2	92.81 (7)	N1—C10—C11	121.47 (19)
C9—N1—C10	119.08 (17)	N1—C10—C13	114.58 (17)
C9—N1—Co1 ⁱ	126.97 (14)	C11—C10—C13	123.95 (18)
C10—N1—Co1 ⁱ	113.75 (13)	C10—C11—C12	118.96 (19)

C14—N2—C16	105.69 (17)	C10—C11—H11	120.5
C14—N2—Co1	127.21 (15)	C12—C11—H11	120.5
C16—N2—Co1	127.09 (14)	C11—C12—C8	120.62 (19)
C14—N3—C15	106.92 (17)	C11—C12—H12	119.7
C14—N3—C18	124.58 (18)	C8—C12—H12	119.7
C15—N3—C18	128.39 (17)	O3—C13—O4	125.53 (19)
C1—O2—Co1	151.15 (14)	O3—C13—C10	118.33 (17)
C13—O4—Co1 ⁱ	117.89 (13)	O4—C13—C10	116.14 (17)
Co1—O5—H5A	124 (2)	N2—C14—N3	111.75 (19)
Co1—O5—H5B	129 (2)	N2—C14—H14	124.1
H5A—O5—H5B	106 (3)	N3—C14—H14	124.1
Co1—O6—H6A	122 (2)	C16—C15—N3	105.86 (19)
Co1—O6—H6B	123 (2)	C16—C15—H15	127.1
H6A—O6—H6B	109 (3)	N3—C15—H15	127.1
O2—C1—O1	126.48 (19)	C15—C16—N2	109.77 (19)
O2—C1—C2	116.35 (18)	C15—C16—H16	125.1
O1—C1—C2	117.17 (17)	N2—C16—H16	125.1
C3—C2—C7	118.53 (18)	C19 ⁱⁱ —C17—C18	120.0 (2)
C3—C2—C1	120.86 (18)	C19 ⁱⁱ —C17—H17	120.0
C7—C2—C1	120.59 (18)	C18—C17—H17	120.0
C2—C3—C4	120.86 (19)	C19—C18—C17	120.91 (19)
C2—C3—H3	119.6	C19—C18—N3	120.09 (19)
C4—C3—H3	119.6	C17—C18—N3	118.99 (19)
C3—C4—C5	120.90 (19)	C18—C19—C17 ⁱⁱ	119.1 (2)
C3—C4—H4	119.6	C18—C19—H19	120.4
C5—C4—H4	119.6	C17 ⁱⁱ —C19—H19	120.4
C6—C5—C4	117.91 (19)		
O5—Co1—N2—C14	76.7 (2)	Co1 ⁱ —N1—C9—C8	172.91 (15)
O6—Co1—N2—C14	-100.1 (2)	C12—C8—C9—N1	-0.4 (3)
O4 ⁱ —Co1—N2—C14	169.2 (2)	C5—C8—C9—N1	-179.25 (19)
N1 ⁱ —Co1—N2—C14	170.6 (3)	C9—N1—C10—C11	2.4 (3)
O2—Co1—N2—C14	-13.3 (2)	Co1 ⁱ —N1—C10—C11	-172.91 (17)
O5—Co1—N2—C16	-103.7 (2)	C9—N1—C10—C13	-176.73 (18)
O6—Co1—N2—C16	79.4 (2)	Co1 ⁱ —N1—C10—C13	8.0 (2)
O4 ⁱ —Co1—N2—C16	-11.3 (2)	N1—C10—C11—C12	-1.0 (3)
N1 ⁱ —Co1—N2—C16	-9.9 (5)	C13—C10—C11—C12	178.0 (2)
O2—Co1—N2—C16	166.21 (19)	C10—C11—C12—C8	-1.2 (4)
O5—Co1—O2—C1	-82.1 (3)	C9—C8—C12—C11	1.9 (3)
N2—Co1—O2—C1	12.6 (3)	C5—C8—C12—C11	-179.4 (2)
O6—Co1—O2—C1	104.7 (3)	Co1 ⁱ —O4—C13—O3	177.72 (17)
O4 ⁱ —Co1—O2—C1	178.5 (3)	Co1 ⁱ —O4—C13—C10	-2.5 (2)
N1 ⁱ —Co1—O2—C1	-168.2 (3)	N1—C10—C13—O3	175.91 (18)
Co1—O2—C1—O1	-7.1 (5)	C11—C10—C13—O3	-3.2 (3)
Co1—O2—C1—C2	172.7 (2)	N1—C10—C13—O4	-3.9 (3)
O2—C1—C2—C3	20.7 (3)	C11—C10—C13—O4	177.0 (2)
O1—C1—C2—C3	-159.4 (2)	C16—N2—C14—N3	0.0 (3)
O2—C1—C2—C7	-157.4 (2)	Co1—N2—C14—N3	179.63 (15)

O1—C1—C2—C7	22.4 (3)	C15—N3—C14—N2	0.4 (3)
C7—C2—C3—C4	0.1 (3)	C18—N3—C14—N2	-176.1 (2)
C1—C2—C3—C4	-178.12 (19)	C14—N3—C15—C16	-0.7 (3)
C2—C3—C4—C5	2.3 (3)	C18—N3—C15—C16	175.6 (2)
C3—C4—C5—C6	-2.2 (3)	N3—C15—C16—N2	0.7 (3)
C3—C4—C5—C8	177.2 (2)	C14—N2—C16—C15	-0.5 (3)
C4—C5—C6—C7	-0.3 (3)	Co1—N2—C16—C15	179.93 (16)
C8—C5—C6—C7	-179.7 (2)	C19 ⁱⁱ —C17—C18—C19	0.5 (4)
C5—C6—C7—C2	2.7 (3)	C19 ⁱⁱ —C17—C18—N3	-179.3 (2)
C3—C2—C7—C6	-2.5 (3)	C14—N3—C18—C19	-143.7 (2)
C1—C2—C7—C6	175.7 (2)	C15—N3—C18—C19	40.6 (4)
C6—C5—C8—C12	-19.3 (3)	C14—N3—C18—C17	36.1 (3)
C4—C5—C8—C12	161.3 (2)	C15—N3—C18—C17	-139.6 (3)
C6—C5—C8—C9	159.4 (2)	C17—C18—C19—C17 ⁱⁱ	-0.5 (4)
C4—C5—C8—C9	-19.9 (3)	N3—C18—C19—C17 ⁱⁱ	179.3 (2)
C10—N1—C9—C8	-1.7 (3)		

Symmetry codes: (i) $-x, -y+1, -z+1$; (ii) $-x+1, -y+2, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O5—H5 <i>A</i> \cdots O3 ⁱⁱⁱ	0.81 (2)	1.89 (2)	2.697 (3)	178 (4)
O5—H5 <i>B</i> \cdots O1 ^{iv}	0.82 (2)	2.01 (2)	2.825 (4)	175 (3)
O6—H6 <i>A</i> \cdots O3 ^v	0.81 (3)	1.98 (3)	2.769 (3)	165 (3)
O6—H6 <i>B</i> \cdots O1 ^{vi}	0.81 (2)	1.99 (2)	2.795 (3)	171 (3)

Symmetry codes: (iii) $-x, -y+2, -z+1$; (iv) $x-1, y, z$; (v) $-x+1, -y+1, -z+1$; (vi) $x, y-1, z$.