



Review

Human $\gamma\delta$ T Cell Subsets and Their Clinical Applications for Cancer Immunotherapy

Derek Lee ^{1,2}, Carl J. Rosenthal ³, Natalie E. Penn ¹, Zachary Spencer Dunn ^{1,4} , Yang Zhou ^{1,2} 
and Lili Yang ^{1,2,5,6,*}

¹ Department of Microbiology, Immunology & Molecular Genetics, University of California, Los Angeles, CA 90095, USA; ylee932@g.ucla.edu (D.L.); nepenn168@g.ucla.edu (N.E.P.); zacharsd@usc.edu (Z.S.D.); zzydcat@ucla.edu (Y.Z.)

² Molecular Biology Institute, University of California, Los Angeles, CA 90095, USA

³ Department of Molecular, Cell and Developmental Biology, University of California, Los Angeles, CA 90095, USA; cjr@dt3.org

⁴ Mork Family Department of Chemical Engineering and Materials Science, University of Southern California, Los Angeles, CA 90089, USA

⁵ Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, CA 90095, USA

⁶ Jonsson Comprehensive Cancer Center, David Geffen School of Medicine, University of California, Los Angeles, CA 90095, USA

* Correspondence: lililyang@ucla.edu; Tel.: +1-310-825-8609

Simple Summary: Research into the immunotherapeutic potential of T cells has predominantly focused on conventional alpha beta ($\alpha\beta$) T cells, which recognize peptide antigens presented by polymorphic major histocompatibility complex (MHC) class I and class II molecules. However, innate-like T cells, such as gamma delta ($\gamma\delta$) T cells, also play important roles in antitumor immunity. Here, we review the current understanding of $\gamma\delta$ T cells in antitumor immunity and discuss strategies that could potentially maximize their potential in cancer immunotherapy.



Citation: Lee, D.; Rosenthal, C.J.; Penn, N.E.; Dunn, Z.S.; Zhou, Y.; Yang, L. Human $\gamma\delta$ T Cell Subsets and Their Clinical Applications for Cancer Immunotherapy. *Cancers* **2022**, *14*, 3005. <https://doi.org/10.3390/cancers14123005>

Academic Editor: Véronique Décot

Received: 5 May 2022

Accepted: 16 June 2022

Published: 18 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Gamma delta ($\gamma\delta$) T cells are a minor population of T cells that share adaptive and innate immune properties. In contrast to MHC-restricted alpha beta ($\alpha\beta$) T cells, $\gamma\delta$ T cells are activated in an MHC-independent manner, making them ideal candidates for developing allogeneic, off-the-shelf cell-based immunotherapies. As the field of cancer immunotherapy progresses rapidly, different subsets of $\gamma\delta$ T cells have been explored. In addition, $\gamma\delta$ T cells can be engineered using different gene editing technologies that augment their tumor recognition abilities and antitumor functions. In this review, we outline the unique features of different subsets of human $\gamma\delta$ T cells and their antitumor properties. We also summarize the past and the ongoing pre-clinical studies and clinical trials utilizing $\gamma\delta$ T cell-based cancer immunotherapy.

Keywords: gamma delta T ($\gamma\delta$ T) cells; cancer immunotherapy; chimeric antigen receptor T (CAR-T) cells; allogeneic cell therapy; butyrophilins (BTN); zoledronate (ZOL)

1. Introduction

Chimeric antigen receptor (CAR) engineered T cell (CAR-T) therapy has revolutionized the treatment of hematological malignancies over the last decade with approved therapies for leukemias, lymphomas, and multiple myeloma (MM). Despite significant improvements in clinical outcomes, CAR-T therapy has faced some hurdles, particularly in treating solid tumors [1]. In this regard, cells with distinct immunological features, such as natural killer (NK) cells, NKT cells, and $\gamma\delta$ T cells, offer alternative cell sources for CAR engineering to treat cancer. Based on the surface expression of $\alpha\beta$ or $\gamma\delta$ T cell receptors (TCRs), T cells are classified into two primary subpopulations— $\alpha\beta$ T and $\gamma\delta$ T cells. While the majority of T cell research and therapeutic application has focused on $\alpha\beta$ T cells, $\gamma\delta$ T

cells are also significant participants in cancer immunity. In fact, Gentles et al. studied complex associations between 22 different leukocyte subsets and cancer survival across 5782 clinical tumor samples and found that intra-tumoral $\gamma\delta$ T cells were the best predictors of overall patient survival [2]. $\alpha\beta$ T cells are adaptive immune cells that recognize their target antigens presented on the major histocompatibility complex (MHC) molecules. $\gamma\delta$ T cells, on the other hand, are innate immune cells that function in an MHC-unrestricted manner and are able to recognize surface proteins such as butyrophilin (BTN) and CD1 molecules. In addition, $\gamma\delta$ T cells have a low risk of developing graft-versus-host disease (GvHD) in allogeneic adoptive transfer settings [3,4], and thus provide a new opportunity for “off-the-shelf” cellular therapeutics. Past studies have shown that $\gamma\delta$ T cells exert potent cytotoxicity against various types of tumors [5–8]. However, compared with their $\alpha\beta$ T cell counterparts, they only comprise 1–5% of total T cells [9]. In this review, we will first provide an introduction to $\gamma\delta$ T cell biology and then summarize past and current clinical trials utilizing $\gamma\delta$ T based-immunotherapy. We will also discuss the challenges of $\gamma\delta$ T based-immunotherapy and the future directions that may overcome these obstacles.

2. $\gamma\delta$ T Cell Subsets and Their Key Features

Recognizing the physiological differences between murine and human $\gamma\delta$ T cells has been critical for understanding their immunotherapeutic potential. Rather than an orthologous conservation between species, mice and humans each possess distinct $\gamma\delta$ T cell subtypes with analogous but not precisely identical functions [10]. Mouse subtypes are typically defined by the γ chain of the $\gamma\delta$ TCR, while human $\gamma\delta$ T cell subtypes are typically defined by the δ chain. The most relevant subtypes for mice are accordingly $V\gamma 1-7$, and those for humans are $V\delta 1-3$ [10].

The human $\gamma\delta$ T subsets that have been most comprehensively studied are $V\delta 1+$ and $V\delta 2+$ T cells, both of which arise during fetal development. Shortly after birth, $V\delta 1+$ T cells are the primary subtype, and occupy the cord blood, postnatal thymus, gut, and skin [11–15]. Adults also maintain a population of $V\delta 1+$ T cells. Although they make up only about 10–15% of $\gamma\delta$ T cells in the blood, they reside primarily in peripheral tissues such as gut epithelia, dermis, spleen, and liver [16,17]. While the development of $V\delta 1+$ T cells is not fully elucidated, $V\delta 2+$ T cell development has been characterized in greater detail. TCR $V\delta 2$ chains are frequently paired with $V\gamma 9$ chains during development [18]. The cells first arise in the fetus, at around five to seven weeks, then migrate from the fetal liver to the thymus at 8 weeks [19]. Though $V\delta 1+$ T cells outnumber $V\delta 2+$ cells immediately after birth, the $V\delta 2+$ cells expand to predominance in the blood and spleen by adulthood, comprising 80–85% of total peripheral $\gamma\delta$ T cells [16]. The $V\gamma 9$ chain is germline-encoded to recognize target BTN proteins, suggesting that BTN family molecule exposure may not be necessary during maturation [20,21]. However, variation in the $J\delta$ chain of fetal and adult $V\delta 2$ cells indicates that $\gamma\delta$ T cell expansion during maturation to adulthood also requires antigen stimulation and selection [20,21]. $V\delta 3+$ T cells comprise the majority of the remaining $\gamma\delta$ T population [22]. Their lineage and rearrangement are less well described than the other subtypes, though they are known to complex with $V\gamma 2$ and $V\gamma 3$ for recognizing CD1d-presented antigens [22]. These cells are found in peripheral blood and the liver, and have been detected in patients with B-cell leukemia as well as in patients with human immunodeficiency virus (HIV) and cytomegalovirus (CMV) infections [22]. Other human $\gamma\delta$ T cell subsets such as $V\delta 4$, $V\delta 6$, $V\delta 7$, and $V\delta 8$ T cells have been found in the peripheral blood of lymphoma patients, but their functions have yet to be determined [23].

3. $\gamma\delta$ T Cell Recognition and Killing of Tumor Cells

$\gamma\delta$ T cells that express the $\gamma 9$ chains paired with the $\delta 2$ chains ($V\gamma 9V\delta 2$ T cells) are the major $\gamma\delta$ T cell population in human peripheral blood. The $V\gamma 9V\delta 2$ T cells react to cells with accumulated intracellular phosphoantigens (pAgs), intermediate metabolites produced by infected or transformed cells. These metabolites include isopentenyl pyrophosphate

(IPP) formed by the mevalonate (MVA) pathway of tumor cells, and (E)-4-Hydroxy-3-methyl-but-2-enyl pyrophosphate (HMB-PP) produced by microbial isoprenoid biosynthesis [24]. Overproduction of IPP in cancer cells as a result of dysregulated MVA pathway leads to activation of V γ 9V δ 2 T cells. A past study showed that oncogenic mutations cause the tumor suppressor protein p53 to interact with sterol regulatory element-binding protein-2 (SREBP-2) nuclear transcription factor, which augments the expression of MVA pathway genes [25,26]. While the precise TCR targeting mechanism of V γ 9V δ 2 T cells has been controversial, recent evidence has shown that the TCR V γ 9 chain directly recognizes BTN2A1, which is associated with BTN3A1 (CD277), another ligand recognized by the TCR V δ 2 chain [27] (Figure 1). BTNs belong to the immunoglobulin superfamily and are structurally similar to B7 family molecules, which include CD80, CD86, and programmed cell death-1 ligand (PD-L1) [28]. pAgs bind the intracellular B30.2 domain of BTN3A1, which enables the BTN2A1-BTN3A1 complex to bind the V γ 9V δ 2 TCR [27]. Strategies to expand V γ 9V δ 2 T cells thus focus on inducing pAg accumulation for TCR activation. Synthetic phosphoantigen analogues mimicking the pAg accumulated state, such as bromohydrin pyrophosphate (BrHPP) and 2-methyl-3-butenyl-1-pyrophosphate (2M3B1PP), can directly stimulate V γ 9V δ 2 T cells [29,30]. Stimulation can also be achieved with bisphosphonates, a class of drugs that prevent or slow down bone loss, such as zoledronate (ZOL) and pamidronate (PAM) [30–34]. These small molecule compounds inhibit farnesyl pyrophosphate synthase in the MVA metabolic pathway, leading to pAg accumulation in treated cells which activates the V γ 9V δ 2 TCR [24]. Notably, the V γ 9V δ 2 T cell subpopulation still retains diversity between and within individuals, and therefore cloning of a single, high-affinity V γ 9V δ 2 TCR to create an effector population could produce more effective therapies than the use of a polyclonal existing population [35].

Non-V γ 9V δ 2 T cells have more diverse TCRs without pAg-mediated BTN recognition capacity. Instead, these non-V γ 9V δ 2 TCRs recognize other molecules such as the CD1 family of surface glycoproteins presenting lipid antigens [36] (Figure 1). V δ 1+ and V δ 3+ subsets have been found to recognize CD1 proteins, although recognition of lipid antigens on CD1 molecules is typically associated with NKT cells [36–40]. However, unlike BTN molecules, which are commonly expressed on both hematologic and solid tumors [41], CD1+ tumor cells mostly arise from myelomonocytic and B-cell lineages, and only a few solid tumors have been discovered to be CD1+ [42]. While it is known that CD1d recognition by NKT cells enables their antitumor response, this reactivity in non-V γ 9V δ 2 T cells remains to be demonstrated [43].

Besides their TCRs, human $\gamma\delta$ T cells also express other surface molecules that can recognize tumor cells and provide additional stimulatory signals. Natural killer receptors expressed by $\gamma\delta$ T cells—including Natural Killer Group 2D (NKG2D), DNAX Accessory Molecule-1 (DNAM-1), NKp30, NKp44, and NKp46—enable them to be activated by and target malignantly transformed cells, which often overexpress ligands for these receptors [44–49] (Figure 1). Both hematologic and solid tumors typically express MHC class I chain-related polypeptide A (MICA) and MICB, as well as members of the UL16 binding protein (ULBP) family (ULBP1–6), which activate NKG2D-expressing V δ 1+ and V δ 2+ T cells. The binding of DNAM-1 to its target CD155 (poliovirus receptor, PVR) [50] or CD112 (Nectin-2) [51] can promote the NK cell-mediated elimination of transformed and virus-infected cells. Although both NKG2D and DNAM-1 are expressed on V δ 1+ and V δ 2+ T cells, NKp30, NKp44, and NKp46 are only expressed on V δ 1+ T cells [52].




Figure 1. The tumor cell recognition of CAR- $\gamma\delta$ T cells. CAR- $\gamma\delta$ T cells recognize tumor cells through multiple receptor pathways, increasing their cytolytic potential and resistance to immunosuppression. Both V δ 1+ and V γ 9V δ 2 TCR possess MHC-nonrestricted cytotoxicity. The V δ 1 TCR recognizes lipid antigens presented on CD1 molecules, and the V γ 9V δ 2 TCR recognizes intracellular pAg-bound BTN3A1/2A1 complex. V γ 9V δ 2 T cells express CD16 (Fc γ RIII), which binds the Fc region of IgG antibodies to induce antibody-dependent cellular cytotoxicity (ADCC). TLRs (Toll-Like Receptors) form another axis of immune cell coordination with each binding pathogen- or damage-associated molecular patterns (PAMPs/DAMPs), simultaneously increasing $\gamma\delta$ T cell cytotoxicity and coordinating with surrounding dendritic cells (DCs). NK receptor pathways expressed on the cells enable cytolytic activity in response to expressed markers of DNA damage, stress, or infection. Both V δ 1+ and V γ 9V δ 2 T cells express NKG2D, which recognizes cellular distress ligands presented on MIC(A/B) and ULBP1-6, and DNAM-1, which binds CD155 and CD112. In addition, V δ 1+ T cells express NKp30, NKp44, and NKp46.

The antitumor cytotoxicity of $\gamma\delta$ T cells occurs through activation induced perforin and granzyme B release, which causes pore formation, granzyme protease entry, and subsequent apoptosis of the tumor cells [53] (Figure 2). In addition to the perforin-granzyme pathway, $\gamma\delta$ T cells also kill tumor cells by expressing TNF-related apoptosis-inducing ligand (TRAIL) and Fas ligand (FasL), which engage their respective receptors expressed by tumor cells to induce apoptosis [53–55]. Furthermore, $\gamma\delta$ T cells can target tumors through antibody dependent cellular cytotoxicity (ADCC), whereby CD16 (Fc γ receptor III) expressed on $\gamma\delta$ T cells binds the Fc region of antibodies bound to a target cell, leading to tumor lysis [53,56,57] (Figure 1). The V γ 9V δ 2 subset in particular has been shown to upregulate CD16 after stimulation, and effectively performs ADCC against tumors coordinated by monoclonal antibodies against HER2 (Trastuzumab) and CD20 (Rituximab) [57].




Figure 2. Regulation of antitumor immunity by $V\gamma 9V\delta 2$ T cells. $V\gamma 9V\delta 2$ T cells perform diverse immunological functions through interactions with other cells. They are directly cytotoxic to tumor cells through multiple mechanisms including perforin and granzyme B production, and express CD36 to facilitate tumor antigen phagocytosis. Upon incorporating antigens, $V\gamma 9V\delta 2$ T cells act as antigen presenting cells (APCs) and both directly and cross present antigens on MHC molecules to $\alpha\beta$ T cells. Combined with expression of the co-stimulatory molecules CD80 and CD86, $V\gamma 9V\delta 2$ T cells effectively induce $\alpha\beta$ T cells to differentiate and proliferate. CD137L expressed on $V\gamma 9V\delta 2$ T cells activates NK cells by engaging CD137, increasing their NKG2D expression, direct cytotoxicity, and ADCC activity. $V\gamma 9V\delta 2$ T cells induce DC maturation through $TNF\alpha$ and $IFN-\gamma$ production, and engage CD40 to allow DCs to generate CD8+ T cell responses. Conversely, DCs activate $V\gamma 9V\delta 2$ T cells by providing CD86 co-stimulation and producing cytokines including IL-12, IL-18, IL-1 β , $TNF\alpha$, and $IFN-\gamma$. Furthermore, mature DCs treated with ZOL upregulate CD40L on $V\gamma 9V\delta 2$ T cells.

4. $\gamma\delta$ T Cell Activation via Toll-like Receptors

Another co-stimulator is the class of toll-like receptors (TLRs), which can activate $\gamma\delta$ T cells directly or by participating in DC activation [58]. TLRs are integral membrane glycoproteins which function as prototype pattern recognition receptors, dimerizing in response to specific lipid, peptide, or nucleic acid PAMPs (pathogen-associated molecular patterns) and DAMPs (danger-associated molecular patterns from damaged tissues) [58,59]. Binding induces signal transduction pathways upregulating the expression of inflammatory cytokines such as $TNF-\alpha$, IL-6, IL-1 β , IL-12, and $IFN-\gamma$ [58,59]. However, TLR signaling can activate or suppress the immune system: with nine TLRs expressed in humans, the result of signaling can depend on the specific TLR and surrounding circumstances. For instance, several studies [60,61] indicated that TLR2 stimulation suppressed Treg potential by suppressing FOXP3, whereas TLR5 stimulation increased FOXP3 expression.

In immunotherapy, the use of TLR agonists enhances $\gamma\delta$ T cell cytotoxicity and proliferation in a DC-dependent and independent manner [58] (Figure 1). $\gamma\delta$ T cells have been shown to respond to a combination of TLR and TCR stimulation while influencing the antigen-presenting property of DCs in return [62,63]. DCs incubated and activated with $\gamma\delta$ T cells increased expression of CD86 and MHC I, enhanced the production of IFN- γ by alloreactive T cells, and produce IL12p70 [62], a messenger which activates NK cells and promotes $\gamma\delta$ T cell proliferation. When activating DCs through TLR signaling, $\gamma\delta$ T cells are co-activated, leading to an increase in CD25 and CD69 expression as well as the production of inflammatory cytokines [64]. V δ 2+ cells, more so than V δ 1+ cells, have a strong IFN- γ production response to a variety of TLRs [65]. V δ 2+ T cells expanded using IPP plus resiquimod, an agonist for TLRs 7 and 8, and were shown to have increased antitumor cytotoxicity and lower PD-1 expression than V δ 2+ T cells expanded with IPP or ZOL [66]. Resiquimod led to the expansion of V δ 2+ T-cells with and without the aid of antigen-presenting cells (APCs), and decreased the presence of inhibitory APCs and downregulated their expression of immunosuppressive PD-L1 and CTLA-4 [66].

5. $\gamma\delta$ T Cell Modulation of Tumor Microenvironment

The tumor microenvironment (TME) is composed of cancer cells, surrounding stroma including fibroblasts and vasculature, and infiltrating immune cells [67]. Although hypoxia and immunosuppressive conditions in the TME hinder many forms of cell therapy [67], $\gamma\delta$ T cells may be more effective in this challenging environment due to their recognition of stress markers expressed on many tumors, MHC independent activation, production of inflammatory cytokines, and the ability to stimulate other immune cells for a coordinated antitumor response. In addition, it was reported that $\gamma\delta$ T cells can target M1 and M2 macrophages [68]. Macrophages that infiltrate the TME may encounter microenvironment specific chemokines, growth factors, and other signals that polarize them away from an antitumor phenotype and towards a tumor promoting M2 state (tumor-associated macrophages, TAMs) [69]. TAMs are a significant component of the TME that support tumor growth and induce extracellular matrix (ECM) remodeling facilitating metastasis through factors such as VEGF and MMP-9 [70]. TAMs also inhibit immune cell antitumor activity by secreting immunosuppressive cytokines such as IL-10 and TGF- β and engaging inhibitory receptors on infiltrating lymphocytes [71]. In treating solid tumors, it is therefore beneficial to repolarize TAMs to an antitumor state, or to directly target and deplete TAMs.

While typically used to stimulate V γ 9V δ 2 T cells, ZOL is also useful against TAMs: it reduces TAM's pro-tumoral activities, repolarizes TAMs to an antitumor state, and may guide V γ 9V δ 2 T cell killing of TAMs. Macrophages have been shown to respond to ZOL in vitro and in vivo [72,73], with increased unphosphorylated Rap1A levels indicating MVA pathway inhibition [72]. Tumors from ZOL treated mice had reduced vascularization and TAM numbers, as well as decreased VEGF and IL-10 and increased proinflammatory IFN- γ [73]. Furthermore, macrophages from these ZOL treated mice repolarized to an antitumor phenotype [73]. In other investigations, ZOL similarly reduced TAM MMP-9 expression while increasing TAM expression of cytokines that restore an antitumor macrophage state [74]. While ZOL alone has promising effects on TAMs, it offers further benefits when administered in combination with V γ 9V δ 2 T cells, where it facilitates direct macrophage targeting; phosphoantigen accumulation in treated macrophages enables V γ 9V δ 2 T cells to respond to these cells and kill them in a perforin-dependent manner [68].

6. $\gamma\delta$ T Cell Interaction with Other Immune Cells

In addition to altering the TME, $\gamma\delta$ T cells have further immunomodulatory roles in interactions with other immune cell types (Figure 2). V γ 9V δ 2 T cells act as antigen presenting cells (APCs) in a similar capacity to DCs [75]. Mature DCs (mDCs) are the canonical professional APC, and their ability to present tumor associated antigens to T cells makes them a crucial component of the anti-cancer immune response [76]. V γ 9V δ 2 T cells similarly act as APCs upon activation, upregulating MHC I and MHC II molecules

and expressing APC associated adhesion and costimulatory molecules [75,77]. These cells also express the scavenger receptor CD36, which may facilitate their ability to uptake antigens from both apoptotic and live tumor cells [78]. After taking in tumor antigens, V γ 9V δ 2 T cells present those antigens to CD4+ $\alpha\beta$ T cells and also utilize insulin-regulated aminopeptidase (IRAP) [79] to cross present them on MHC I molecules to prime CD8+ $\alpha\beta$ T cells—a hallmark of professional APCs [75,77]. Antigen presentation by V γ 9V δ 2 T cells induces naive CD4+ and CD8+ $\alpha\beta$ T cell differentiation and proliferation to a similar degree as DCs [75,77].

Beyond acting as APCs to activate T cells, $\gamma\delta$ T cells have diverse roles interacting with other immune cells. One well characterized set of interactions is the bidirectional crosstalk between $\gamma\delta$ T cells and DCs [80]. V γ 9V δ 2 T cells assist with DC maturation by producing TNF- α and IFN- γ , yielding DCs with upregulated co-stimulatory molecules and enhanced cytokine secretion which effectively induce $\alpha\beta$ T cell proliferation and IFN- γ production [81]. Conversely, DCs can activate $\gamma\delta$ T cells. CD86-CD28 cell contact interactions [64], as well as production of stimulatory cytokines such as IL-12, IL-1 β , TNF- α , and type 1 IFNs [82–85], enable DCs to activate $\gamma\delta$ T cell cytokine production and cytotoxicity. In particular, DCs treated with ZOL are especially effective at activating V γ 9V δ 2 T cells. ZOL pulsed DCs induced greater $\gamma\delta$ T cell proliferation and expanded CD62L+ populations compared to untreated DCs [86,87]. Furthermore, V γ 9V δ 2 T cells activated by ZOL treated immature DCs had upregulated CD40L expression and IFN- γ production, and subsequently enhanced antitumor CD8+ $\alpha\beta$ T cell proliferation [88].

$\gamma\delta$ T cells also interact with and activate NK cells. Maniar et al. showed that IPP expanded V γ 9V δ 2 T cells co-cultured with NK cells increased NK cell cytotoxicity against tumors that are usually resistant to NK cytolysis [89]. These $\gamma\delta$ T cells activate NK cells through co-stimulatory interactions, with CD137L presented on activated $\gamma\delta$ T cells and CD137 expressed on activated NK cells. Activated NK cells augment their NKG2D expression and increase their tumor cell cytotoxicity. In addition, when stimulated with ZOL, V γ 9V δ 2 T cells also enhance NK tumor cell killing through ADCC pathways [89]. Further evidence that $\gamma\delta$ T cells support NK cells comes from in vivo results suggesting a critical role for $\gamma\delta$ T cells in enabling NK cell IFN- γ production during early antibacterial responses [90]. Additional work found that $\gamma\delta$ T cells enhanced NK cell mediated IFN- γ production in early stages after ZOL stimulation, and these effects were formed by interactions between $\gamma\delta$ T cells, DC-like cells, and NK cells [91].

7. Autologous $\gamma\delta$ T Cell Adoptive Transfer Therapy

Early clinical investigations of $\gamma\delta$ T cells predominantly used them autologously (Tables 1 and 2). The first pilot clinical study of $\gamma\delta$ T cells was conducted by Wilhelm et al. in 2003 [92]. Nineteen patients with relapsed or refractory (R/R) MM and low-grade non-Hodgkin lymphoma (NHL) were given intravenous (i.v.) IL-2 and PAM. Several patients experienced low-grade adverse responses to IL-2 infusion, and two experienced a grade 3 response, but none were severe enough to stop treatment. $\gamma\delta$ T cells did not measurably proliferate in an initial cohort of ten patients, and the group did not experience any objective tumor response. An additional nine patients were selected based on in vitro $\gamma\delta$ T cell proliferation in response to PAM/IL-2. In this cohort, five patients were found to have significant in vivo $\gamma\delta$ T cell proliferation after treatment. Of the nine patients, three had partial remission and two additional patients had stable disease. This study reveals the interpatient heterogeneity of $\gamma\delta$ T cell responsiveness to bisphosphonate compound in vivo and highlights the value of prescreening $\gamma\delta$ T cells in vitro before autologous therapeutic use. A later phase I clinical trial conducted by Dieli et al. was the first to evaluate combined ZOL and IL-2 and confirmed the importance of IL-2 for maintaining $\gamma\delta$ T cells in vivo [55]. They compared the effect of giving ZOL alone or in combination with IL-2 on $\gamma\delta$ T cells in eighteen patients with metastatic hormone-refractory prostate cancer. All patients tolerated treatment well: eight patients across both groups developed expected, mild, and transient flu-like symptoms controlled by paracetamol, and two developed mild erythema around

the IL-2 injection site, but no further adverse effects were observed. In nine patients receiving ZOL/IL-2, $\gamma\delta$ T cells proliferated and had an effector memory phenotype, with IFN- γ , TRAIL, and perforin production. Three of these patients had partial remission, and five experienced stable disease. In contrast, $\gamma\delta$ T cells from nine patients receiving only ZOL generally did not proliferate and had lower TRAIL serum concentrations. In this group, only one patient experienced partial remission, and one patient had stable disease.

The benefit of combined ZOL/IL-2 in vivo was further supported by another phase 1 clinical trial in 2010 by Meraviglia et al., who evaluated this treatment in ten terminal patients with advanced metastatic breast cancer [93]. The combination was well-tolerated: five patients developed expected, mild, and transient flu-like symptoms controlled by paracetamol, and two developed mild erythema around the IL-2 injection site, but no further adverse effects were observed. $\gamma\delta$ T cells in three patients proliferated into an effector population with increased IFN- γ production. These patients had reduced CA15-3 tumor marker levels, concomitant with one instance of partial remission and two cases of stable disease. Patients without sustained $\gamma\delta$ T cell numbers had progressively worsening disease. This study, taken in combination with previous results, supports a correlation between $\gamma\delta$ T cell responsiveness and cancer prognosis.

Beyond supplementing endogenous $\gamma\delta$ T cells with stimulatory molecules and cytokines, clinical studies have also assessed the autologous use of $\gamma\delta$ T cells expanded ex vivo (Table 2). The first such investigation, a 2007 pilot study by Kobayashi et al., expanded $\gamma\delta$ T cells from the peripheral blood of seven advanced renal cell carcinoma (RCC) patients [94]. $\gamma\delta$ T cells were cultured with pAg stimulation, and then reinfused into patients along with IL-2. Antitumor effects were reported in five patients, three of which showed demonstrably slower tumor growth, but the other two showed increased growth speed. No patients experienced high-grade adverse effects, though many of them had low-grade (mild or bothersome) adverse effects from IL-2 infusions. Following up their pilot study, Kobayashi et al. took the ex vivo expansion, IL-2-supplemented system and added injection with ZOL several hours before cell infusion for a phase I/II study [95]. Of eleven patients with advanced RCC, one experienced a complete response and five experienced stable disease, but the rest continued to show disease progression. All patients experienced adverse effects spanning grades 1 through 4, but only two patients dropped out of the study, and all others saw a reduction in adverse events as their treatment continued. Compared to previous results, ex vivo stimulation in combination with ZOL and IL-2 significantly improved $\gamma\delta$ T cytotoxicity and therapeutic effects.

One autologous study by Wada et al. identified cell localization as a potential issue to address, and therefore attempted localized injection of the therapeutic product [96]. Seven patients with gastric adenocarcinoma and secondary malignant ascites received ex vivo expanded $\gamma\delta$ T cells injected directly into their peritoneal cavities. Adverse effects were limited to fever and ZOL-induced hypocalcemia, both of which were reversible and mild. One patient in the study experienced reduced ascites volume, and for another patient ascites almost completely disappeared from the peritoneal cavity.

Table 1. Autologous, in vivo expanded $\gamma\delta$ T cell clinical trials.

Year	Author	Phase	Tumor Type	Treatment	Clinical Outcome
2003	Wilhelm et al. [92]	Pilot clinical study	MM, CLL, MZL	PAM + IL-2	<p>Cohort A SD: 1/10, PD: 8/10</p> <ul style="list-style-type: none"> - 1 patient dropped out of the study. - Little to no $\gamma\delta$T proliferation was observed. <p>Cohort B PR: 3/9, SD: 2/9, PD: 4/9</p> <ul style="list-style-type: none"> - $\gamma\delta$T proliferation was observed in 5/9 patients.
2007	Dieli et al. [55]	I	Prostate cancer	ZOL or ZOL + IL-2	<p>ZOL only PD: 3/9, SD: 1/9, PR: 1/9</p> <ul style="list-style-type: none"> - 4 patients dropped out of the study. - 1/4 that could be evaluated had a sustained effector population, and was the only patient with a PR. <p>ZOL + IL-2. PD: 1/9, SD: 4/9, PR: 2/9</p> <ul style="list-style-type: none"> - 2 patients dropped out of the study. - 5/7 had an increased effector population.
2010	Meraviglia et al. [93]	I	Breast cancer	ZOL + IL-2	<p>PD: 4/10, SD: 2/10, PR: 1/10</p> <ul style="list-style-type: none"> - 3 patients dropped out of the study. - Each patient with SD or PD showed sharp decline in their $\gamma\delta$ T cell population either before death or before reduction to SD.
2010	Bennouna et al. [97]	I	Solid tumor variety	BrHPP + IL-2	PD: 16/28, SD: 12/28
2012	Kunzmann et al. [98]	I/II	RCC, MM, AML	ZOL + IL-2	<p>PR: 2/21, SD: 6/21, PD: 12/21</p> <ul style="list-style-type: none"> - 1 patient dropped out of the study.
2011	Lang et al. [99]	Pilot clinical study	RCC	ZOL + IL-2	<p>SD: 5/12</p> <ul style="list-style-type: none"> - 3 patients dropped out of the study.
2016	Pressey et al. [100] (NCT01404702)	I	Neuroblastoma	ZOL + IL-2	PD: 4/4
2022	LAVA Therapeutics (NCT04887259)	I/IIa	CLL, MM	Bispecific Ab for $V\gamma 9V\delta 2$ TCR and CD1d	<p>No patient response has yet been reported.</p> <ul style="list-style-type: none"> - The therapy was safe and well-tolerated.

Abbreviations are as follows: **PD**: progressive disease; **SD**: stable disease; **PR**: partial response; **Ab**: antibody; **ZOL**: zoledronate/zoledronic acid; **PAM**: pamidronate; **BrHPP**: bromohydrin pyrophosphate; **AML**: acute myeloid leukemia; **CLL**: chronic lymphocytic leukemia; **MM**: multiple myeloma; **MZL**: marginal zone lymphoma; **RCC**: renal cell carcinoma.

Table 2. Autologous, ex vivo expanded $\gamma\delta$ T cell clinical trials.

Year	Author	Phase	Tumor Type	Effector Cells	Clinical Outcome
2007	Kobayashi et al. [94] (NCT00588913)	Pilot clinical study	RCC	V γ 9V δ 2 T + IL-2	<i>PD: 7/7</i> 5/7 showed an increased $\gamma\delta$ T cell population.
2008	Bennouna et al. [101]	I	RCC	V γ 9V δ 2 T + IL-2	At one time point, <i>SD: 6/10, PD: 4/10</i> . However, 4 patients were later withdrawn prematurely.
2009	Abe et al. [102]	Pilot clinical study	MM	V γ 9V δ T	<i>PD: 6/6</i> <ul style="list-style-type: none"> - The effector memory Vγ9+ T population increased, as well as the overall Vγ9+ T percentage in bone marrow and number in peripheral blood. - Expression of NKG2D and CD69 increased on Vγ9+ T cells after stimulation. - Increased IFN-γ production.
2010	Nakajima et al. [103] (C000000336)	I	NSCLC	V γ 9V δ 2 T	At one time point, <i>SD: 3/10, PD: 5/10</i> . However, only 6 patients remained at the end of the study.
2011	Kobayashi et al. [95] (NCT00588913)	I/II	RCC	V γ 9V δ 2 T + ZOL + IL-2	<i>PD: 5/11, SD: 5/11, CR: 1/11</i>
2011	Sakamoto et al. [104] (C000000336)	I	NSCLC	V γ 9V δ 2 T	<i>PD: 6/15, SD: 6/15</i> <ul style="list-style-type: none"> - 3 patients dropped out of the study.
2011	Nicol et al. [105]	I	Solid tumors	V γ 9V δ 2 T + ZOL / V γ 9V δ 2 T + ZOL + conventional therapy	V γ 9V δ 2 T+ ZOL only: <i>SD: 3/15, PD: 11/15</i> V γ 9V δ 2 T+ ZOL + other treatments: <i>CR: 1/3, PR: 2/3</i> <ul style="list-style-type: none"> - 1 patient dropped out of the study.
2013	Izumi et al. [106] (UMIN000000854)	-	CRC	V γ 9V δ 2 T	<i>PD: 6/6</i> <ul style="list-style-type: none"> - All patients demonstrated expansion of the Vγ9Vδ2 T population.
2014	Kakimi et al. [107] (C000000336)	I	NSCLC	V γ 9V δ 2 T	<i>PD: 6/15, SD: 6/15</i> <ul style="list-style-type: none"> - 3 patients dropped out of the study. - Increased IFN-γ release was observed. - Interestingly, Vγ9Vδ2 T population expansion did not correlate with prognosis.
2014	Wada et al. [96] (UMIN000004130)	Pilot clinical study	Gastric cancer	V γ 9V δ 2 T + ZOL	<i>PD: 7/7</i> <ul style="list-style-type: none"> - Reduction in tumor ascites.
2015	Cui et al. [108]	Pilot clinical study	Gastric cancer	Chemotherapy + $\gamma\delta$ T/NK/Killer cells	Progression-free survival of the group infused with effectors was significantly higher than those treated with chemotherapy alone: 70% vs. 46.4% survived over the two years, respectively.
2017	Gadeta (NL6357)	I	AML, MM, MDS	V γ 9V δ 2 TCR transduced $\alpha\beta$ T cells	-

Abbreviations are as follows: **PD**: progressive disease; **SD**: stable disease; **PR**: partial response; **CR**: complete response/remission; **ZOL**: zoledronate/zoledronic acid; **AML**: acute myeloid leukemia; **CRC**: colorectal cancer; **MDS**: myelodysplastic syndrome; **MM**: multiple myeloma; **NSCLC**: non-small cell lung cancer; **RCC**: renal cell carcinoma.

8. Allogeneic $\gamma\delta$ T Cell Adoptive Transfer Therapy

In almost all autologous studies, treatments showed some promise and reduced tumor burden in at least a few patients, but the effects were inconsistent. Researchers looked for a way to standardize the cell product: introducing allogeneic therapeutic cells harnessed from healthy donors into multiple patients (Tables 3 and 4). In a pilot study, Wilhelm et al. took a partial step toward this by selecting donors who were half-matched (HLA-haploidentical) family members of four patients with advanced hematological malignancies [109]. These patients had been judged ineligible for allogeneic stem cell therapy due to its high toxicity. Prior to infusion, all patients underwent lymphocyte-depleting chemotherapy. They were then injected with 99% pure $\gamma\delta$ T cells from the family donors, along with ZOL on day 0 and repeated IL-2 infusions afterwards to promote cell expansion. Neutropenia occurred within the expected time range, with hematopoiesis recovering several weeks after the chemotherapy, and no patient experienced GvHD or organ injury. Patients experienced a mean 68-fold increase of donor $\gamma\delta$ T cells, peaking around day eight, but persisting up to 28 days within the body. One patient passed away due to an infection unrelated to treatment, but the remaining three patients all experienced complete remission as a result of the therapy. This pilot study indicated that the use of allogeneic $\gamma\delta$ T cells is feasible and safe, and that ZOL/IL-2 infusions can activate and expand allogeneic $\gamma\delta$ T cells in vivo to achieve promising therapeutic response. Even with non-family donors, the safety of allogeneic $\gamma\delta$ T cell therapy remained. A case study by Alnaggar et al. gave eight infusions of $\gamma\delta$ T cells expanded from a healthy donor to a patient with cholangiocarcinoma over the course of one year [110]. The patient experienced no adverse effects. And even without ZOL or IL-2 supplementation, he experienced depleted tumor activity, tumor size reduction, and improved quality of life during and after the treatment.

Table 3. Allogeneic, ex vivo expanded $\gamma\delta$ T cell clinical trials.

Year	Author	Phase	Tumor Type	Effector Cells	Clinical Outcome
2014	Wilhelm et al. [109]	Pilot clinical study	NHL, MM, AML, PCL	V γ 9V δ 2 T, with some CD4-/8- $\alpha\beta$ T + ZOL + IL-2	CR: 3/4 - 1 patient dropped out of the study. - Patients experienced an average 68-fold expansion of donated V γ 9V δ 2 T cells.
2019	Alnaggar et al. [110]	Case Study	Cholangiocarcinoma	V γ 9V δ 2 T, 8 infusions	CR: 1/1 - Lack of adverse effects.
2020	Lin et al. [111] (NCT03180437)	I/II	Pancreatic cancer	IRE + V γ 9V δ 2 T	- Patients receiving multiple courses of $\gamma\delta$ T cell infusion had a statistically significant extension in length of life.

Abbreviations are as follows: **CR**: complete response/remission; **ZOL**: zoledronate/zoledronic acid; **IRE**: Irreversible electroporation; **AML**: acute myeloid leukemia; **MM**: multiple myeloma; **NHL**: non-Hodgkin's lymphoma; **PCL**: plasma cell leukemia.

Table 4. Allogeneic, ex vivo expanded CAR- $\gamma\delta$ T cell clinical trials.

Year	Author	Phase	Tumor Type	Effector Cells	Clinical Outcome
2017	Beijing Doing Biomedical (NCT02656147)	I	B-Cell Lymphoma, ALL, CLL	CD19 CAR- $\gamma\delta$ T	-
2019	CytoMed Therapeutics (NCT04107142)	I	Solid tumors	NKG2DL CAR- $\gamma\delta$ T	-
2021	Adicet Bio (NCT04735471)	I	B-cell malignancies	CD20 CAR-V δ 1 T + IL-2	Interim result—CR: 2/4, PR: 1/4, PD: 1/4
2021	PersonGen BioTherapeutics (NCT04702841)	I	CD7+ T lymphoma (ALL)	CD7 CAR- $\gamma\delta$ T	-

Abbreviations are as follows: **PD**: progressive disease; **PR**: partial response; **CR**: complete response/remission; **CAR**: chimeric antigen receptor; **ALL**: acute lymphocytic leukemia; **CLL**: chronic lymphocytic leukemia.

9. CAR-Engineered $\gamma\delta$ T Cell Adoptive Transfer Therapy

With clear evidence that adverse effects would be minor and reversible, and that $\gamma\delta$ T cells had the potential to achieve a complete response, researchers genetically manipulated the cells to increase their antitumor functions. In one of the earliest pre-clinical studies of CAR- $\gamma\delta$ T cells, Deniger et al. electroporated polyclonal $\gamma\delta$ T cells with Sleeping Beauty (SB) transposon and transposase to overexpress a CD19 CAR and expanded them with artificial antigen-presenting cells (aAPCs) engineered to express CD19 [112]. CAR- $\gamma\delta$ T cells and IL-2 were later intravenously injected into NSG mice xenografted with a CD19+ leukemia cell line. The group administered the cells and IL-2 weekly for three weeks, and results indicated that polyclonal CAR- $\gamma\delta$ T cells can suppress tumor growth in this pre-clinical setting.

In 2020, Ang et al. expanded V γ 9V δ 2 T cells with Zometa/IL-2 and engineered the cells with a CAR based on the extracellular domain of NKG2D (NKG2DL) using mRNA electroporation [113]. The NKG2DL CAR- $\gamma\delta$ T cells were capable of killing multiple solid tumor cell lines in vitro, including those resistant to Zometa treatment. In addition, they conducted pre-clinical studies in mice previously intraperitoneally injected with colon tumors or ovarian tumors and demonstrated NKG2DL CAR- $\gamma\delta$ T cells caused tumor regression in mice and extended their survival in both tumor models.

Another group, Rozenbaum et al., expanded CD19 CAR-V γ 9V δ 2 T cells using ZOL/IL-2. They gave CD19+ tumor bearing NSG mice the therapeutic cells plus ZOL and observed tumor reduction [114]. Importantly, the group generated a CD19 knockout tumor model by CRISPR/Cas9 and showed that CD19 CAR-V γ 9V δ 2 T cells in the presence of ZOL had enhanced cytotoxicity against CD19- tumor cells compared to conventional CD19-CAR T cells. This important finding suggests that CAR-V γ 9V δ 2 T cells combined with a bisphosphonate compound could target tumors in an antigen escape situation through CAR-independent pathways, significantly increasing their therapeutic potential in clinics.

Table 4 summarizes past and current CAR- $\gamma\delta$ T cell clinical trials. A phase I clinical trial (NCT02656147) to assess the safety and efficacy of CD19 CAR- $\gamma\delta$ T cells for the treatment of lymphoma and leukemia was conducted by Beijing Doing Biomedical. Another phase I clinical trial (NCT04107142) performed by CytoMed Therapeutics has evaluated the safety and tolerability of haploidentical/allogeneic NKG2DL CAR- $\gamma\delta$ T cells in patients with R/R solid tumors, including colorectal cancer, triple-negative breast cancer, sarcoma, nasopharyngeal cancer, prostate cancer, and gastric cancer. Although both studies have been completed, no formal reports on their results have yet been published. Currently, PersonGen BioTherapeutics is validating its CD7 CAR- $\gamma\delta$ T cells in patients with R/R CD7+ T cell-derived malignant tumors in an early Phase I study (NCT04702841). Adicet Bio is testing CD20 CAR-V δ 1+ T cells (ADI-001) in patients with B cell malignancies in another Phase I trial (NCT04735471). In December 2021, Adicet Bio announced very positive interim clinical data from this study. Of the four efficacy evaluable patients, three received CAR-V δ 1+ T cells at a lower dose (30 million CAR+ cells) and one received a higher dose (100 million CAR+ cells). Among the low dose patients, one achieved a complete response (CR), one achieved a near CR, and one had progressive disease (PD). For the higher dose patient, a CR was achieved. This encouraging result granted the company's lead candidate ADI-001 "Fast Track Designation" from the FDA.

10. Other $\gamma\delta$ T Cell-Based Therapies

Efforts to optimize $\gamma\delta$ T cell-based immunotherapies have resulted in a diversity of approaches to supplement the unique properties of $\gamma\delta$ T cells, increase efficacy, and reduce side effects. One approach is harnessing a high affinity V γ 9V δ 2 TCR from $\gamma\delta$ T cells and overexpressing it in $\alpha\beta$ T cells. The engineered cell products (TEGs) possess the properties of $\alpha\beta$ T cells, such as robust proliferation and longevity, with the additional targeting ability of the V γ 9V δ 2 TCR to BTN molecules on tumor cells without MHC restriction. After verifying these cells' efficacy and safety in a PDX leukemia model [115], Gadeta began a Phase I clinical trial (NTR6541), assessing its lead candidate, TEG001, in patients who had exhausted other treatment options for B cell malignancies, high-risk Myelodysplastic

Syndrome, or MM (Table 2). Conventional CAR constructs combine the CD3 ζ intracellular signaling domain and co-stimulatory domain, enabling one antigen binding event to generate the two activating signals required for T cell stimulation. Fisher et al. sought to avoid overactivation of a therapeutic product against any healthy cells that may express a CAR target antigen [116]. They attempted to minimize reactions against healthy cells by GD2 CAR- $\gamma\delta$ T cells by separating these activating signals. By replacing both CD3 ζ and the CD28 co-stimulatory domain with DAP10 (the NKG2D co-stimulatory domain) in their CAR construct, two independent tumor binding events—V γ 9V δ 2 TCR signaling plus CAR signaling—were required for cell activation. This separated signaling method could develop safer immunotherapies with less potential for off-target effects, allowing broader target selection.

Additional immunotherapy approaches have attempted to improve targeting and efficacy by developing bispecific antibodies featuring two independent domains that link T cells to tumor cells. It is difficult to target an antibody to an inherently variable $\alpha\beta$ TCR, so researchers primarily chose to target CD3 [117]. Two FDA-approved examples are blinatumomab, which targets CD3 and CD19 [118], and catumaxomab, which targets CD3 and EpCAM [119], and can also be recognized by CD16 on the $\gamma\delta$ T cell membrane. However, because CD3 is ubiquitous among different TCRs, anti-CD3 antibodies can indiscriminately activate all T cell subsets and induce cytokine storms, causing significant adverse effects in the first few cycles of the drug. To reduce these side effects, finer effector cell selectivity is ideal. The consistent and unique molecules that comprise the $\gamma\delta$ TCR present a steadier target for selective stimulation of one cell type. A bispecific antibody binding the V γ 9 chain and HER2 tumor antigen enhanced V γ 9V δ 2 T cell in vitro and in vivo killing of pancreatic adenocarcinomas, with superior in vitro killing compared to pAg stimulated $\gamma\delta$ T cells alone or with a CD3/HER2 bispecific antibody [120]. An additional report with a V γ 9/CD123 bispecific antibody in acute myeloid leukemia (AML) provided further support for a V γ 9 targeting strategy over a CD3 targeting strategy [121]. This bispecific antibody selectively stimulated V γ 9+ T cells, with significantly less production of cytokine storm associated IL-6 and IL-10 compared to a CD3/CD123 antibody. The V γ 9/CD123 antibody mediated $\gamma\delta$ T cell conjugation with tumor cells, and induced cytotoxicity against antigen expressing tumors both in vitro and in vivo. Currently, Lava Therapeutics is conducting a Phase I/II trial (NCT04887259) for LAVA-051, an antibody which binds both the V δ 2 chain and CD1d, allowing for selective stimulation of V γ 9V δ 2 T cells and NKT cells to target blood cancers such as chronic lymphocytic leukemia (CLL), MM, and AML (Table 1). In their preliminary data, the V δ 2/CD1d antibody was well-tolerated without cytokine storm syndrome and circulating V γ 9V δ 2 T cells effectively engaged the antibody and displayed a more activated phenotype after treatment.

Other than the approaches discussed above, stem cell derived $\gamma\delta$ T cells for adoptive transfer cell therapy could be another promising direction. Watanabe et al. generated induced pluripotent stem cells (iPSCs) that harbor rearrangements of the TCR gamma chain (V γ 9) and TCR delta chain (V δ 2) gene regions ($\gamma\delta$ T-iPSCs) [122]. They activated PBMC $\gamma\delta$ T cells with IL-2 and ZOL, then transduced those cells with the Sendai virus vector encoded KLF4, c-MYC, OCT3/4, and SOX2 transcription factor genes. They successfully reprogrammed the cells into iPSC lines that were pluripotent and capable of differentiating into embryoid bodies (EB) and hematopoietic progenitor cells. However, they did not continue to differentiate the cells into fully mature $\gamma\delta$ T cells. Zeng et al. also reprogrammed $\gamma\delta$ T-iPSC lines from V γ 9V δ 2 T cells [123]. They further differentiated the cells into NK-like $\gamma\delta$ T cells, which they designated “ $\gamma\delta$ natural killer T” ($\gamma\delta$ NKT) cells. The group used a “NK cell-promoting protocol” and utilized OP9 mouse stromal cells overexpressing DLL1 Notch ligands to support the differentiation process. The iPSC derived $\gamma\delta$ NKT cells exhibited a NK cell-like phenotype and expressed many NK receptors such as NKG2D, DNAM-1, NKp30, NKp44, NKp46, and CD16. An in vitro cytotoxicity assay demonstrated that these cells are powerful killers of various cancer cell lines, especially solid tumors.

11. $\gamma\delta$ T Cell-Based Cancer Immunotherapy: Challenges and Perspectives

The past several decades of immunology research have yielded great progress in our understanding of the molecular mechanisms underlying antigen presentation to $\alpha\beta$ T cells and their subsequent response. However, progress in understanding these mechanisms in $\gamma\delta$ T cells has been comparatively slow and is impeded by limited knowledge of what antigens these cells respond to [124]. $V\gamma9V\delta2$ T cells were initially thought to directly recognize pAgs produced by microbial pathogens and tumor cells with abnormal MVA pathway activation [125]. Increasing structural and biochemical evidence evolved this understanding, indicating instead that the $V\gamma9V\delta2$ TCR reacts to BTN3A1 on the cell surface, which is bound at its intracellular B30.2 domain by pAgs [24]. More recent evidence has refined this paradigm, proposing that a BTN2A1-BTN3A1 complex acts as a dual ligand binding the $V\gamma9V\delta2$ TCR in response to intracellular pAg binding the B30.2 domain [27]. A better understanding of mechanisms of $\gamma\delta$ T cell activation and its TCR targeting may allow us to maximize the potential of $\gamma\delta$ T cell-based therapy. While certain subsets of $\gamma\delta$ T cells have been selected as potential immunotherapy agents, other subsets were identified as pro-tumoral $\gamma\delta$ T cells and have immunosuppressive roles in some cancers [126,127]. Peng et al. showed that a dominant $V\delta1$ T cell population infiltrating breast tumors were associated with immunosuppressive functions by inhibiting naive and effector T cell responses and by impairing DC maturation and function [128]. In 2014, Wu et al. reported IL-17-producing $\gamma\delta$ T cells were found to have a pro-tumoral role [129]. In particular, $V\delta1$ T cells were the primary source of IL-17 in colorectal cancer chronic inflammation. Furthermore, these $V\delta1$ T cells produced TNF, IL-8, and granulocyte-macrophage colony-stimulating factor (GM-CSF), which attracted immunosuppressive myeloid-derived suppressor cells (MDSCs) into the tumor microenvironment to support immunosuppressive activity [129]. Besides these two reports, several studies also found that $\gamma\delta$ T cells may have immunosuppressive roles in different cancers [130–133]. Although $\gamma\delta$ T cells have tremendous potential for therapeutic applications in cell-based immunotherapy, their pro-tumoral mechanisms need to be studied further to prevent their possible deleterious functions for cancer treatment.

Another challenge of $\gamma\delta$ T cell-based cancer immunotherapy is that although several studies have reported that $\gamma\delta$ T cells can modulate immunosuppressive cells in the TME, the lack of nutrients, presence of inhibitory molecules, and hypoxia in the TME may still limit the therapeutic potential of $\gamma\delta$ T cells [67,134]. Inhibitory molecules such as TGF- β [135], prostaglandin-E2 (PGE2) [136], adenosine [137], and soluble NKG2D ligands [138] produced by tumor cells and other cells of the TME can interfere with the proliferation and function of $\gamma\delta$ T cells. Gonnermann et al. reported that Colo357 cells (pancreatic adenocarcinoma) develop resistance to $\gamma\delta$ T cell lysis by enhancing the expression of cyclooxygenase 2 (Cox-2) [139]. This enzyme catalyzes the metabolism of arachidonic acid to prostanoids, such as PGE2, which in turn promotes tumorigenesis, tumor progression, and metastasis. This Cox-2 metabolite contributes to establishing immunosuppressive factors in the TME and inhibits the antitumor activities of immune cells. However, the PGE2-mediated inhibition of $\gamma\delta$ T cell cytotoxicity can be partially overcome with Cox-2 inhibitors [139]. This suggests that the use of Cox-2 inhibitors with $\gamma\delta$ T cells or other cell types may improve the cytotoxic functions of immune cells in the TME and have beneficial effects in clinics.

Exhaustion is a hyporesponsive cell state induced by chronic stimulation from repeated antigen exposure, and manifests as a loss of effector functions including proliferation, cytokine production, and cytotoxicity. These changes are concomitant with the expression of inhibitory receptors such as PD-1, TIM-3, LAG-3, and CTLA-4 [140,141]. Although exhaustion may have homeostatic function in limiting immunopathology in chronic infections, in immunotherapy it can be an obstacle reducing the efficacy of cell therapies. CD8+ T cell exhaustion has been documented in many cancers, and this process also occurs in $\gamma\delta$ T cells [142]. In a nonhuman primate model, $V\gamma9V\delta2$ T cells repeatedly stimulated with pAgs and IL-2 had lower expansion rates upon restimulation, compared to the initial pAg response. $\gamma\delta$ T cells from CLL patients failed to proliferate in response to IPP accumulation

in CLL cells, indicating an exhausted state that could arise from prolonged stimulation from these patients' cancer cells [143]. The TME often has upregulated expression of ligands for inhibitory receptors present on immune cells, which engage to enforce an exhausted state. In one report, MM patients' bone marrow (BM) had upregulated PD-L1 and contained V γ 9V δ 2 T cells with greater PD-1 expression than $\gamma\delta$ T cells from other regions, as well as reduced proliferation response to ZOL [144]. This study found that an anti-PD-1 treatment conferred a fivefold increase in proliferation of MM BM $\gamma\delta$ T cells after ZOL stimulation. Another study found that anti-PD-1 treatment increased $\gamma\delta$ T cell IFN- γ production both with direct ZOL stimulation, and from ZOL-treated primary AML cells [145]. Using $\gamma\delta$ T cells combined with immune checkpoint inhibitors is thus appealing to preemptively guard against exhaustion from sustained interactions with tumors, maintaining cell effector functions for a more efficacious therapy.

$\gamma\delta$ T cells are especially prone to exhaustion when differentiated toward an effector memory (T_{EM}) or terminally differentiated effector memory (T_{EMRA}) state [146,147]. T cells progress to more T_{EM} and T_{EMRA} states in response to activation signals and stimulation, including tumor recognition [148,149]. $\gamma\delta$ T cell memory states have traditionally been defined by the expression of CD45RA and CD27, as well as several lymph node homing receptors [147]. Naive $\gamma\delta$ T cells are the earliest memory state. They are CD45RA+CD27+ and express the lymph node homing markers CD62L and CCR7, and are infrequent in circulation but are the predominant $\gamma\delta$ T cell memory type in lymph nodes [147]. Naive $\gamma\delta$ T cells proliferate in response to pAg but have minimal cytokine releasing activity. These cells have strong self-renewal potential to yield a population with the same memory state, a property which diminishes with greater differentiation. After stimulation, naive $\gamma\delta$ T cells remain CD62L+ and CCR7+ but progress to a CD45RA-CD27+ central memory (T_{CM}) state that is CD45RO+ and mostly found in the peripheral blood [147]. T_{CM} $\gamma\delta$ T cells have greater proliferative potential in response to pAg compared to naive cells, but still have reduced effector functions such as cytokine release. Sites of inflammation contain further differentiated $\gamma\delta$ T cells without CD62L or CCR7, including CD45RA-CD27- T_{EM} cells and CD45RA+CD27- T_{EMRA} cells. T_{EM} $\gamma\delta$ T cells have reduced proliferative potential and minimal self-renewal ability, but vigorously release cytokines including IFN- γ . Although T_{EMRA} $\gamma\delta$ T cells exert potent cytotoxicity against tumor cells, they soon become exhausted when they are excessively restimulated by tumor antigens. Exhausted T cells not only lose the ability to proliferate and self-renew, but also have weakened antitumor functions [150].

$\gamma\delta$ T cell expansion protocols induce $\gamma\delta$ TCR signaling to yield practical cell numbers at the expense of the population's memory state. While the original $\gamma\delta$ T cells being expanded may contain some $\gamma\delta$ T cells in earlier memory states, stimulation differentiates the resultant population toward effector and terminally differentiated effector memories. Therapeutically, these further differentiated populations could initially yield potent antitumor effects, but longer-term tumor control would be limited by their low proliferation/self-renewal ability and persistence. Instead, maintaining the naive or central memory status of $\gamma\delta$ T cells is desirable for clinical use. These populations have long persistence, which is amenable for ongoing tumor surveillance; upon antigen stimulation they proliferate greatly to yield a large population of tumor responsive effector cells while reserving a subset of self-renewing memory cells to generate future waves of effectors against subsequent tumor encounters. Since current $\gamma\delta$ T cell expansion protocols rely on stimulatory signals which induce differentiation away from these memory states, generating a population of naive and central memory $\gamma\delta$ T cells in vitro large enough for therapeutic use could be challenging. However, such direction is crucial for therapeutic efficacy. Fortunately, there have been continuous advances in knowledge about $\gamma\delta$ T cell biology and culture conditions. As $\gamma\delta$ T cells increasingly attract research attention for their immunotherapeutic potential, growing bodies of work in these areas could yield key insights to effectively generate $\gamma\delta$ T cells that have superior persistence and efficacy against tumors.

Author Contributions: D.L. researched data and provided guidance on the outline of this review. D.L., C.J.R., N.E.P. and Z.S.D. all contributed to the writing as well as the review/editing. Y.Z. helped

to organize the content and provided suggestion of manuscript writing. L.Y. reviewed and finalized the article. All authors contributed to the manuscript and approved the submitted version. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by a Partnering Opportunity for Translational Research Projects Award from the California Institute for Regenerative Medicine (TRAN1-12250, to L.Y.), a UCLA BSCRC Innovation Award (to L.Y.), an Ablon Scholars Award (to L.Y.), and a T32 Microbial Pathogenesis Training Grant (Ruth L. Kirschstein National Research Service Award, T32-AI007323) (to D.L.).

Acknowledgments: We want to thank University of California, Los Angeles (UCLA) and members of Lili Yang's lab for providing supports. We also want to thank BioRender.com, both Figure 1 (Licensing Agreement number: AS23VHMVTV, accessed on 3 May 2022) and Figure 2 (Licensing Agreement number: HH23VHMQBUB, accessed on 3 May 2022) were created by this website.

Conflicts of Interest: L.Y. is a scientific advisor to AlzChem and Amberstone Biosciences, and a co-founder, stockholder, and advisory board member of Appia Bio. L.Y., D.L. and Y.Z. are listed on the IP-related documents filed by UCLA. None of the declared companies contributed to or directed any of the research reported in this article. The remaining authors declare that they have no competing interests.

References

1. Marofi, F.; Motavalli, R.; Safonov, V.A.; Thangavelu, L.; Yumashev, A.V.; Alexander, M.; Shomali, N.; Chartrand, M.S.; Pathak, Y.; Jarahian, M.; et al. CAR T cells in solid tumors: Challenges and opportunities. *Stem Cell Res. Ther.* **2021**, *12*, 81. [[CrossRef](#)] [[PubMed](#)]
2. Gentles, A.J.; Newman, A.M.; Liu, C.L.; Bratman, S.V.; Feng, W.; Kim, D.; Nair, V.S.; Xu, Y.; Khuong, A.; Hoang, C.D.; et al. The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nat. Med.* **2015**, *21*, 938–945. [[CrossRef](#)] [[PubMed](#)]
3. Minculescu, L.; Sengelov, H. The role of gamma delta T cells in haematopoietic stem cell transplantation. *Scand. J. Immunol.* **2015**, *81*, 459–468. [[CrossRef](#)] [[PubMed](#)]
4. Handgretinger, R.; Schilbach, K. The potential role of gammadelta T cells after allogeneic HCT for leukemia. *Blood* **2018**, *131*, 1063–1072. [[CrossRef](#)]
5. Parker, C.M.; Groh, V.; Band, H.; Porcelli, S.A.; Morita, C.; Fabbi, M.; Glass, D.; Strominger, J.L.; Brenner, M.B. Evidence for extrathymic changes in the T cell receptor gamma/delta repertoire. *J. Exp. Med.* **1990**, *171*, 1597–1612. [[CrossRef](#)]
6. Kunzmann, V.; Wilhelm, M. Anti-lymphoma effect of gammadelta T cells. *Leuk. Lymphoma* **2005**, *46*, 671–680. [[CrossRef](#)]
7. Di Lorenzo, B.; Simoes, A.E.; Caiado, F.; Tieppo, P.; Correia, D.V.; Carvalho, T.; da Silva, M.G.; Dechanet-Merville, J.; Schumacher, T.N.; Prinz, I.; et al. Broad Cytotoxic Targeting of Acute Myeloid Leukemia by Polyclonal Delta One T Cells. *Cancer Immunol. Res.* **2019**, *7*, 552–558. [[CrossRef](#)]
8. Kabelitz, D.; He, W. The multifunctionality of human Vgamma9Vdelta2 gammadelta T cells: Clonal plasticity or distinct subsets? *Scand. J. Immunol.* **2012**, *76*, 213–222. [[CrossRef](#)]
9. Pistoia, V.; Tumino, N.; Vacca, P.; Veneziani, I.; Moretta, A.; Locatelli, F.; Moretta, L. Human gammadelta T-Cells: From Surface Receptors to the Therapy of High-Risk Leukemias. *Front. Immunol.* **2018**, *9*, 984. [[CrossRef](#)]
10. Silva-Santos, B.; Mensurado, S.; Coffelt, S.B. gammadelta T cells: Pleiotropic immune effectors with therapeutic potential in cancer. *Nat. Rev. Cancer* **2019**, *19*, 392–404. [[CrossRef](#)]
11. Morita, C.T.; Parker, C.M.; Brenner, M.B.; Band, H. TCR usage and functional capabilities of human gamma delta T cells at birth. *J. Immunol.* **1994**, *153*, 3979–3988. [[PubMed](#)]
12. McVay, L.D.; Carding, S.R.; Bottomly, K.; Hayday, A.C. Regulated expression and structure of T cell receptor gamma/delta transcripts in human thymic ontogeny. *EMBO J.* **1991**, *10*, 83–91. [[CrossRef](#)] [[PubMed](#)]
13. Casorati, G.; De Libero, G.; Lanzavecchia, A.; Migone, N. Molecular analysis of human gamma/delta+ clones from thymus and peripheral blood. *J. Exp. Med.* **1989**, *170*, 1521–1535. [[CrossRef](#)] [[PubMed](#)]
14. Bos, J.D.; Teunissen, M.B.; Cairo, I.; Krieg, S.R.; Kapsenberg, M.L.; Das, P.K.; Borst, J. T-cell receptor gamma delta bearing cells in normal human skin. *J. Invest. Dermatol.* **1990**, *94*, 37–42. [[CrossRef](#)]
15. Deusch, K.; Luling, F.; Reich, K.; Classen, M.; Wagner, H.; Pfeffer, K. A major fraction of human intraepithelial lymphocytes simultaneously expresses the gamma/delta T cell receptor, the CD8 accessory molecule and preferentially uses the V delta 1 gene segment. *Eur. J. Immunol.* **1991**, *21*, 1053–1059. [[CrossRef](#)]
16. Sandberg, Y.; Almeida, J.; Gonzalez, M.; Lima, M.; Barcena, P.; Szczepanski, T.; van Gestel-Mol, E.J.; Wind, H.; Balanzategui, A.; van Dongen, J.J.; et al. TCRgammadelta+ large granular lymphocyte leukemias reflect the spectrum of normal antigen-selected TCRgammadelta+ T-cells. *Leukemia* **2006**, *20*, 505–513. [[CrossRef](#)]
17. Qi, C.; Wang, Y.; Li, P.; Zhao, J. Gamma Delta T Cells and Their Pathogenic Role in Psoriasis. *Front. Immunol.* **2021**, *12*, 627139. [[CrossRef](#)]

18. Davey, M.S.; Willcox, C.R.; Hunter, S.; Kasatskaya, S.A.; Remmerswaal, E.B.M.; Salim, M.; Mohammed, F.; Bemelman, F.J.; Chudakov, D.M.; Oo, Y.H.; et al. The human Vdelta2⁺ T-cell compartment comprises distinct innate-like Vgamma9⁺ and adaptive Vgamma9⁻ subsets. *Nat. Commun.* **2018**, *9*, 1760. [[CrossRef](#)]
19. McVay, L.D.; Carding, S.R. Extrathymic origin of human gamma delta T cells during fetal development. *J. Immunol.* **1996**, *157*, 2873–2882.
20. Papadopoulou, M.; Tieppo, P.; McGovern, N.; Gosselin, F.; Chan, J.K.Y.; Goetzeluk, G.; Dauby, N.; Cogan, A.; Donner, C.; Ginhoux, F.; et al. TCR Sequencing Reveals the Distinct Development of Fetal and Adult Human Vgamma9Vdelta2 T Cells. *J. Immunol.* **2019**, *203*, 1468–1479. [[CrossRef](#)]
21. Willcox, C.R.; Davey, M.S.; Willcox, B.E. Development and Selection of the Human Vgamma9Vdelta2⁺ T-Cell Repertoire. *Front. Immunol.* **2018**, *9*, 1501. [[CrossRef](#)] [[PubMed](#)]
22. Kenna, T.; Golden-Mason, L.; Norris, S.; Hegarty, J.E.; O’Farrelly, C.; Doherty, D.G. Distinct subpopulations of gamma delta T cells are present in normal and tumor-bearing human liver. *Clin. Immunol.* **2004**, *113*, 56–63. [[CrossRef](#)] [[PubMed](#)]
23. Wang, L.; Xu, M.; Wang, C.; Zhu, L.; Hu, J.; Chen, S.; Wu, X.; Li, B.; Li, Y. The feature of distribution and clonality of TCR gamma/delta subfamilies T cells in patients with B-cell non-Hodgkin lymphoma. *J. Immunol. Res.* **2014**, *2014*, 241246. [[CrossRef](#)]
24. Rigau, M.; Uldrich, A.P.; Behren, A. Targeting butyrophilins for cancer immunotherapy. *Trends Immunol.* **2021**, *42*, 670–680. [[CrossRef](#)] [[PubMed](#)]
25. Guerra, B.; Recio, C.; Aranda-Tavio, H.; Guerra-Rodriguez, M.; Garcia-Castellano, J.M.; Fernandez-Perez, L. The Mevalonate Pathway, a Metabolic Target in Cancer Therapy. *Front. Oncol.* **2021**, *11*, 626971. [[CrossRef](#)]
26. Freed-Pastor, W.A.; Mizuno, H.; Zhao, X.; Langerod, A.; Moon, S.H.; Rodriguez-Barrueco, R.; Barsotti, A.; Chicas, A.; Li, W.; Polotskaia, A.; et al. Mutant p53 disrupts mammary tissue architecture via the mevalonate pathway. *Cell* **2012**, *148*, 244–258. [[CrossRef](#)]
27. Rigau, M.; Ostrouska, S.; Fulford, T.S.; Johnson, D.N.; Woods, K.; Ruan, Z.; McWilliam, H.E.G.; Hudson, C.; Tutuka, C.; Wheatley, A.K.; et al. Butyrophilin 2A1 is essential for phosphoantigen reactivity by gammadelta T cells. *Science* **2020**, *367*, eaay5516. [[CrossRef](#)]
28. Malinowska, M.; Tokarz-Deptula, B.; Deptula, W. Butyrophilins: An important new element of resistance. *Cent. Eur. J. Immunol.* **2017**, *42*, 399–403. [[CrossRef](#)]
29. Salot, S.; Laplace, C.; Saiagh, S.; Bercegeay, S.; Tenaud, I.; Cassidanius, A.; Romagne, F.; Dreno, B.; Tiollier, J. Large scale expansion of gamma 9 delta 2 T lymphocytes: Innacell gamma delta cell therapy product. *J. Immunol. Methods* **2007**, *326*, 63–75. [[CrossRef](#)]
30. Deniger, D.C.; Moyes, J.S.; Cooper, L.J. Clinical applications of gamma delta T cells with multivalent immunity. *Front. Immunol.* **2014**, *5*, 636. [[CrossRef](#)]
31. Kondo, M.; Sakuta, K.; Noguchi, A.; Ariyoshi, N.; Sato, K.; Sato, S.; Sato, K.; Hosoi, A.; Nakajima, J.; Yoshida, Y.; et al. Zoledronate facilitates large-scale ex vivo expansion of functional gammadelta T cells from cancer patients for use in adoptive immunotherapy. *Cytotherapy* **2008**, *10*, 842–856. [[CrossRef](#)] [[PubMed](#)]
32. Kondo, M.; Izumi, T.; Fujieda, N.; Kondo, A.; Morishita, T.; Matsushita, H.; Kakimi, K. Expansion of human peripheral blood $\gamma\delta$ T cells using zoledronate. *J. Vis. Exp.* **2011**, *55*, e3182. [[CrossRef](#)]
33. Nussbaumer, O.; Gruenbacher, G.; Gander, H.; Komuczki, J.; Rahm, A.; Thurnher, M. Essential requirements of zoledronate-induced cytokine and gammadelta T cell proliferative responses. *J. Immunol.* **2013**, *191*, 1346–1355. [[CrossRef](#)] [[PubMed](#)]
34. Kunzmann, V.; Bauer, E.; Wilhelm, M. Gamma/delta T-cell stimulation by pamidronate. *N. Engl. J. Med.* **1999**, *340*, 737–738. [[CrossRef](#)]
35. Marcu-Malina, V.; Heijhuurs, S.; van Buuren, M.; Hartkamp, L.; Strand, S.; Sebestyen, Z.; Scholten, K.; Martens, A.; Kuball, J. Redirecting alphabeta T cells against cancer cells by transfer of a broadly tumor-reactive gammadeltaT-cell receptor. *Blood* **2011**, *118*, 50–59. [[CrossRef](#)]
36. Luoma, A.M.; Castro, C.D.; Adams, E.J. gammadelta T cell surveillance via CD1 molecules. *Trends Immunol.* **2014**, *35*, 613–621. [[CrossRef](#)]
37. Bai, L.; Picard, D.; Anderson, B.; Chaudhary, V.; Luoma, A.; Jabri, B.; Adams, E.J.; Savage, P.B.; Bendelac, A. The majority of CD1d-sulfatide-specific T cells in human blood use a semiinvariant Vdelta1 TCR. *Eur. J. Immunol.* **2012**, *42*, 2505–2510. [[CrossRef](#)]
38. Russano, A.M.; Agea, E.; Corazzi, L.; Postle, A.D.; De Libero, G.; Porcelli, S.; de Benedictis, F.M.; Spinozzi, F. Recognition of pollen-derived phosphatidyl-ethanolamine by human CD1d-restricted gamma delta T cells. *J. Allergy Clin. Immunol.* **2006**, *117*, 1178–1184. [[CrossRef](#)]
39. Agea, E.; Russano, A.; Bistoni, O.; Mannucci, R.; Nicoletti, I.; Corazzi, L.; Postle, A.D.; De Libero, G.; Porcelli, S.A.; Spinozzi, F. Human CD1-restricted T cell recognition of lipids from pollens. *J. Exp. Med.* **2005**, *202*, 295–308. [[CrossRef](#)]
40. Mangan, B.A.; Dunne, M.R.; O’Reilly, V.P.; Dunne, P.J.; Exley, M.A.; O’Shea, D.; Scotet, E.; Hogan, A.E.; Doherty, D.G. Cutting edge: CD1d restriction and Th1/Th2/Th17 cytokine secretion by human Vdelta3 T cells. *J. Immunol.* **2013**, *191*, 30–34. [[CrossRef](#)]
41. Liang, F.; Zhang, C.; Guo, H.; Gao, S.H.; Yang, F.Y.; Zhou, G.B.; Wang, G.Z. Comprehensive analysis of BTN3A1 in cancers: Mining of omics data and validation in patient samples and cellular models. *FEBS Open Bio* **2021**, *11*, 2586–2599. [[CrossRef](#)] [[PubMed](#)]
42. Nair, S.; Dhodapkar, M.V. Natural Killer T Cells in Cancer Immunotherapy. *Front. Immunol.* **2017**, *8*, 1178. [[CrossRef](#)] [[PubMed](#)]
43. Sebestyen, Z.; Prinz, I.; Dechanet-Merville, J.; Silva-Santos, B.; Kuball, J. Translating gammadelta (gammadelta) T cells and their receptors into cancer cell therapies. *Nat. Rev. Drug Discov.* **2020**, *19*, 169–184. [[CrossRef](#)] [[PubMed](#)]

44. Bauer, S.; Groh, V.; Wu, J.; Steinle, A.; Phillips, J.H.; Lanier, L.L.; Spies, T. Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science* **1999**, *285*, 727–729. [[CrossRef](#)]
45. Simoes, A.E.; Di Lorenzo, B.; Silva-Santos, B. Molecular Determinants of Target Cell Recognition by Human gammadelta T Cells. *Front. Immunol.* **2018**, *9*, 929. [[CrossRef](#)]
46. Groh, V.; Rhinehart, R.; Secrist, H.; Bauer, S.; Grabstein, K.H.; Spies, T. Broad tumor-associated expression and recognition by tumor-derived gamma delta T cells of MICA and MICB. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 6879–6884. [[CrossRef](#)]
47. Kong, Y.; Cao, W.; Xi, X.; Ma, C.; Cui, L.; He, W. The NKG2D ligand ULBP4 binds to TCRgamma9/delta2 and induces cytotoxicity to tumor cells through both TCRgammadelta and NKG2D. *Blood* **2009**, *114*, 310–317. [[CrossRef](#)]
48. Lanca, T.; Correia, D.V.; Moita, C.F.; Raquel, H.; Neves-Costa, A.; Ferreira, C.; Ramalho, J.S.; Barata, J.T.; Moita, L.F.; Gomes, A.Q.; et al. The MHC class Ib protein ULBP1 is a nonredundant determinant of leukemia/lymphoma susceptibility to gammadelta T-cell cytotoxicity. *Blood* **2010**, *115*, 2407–2411. [[CrossRef](#)]
49. Kabelitz, D.; Serrano, R.; Kouakanou, L.; Peters, C.; Kalyan, S. Cancer immunotherapy with gammadelta T cells: Many paths ahead of us. *Cell Mol. Immunol.* **2020**, *17*, 925–939. [[CrossRef](#)]
50. Chan, C.J.; Andrews, D.M.; McLaughlin, N.M.; Yagita, H.; Gilfillan, S.; Colonna, M.; Smyth, M.J. DNAM-1/CD155 interactions promote cytokine and NK cell-mediated suppression of poorly immunogenic melanoma metastases. *J. Immunol.* **2010**, *184*, 902–911. [[CrossRef](#)]
51. Pende, D.; Bottino, C.; Castriconi, R.; Cantoni, C.; Marcenaro, S.; Rivera, P.; Spaggiari, G.M.; Dondero, A.; Carnemolla, B.; Reymond, N.; et al. PVR (CD155) and Nectin-2 (CD112) as ligands of the human DNAM-1 (CD226) activating receptor: Involvement in tumor cell lysis. *Mol. Immunol.* **2005**, *42*, 463–469. [[CrossRef](#)] [[PubMed](#)]
52. Hudspeth, K.; Silva-Santos, B.; Mavilio, D. Natural cytotoxicity receptors: Broader expression patterns and functions in innate and adaptive immune cells. *Front. Immunol.* **2013**, *4*, 69. [[CrossRef](#)] [[PubMed](#)]
53. Zhao, Y.; Niu, C.; Cui, J. Gamma-delta (gammadelta) T cells: Friend or foe in cancer development? *J. Transl. Med.* **2018**, *16*, 3. [[CrossRef](#)] [[PubMed](#)]
54. Todaro, M.; D’Asaro, M.; Caccamo, N.; Iovino, F.; Francipane, M.G.; Meraviglia, S.; Orlando, V.; La Mendola, C.; Gulotta, G.; Salerno, A.; et al. Efficient killing of human colon cancer stem cells by gammadelta T lymphocytes. *J. Immunol.* **2009**, *182*, 7287–7296. [[CrossRef](#)] [[PubMed](#)]
55. Dieli, F.; Vermijlen, D.; Fulfaro, F.; Caccamo, N.; Meraviglia, S.; Cicero, G.; Roberts, A.; Buccheri, S.; D’Asaro, M.; Gebbia, N.; et al. Targeting human $\gamma\delta$ T cells with zoledronate and interleukin-2 for immunotherapy of hormone-refractory prostate cancer. *Cancer Res.* **2007**, *67*, 7450–7457. [[CrossRef](#)]
56. Couzi, L.; Pitard, V.; Sicard, X.; Garrigue, I.; Hawchar, O.; Merville, P.; Moreau, J.F.; Dechanet-Merville, J. Antibody-dependent anti-cytomegalovirus activity of human gammadelta T cells expressing CD16 (FcgammaRIIIa). *Blood* **2012**, *119*, 1418–1427. [[CrossRef](#)]
57. Tokuyama, H.; Hagi, T.; Mattarollo, S.R.; Morley, J.; Wang, Q.; So, H.F.; Moriyasu, F.; Nieda, M.; Nicol, A.J. V gamma 9 V delta 2 T cell cytotoxicity against tumor cells is enhanced by monoclonal antibody drugs—rituximab and trastuzumab. *Int. J. Cancer* **2008**, *122*, 2526–2534. [[CrossRef](#)]
58. Dar, A.A.; Patil, R.S.; Chiplunkar, S.V. Insights into the Relationship between Toll Like Receptors and Gamma Delta T Cell Responses. *Front. Immunol.* **2014**, *5*, 366. [[CrossRef](#)]
59. O’Neill, L.A.; Golenbock, D.; Bowie, A.G. The History of Toll-like Receptors—Redefining Innate Immunity. *Nat. Rev. Immunol.* **2013**, *13*, 453–460. [[CrossRef](#)]
60. Liu, H.; Komai-Koma, M.; Xu, D.; Liew, F.Y. Toll-like receptor 2 signaling modulates the functions of CD4+ CD25+ regulatory T cells. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 7048–7053. [[CrossRef](#)]
61. Suttmuller, R.P.; den Brok, M.H.; Kramer, M.; Bennink, E.J.; Toonen, L.W.; Kullberg, B.J.; Joosten, L.A.; Akira, S.; Netea, M.G.; Adema, G.J. Toll-like receptor 2 controls expansion and function of regulatory T cells. *J. Clin. Investig.* **2006**, *116*, 485–494. [[CrossRef](#)] [[PubMed](#)]
62. Leslie, D.S.; Vincent, M.S.; Spada, F.M.; Das, H.; Sugita, M.; Morita, C.T.; Brenner, M.B. CD1-mediated gamma/delta T cell maturation of dendritic cells. *J. Exp. Med.* **2002**, *196*, 1575–1584. [[CrossRef](#)] [[PubMed](#)]
63. Fang, H.; Welte, T.; Zheng, X.; Chang, G.J.; Holbrook, M.R.; Soong, L.; Wang, T. gammadelta T cells promote the maturation of dendritic cells during West Nile virus infection. *FEMS Immunol. Med. Microbiol.* **2010**, *59*, 71–80. [[CrossRef](#)] [[PubMed](#)]
64. Conti, L.; Casetti, R.; Cardone, M.; Varano, B.; Martino, A.; Belardelli, F.; Poccia, F.; Gessani, S. Reciprocal activating interaction between dendritic cells and pamidronate-stimulated gammadelta T cells: Role of CD86 and inflammatory cytokines. *J. Immunol.* **2005**, *174*, 252–260. [[CrossRef](#)] [[PubMed](#)]
65. Pietschmann, K.; Beetz, S.; Welte, S.; Martens, I.; Gruen, J.; Oberg, H.H.; Wesch, D.; Kabelitz, D. Toll-like receptor expression and function in subsets of human gammadelta T lymphocytes. *Scand. J. Immunol.* **2009**, *70*, 245–255. [[CrossRef](#)]
66. Wang, H.; Chen, H.; Liu, S.; Zhang, J.; Lu, H.; Somasundaram, R.; Choi, R.; Zhang, G.; Ou, L.; Scholler, J.; et al. Costimulation of gammadeltaTCR and TLR7/8 promotes Vdelta2 T-cell antitumor activity by modulating mTOR pathway and APC function. *J. Immunother. Cancer* **2021**, *9*. [[CrossRef](#)]
67. Corsale, A.M.; Di Simone, M.; Lo Presti, E.; Picone, C.; Dieli, F.; Meraviglia, S. Metabolic Changes in Tumor Microenvironment: How Could They Affect gammadelta T Cells Functions? *Cells* **2021**, *10*, 2896. [[CrossRef](#)]

68. Fowler, D.W.; Copier, J.; Dalglish, A.G.; Bodman-Smith, M.D. Zoledronic acid renders human M1 and M2 macrophages susceptible to Vdelta2⁺ gammadelta T cell cytotoxicity in a perforin-dependent manner. *Cancer Immunol. Immunother.* **2017**, *66*, 1205–1215. [[CrossRef](#)]
69. Lin, Y.; Xu, J.; Lan, H. Tumor-associated macrophages in tumor metastasis: Biological roles and clinical therapeutic applications. *J. Hematol. Oncol.* **2019**, *12*, 76. [[CrossRef](#)]
70. Foda, H.D.; Zucker, S. Matrix metalloproteinases in cancer invasion, metastasis and angiogenesis. *Drug Discov. Today* **2001**, *6*, 478–482. [[CrossRef](#)]
71. Chen, Y.; Song, Y.; Du, W.; Gong, L.; Chang, H.; Zou, Z. Tumor-associated macrophages: An accomplice in solid tumor progression. *J. Biomed. Sci.* **2019**, *26*, 78. [[CrossRef](#)] [[PubMed](#)]
72. Rogers, T.L.; Wind, N.; Hughes, R.; Nutter, F.; Brown, H.K.; Vasiliadou, I.; Ottewill, P.D.; Holen, I. Macrophages as potential targets for zoledronic acid outside the skeleton-evidence from in vitro and in vivo models. *Cell Oncol.* **2013**, *36*, 505–514. [[CrossRef](#)] [[PubMed](#)]
73. Coscia, M.; Quaglino, E.; Iezzi, M.; Curcio, C.; Pantaleoni, F.; Riganti, C.; Holen, I.; Monkkonen, H.; Boccardo, M.; Forni, G.; et al. Zoledronic acid repolarizes tumour-associated macrophages and inhibits mammary carcinogenesis by targeting the mevalonate pathway. *J. Cell Mol. Med.* **2010**, *14*, 2803–2815. [[CrossRef](#)] [[PubMed](#)]
74. Melani, C.; Sangaletti, S.; Barazzetta, F.M.; Werb, Z.; Colombo, M.P. Amino-biphosphonate-mediated MMP-9 inhibition breaks the tumor-bone marrow axis responsible for myeloid-derived suppressor cell expansion and macrophage infiltration in tumor stroma. *Cancer Res.* **2007**, *67*, 11438–11446. [[CrossRef](#)] [[PubMed](#)]
75. Brandes, M.; Willmann, K.; Moser, B. Professional antigen-presentation function by human gammadelta T Cells. *Science* **2005**, *309*, 264–268. [[CrossRef](#)] [[PubMed](#)]
76. Sabado, R.L.; Balan, S.; Bhardwaj, N. Dendritic cell-based immunotherapy. *Cell Res.* **2017**, *27*, 74–95. [[CrossRef](#)]
77. Brandes, M.; Willmann, K.; Bioley, G.; Levy, N.; Eberl, M.; Luo, M.; Tampe, R.; Levy, F.; Romero, P.; Moser, B. Cross-presenting human gammadelta T cells induce robust CD8⁺ alpha T cell responses. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 2307–2312. [[CrossRef](#)]
78. Muto, M.; Baghdadi, M.; Maekawa, R.; Wada, H.; Seino, K. Myeloid molecular characteristics of human gammadelta T cells support their acquisition of tumor antigen-presenting capacity. *Cancer Immunol. Immunother.* **2015**, *64*, 941–949. [[CrossRef](#)]
79. Saveanu, L.; Carroll, O.; Weimershaus, M.; Guermonprez, P.; Firat, E.; Lindo, V.; Greer, F.; Davoust, J.; Kratzer, R.; Keller, S.R.; et al. IRAP identifies an endosomal compartment required for MHC class I cross-presentation. *Science* **2009**, *325*, 213–217. [[CrossRef](#)]
80. Scotet, E.; Nedellec, S.; Devilder, M.C.; Allain, S.; Bonneville, M. Bridging innate and adaptive immunity through gammadelta T-dendritic cell crosstalk. *Front. Biosci.* **2008**, *13*, 6872–6885. [[CrossRef](#)]
81. van Beek, J.J.; Wimmers, F.; Hato, S.V.; de Vries, I.J.; Skold, A.E. Dendritic cell cross talk with innate and innate-like effector cells in antitumor immunity: Implications for DC vaccination. *Crit. Rev. Immunol.* **2014**, *34*, 517–536. [[CrossRef](#)] [[PubMed](#)]
82. Devilder, M.C.; Allain, S.; Dousset, C.; Bonneville, M.; Scotet, E. Early triggering of exclusive IFN-gamma responses of human Vgamma9Vdelta2 T cells by TLR-activated myeloid and plasmacytoid dendritic cells. *J. Immunol.* **2009**, *183*, 3625–3633. [[CrossRef](#)] [[PubMed](#)]
83. Kunzmann, V.; Kretzschmar, E.; Herrmann, T.; Wilhelm, M. Polyinosinic-polycytidylic acid-mediated stimulation of human gammadelta T cells via CD11c dendritic cell-derived type I interferons. *Immunology* **2004**, *112*, 369–377. [[CrossRef](#)] [[PubMed](#)]
84. Fowler, D.W.; Copier, J.; Wilson, N.; Dalglish, A.G.; Bodman-Smith, M.D. Mycobacteria activate gammadelta T-cell anti-tumour responses via cytokines from type 1 myeloid dendritic cells: A mechanism of action for cancer immunotherapy. *Cancer Immunol. Immunother.* **2012**, *61*, 535–547. [[CrossRef](#)]
85. Li, W.; Kubo, S.; Okuda, A.; Yamamoto, H.; Ueda, H.; Tanaka, T.; Nakamura, H.; Yamanishi, H.; Terada, N.; Okamura, H. Effect of IL-18 on expansion of gammadelta T cells stimulated by zoledronate and IL-2. *J. Immunother.* **2010**, *33*, 287–296. [[CrossRef](#)]
86. Fiore, F.; Castella, B.; Nuschak, B.; Bertieri, R.; Mariani, S.; Bruno, B.; Pantaleoni, F.; Foglietta, M.; Boccardo, M.; Massaia, M. Enhanced ability of dendritic cells to stimulate innate and adaptive immunity on short-term incubation with zoledronic acid. *Blood* **2007**, *110*, 921–927. [[CrossRef](#)]
87. Su, X.; Zhang, L.; Jin, L.; Ye, J.; Guan, Z.; Chen, R. Coculturing dendritic cells with zoledronate acid efficiently enhance the anti-tumor effects of cytokine-induced killer cells. *J. Clin. Immunol.* **2010**, *30*, 766–774. [[CrossRef](#)]
88. Takahara, M.; Miyai, M.; Tomiyama, M.; Mutou, M.; Nicol, A.J.; Nieda, M. Copulsing tumor antigen-pulsed dendritic cells with zoledronate efficiently enhance the expansion of tumor antigen-specific CD8⁺ T cells via Vgamma9gammadelta T cell activation. *J. Leukoc. Biol.* **2008**, *83*, 742–754. [[CrossRef](#)]
89. Maniar, A.; Zhang, X.; Lin, W.; Gastman, B.R.; Pauza, C.D.; Strome, S.E.; Chapoval, A.I. Human gammadelta T lymphocytes induce robust NK cell-mediated antitumor cytotoxicity through CD137 engagement. *Blood* **2010**, *116*, 1726–1733. [[CrossRef](#)]
90. Ladel, C.H.; Blum, C.; Kaufmann, S.H. Control of natural killer cell-mediated innate resistance against the intracellular pathogen *Listeria monocytogenes* by gamma/delta T lymphocytes. *Infect. Immun.* **1996**, *64*, 1744–1749. [[CrossRef](#)]
91. Nussbaumer, O.; Gruenbacher, G.; Gander, H.; Thurnher, M. DC-like cell-dependent activation of human natural killer cells by the bisphosphonate zoledronic acid is regulated by gammadelta T lymphocytes. *Blood* **2011**, *118*, 2743–2751. [[CrossRef](#)] [[PubMed](#)]
92. Wilhelm, M.; Kunzmann, V.; Eckstein, S.; Reimer, P.; Weissinger, F.; Ruediger, T.; Tony, H.P. Gammadelta T cells for immune therapy of patients with lymphoid malignancies. *Blood* **2003**, *102*, 200–206. [[CrossRef](#)] [[PubMed](#)]

93. Meraviglia, S.; Eberl, M.; Vermijlen, D.; Todaro, M.; Buccheri, S.; Cicero, G.; La Mendola, C.; Guggino, G.; D'Asaro, M.; Orlando, V.; et al. In vivo manipulation of Vgamma9Vdelta2 T cells with zoledronate and low-dose interleukin-2 for immunotherapy of advanced breast cancer patients. *Clin. Exp. Immunol.* **2010**, *161*, 290–297. [[CrossRef](#)] [[PubMed](#)]
94. Kobayashi, H.; Tanaka, Y.; Yagi, J.; Osaka, Y.; Nakazawa, H.; Uchiyama, T.; Minato, N.; Toma, H. Safety profile and anti-tumor effects of adoptive immunotherapy using gamma-delta T cells against advanced renal cell carcinoma: A pilot study. *Cancer Immunol. Immunother.* **2007**, *56*, 469–476. [[CrossRef](#)] [[PubMed](#)]
95. Kobayashi, H.; Tanaka, Y.; Yagi, J.; Minato, N.; Tanabe, K. Phase I/II study of adoptive transfer of gammadelta T cells in combination with zoledronic acid and IL-2 to patients with advanced renal cell carcinoma. *Cancer Immunol. Immunother.* **2011**, *60*, 1075–1084. [[CrossRef](#)]
96. Wada, I.; Matsushita, H.; Noji, S.; Mori, K.; Yamashita, H.; Nomura, S.; Shimizu, N.; Seto, Y.; Kakimi, K. Intraperitoneal injection of in vitro expanded Vgamma9Vdelta2 T cells together with zoledronate for the treatment of malignant ascites due to gastric cancer. *Cancer Med.* **2014**, *3*, 362–375. [[CrossRef](#)]
97. Bennouna, J.; Levy, V.; Sicard, H.; Senellart, H.; Audrain, M.; Hiret, S.; Rolland, F.; Bruzzoni-Giovanelli, H.; Rimbart, M.; Galea, C.; et al. Phase I study of bromohydrin pyrophosphate (BrHPP, IPH 1101), a Vgamma9Vdelta2 T lymphocyte agonist in patients with solid tumors. *Cancer Immunol. Immunother.* **2010**, *59*, 1521–1530. [[CrossRef](#)]
98. Kunzmann, V.; Smetak, M.; Kimmel, B.; Weigang-Koehler, K.; Goebeler, M.; Birkmann, J.; Becker, J.; Schmidt-Wolf, I.G.; Einsele, H.; Wilhelm, M. Tumor-promoting versus tumor-antagonizing roles of gammadelta T cells in cancer immunotherapy: Results from a prospective phase I/II trial. *J. Immunother.* **2012**, *35*, 205–213. [[CrossRef](#)]
99. Lang, J.M.; Kaikobad, M.R.; Wallace, M.; Staab, M.J.; Horvath, D.L.; Wilding, G.; Liu, G.; Eickhoff, J.C.; McNeel, D.G.; Malkovsky, M. Pilot trial of interleukin-2 and zoledronic acid to augment gammadelta T cells as treatment for patients with refractory renal cell carcinoma. *Cancer Immunol. Immunother.* **2011**, *60*, 1447–1460. [[CrossRef](#)]
100. Pressey, J.G.; Adams, J.; Harkins, L.; Kelly, D.; You, Z.; Lamb, L.S., Jr. In vivo expansion and activation of gammadelta T cells as immunotherapy for refractory neuroblastoma: A phase 1 study. *Medicine* **2016**, *95*, e4909. [[CrossRef](#)]
101. Bennouna, J.; Bompas, E.; Neidhardt, E.M.; Rolland, F.; Philip, I.; Galea, C.; Salot, S.; Saiagh, S.; Audrain, M.; Rimbart, M.; et al. Phase-I study of Innacell gammadelta, an autologous cell-therapy product highly enriched in gamma9delta2 T lymphocytes, in combination with IL-2, in patients with metastatic renal cell carcinoma. *Cancer Immunol. Immunother.* **2008**, *57*, 1599–1609. [[CrossRef](#)] [[PubMed](#)]
102. Abe, Y.; Muto, M.; Nieda, M.; Nakagawa, Y.; Nicol, A.; Kaneko, T.; Goto, S.; Yokokawa, K.; Suzuki, K. Clinical and immunological evaluation of zoledronate-activated Vgamma9gammadelta T-cell-based immunotherapy for patients with multiple myeloma. *Exp. Hematol.* **2009**, *37*, 956–968. [[CrossRef](#)] [[PubMed](#)]
103. Nakajima, J.; Murakawa, T.; Fukami, T.; Goto, S.; Kaneko, T.; Yoshida, Y.; Takamoto, S.; Kakimi, K. A phase I study of adoptive immunotherapy for recurrent non-small-cell lung cancer patients with autologous gammadelta T cells. *Eur. J. Cardiothorac. Surg.* **2010**, *37*, 1191–1197. [[CrossRef](#)] [[PubMed](#)]
104. Sakamoto, M.; Nakajima, J.; Murakawa, T.; Fukami, T.; Yoshida, Y.; Murayama, T.; Takamoto, S.; Matsushita, H.; Kakimi, K. Adoptive immunotherapy for advanced non-small cell lung cancer using zoledronate-expanded gammadelta T cells: A phase I clinical study. *J. Immunother.* **2011**, *34*, 202–211. [[CrossRef](#)] [[PubMed](#)]
105. Nicol, A.J.; Tokuyama, H.; Mattarollo, S.R.; Hagi, T.; Suzuki, K.; Yokokawa, K.; Nieda, M. Clinical evaluation of autologous gamma delta T cell-based immunotherapy for metastatic solid tumours. *Br. J. Cancer* **2011**, *105*, 778–786. [[CrossRef](#)]
106. Izumi, T.; Kondo, M.; Takahashi, T.; Fujieda, N.; Kondo, A.; Tamura, N.; Murakawa, T.; Nakajima, J.; Matsushita, H.; Kakimi, K. Ex vivo characterization of gammadelta T-cell repertoire in patients after adoptive transfer of Vgamma9Vdelta2 T cells expressing the interleukin-2 receptor beta-chain and the common gamma-chain. *Cytotherapy* **2013**, *15*, 481–491. [[CrossRef](#)]
107. Kakimi, K.; Matsushita, H.; Murakawa, T.; Nakajima, J. gammadelta T cell therapy for the treatment of non-small cell lung cancer. *Transl. Lung Cancer Res.* **2014**, *3*, 23–33. [[CrossRef](#)]
108. Cui, J.; Li, L.; Wang, C.; Jin, H.; Yao, C.; Wang, Y.; Li, D.; Tian, H.; Niu, C.; Wang, G.; et al. Combined cellular immunotherapy and chemotherapy improves clinical outcome in patients with gastric carcinoma. *Cytotherapy* **2015**, *17*, 979–988. [[CrossRef](#)]
109. Wilhelm, M.; Smetak, M.; Schaefer-Eckart, K.; Kimmel, B.; Birkmann, J.; Einsele, H.; Kunzmann, V. Successful adoptive transfer and in vivo expansion of haploidentical gammadelta T cells. *J. Transl. Med.* **2014**, *12*, 45. [[CrossRef](#)]
110. Alnaggar, M.; Xu, Y.; Li, J.; He, J.; Chen, J.; Li, M.; Wu, Q.; Lin, L.; Liang, Y.; Wang, X.; et al. Allogenic Vgamma9Vdelta2 T cell as new potential immunotherapy drug for solid tumor: A case study for cholangiocarcinoma. *J. Immunother. Cancer* **2019**, *7*, 36. [[CrossRef](#)]
111. Lin, M.; Zhang, X.; Liang, S.; Luo, H.; Alnaggar, M.; Liu, A.; Yin, Z.; Chen, J.; Niu, L.; Jiang, Y. Irreversible electroporation plus allogenic Vgamma9Vdelta2 T cells enhances antitumor effect for locally advanced pancreatic cancer patients. *Signal Transduct Target Ther.* **2020**, *5*, 215. [[CrossRef](#)] [[PubMed](#)]
112. Deniger, D.C.; Switzer, K.; Mi, T.; Maiti, S.; Hurton, L.; Singh, H.; Huls, H.; Olivares, S.; Lee, D.A.; Champlin, R.E.; et al. Bispecific T-cells expressing polyclonal repertoire of endogenous gammadelta T-cell receptors and introduced CD19-specific chimeric antigen receptor. *Mol. Ther.* **2013**, *21*, 638–647. [[CrossRef](#)]
113. Ang, W.X.; Ng, Y.Y.; Xiao, L.; Chen, C.; Li, Z.; Chi, Z.; Tay, J.C.; Tan, W.K.; Zeng, J.; Toh, H.C.; et al. Electroporation of NKG2D RNA CAR Improves Vgamma9Vdelta2 T Cell Responses against Human Solid Tumor Xenografts. *Mol. Ther. Oncolytics* **2020**, *17*, 421–430. [[CrossRef](#)] [[PubMed](#)]

114. Rozenbaum, M.; Meir, A.; Aharoni, Y.; Itzhaki, O.; Schachter, J.; Bank, I.; Jacoby, E.; Besser, M.J. Gamma-Delta CAR-T Cells Show CAR-Directed and Independent Activity Against Leukemia. *Front. Immunol.* **2020**, *11*, 1347. [[CrossRef](#)]
115. Johanna, I.; Straetmans, T.; Heijhuurs, S.; Aarts-Riemens, T.; Norell, H.; Bongiovanni, L.; de Bruin, A.; Sebestyen, Z.; Kuball, J. Evaluating in vivo efficacy-toxicity profile of TEG001 in humanized mice xenografts against primary human AML disease and healthy hematopoietic cells. *J. Immunother. Cancer* **2019**, *7*, 69. [[CrossRef](#)]
116. Fisher, J.; Abramowski, P.; Wisidagamage Don, N.D.; Flutter, B.; Capsomidis, A.; Cheung, G.W.; Gustafsson, K.; Anderson, J. Avoidance of On-Target Off-Tumor Activation Using a Co-stimulation-Only Chimeric Antigen Receptor. *Mol. Ther.* **2017**, *25*, 1234–1247. [[CrossRef](#)]
117. Middelburg, J.; Kemper, K.; Engelberts, P.; Labrijn, A.F.; Schuurman, J.; van Hall, T. Overcoming Challenges for CD3-Bispecific Antibody Therapy in Solid Tumors. *Cancers* **2021**, *13*, 287. [[CrossRef](#)]
118. Goebeler, M.E.; Bargou, R. Blinatumomab: A CD19/CD3 bispecific T cell engager (BiTE) with unique anti-tumor efficacy. *Leuk. Lymphoma* **2016**, *57*, 1021–1032. [[CrossRef](#)]
119. Wang, L.; Qiao, Y.; Zong, H.; Han, L.; Ke, Y.; Pan, Z.; Chen, J.; Lu, J.; Li, J.; Ying, T.; et al. IgG-like Bispecific Antibody CD3xEpCAM Generated by Split Intein Against Colorectal Cancer. *Front. Pharmacol.* **2022**, *13*, 803059. [[CrossRef](#)]
120. Oberg, H.H.; Peipp, M.; Kellner, C.; Sebens, S.; Krause, S.; Petrick, D.; Adam-Klages, S.; Rocken, C.; Becker, T.; Vogel, I.; et al. Novel bispecific antibodies increase gammadelta T-cell cytotoxicity against pancreatic cancer cells. *Cancer Res.* **2014**, *74*, 1349–1360. [[CrossRef](#)]
121. Ganesan, R.; Chennupati, V.; Ramachandran, B.; Hansen, M.R.; Singh, S.; Grewal, I.S. Selective recruitment of gammadelta T cells by a bispecific antibody for the treatment of acute myeloid leukemia. *Leukemia* **2021**, *35*, 2274–2284. [[CrossRef](#)] [[PubMed](#)]
122. Watanabe, D.; Koyanagi-Aoi, M.; Taniguchi-Ikeda, M.; Yoshida, Y.; Azuma, T.; Aoi, T. The Generation of Human gammadelta T Cell-Derived Induced Pluripotent Stem Cells from Whole Peripheral Blood Mononuclear Cell Culture. *Stem Cells Transl. Med.* **2018**, *7*, 34–44. [[CrossRef](#)] [[PubMed](#)]
123. Zeng, J.; Tang, S.Y.; Wang, S. Derivation of mimetic gammadelta T cells endowed with cancer recognition receptors from reprogrammed gammadelta T cell. *PLoS ONE* **2019**, *14*, e0216815. [[CrossRef](#)] [[PubMed](#)]
124. Vermijlen, D.; Gatti, D.; Kouzeli, A.; Rus, T.; Eberl, M. gammadelta T cell responses: How many ligands will it take till we know? *Semin. Cell Dev. Biol.* **2018**, *84*, 75–86. [[CrossRef](#)]
125. Vavassori, S.; Kumar, A.; Wan, G.S.; Ramanjaneyulu, G.S.; Cavallari, M.; El Daker, S.; Beddoe, T.; Theodossis, A.; Williams, N.K.; Gostick, E.; et al. Butyrophilin 3A1 binds phosphorylated antigens and stimulates human gammadelta T cells. *Nat. Immunol.* **2013**, *14*, 908–916. [[CrossRef](#)]
126. Park, J.H.; Lee, H.K. Function of $\gamma\delta$ T cells in tumor immunology and their application to cancer therapy. *Exp. Mol. Med.* **2021**, *53*, 318–327. [[CrossRef](#)]
127. Chabab, G.; Barjon, C.; Bonnefoy, N.; Lafont, V. Pro-tumor gammadelta T Cells in Human Cancer: Polarization, Mechanisms of Action, and Implications for Therapy. *Front. Immunol.* **2020**, *11*, 2186. [[CrossRef](#)]
128. Peng, G.; Wang, H.Y.; Peng, W.; Kiniwa, Y.; Seo, K.H.; Wang, R.F. Tumor-infiltrating gammadelta T cells suppress T and dendritic cell function via mechanisms controlled by a unique toll-like receptor signaling pathway. *Immunity* **2007**, *27*, 334–348. [[CrossRef](#)]
129. Wu, P.; Wu, D.; Ni, C.; Ye, J.; Chen, W.; Hu, G.; Wang, Z.; Wang, C.; Zhang, Z.; Xia, W.; et al. gammadelta T17 cells promote the accumulation and expansion of myeloid-derived suppressor cells in human colorectal cancer. *Immunity* **2014**, *40*, 785–800. [[CrossRef](#)]
130. Rong, L.; Li, K.; Li, R.; Liu, H.M.; Sun, R.; Liu, X.Y. Analysis of tumor-infiltrating gamma delta T cells in rectal cancer. *World J. Gastroenterol.* **2016**, *22*, 3573–3580. [[CrossRef](#)]
131. Patil, R.S.; Shah, S.U.; Shrikhande, S.V.; Goel, M.; Dikshit, R.P.; Chiplunkar, S.V. IL17 producing gammadelta T cells induce angiogenesis and are associated with poor survival in gallbladder cancer patients. *Int. J. Cancer* **2016**, *139*, 869–881. [[CrossRef](#)] [[PubMed](#)]
132. Chen, X.; Shang, W.; Xu, R.; Wu, M.; Zhang, X.; Huang, P.; Wang, F.; Pan, S. Distribution and functions of gammadelta T cells infiltrated in the ovarian cancer microenvironment. *J. Transl. Med.* **2019**, *17*, 144. [[CrossRef](#)] [[PubMed](#)]
133. Daley, D.; Zambirinis, C.P.; Seifert, L.; Akkad, N.; Mohan, N.; Werba, G.; Barilla, R.; Torres-Hernandez, A.; Hundeyin, M.; Mani, V.R.K.; et al. gammadelta T Cells Support Pancreatic Oncogenesis by Restraining alphabeta T Cell Activation. *Cell* **2016**, *166*, 1485–1499.e15. [[CrossRef](#)] [[PubMed](#)]
134. Lo Presti, E.; Pizzolato, G.; Corsale, A.M.; Caccamo, N.; Sireci, G.; Dieli, F.; Meraviglia, S. gammadelta T Cells and Tumor Microenvironment: From Immunosurveillance to Tumor Evasion. *Front. Immunol.* **2018**, *9*, 1395. [[CrossRef](#)] [[PubMed](#)]
135. Li, M.O.; Wan, Y.Y.; Sanjabi, S.; Robertson, A.K.; Flavell, R.A. Transforming growth factor-beta regulation of immune responses. *Annu. Rev. Immunol.* **2006**, *24*, 99–146. [[CrossRef](#)] [[PubMed](#)]
136. Wang, J.; Zhang, L.; Kang, D.; Yang, D.; Tang, Y. Activation of PGE2/EP2 and PGE2/EP4 signaling pathways positively regulate the level of PD-1 in infiltrating CD8⁺ T cells in patients with lung cancer. *Oncol. Lett.* **2018**, *15*, 552–558. [[CrossRef](#)]
137. Vigano, S.; Alatzoglou, D.; Irving, M.; Menetrier-Caux, C.; Caux, C.; Romero, P.; Coukos, G. Targeting Adenosine in Cancer Immunotherapy to Enhance T-Cell Function. *Front. Immunol.* **2019**, *10*, 925. [[CrossRef](#)]
138. Marten, A.; von Lilienfeld-Toal, M.; Buchler, M.W.; Schmidt, J. Soluble MIC is elevated in the serum of patients with pancreatic carcinoma diminishing gammadelta T cell cytotoxicity. *Int. J. Cancer* **2006**, *119*, 2359–2365. [[CrossRef](#)]

139. Gonnermann, D.; Oberg, H.H.; Kellner, C.; Peipp, M.; Sebens, S.; Kabelitz, D.; Wesch, D. Resistance of cyclooxygenase-2 expressing pancreatic ductal adenocarcinoma cells against gammadelta T cell cytotoxicity. *Oncoimmunology* **2015**, *4*, e988460. [[CrossRef](#)]
140. Blackburn, S.D.; Shin, H.; Haining, W.N.; Zou, T.; Workman, C.J.; Polley, A.; Betts, M.R.; Freeman, G.J.; Vignali, D.A.; Wherry, E.J. Coregulation of CD8+ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat. Immunol.* **2009**, *10*, 29–37. [[CrossRef](#)]
141. Andrews, L.P.; Marciscano, A.E.; Drake, C.G.; Vignali, D.A. LAG3 (CD223) as a cancer immunotherapy target. *Immunol. Rev.* **2017**, *276*, 80–96. [[CrossRef](#)] [[PubMed](#)]
142. Lee, H.W.; Chung, Y.S.; Kim, T.J. Heterogeneity of Human gammadelta T Cells and Their Role in Cancer Immunity. *Immune Netw.* **2020**, *20*, e5. [[CrossRef](#)] [[PubMed](#)]
143. Coscia, M.; Vitale, C.; Peola, S.; Foglietta, M.; Rigoni, M.; Griggio, V.; Castella, B.; Angelini, D.; Chiaretti, S.; Riganti, C.; et al. Dysfunctional Vgamma9Vdelta2 T cells are negative prognosticators and markers of dysregulated mevalonate pathway activity in chronic lymphocytic leukemia cells. *Blood* **2012**, *120*, 3271–3279. [[CrossRef](#)] [[PubMed](#)]
144. Castella, B.; Foglietta, M.; Sciancalepore, P.; Rigoni, M.; Coscia, M.; Griggio, V.; Vitale, C.; Ferracini, R.; Saraci, E.; Omede, P.; et al. Anergic bone marrow Vgamma9Vdelta2 T cells as early and long-lasting markers of PD-1-targetable microenvironment-induced immune suppression in human myeloma. *Oncoimmunology* **2015**, *4*, e1047580. [[CrossRef](#)] [[PubMed](#)]
145. Hoeres, T.; Holzmann, E.; Smetak, M.; Birkmann, J.; Wilhelm, M. PD-1 signaling modulates interferon-gamma production by Gamma Delta (gammadelta) T-Cells in response to leukemia. *Oncoimmunology* **2019**, *8*, 1550618. [[CrossRef](#)] [[PubMed](#)]
146. Dieli, F.; Poccia, F.; Lipp, M.; Sireci, G.; Caccamo, N.; Di Sano, C.; Salerno, A. Differentiation of effector/memory Vdelta2 T cells and migratory routes in lymph nodes or inflammatory sites. *J. Exp. Med.* **2003**, *198*, 391–397. [[CrossRef](#)]
147. Battistini, L.; Caccamo, N.; Borsellino, G.; Meraviglia, S.; Angelini, D.F.; Dieli, F.; Cencioni, M.T.; Salerno, A. Homing and memory patterns of human gammadelta T cells in physiopathological situations. *Microbes Infect.* **2005**, *7*, 510–517. [[CrossRef](#)]
148. Martin, M.D.; Badovinac, V.P. Defining Memory CD8 T Cell. *Front. Immunol.* **2018**, *9*, 2692. [[CrossRef](#)]
149. Xu, L.; Yao, D.; Tan, J.; He, Z.; Yu, Z.; Chen, J.; Luo, G.; Wang, C.; Zhou, F.; Zha, X.; et al. Memory T cells skew toward terminal differentiation in the CD8+ T cell population in patients with acute myeloid leukemia. *J. Hematol. Oncol.* **2018**, *11*, 93. [[CrossRef](#)]
150. Cui, W.; Kaech, S.M. Generation of effector CD8+ T cells and their conversion to memory T cells. *Immunol. Rev.* **2010**, *236*, 151–166. [[CrossRef](#)]