



## Original Article

## Role of osteopontin in the process of pulpal healing following tooth replantation in mice



Kiyoko Suzuki-Barrera<sup>a</sup>, Sanako Makishi<sup>a</sup>, Mitsushiro Nakatomi<sup>b</sup>, Kotaro Saito<sup>a</sup>, Hiroko Ida-Yonemochi<sup>a</sup>, Hayato Ohshima<sup>a,\*</sup>

<sup>a</sup> Division of Anatomy and Cell Biology of the Hard Tissue, Department of Tissue Regeneration and Reconstruction, Niigata University Graduate School of Medical and Dental Sciences, 2-5274 Gakkocho-dori, Chuo-ku, Niigata, 951-8514, Japan

<sup>b</sup> Department of Human, Information and Life Sciences, School of Health Sciences, University of Occupational and Environmental Health, 1-1 Iseigaoka, Yahatanishi-ku, Kitakyushu, 807-8555, Japan

## ARTICLE INFO

## Article history:

Received 24 June 2022

Received in revised form

24 September 2022

Accepted 29 September 2022

## Keywords:

Animal model

Blood supply

Dentinogenesis

Innervation

Osteopontin

Tooth replantation

## ABSTRACT

**Introduction:** The role of osteopontin (OPN) following severe injury remains to be elucidated, especially its relationship with type I collagen (encoded by the *Col1a1* gene) secretion by newly-differentiated odontoblast-like cells (OBLCs). In this study, we examined the role of OPN in the process of reparative dentin formation with a focus on reinnervation and revascularization after tooth replantation in *Opn* knockout (KO) and wild-type (WT) mice.

**Methods:** Maxillary first molars of 2- and 3-week-old-*Opn* KO and WT mice (*Opn* KO 2W, *Opn* KO 3W, WT 2W, and WT 3W groups) were replanted, followed by fixation 3–56 days after operation. Following micro-computed tomography analysis, the decalcified samples were processed for immunohistochemistry for Ki67, Nestin, PGP 9.5, and CD31 and *in situ* hybridization for *Col1a1*.

**Results:** An intense inflammatory reaction occurred to disrupt pulpal healing in the replanted teeth of the *Opn* KO 3W group, whereas dental pulp achieved healing in the *Opn* KO 2W and WT groups. The tertiary dentin in the *Opn* KO 3W group was significantly decreased in area compared with the *Opn* KO 2W and WT groups, with a significantly low percentage of Nestin-positive, newly-differentiated OBLCs during postoperative days 7–14. In the *Opn* KO 3W group, the blood vessels were significantly decreased in area and pulp healing was disturbed with a failure of pulpal revascularization and reinnervation.

**Conclusions:** OPN is necessary for proper reinnervation and revascularization to deposit reparative dentin following severe injury within the dental pulp of erupted teeth with advanced root development. © 2022, The Japanese Society for Regenerative Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Dental pulp is a highly specialized mesenchymal tissue with a remarkable capacity for repair and regeneration [1]. Pulpal healing with tertiary dentin indicates that the original pulp tissue is

replaced by a different tissue with the deposition of a calcified scar [2]. After tooth injury, the formation of tertiary dentin adjacent to preexisting dentin is a secretory response regulated by odontoblasts or odontoblast-like cells (OBLCs). There is an intimate relationship between reactionary dentinogenesis with nerve fiber sprouting following injuries, such as dental caries [3]. This indicates that crosstalk exists between matrix secretory activity by odontoblasts and dynamic neuronal responses. Pulpal nerve fibers exhibit neuroplasticity with sprouting, degenerative, or regenerative processes depending on various scenarios, such as injury or physiological events [4–8]. The response of blood vessels is to initiate angiogenic events as a consequence of hypoxia, because the secretory activity of the odontoblasts is associated with the positional and ultrastructural changes of pulpal capillaries [9]. The mechanisms that regulate multi-event responses with tertiary

**Abbreviations:** M1, first molars; GFP, green fluorescent protein; H&E, hematoxylin and eosin; H2B, histone 2B; KO, knockout; MSCs, mesenchymal stem cells;  $\mu$ CT, micro-computed tomography; OPN, osteopontin; OBLCs, odontoblast-like cells; SCs, Schwann cells; SCAP, stem cells derived from the apical papilla; VEGF, vascular endothelial growth factor; WT, wild-type.

\* Corresponding author. Fax: +81-25-227-0804.

E-mail address: [histoman@dent.niigata-u.ac.jp](mailto:histoman@dent.niigata-u.ac.jp) (H. Ohshima).

Peer review under responsibility of the Japanese Society for Regenerative Medicine.

<https://doi.org/10.1016/j.reth.2022.09.011>

2352-3204/© 2022, The Japanese Society for Regenerative Medicine. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

dentinogenesis, reinnervation, and revascularization within the dental pulp following exogenous injury remain to be fully elucidated.

Tooth replantation is a severe injury that induces the death of most odontoblasts because of the interruption of the neurovascular supply to the dental pulp. Although pulpal healing with reinnervation and revascularization has been shown to occur in humans [10] and experimental animals [11,12], there are many factors that affect this successful outcome. The lack of proper oxygenated medium is decisive for the survival of odontoblast-lineage cells and the occlusal force during and/or after the operation is detrimental to the cells [12]. Furthermore, the width and length of the roots are important predictors of pulpal healing with a significant association between the stage of root development or a favorable ratio of the broad apical foramen/short root canal and proper pulpal healing [10,13]. Therefore, the evaluation and comparison of these factors are important to provide insight into the manner in which dental pulp can (or not) succeed in its own proper healing following tooth replantation.

Understanding how tertiary dentinogenesis and OBLC differentiation occur after tooth replantation is fundamental for evaluating strategies and predicting outcomes for dentin-pulp complex regeneration. The regulation of matrix mineralization and cell adhesion is associated with the presence of non-collagenous proteins, especially small integrin-binding ligand N-linked glycoproteins, which are composed of bone sialoprotein, dentin sialophosphoprotein, dentin matrix protein-1, matrix extracellular phosphoprotein, and osteopontin (OPN) [14]. OPN is localized in the boundary between tertiary dentin and preexisting dentin in mice [15]. Immunocompetent cells, such as macrophages and dendritic cells, secrete OPN, which is deposited at the dentin-predentin interface before OBLC differentiation and after tooth transplantation [16]. In addition, OPN is essential for type I collagen secretion by new OBLCs to form reparative dentin after an injury, such as cavity preparation [17]. However, the role of OPN during reparative dentinogenesis and its interplay with the neurovascular response after severe injury, such as tooth replantation, is not yet fully understood. In this study, we determined the role of OPN in the process of reparative dentin formation with special focus on reinnervation, revascularization, and different root development stages following tooth replantation in *Opn* knockout (KO) and wild-type (WT) mice.

## 2. Methods

### 2.1. Mice

All animal experiments were conducted in compliance with ARRIVE guidelines and a protocol that was reviewed by the Institutional Animal Care and Use Committee and approved by the President of Niigata University (SA00780). Information regarding the animals is available from a previous study [17]. Male and female *Opn*<sup>-/-</sup> (B6. Cg-Spp1<sup>tm1Blh</sup>/J) and wild-type (WT: C57BL/6J) mice were obtained from the Jackson Laboratories (Bar Harbor, ME) and Charles River Laboratories of Japan (Yokohama, Japan), respectively.

### 2.2. Tooth replantation

Two- and three-week-old animals were used for tooth replantation, which were referred to as 2W and 3W groups, respectively to clarify the role of OPN in the process of reparative dentin formation following tooth replantation using the unerupted teeth with short roots in the 2W group as well as the erupted teeth with

long roots in the 3W group. The upper right first molars (M1) of *Opn* KO and WT mice were extracted under deep anesthesia following an intraperitoneal injection of a mixed solution (0.05–0.1 mL/10 g) of Domitor® (1.875 mL: Nippon Zenyaku Kogyo Co, Ltd, Koriyama, Japan), midazolam (2 mL: Sandoz KK, Tokyo, Japan), Vetorphale® (2.5 mL: Meiji Seika Pharma Co, Ltd, Tokyo, Japan), and physiological saline (18.625 mL), using a pair of modified dental tweezers. The tooth was immediately repositioned in the original socket. The animals were divided into four groups: *Opn* KO 2W, *Opn* KO 3W, WT 2W, and WT 3W groups.

### 2.3. Micro-computed tomography ( $\mu$ CT) analysis

$\mu$ CT analysis (Elescan; Nippon Steel Texeng. Co., Ltd, Tokyo, Japan) was used to examine the stages of root development and the morphological changes of the replants in the *Opn* KO 2W and WT 2W groups at 3, 5, 7, and 14 days following tooth replantation. The CT settings were as follows: pixel matrix, 512 × 512 × 256; slice thickness, 20.67  $\mu$ m; projection number, 900 × 32; magnification, × 5.3; voltage, 63 kV; and electrical current, 101  $\mu$ A. The maxillae were reconstructed using a software program (TRI/3D Bon, Ratoc System Engineering, Tokyo, Japan) to evaluate the three-dimensionally reconstructed views of the maxillae including the upper first molars (M1). The untreated teeth from the *Opn* KO 2W and WT 2W groups were used as the control groups.

### 2.4. Tissue preparation

Tissues were collected from three to ten animals at intervals of 3, 5, 7, 14, and 56 days after tooth replantation (n = 84) as shown in Table 1. At each stage, the animals were perfused with 4% paraformaldehyde in 0.1M phosphate buffer (pH 7.4) under deep anesthesia following an intraperitoneal injection of a mixed solution (0.05–0.1 mL/10 g) of Domitor®, midazolam, Vetorphale®, and physiological saline and each maxilla was immersed in the same fixative. The procedure for tissue preparation was previously described [17].

### 2.5. Immunohistochemical analysis

Immunohistochemistry was performed essentially as described in our previous report [17] with a mouse anti-Nestin monoclonal antibody diluted 1:200 (Millipore, Temecula, CA; catalog number: MAB353). The Envision + Horseradish Peroxidase System (Dako Japan, catalog number: K5027) was used along with a rabbit anti-human PGP9.5 antibody diluted to 1:400 (Ultra Clone Ltd., Isle of Wight, UK; catalog number: RA95101), a rat anti-mouse CD31 antibody diluted to 1:50 (BD Pharmingen, San Diego, CA; catalog number: 553,370), and a rat anti-Ki67 monoclonal antibody diluted to 1:100 for the cell proliferation assay (Dako Japan, Tokyo, Japan; catalog number: M7249). The avidin-biotin peroxidase complex (Vectastain ABC Kit; Vector Laboratories, Burlingame, CA) formation was determined using biotinylated anti-rabbit or anti-rat immunoglobulin G (Vector Laboratories; catalog number: BA-4000).

### 2.6. In situ hybridization

Section *in situ* hybridization was performed as previously described [18]. Digoxigenin-labeled probe for *Col1a1* [19], which encodes type I collagen protein, was prepared according to the manufacturer's instructions (Roche Diagnostics Corp, Indianapolis, IN).

**Table 1**  
Number of animals in each experiment and for statistical analysis after tooth replantation.

		Experiment	3D	5D	7D	14D	56D	Total
Opn KO	2W Replantation	μCT	(3)	(3)	(3)	(3)	0	(12)
		H&E	4	4	3	3	6	20
		Nestin (IHC)	(4)	(4)	(4)	(3)	(3)	(18)
		<i>Col1a1</i> (ISH)	1	1	1	0	0	3
		Ki67 (IHC)	(4)	(4)	(3)	(3)	0	(14)
		CD31 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		PGP 9.5 (IHC)	(3)	(3)	(3)	(3)	0	(12)
	3W Replantation	μCT	(3)	(3)	(3)	(3)	0	(12)
		H&E	6	10	10	7	0	33
		Nestin (IHC)	(3)	(4)	(6)	(3)	0	(16)
		Ki67 (IHC)	(3)	(4)	(4)	(3)	0	(14)
		CD31 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		PGP 9.5 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		μCT	(3)	(3)	(3)	(3)	0	(12)
WT	2W Replantation	H&E	4	3	3	3	0	13
		Nestin (IHC)	(4)	(4)	(4)	(3)	0	(15)
		<i>Col1a1</i> (ISH)	1	1	1	0	0	3
		Ki67 (IHC)	(4)	(4)	(3)	(3)	0	(14)
		CD31 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		PGP 9.5 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		μCT	(3)	(3)	(3)	(3)	0	(12)
	3W Replantation	H&E	3	3	3	3	0	12
		Nestin (IHC)	(3)	(3)	(3)	(3)	0	(12)
		Ki67 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		CD31 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		PGP 9.5 (IHC)	(3)	(3)	(3)	(3)	0	(12)
		μCT	(3)	(3)	(3)	(3)	0	(12)

## 2.7. Statistical analysis

The root length of the upper first molar was measured in μCT images using TRI/3D Bon Software and Image J software (Image J 1.45s; National Institutes of Health, Bethesda, MD). The percentage of Nestin-positive perimeters in the total perimeter of the pulp-dentin border was calculated using Image J software. Data were obtained from samples of *Opn* KO and WT teeth (Table 1) and one optimal section was selected from each tooth. For the cell proliferation assay based on Ki67 immunohistochemistry, one optimal section was selected from each tooth (Table 1). For the analysis of hard tissue and blood vessel areas, hematoxylin and eosin (H&E) stained sections were prepared from the samples of *Opn* KO and WT teeth (Table 1) and analyzed using Image J and WinRoof software (WinRoof Version 7.4; Mitani Corporation, Tokyo, Japan). The root length, the percentage of hard tissue and blood vessel areas, Nestin-positive perimeter, and Ki67-positive cell density among different groups were compared using a one-way analysis of variance followed by Bonferroni's multiple comparisons test after the confirmation of data normality and homogeneity of variance (SPSS 21.0.0.0 for Windows; SPSS Japan, Tokyo, Japan). The samples showing no normal distribution were compared using the Kruskal–Wallis test followed by Bonferroni's test for multiple comparisons. Data were reported as the mean + SD and p denotes the p-value.

## 3. Results


### 3.1. Pulpal healing was achieved in the *Opn* KO 2W group, exhibiting newly-differentiated OBLC differentiation and reparative dentin formation

The root formation of replants progressed for 3–14 days after tooth replantation in the *Opn* KO 2W and WT 2W groups (Fig. 1a), although the length of the roots of the replants was shorter than that of untreated control developing teeth (Fig. 1b).


A significant difference was observed in the length of the mesial root between the control and replanted teeth in the *Opn* KO 2W group on day 5 (Fig. 1b). On day 3 in the *Opn* KO 2W and WT 2W groups, the odontoblasts exhibited degenerated features or an exudative lesion occurred beneath the coronal dentin. On day 7 in the *Opn* KO 2W and WT 2W groups, the revascularization was completed and the inflammatory reaction ceased throughout the dental pulp to achieve pulpal healing (Fig. 1c, d, g, h). In the WT 2W and *Opn* KO 2W groups, Nestin-positive newly-differentiated OBLCs were arranged beneath the reparative dentin (Fig. 1k–r). The filamentous structures also showed Nestin-positive reaction in addition to the OBLCs during days 3–5 (Fig. 1k and l). The timing of *Col1a1* mRNA expression in the root pulp in the *Opn* KO 2W group was faster compared with that of the WT 2W group on day 3 (Fig. 1e, i), and the pulp-dentin border of the whole pulp expressed *Col1a1* in both the WT 2W and *Opn* KO 2W groups on day 7 (Fig. 1f, j). On day 56 in the *Opn* KO 2W group, Nestin-positive OBLCs were arranged along the pulp-dentin border, although the pulp cavity was reduced in size (Fig. 2a–d).

### 3.2. Pulpal healing was not achieved in the *Opn* KO 3W group

The occurrence of hard tissue was calculated based on Nestin-positive (tertiary dentin) or Nestin negative hard tissue areas in the total pulp area (Fig. 2e). The tertiary dentin area of the *Opn* KO 3W group significantly decreased compared with that of the *Opn* KO 2W and WT groups. In contrast, a severe inflammatory reaction occurred to disturb pulpal healing in the *Opn* KO 3W group (Fig. 2f and g). On day 3 in the WT 3W group, the odontoblasts exhibited degenerated features beneath the coronal dentin (Fig. 2h). On day 7 in the WT 3W group, the revascularization was completed and the inflammatory reaction ceased throughout the dental pulp to achieve pulpal healing (Fig. 2i). The *Opn* KO 3W group exhibited impaired reparative dentin formation with a significantly low percentage of Nestin-positive newly-differentiated OBLCs during



**Fig. 1.** μCT images (a), quantitative analyses of mesial and distal root length in μCT images (b), H&E-stained sections (c, d, g, h), *Col1a1* in situ hybridization (e, f, i, j), and Nestin immunohistochemistry (k–r) of replanted teeth 3 (a-1, a-5, c, g, e, i, k, o), 5 (a-2, a-6, l, p), 7 (a-3, a-7, d, h, f, j, m, q), and 14 days (a-4, a-8, n, r) after operation in *Opn* KO 2W (a-1–4, c–f, k–n) and WT 2W (a-5–8, g–j, o–r) groups. (a) The root formation of replants progresses during days 3–14. (b) A significant difference was observed in the length of the mesial root between the control and replanted teeth in the *Opn* KO 2W group on day 5. (c, d, g, h) On day 3 in the *Opn* KO 2W and WT 2W groups, the odontoblasts show degenerated features (\*) or an exudative lesion (E) occurs beneath the coronal dentin, with an artificial detached distribution of their layer from the predentin. On day 7 in the *Opn* KO 2W and WT 2W groups, the revascularization (arrows) is completed and the inflammatory reaction ceases throughout the dental pulp to achieve pulpal healing. (e, f, i, j) The timing of *Col1a1* mRNA expression in the root pulp of the *Opn* KO 2W group is faster compared with that of the WT 2W group on day 3. The pulp-dentin border of the whole pulp expresses *Col1a1* in both the WT 2W and *Opn* KO 2W groups on day 7. (k–r) In the *Opn* KO 2W and WT 2W groups, Nestin-positive newly-differentiated OBLCs are arranged beneath the reparative dentin. Scale bars = (a) 1000 μm, (e, f, i, j, k–r) 250 μm, (c, d, g, h) 100 μm. B, bone; D, dentin; DP, dental pulp; OBLC, odontoblast-like cells.



**Fig. 2.** H&E-stained sections (a, c, f–i) and Nestin immunohistochemistry (b, d, k–r) in replanted teeth 3 (f, h, k, o), 5 (l, p), 7 (g, i, m, q), 14 (n, r), and 56 days (a–d) after operation in the *Opn* KO 2W (a–d), *Opn* KO 3W (f, g, k–n), and WT 3W (h, i, o–r) groups. Quantitative analysis of the hard tissue area percentage at 14 and 56 days in Nestin-stained sections (e) and Nestin-positive perimeter (j). (a–d) The *Opn* KO 2W shows continuous reparative dentin formation, resulting in a reduced size of pulp cavity. (e) The tertiary dentin of the *Opn* KO 3W group significantly decreases in area compared with the *Opn* KO 2W and WT groups. (f–i) A severe inflammatory reaction occurs which disturbs pulpal healing in the *Opn* KO 3W group. In contrast, in the WT 2W group the dental pulp achieves pulpal healing. (j) Quantitative analysis of the Nestin-positive perimeter shows a significant lower percentage in the *Opn* KO 3W group compared with all other groups during days 7–14. (k–r) The *Opn* KO 3W group displays impaired reparative dentin formation with a low percentage of Nestin-positive newly-differentiated OBLCs during postoperative days 7–14. In the WT 3W group, Nestin-positive newly-differentiated OBLCs are arranged beneath the reparative dentin. Scale bars = (a, b, k–r) 250  $\mu$ m, (f–i) 100  $\mu$ m, (c, d) 50  $\mu$ m. B, bone; D, dentin; DP, dental pulp; OBLC, odontoblast-like cells; \*, degenerating odontoblasts; arrows, revascularization.

postoperative days 7–14. In contrast, in the WT 3W group, Nestin-positive newly-differentiated OBLCs were arranged beneath the reparative dentin (Fig. 2k–r). The *Opn* KO 2W and WT 2W groups showed a significantly higher occurrence of Nestin-positive newly-differentiated OBLCs along the pulp-dentin border on day 5 compared with the *Opn* KO 3W group. Subsequently, there were no significant differences among the *Opn* KO 2W and WT groups during days 7–14 (Fig. 2j).

### 3.3. The *Opn* KO 2W group displayed cell proliferation within the dental pulp after replantation

The analysis of cell proliferation by Ki67 immunostaining confirmed a lack of proper pulp healing in the *Opn* KO 3W group, with significantly decreased cell proliferation activity compared with the *Opn* KO 2W and WT 2W groups on day 3 and the *Opn* KO 2W and WT 3W groups on day 5 (Fig. 3a–c). The cell proliferation

activity of the *Opn* KO 2W group tended to be higher compared with that of the WT 2W group. The proliferative activity reached its peak on day 5 in the *Opn* KO 2W group, whereas it occurred on day 3 in the WT 2W group (Fig. 3c).

### 3.4. Vascularization and innervation were reestablished within the dental pulp in the *Opn* KO 2W group

CD31 is a cell adhesion molecule highly expressed on endothelial cells and concentrated at the junctions between them to be used as a general marker for blood vessels (Fig. 4a, c, e, g, i, k, m, o), whereas PGP9.5 is highly expressed in neurons to be used as a general marker for neuronal tissue including peripheral nerves (Fig. 4b, d, f, h, j, l, n, p). In the *Opn* KO 3W group, pulp healing was disturbed with a failure of pulpal revascularization and reinnervation (Fig. 4i–l). The area of blood vessels in the *Opn* KO 3W group was significantly decreased compared with that in the WT groups on day 5, WT 3W group on day 7, and the *Opn* KO 2W and WT 3W groups on day 14 (Fig. 4q). In contrast, the *Opn* KO 2W group achieved revascularization and reinnervation within the dental pulp (Fig. 4a–d). The existence of reestablished blood vessels was confirmed by the presence of CD31-positive endothelial cells during days 3–7 (Fig. 4a, c, e, g, m, o).


## 4. Discussion

### 4.1. The significance of the experimental design


The main difference between the experimental groups is that the maxillary first molar of the *Opn* KO 2W group is characterized by early root development and non-erupted tooth, whereas the *Opn* KO 3W group contained the erupted tooth with advanced root development. Compared the *Opn* KO 2W with WT 2W control groups, root development in the former group was delayed compared with the latter group, indicating a different baseline between these groups. Moreover, a significant difference was temporarily observed in the length of the mesial root between the control and replanted teeth in the *Opn* KO 2W group on day 5. The experiment for tooth replantation using 2-week-old mice provides a better environment for replant without serious deleterious effects on Hertwig’s epithelial root sheath because of the short length of the replant roots and gingival covering.

### 4.2. The role of OPN in the process of reparative dentinogenesis

The tertiary dentin in the *Opn* KO 3W group was significantly decreased in area compared with the *Opn* KO 2W and WT groups,



**Fig. 3.** Ki67 immunohistochemistry (a) in the dental pulp 3 (a-1, a-3, a-5, a-7) and 5 days (a-2, a-4, a-6, a-8) after operation in the *Opn* KO 2W (a-1, a-2), *Opn* KO 3W (a-5, a-6), WT 2W (a-3, a-4), and WT 3W (a-7, a-8) groups. Representative images (a) are higher magnified views of the boxed area in the coronal pulp (b). Quantitative analysis of Ki67-positive cell density in the dental pulp (c). (a) The *Opn* KO 3W group lacks proliferative activity in the dental pulp, whereas all other groups show active cell proliferation. (c) The *Opn* KO 3W group exhibits a significant decreased in cell proliferative activity compared with the *Opn* KO 2W and WT 2W groups on day 3 and the *Opn* KO 2W and WT 3W groups on day 5. Scale bars = (b) 500  $\mu$ m, (a) 50  $\mu$ m.



**Fig. 4.** CD31 (a, c, e, g, i, k, m, o) and PGP 9.5 (b, d, f, h, j, l, n, p) immunohistochemistry in replanted teeth 3 (a, b, e, f, i, j, m, n) and 7 days (c, d, g, h, k, l, o, p) after operation in the *Opn* KO 2W (a–d), WT 2W (e–h), *Opn* KO 3W (i–l), and WT 3W (m–p) groups. Quantitative analysis of blood vessel area percentage in H&E-stained sections (q). (a–p) In the *Opn* KO 3W group, pulp healing is disturbed with a failure of pulpal revascularization and reinnervation. The *Opn* KO 2W group succeeded in revascularization (\*) and reinnervation (arrows) within the dental pulp. (q) The area of blood vessels in the *Opn* KO 3W group significantly decreased compared with that in the WT groups on day 5, the WT 3W group on day 7, and the *Opn* KO 2W and WT 3W groups after 14 days. Scale bars = 100  $\mu$ m.

with a significantly low percentage of Nestin-positive newly-differentiated OBLCs present during postoperative days 7–14. OPN deficiency inhibits OBLC differentiation to induce no hard tissue formation in the replants with advanced root formation. OPN is deposited at the boundary between the preexisting and reparative dentin after cavity preparation in mice, resulting in a role for OPN in type I collagen secretion [17]. However, the *Opn* KO 2W group showed OBLC differentiation. Furthermore, the Nestin-positive filamentous structures appear in the pulp in which increasing numbers of proliferative cells and rich revascularization occur. These findings may indicate a correlation between the up-regulation of Nestin expression and activation of the proliferative potential of dental pulp stem/progenitor cells. A reduction in root length can accelerate pulp regeneration by improving the

revascularization of the replanted teeth [20]. The favorable ratio of broad apical foramen/short root canal is important, since the foramen is the primary access route for blood vessels and nerves to the coronal dental pulp. If the apical foramen is too small, it may prevent cell migration, revascularization, and reinnervation. Moreover, ischemic pulp healing depends on the prevalence of reinnervation and revascularization and the absence of bacterial invasion [21]. This is based on the difference between the experimental groups (non-erupted versus erupted replants) and the gingival covering that protects from oral microflora in the case of the *Opn* KO 2W group. The *Opn* KO 2W and WT 2W groups exhibited significantly higher numbers of Nestin-positive newly-differentiated OBLCs along the pulp-dentin border on day 5 compared with the *Opn* KO 3W group. Subsequently, there were no

significant differences between the *Opn* KO 2W and WT groups during days 7–14. With respect to cavity preparation, reparative dentin formation is disturbed because of the loss of *Col1a1* mRNA expression in the *Opn* KO mice. In contrast, reactionary dentin formation is quite normal even in these mice, indicating that OBLCs require OPN for the secretion of type I collagen, whereas OPN is not necessary for the surviving odontoblasts responsible for reactionary dentin [17]. As mentioned above, reparative dentin formation occurred in the *Opn* KO 2W group in the present study. The stem/progenitor cells for OBLCs may be different between the mild injury models, such as cavity preparation, and severe injury models, such as tooth replantation. The former model provides localized injury beneath the disrupted dentin, whereas the latter model results in the total death of odontoblasts throughout the dental pulp. Furthermore, tooth replantation results in the degeneration of the subodontoblastic layer and the central pulp tissue in addition to the odontoblasts. Dental pulp stem/progenitor cells are localized within the perivascular niche in the subodontoblastic layer and the central pulp tissue. In fact, our previous studies using prenatal labeling methods with BrdU and Tet-OP-histone 2B (H2B)-green fluorescent protein (GFP) mice demonstrated the occurrence of label-retaining cells in these niches [2]. These findings suggest that other types of stem/progenitor cells commit themselves to OBLCs in the *Opn* KO 2W group.

During early root development, the apical papilla harbors mesenchymal stem cells (MSCs), which are stem cells derived from the apical papilla (SCAP), which are capable of becoming OBLCs to produce dentin *in vivo* [22], enlightening the potential role of SCAP in pulpal healing and regeneration. The first molar of the *Opn* KO 2W group is the developing tooth with immature roots containing many SCAP at the apical portion. SCAP-derived OBLCs may not need OPN for type I collagen secretion. This notion is supported by the previous evidence that the coronal subodontoblastic layer is different from the other pulp cells with respect to their origin. The former is of neuronal ectodermal origin, whereas the latter is of neural crest origin [23]. SCAP can easily reach the coronal pulp chamber because of the short distance. Further studies are needed to identify the multiple types of stem/progenitor cells for OBLCs. The timing of *Col1a1* mRNA expression in the root pulp of the *Opn* KO 2W group is faster compared with that of the WT 2W group on day 3, and the pulp-dentin border of the whole pulp expressed *Col1a1* mRNA in both the WT 2W and *Opn* KO 2W groups on day 7. These findings support the above notion. In the early stage of healing, the activity of *Col1a1* mRNA in the *Opn* 2W group suggests high SCAP activity. The cell proliferative activity of the *Opn* KO 2W group tended to be higher compared with that of the WT 2W group, however, peak proliferative activity in the WT 2W group is faster (on day 3) compared with that in the *Opn* KO 2W group (on day 5). The delayed healing process in the *Opn* KO 2W group compared with that in the WT 2W group may be attributed to an abnormal immune response resulting from *Opn* deficiency [24].

#### 4.3. Relationship between OPN and angiogenesis/axonal regeneration

In the *Opn* KO 3W group, the blood vessels were significantly decreased in area and the pulp healing was disturbed with a failure of pulpal revascularization and reinnervation. Angiogenesis is important for organ growth and repair, as hypoxia following injury induces complex and specific pro-angiogenic responses within the dental pulp where vascular endothelial growth factor (VEGF) participates in the revascularization process [25]. The consequence of vascular disruption during hypoxia triggers an increase of MSC migration and expression of OPN in osteocytes [26]. In the *Opn* KO mice with advanced root development, pulp healing is disturbed

with a failure of pulpal revascularization. OPN enhances angiogenesis directly through the activation of the PI3K/Akt pathway and improves the expression of VEGF [27], confirming an important role of OPN during angiogenesis in tissue regenerative events. Revascularization is an important event during reinnervation, in which the distal stump becomes vascularized in response to the macrophage-induced VEGF signal with Schwann cells (SCs) migrating along the vasculature [28]. A peripheral nerve injury triggers a Wallerian degeneration process, which includes a multicellular response primarily from SCs, blood vessels, and immunocompetent cells [29]. The dental pulp also exhibits neuroplasticity, and nerve fibers display a prominent and progressive axonal degeneration in a Wallerian-like scheme under physiological root resorption [7]. In the present study, the axonal regeneration was impaired within dental pulp of the *Opn* KO 3W group with advanced root development. Previous studies demonstrated that SCs express OPN in the degenerating distal nerve stump during the first days following injury [30]. OPN was significantly upregulated after peripheral nerve injury to promote proliferation and inhibition of SC apoptosis [31]. These findings indicate the importance of OPN during axonal regeneration in peripheral nerves with a special role in the survival of SCs, which are fundamental in providing the necessary signals and spatial cues to the injured nerves.

## 5. Conclusions

OPN has an important role during pulp regeneration, since the re-establishment of innervation and vascularization are major processes during pulp healing. The present study also demonstrated that OPN is necessary for proper reinnervation and revascularization to deposit reparative dentin after severe injury within the dental pulp in erupted teeth with advanced root development.

### Author contributions

KS-B contributed to data curation; formal analysis; investigation; methodology; validation; writing—original draft preparation. SM, KS, MN, and HI-Y contributed to investigation; validation; writing - review & editing. HO contributed to conceptualization; data curation; investigation; methodology; supervision; validation; writing—original draft preparation.

### Declaration of competing interest

The authors declare no conflicts of interest related to this study.

### Acknowledgments

The authors cordially thank Dr. T. Komori for providing *Col1a1* plasmids, Ms. Y. Abe and Ms. M. Kawachi for their technical assistance, and Enago ([www.enago.jp](http://www.enago.jp)) for the English language review. Dr. K. Suzuki-Barrera was supported by Scholarship for Japanese Emigrants and their Descendants in Latin America and the Caribbean: Program for Developing Leaders in Nikkei Communities, JICA-Japan. This work was supported by Grants-in-Aid for Scientific Research (B) (no. 17H04366 to HO) and Challenging Research (Pioneering) (no. 20K21672 to HO) from the Japan Society for the Promotion of Science.

### References

- [1] Hargreaves KM, Goodis HE, Tay FR. Seltzer and Bender's dental pulp. 2d ed. Quintessence Publishing; 2012.
- [2] Saito K, Ohshima H. Differentiation capacity and maintenance of dental pulp stem/progenitor cells in the process of pulpal healing following tooth injuries. *J Oral Biosci* 2017;59:63–70.

- [3] Couve E, Osorio R, Schmachtenberg O. Reactionary dentinogenesis and neuroimmune response in dental caries. *J Dent Res* 2014;93:788–93.
- [4] Byers MR, Kvinnsland I, Bothwell M. Analysis of low affinity nerve growth factor receptor during pulpal healing and regeneration of myelinated and unmyelinated axons in replanted teeth. *J Comp Neurol* 1992;326:470–84.
- [5] Luukko K, Kettunen P. Coordination of tooth morphogenesis and neuronal development through tissue interactions: lessons from mouse models. *Exp Cell Res* 2014;325:72–7.
- [6] Saito K, Ida-Yonemochi H, Ushiki T, Ohshima H. Responses of pulp vasculature after cavity preparation in rat molars. *J Oral Biosci* 2015;57:157–64.
- [7] Suzuki K, Lovera M, Schmachtenberg O, Couve E. Axonal degeneration in dental pulp precedes human primary teeth exfoliation. *J Dent Res* 2015;94:1446–53.
- [8] Couve E, Lovera M, Suzuki K, Schmachtenberg O. Schwann cell phenotype changes in aging human dental pulp. *J Dent Res* 2018;97:347–55.
- [9] Seino Y, Takano Y, Ohshima H. Positional and ultrastructural changes in peripheral pulp capillaries correlate with the active phase of dentin deposition and mineralization in rat molars. *J Oral Biosci* 2017;59:163–71.
- [10] Andreasen JO, Borum MK, Jacobsen HL, Andreasen FM. Replantation of 400 avulsed permanent incisors. 2. Factors related to pulpal healing. *Endod Dent Traumatol* 1995;11:59–68.
- [11] Shimizu A, Nakakura-Ohshima K, Noda T, Maeda T, Ohshima H. Responses of immunocompetent cells in the dental pulp to replantation during the regeneration process in rat molars. *Cell Tissue Res* 2000;302:221–33.
- [12] Hasegawa T, Suzuki H, Yoshie H, Ohshima H. Influence of extended operation time and of occlusal force on determination of pulpal healing pattern in replanted mouse molars. *Cell Tissue Res* 2007;329:259–72.
- [13] Kafourou V, Tong HJ, Day P, Houghton N, Spencer RJ, Duggal M. Outcomes and prognostic factors that influence the success of tooth autotransplantation in children and adolescents. *Dent Traumatol* 2017;33:393–9.
- [14] Nikoloudaki G. Functions of matricellular proteins in dental tissues and their emerging roles in orofacial tissue development, maintenance, and disease. *Int J Mol Sci* 2021:22.
- [15] Takamori Y, Suzuki H, Nakakura-Ohshima K, Cai J, Cho SW, Jung HS, et al. Capacity of dental pulp differentiation in mouse molars as demonstrated by allogenic tooth transplantation. *J Histochem Cytochem* 2008;56:1075–86.
- [16] Saito K, Nakatomi M, Ida-Yonemochi H, Kenmotsu S, Ohshima H. The expression of GM-CSF and osteopontin in immunocompetent cells precedes the odontoblast differentiation following allogenic tooth transplantation in mice. *J Histochem Cytochem* 2011;59:518–29.
- [17] Saito K, Nakatomi M, Ida-Yonemochi H, Ohshima H. Osteopontin is essential for type I collagen secretion in reparative dentin. *J Dent Res* 2016;95:1034–41.
- [18] Nakatomi M, Morita I, Eto K, Ota MS. Sonic hedgehog signaling is important in tooth root development. *J Dent Res* 2006;85:427–31.
- [19] Metsaranta M, Toman D, De Crombrughe B, Vuorio E. Specific hybridization probes for mouse type I, II, III and IX collagen mRNAs. *Biochim Biophys Acta* 1991;1089:241–3.
- [20] Nakakura-Ohshima K, Quispe-Salcedo A, Sano H, Hayasaki H, Ohshima H. The effects of reducing the root length by apicoectomy on dental pulp revascularization following tooth replantation in mice. *Dent Traumatol* 2021;37:677–90.
- [21] Andreasen FM, Kahler B. Pulpal response after acute dental injury in the permanent dentition: clinical implications—a review. *J Endod* 2015;41:299–308.
- [22] Sonoyama W, Liu Y, Yamaza T, Tuan RS, Wang S, Shi S, et al. Characterization of the apical papilla and its residing stem cells from human immature permanent teeth: a pilot study. *J Endod* 2008;34:166–71.
- [23] Nakatomi M, Quispe-Salcedo A, Sakaguchi M, Ida-Yonemochi H, Okano H, Ohshima H. Nestin expression is differently regulated between odontoblasts and the subodontoblastic layer in mice. *Histochem Cell Biol* 2018;149:383–91.
- [24] Weber GF, Cantor H. The immunology of Eta-1/osteopontin. *Cytokine Growth Factor Rev* 1996;7:241–8.
- [25] Aranha AM, Zhang Z, Neiva KG, Costa CA, Hebling J, Nor JE. Hypoxia enhances the angiogenic potential of human dental pulp cells. *J Endod* 2010;36:1633–7.
- [26] Raheja LF, Genetos DC, Yellowley CE. Hypoxic osteocytes recruit human MSCs through an OPN/CD44-mediated pathway. *Biochem Biophys Res Commun* 2008;366:1061–6.
- [27] Dai J, Peng L, Fan K, Wang H, Wei R, Ji G, et al. Osteopontin induces angiogenesis through activation of PI3K/AKT and ERK1/2 in endothelial cells. *Oncogene* 2009;28:3412–22.
- [28] Cattin AL, Lloyd AC. The multicellular complexity of peripheral nerve regeneration. *Curr Opin Neurobiol* 2016;39:38–46.
- [29] Jessen KR, Mirsky R. The repair Schwann cell and its function in regenerating nerves. *J Physiol* 2016;594:3521–31.
- [30] Jander S, Bussini S, Neuen-Jacob E, Bosse F, Menge T, Muller HW, et al. Osteopontin: a novel axon-regulated Schwann cell gene. *J Neurosci Res* 2002;67:156–66.
- [31] Wang JB, Zhang Z, Li JN, Yang T, Du S, Cao RJ, et al. SPP1 promotes Schwann cell proliferation and survival through PKC $\alpha$  by binding with CD44 and  $\alpha$ v $\beta$ 3 after peripheral nerve injury. *Cell Biosci* 2020;10:98.