





Strong Association of Metabolic Parameters with ADMA and VCAM-I in Normo-Weight Subjects with Metabolic Syndrome

Iis Inayati Rakhmat ¹⁻³, Gaga Irawan Nugraha^{3,4}, Eko Fuji Ariyanto ^{3,4}, Yuni Susanti Pratiwi^{3,4}, Desy Linasari ⁵, Siti Nur Fatimah^{3,6}, Mohammad Ghozali ^{3,4}, Mas Rizky AA Syamsunarno^{3,4}, Mohammad Rizki Akbar^{3,7}, Tri Hanggono Achmad^{3,4}

¹Doctoral Study Program Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia; ²Department of Biochemistry, Faculty of Medicine, Universitas Jenderal Achmad Yani, Cimahi, West Java, Indonesia; ³Cardiometabolic Working Group Study, Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia; ⁴Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia; ⁵Department of Community Medicine, Faculty of Medicine, Universitas Jenderal Achmad Yani, Cimahi, West Java, Indonesia; ⁶Department of Public Health, Faculty of Medicine, Universitas Padjadjaran, Bandung, West Java, Indonesia; ⁷Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Padjadjaran-Hasan Sadikin General Hospital, Bandung, West Java, Indonesia

Correspondence: Gaga Irawan Nugraha, Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran, Eymann 38, Bandung, West Java, 40161, Indonesia, Tel/Fax +6222 2032170, Email gaga@unpad.ac.id

Background: Metabolic syndrome (MetS) is a risk factor for cardiovascular disease and is linked to obesity. Subjects with MetS who have normo-weight potentially show higher mortality and morbidity.

Purpose: This study aims to reveal the critical essential metabolic parameters associated with endothelial dysfunction in MetS subjects with normo-weight compared to obese.

Patients and Methods: The study was designed using a case-control approach. Ninety-nine MetS subjects (34 Normo-weight and 65 obese) from the urban population were enrolled in this study. The components of MetS are based on NCEP/ATP III criteria. Asymmetric dimethylarginine (ADMA) and vascular cell adhesion molecule 1 (VCAM-1) as markers for endothelial dysfunction were measured in both groups.

Results: Fasting blood glucose (FBG) levels were higher in the normo-weight group (143.38 ± 79.8 mg/dL) compared to the obese group (120.89 ± 46.5 mg/dL). High-density lipoprotein cholesterol (HDL-c) levels in the normo-weight group were lower (42.82 ± 10.1 mg/dL) compared to obesity (45.74 ± 9.3 mg/dL), while triacylglycerol (TAG) levels were higher in the obese (197.25 ± 110.5 mg/dL) compared to the normo-weight group (167.03 ± 98.4 mg/dL), although the differences were statistically not significant (all $p > 0.05$). The difference between ADMA and VCAM-1 levels was statistically not significant in both groups. Correlation between MetS components with endothelial dysfunction parameters shows that metabolic parameters correlate strongly. Interestingly, a stronger correlation between FBG and ADMA was observed in normo-weight ($r = 0.519$) compared to obese groups ($r = 0.445$). In addition, TAG consistently shows a significant correlation with ADMA and VCAM-1 in normo-weight groups.

Conclusion: Metabolic parameters, especially FBG and TAG, correlate strongly with endothelial dysfunction parameters in normo-weight subjects with metabolic syndrome.

Keywords: ADMA, normal-weight metabolically unhealthy, VCAM-1

Introduction

Cardiovascular disease remains a significant global threat, with 17.9 million people dying from it in 2016, accounting for 31% of all global deaths.¹ Metabolic syndrome (MetS) is a common condition worldwide linked to an increased risk of cardiovascular events and diabetes mellitus.²⁻⁴ Metabolic syndrome, initially linked to obesity, can also occur in normo-weight adults.⁵ Interestingly, recent studies in the US revealed that the mortality rate in normal-weight MetS patients was even higher than in

obese MetS patients.⁶ These findings raise the issue of which MetS criteria influence more to cardiovascular events rather than obesity.^{7–10}

Several criteria have been developed to diagnose MetS, including central obesity, elevated TAG, low HDL-c, elevated fasting blood sugar, and hypertension.^{2,3} These criteria can also be found in normo-weight subjects.^{9,10} Several studies have reported metabolically unhealthy subjects with cardiovascular disease,^{5,6,8,10,11} associated with endothelial dysfunction, and alteration in ADMA and VCAM-1 used as surrogate biomarkers.^{12–15} This study aims to reveal the metabolic parameters primarily related to endothelial dysfunction, represented by ADMA and VCAM-1 elevation in metabolically unhealthy subjects with normal weight compared to obese.

Materials and Methods

Study Design and Subjects

Analytical descriptive design with a case–control approach was conducted in the Cibeber subdistrict, Cimahi, West Java, Indonesia. A total of 256 participated in this study after signing informed consent. Eight subjects were excluded due to having been diagnosed and treated for cardiometabolic diseases. Of 248 subjects examined for MetS based on NCEP/ATP III criteria, 107 subjects were categorized as MetS. Eight people withdrew from the study, enrolling 99 subjects (34 are normal weight). The Research Ethics Committee of Universitas Padjadjaran approved the research protocol No. 1123/UN6.KEP/EC/2022 and confirmed that this study complies with the Declaration of Helsinki.

Anthropometric and Blood Pressure Measurements

The body mass index (BMI) was determined by the formula weight (kg) divided by squared height (m²) based on the WHO task-forced Asia Pacific criteria.¹⁶ The weight was measured using body weight and body composition OMRON HBF-214 (Singapore), and the height was measured using a portable stadiometer ADE MZ10042 (Hamburg, Germany). The waist circumference (WC) measured at the top of the iliac crests was taken with light clothes using non-stretchable tape. Blood pressure was measured using a digital sphygmomanometer OMRON hem 8712 (Singapore).

Laboratory Assays

Blood samples were collected after a 12-hour overnight fasting. Fasting blood glucose (FBG) is measured using spectrophotometry using the GOD-PAP method (Proline Glucose GOD FS). HDL cholesterol is measured by the PEG Trinder precipitation method (SEKISUI MEDICAL CO, LTD, Japan). Serum triglycerides were measured using the colorimetric enzymatic test Glycerol Peroxidase Phosphate Acid (GPO-PAP) (Proline triglycerides FS 10'). ADMA was measured using Agilent 6460 Triple Quad with 1290 UPLC with LC/MS-MS method. VCAM-1 was measured using Biorad Microplate Reader 680 with an Enzyme-linked immunosorbent assay (ELISA) method (Human VCAM-1/CD106 Immunoassay). ADMA levels are elevated when ≥ 100 ug/dl^{12,17} while VCAM-1 levels increase when ≥ 714 ug/dl.¹³

Statistical Analysis

Data analysis was performed by SPSS v26 (SPSS Inc., Chicago, Illinois, USA). Comparisons were performed by Student's unpaired *t*-test or Mann–Whitney test. The Pearson correlation was used to determine the relationship between metabolic syndrome components with ADMA and VCAM-1 serum. A value of $p < 0.05$ was considered statistically significant.

Results

The Characteristic of Participants

The characteristics of participants who were metabolic syndrome patients are presented in Table 1. They were divided into normo-weight and obese groups. The mean BMI of the normo-weight group was 23.39 ± 1.3 kg/m², and the obese group was 30.54 ± 4.7 kg/m² ($p = 0.000$). It was also observed that WC in the obese group was significantly higher than in the normo-weight group (96.99 ± 13.7 cm vs 84.85 ± 6.6 cm, respectively, $p = 0.000$). Average fasting blood glucose levels were higher in the normal weight group (143.38 ± 79.8 mg/dL) compared to the obese group (120.89 ± 46.5 mg/dL), although statistically, there was no significant between the two groups ($p = 0.079$). Average HDL-c levels in the normal weight group were lower

Table 1 Study Group Data Based on Body Mass Index (N=99)

	Metabolic Syndrome Patient		p-value
	Normo-Weight (n=34)	Obese (n=65)	
	Mean ± SD	Mean ± SD	
Age (years)	55.32 ± 8.6	51.17 ± 10.5	0.051
BMI (kg/m ²)	23.39 ± 1.3	30.54 ± 4.7	0.000*
Waist circumference (cm)	84.85 ± 6.6	96.99 ± 13.7	0.000*
SBP (mmHg)	147.00 ± 18.6	154.91 ± 23.3	0.090
DBP (mmHg)	89.97 ± 10.9	97.78 ± 14.8	0.098
FBG (mg/dl)	143.38 ± 79.8	120.89 ± 46.5	0.079
HDL-c (mg/dl)	42.82 ± 10.1	45.74 ± 9.3	0.158
TAG (mg/dl)	167.03 ± 98.4	197.25 ± 110.5	0.183
ADMA (ug/dl)	93.79 ± 17.1	92.38 ± 20.8	0.735
VCAM-I (ug/dl)	771.73 ± 151.53	780.44 ± 285.3	0.869

Notes: Data is shown with mean ± SD, *significantly different if p-value <0.05.

Abbreviations: BMI, Body Mass Index; WP, Waist Circumference; SBP, Systolic Blood Pressure; DBP, Diastolic Blood Pressure; FBG, Fasting Blood Glucose; HDL-c, High Density Lipoprotein-cholesterol; TAG, Triglycerides; ADMA, Asymmetric dimethylarginine; VCAM-I, Vascular cells adhesion molecules-I.

(42.82 ± 10.1 mg/dL) compared with obesity (45.74 ± 9.3 mg/dL), although statistically also not significant ($p = 0.158$). Meanwhile, the average TAG levels were higher in the obese group (197.25 ± 110.5 mg/dL) compared to the normal weight group (167.03 ± 98.4 mg/dL), although statistically, there was also no significant correlation between the two groups ($p=0.183$). The mean serum VCAM-1 level was higher than normal (771.73 ± 151.53 ug/dL and 780.44 ± 285.3 ug/dL), while the average ADMA level is close to the upper limit of normal values in both groups (93.79 ± 17.1 ug/dL and 92.38 ± 20.8 ug/dL), although there was no significant difference.

The Association Between ADMA and VCAM-I Level with Age, BMI, and MetS Components

Table 2 shows the association between ADMA and VCAM-1 levels with age, BMI, and MetS components in both groups. ADMA levels positively correlate with age, FBG, and TAG (r range is 0.223–0.652) in both groups. VCAM-1 is positively correlated with FBG and TAG (range of r is 0.329–0.485) in the normo-weight group, while in the obese group, it is positively correlated with age and FBG (range of r is 0.209–0.581) and negatively correlated with HDL-c ($r=-0.323$). FBG more strongly correlates with ADMA in normo-weight than in obese groups, while FBG correlates with VCAM-1 more strongly in obese than with normo-weight groups. TAG consistently correlates significantly with ADMA and VCAM-1 in normo-weight groups.

Discussion

Our study aims to elaborate the association between metabolic parameters and endothelial dysfunction in normo-weight and obese subjects with metabolic syndrome and revealed that FBG is strongly associated with ADMA and VCAM-1.

Previous studies have reported that high FBG level is strongly associated with obesity.^{18–20} Our study, which is shown in Table 1, uncovered a fascinating outcome where the FBG levels were noticeably greater in the normo-weight group compared to the obese group, despite the difference not being statistically significant. Additionally, our results showed that the concentration of HDL-c was decreased in the normo-weight group when compared to the obese group, although

Table 2 Association Between Age, BMI, and MetS Components with ADMA and VCAM-I Level (N=99)

Variable	ADMA Level (ug/mL)				VCAM-I Level (ug/mL)			
	Normo-Weight (n=34)		Obese (n=65)		Normo-Weight (n=34)		Obese (n=65)	
	r	p-value	r	p-value	r	p-value	r	p-value
Age (years)	0.346	0.002*	0.223	0.020*	0.148	0.086	0.209	0.027*
BMI (kg/m ²)	-0.196	0.882	0.181	0.510	0.045	0.990	0.040	0.370
WC (cm)	-0.037	0.301	0.283	0.597	0.008	0.576	0.075	0.501
SBP (mmHg)	0.301	0.179	0.134	0.303	0.027	0.052	0.022	0.137
DBP (mmHg)	-0.014	0.524	0.045	0.346	0.129	0.251	-0.037	0.341
FBG (mg/dl)	0.519	0.000*	0.445	0.045*	0.329	0.009*	0.581	0.000*
HDL-c (mg/dl)	0.232	0.175	-0.242	0.139	0.140	0.131	-0.323	0.008*
TAG (mg/dl)	0.652	0.000*	0.522	0.006*	0.485	0.001*	0.399	0.270

Notes: Data is shown with r (coefficient correlation) and p-value (*significantly correlated if p-value <0.05).

Abbreviations: BMI, Body Mass Index; WP, Waist Circumference; SBP, Systolic Blood Pressure; DBP, Diastolic Blood Pressure; FBG, Fasting Blood Glucose; HDL-c, High Density Lipoprotein-cholesterol; TAG, Triglycerides; ADMA, Asymmetric dimethylarginine; VCAM-I, Vascular cells adhesion molecules-I.

the difference was not statistically significant. Taken together, these data indicated that the insulin function of normo-weight was worse than those in the obese group.

This study analyzed ADMA and VCAM-1 as they can be used as markers of the early phase of cardiovascular events.^{14,21,22} Average ADMA levels were still normal in both groups, while average VCAM-1 levels in both groups had increased. However, no significant difference was observed between normo-weight and obese groups. This phenomenon shows that endothelial injury has occurred in both groups, where the endothelial tone is still good, while the adhesive molecules have already been disrupted.²³

Endothelial injury in both groups, characterized by elevated VCAM-1 levels, may have been due to high systolic and diastolic levels in both groups. A previous study reported that in subjects with hypertension, VCAM-1 levels increased following high endothelin-1 (ET-1) levels. ET-1 stimulates arterial VCAM-1 expression by producing O₂⁻ from an ETA receptor/NADPH oxidase pathway in low-renin mineralocorticoid hypertension.^{24,25} Previous research has shown that MetS is linked to accelerated aortic stiffening, which is primarily associated with changes in blood pressure. Furthermore, individuals with hypertension have significantly higher levels of aortic stiffness, as measured by pulse wave velocity (PWV), compared to those without hypertension. The arterial structural and functional properties are thought to be related to this increased stiffness.^{26,27} In our study, this mechanism might undergo in our subjects, which leads to endothelial dysfunction represented by an increase in VCAM-1 in both groups, as shown in Table 2.

Table 2 shows that the correlation coefficient of endothelial dysfunction in normo-weight is stronger compared to obese, indicated by the r value of FBG and TAG in the ADMA value. This may be related to the “obesity paradox”, which explains that the prognosis of obese patients is better than that of normal-weight patients and that metabolic disorders are better indicators than BMI. Obese patients have more metabolic reserves. In addition, the cytokine and neurohormone profiles of obese patients are also cardioprotective. Adipose tissue produces tumor necrosis factor-alpha receptors that can counteract the adverse effects of tumor necrosis factor-alpha. Higher levels of lipoproteins in obese patients can bind to and neutralize lipopolysaccharides that play a role in stimulating the release of inflammatory cytokines. In addition, obese people tend to check themselves early, even in mild pain and get more aggressive treatment because they are considered more risky.²⁸⁻³⁰

Our study reveals that not all MetS components correlate significantly with ADMA and VCAM-1 as markers of endothelial dysfunction. This might be related to the conditions under which this study was conducted in populations

where MetS may still be in its early phase. This is an advantage because this study can detect early the incidence of endothelial dysfunction in people who experience metabolic disorders.

It is known that lipid disorders play a more important role in the incidence of endothelial dysfunction. Still, in this study, it is shown that FBG is consistently associated with VCAM-1. This suggests that carbohydrate metabolic disorders play a more important role than lipids, which may also be related to the paradoxical mechanism of obesity, as a result of a progressive catabolic state and lean mass loss.³⁰ The results of the study also showed that hyperglycemia triggers an inflammatory response and upregulation of VCAM-1.^{31,32}

The MetS components positively correlated with VCAM-1 as markers of endothelial dysfunction were FBG in both groups. Hyperglycemia is the major causal factor in the development of endothelial dysfunction in diabetes mellitus, although the mechanisms underlying this phenomenon are likely to be multifactorial.^{33,34} Increasing evidence suggests that the progression of insulin resistance to type 2 diabetes parallels the progression of endothelial dysfunction to atherosclerosis. Insulin resistance is closely linked to visceral adiposity, and early data suggested that free fatty acids were responsible for this association.^{33,35,36} This shows that metabolic parameters are more sensitive and play an important role in endothelial dysfunction than the functional disorders represented by SBP and DBP. This means that even if a person is of normal weight, they still have the same risk as an obese person for metabolic syndrome.³⁷

Metabolic syndrome or metabolically unhealthy patients were usually diagnosed late and were found in hospitals. In order to gain early awareness in preventing cardiovascular event in the community, this study was done in population setting. Urban areas in Indonesia are known to have high MetS data, including in the Cimahi area, West Java, which is the location of this study.^{38–40} The average age of MetS sufferers in this study was more than 40 years, consistent with previous studies.^{11,41,42} However, some studies report that MetS can occur at a younger age because the incidence of MetS often parallels the incidence of obesity and type-2 diabetes.^{38,43–45} The suggestion for future research is to elaborate molecular mechanisms in vitro regarding how hyperglycemia induced VCAM-1 increase. The results of this study can be used to formulate more precise strategies for early prevention of metabolically unhealthy state.

Conclusion

Among MetS components, metabolic parameters, especially FBG and TAG, have stronger correlation with endothelial dysfunction parameters (ADMA and VCAM-1) in normo-weight subjects with metabolic syndrome. Moreover, significant correlation between metabolic and endothelial dysfunction parameters was consistently observed in normo-weight metabolically unhealthy subjects.

Acknowledgments

We gratefully acknowledge support from the Universitas Padjadjaran Academic Leadership Grant for THA and Universitas Jenderal Achmad Yani research grant for IIR.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Chigom E. *Non-communicable diseases*. World Health. 1988.
2. Grundy SM, Brewer HB, Cleeman JI, Smith SC, Lenfant C. Definition of Metabolic Syndrome: report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation*. 2004;109(3):433–438. doi:10.1161/01.CIR.0000111245.75752.C6
3. Kaur J. A comprehensive review on metabolic syndrome. *Cardiol Res Pract*. 2014;2014:1–21. doi:10.1155/2014/943162
4. Pullaiah CP, Reddy DG. Metabolic Syndrome: a Rising Epidemic: comprehensive Review. *Int J Res Ayurveda Pharm*. 2017;8(3):36–41. doi:10.7897/2277-4343.083140
5. Dearborn JL, Viscoli CM, Inzucchi SE, Young LH, Kernan WN. Metabolic syndrome identifies normal weight insulin-resistant stroke patients at risk for recurrent vascular disease. *Int J Stroke*. 2019;14(6):639–645.
6. Shi TH, Wang B, Natarajan S. The influence of metabolic syndrome in predicting mortality risk among US adults: importance of metabolic syndrome even in adults with normal weight. *Prev Chronic Dis*. 2020;17:1–10. doi:10.5888/pcd17.200020

7. Mata AJ, Jasul G. Prevalence of metabolic syndrome and its individual features across different (Normal, overweight, pre-obese and obese) body mass index (BMI) categories in a tertiary hospital in the Philippines. *J ASEAN Federation Endocrine Societies*. 2017;32:117–122. doi:10.15605/jafes.032.02.04
8. Ding C, Chan Z, Magkos F. Lean, but not healthy. *Curr Opin Clin Nutr Metab Care*. 2016;19(6):408–417. doi:10.1097/MCO.0000000000000317
9. Ruderman N, Chisholm D, Pi-Sunyer X, Schneider S. The metabolically obese, normal-weight individual revisited. *Diabetes*. 1998;47(5):699–713. doi:10.2337/diabetes.47.5.699
10. Kim NH, Kim KJ, Choi J, Kim SG. Metabolically unhealthy individuals, either with obesity or not, have a higher risk of critical coronavirus disease 2019 outcomes than metabolically healthy individuals without obesity. *Metabolism*. 2022;128(January):154894. doi:10.1016/j.metabol.2021.154894
11. Suliga E, Koziel D, Głuszek S. Prevalence of metabolic syndrome in normal weight individuals. *Ann Agric Environ Med*. 2016;23(4):631–635. doi:10.5604/12321966.1226858
12. Eliana F, Suwondo P, Makmun LH, Kunci K. Faktor risiko disfungsi endotel pada prediabetes The risk factor of endothelial dysfunction in prediabetes. *J Kedokt Yars*. 2009;17(3):160–168.
13. Hardiman D, Medika M. Metformin Dosis Rendah Jangka Pendek dan Risiko Timbulnya Aterosklerosis pada Prediabetes. *Int J Med*. 2010;43(2):69–74.
14. Sibal L, Agarwal C, Sd H, Boger PH. The Role of Asymmetric Dimethylarginine (ADMA) in Endothelial Dysfunction and Cardiovascular Disease. *Curr Cardiol Rev*. 2010;6(2):82–90. doi:10.2174/157340310791162659
15. Palomo I, Contreras A, Alarcón LM, et al. Elevated concentration of asymmetric dimethylarginine (ADMA) in individuals with metabolic syndrome. *Nitric Oxide - Biol Chem*. 2011;24(4):224–228. doi:10.1016/j.niox.2011.03.002
16. Kemenkes. Epidemi Obesitas. FactSheet_Obesitas_Kit_Informasi_Obesitas.pdf. 2021. 1–8.
17. Fathurrahman M. Hubungan Asymmetric Dimethylarginine (Adma) Dengan Hasil Luanan Bayi Pada Pasien Dengan Preeklamsia. 2017.
18. Gupta G, Wadhwa R, Pandey P, et al. Obesity and Diabetes: Pathophysiology of Obesity-Induced Hyperglycemia and Insulin Resistance. *Pathophysiology of Obesity-Induced Health Complications*. Cham. Springer International Publishing. 2020 81–97. Available from: <http://www.springer.com/series/7064>. Accessed February 13, 2024.
19. Martyn JAJ, Kaneki M, Yasuhara S. Obesity-induced insulin resistance and hyperglycemia: etiologic factors and molecular mechanisms. *Anesthesiology*. 2008;109(1):137–148. doi:10.1097/ALN.0b013e3181799d45
20. Toelsie JR, Morpurgo F, Krishnadath I, Bipat R. Obesity, overweight and hyperglycemia among primary school children in a low-middle income country with a multiethnic population. *Obes Pillars*. 2023;5(November 2022):100053. doi:10.1016/j.obpill.2022.100053
21. Sibal L, Agarwal SC, Schwedhelm E, Lüneburg N, Böger RH, Home PD. A study of endothelial function and circulating asymmetric dimethylarginine levels in people with Type 1 diabetes without macrovascular disease or microalbuminuria. *Cardiovasc Diabetol*. 2009;8(1):1–7. doi:10.1186/1475-2840-8-27
22. Couffinhal T, Duplâa C, Moreau C, Lamazière JMD, Bonnet J. Regulation of vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 in human vascular smooth muscle cells. *Circ Res*. 1994;74(2):225–234. doi:10.1161/01.RES.74.2.225
23. Medina-leyte DJ, Zepeda-garc O, Dom M, González-Garrido A, Villarreal-Molina T, Jacobo-Albavera L. Endothelial Dysfunction, Inflammation and Coronary Artery Disease: potential Biomarkers and Promising Therapeutical Approaches. *Int J Mol Sci*. 2021;22(8):1–28. doi:10.3390/ijms22083850
24. Li L, Chu Y, Fink GD, Engelhardt JF, Heistad DD, Chen AF. Endothelin-1 Stimulates Arterial VCAM-1 Expression Via NADPH Oxidase-Derived Superoxide in Mineralocorticoid Hypertension. *Hypertension*. 2003;42(5):997–1003. doi:10.1161/01.HYP.0000095980.43859.59
25. Abraham D, Distler O. How does endothelial cell injury start? The role of endothelin in systemic sclerosis. *Arthritis Res Ther*. 2007;9(SUPPL. 2):S2. doi:10.1186/ar2186
26. Maloberti A, Bruno RM, Facchetti R, et al. THE role of metabolic syndrome in blood pressure control and pulse wave velocity progression over a 3.5 years in treated hypertensive PATIENTS. *Eur J Intern Med*. 2020;76(February):107–109. doi:10.1016/j.ejim.2020.02.005
27. Maloberti A, Bombelli M, Vallerio P, et al. Metabolic syndrome is related to vascular structural alterations but not to functional ones both in hypertensives and healthy subjects. *Nutr Metab Cardiovasc Dis*. 2021;31(4):1044–1052. doi:10.1016/j.numecd.2020.11.011
28. Nursalim A, Yuniadi Y. Paradox Obesitas pada Pasien Gagal Jantung. *J Kardiol Indones*. 2011;32(4):207–215.
29. Cheung Y-M, Joham A, Marks S, Teede H. The obesity paradox: an endocrine perspective. *Intern Med J*. 2017;47(7):727–733. doi:10.1111/imj.13257
30. Elagizi A, Kachur S, Lavie CJ, et al. An Overview and Update on Obesity and the Obesity Paradox in Cardiovascular Diseases. *Prog Cardiovasc Dis*. 2018;61(2):142–150. doi:10.1016/j.pcad.2018.07.003
31. Zhang LF, Zalewski A, Liu Y, et al. Diabetes-induced oxidative stress and low-grade inflammation in porcine coronary arteries. *Circulation*. 2003;108(4):472–478. doi:10.1161/01.CIR.0000080378.96063.23
32. Gustavsson C, Agardh CD, Zetterqvist AV, Nilsson J, Agardh E, Gomez MF. Vascular cellular adhesion molecule-1 (VCAM-1) expression in mice retinal vessels is affected by both hyperglycemia and hyperlipidemia. *PLoS One*. 2010;5(9):1–12. doi:10.1371/journal.pone.0012699
33. Kolluru GK, Bir SC, Kevil CG. Endothelial dysfunction and diabetes: effects on angiogenesis, vascular remodeling, and wound healing. *Int J Vasc Med*. 2012;2012:8267. doi:10.1155/2012/918267
34. Hadi HAR, Al Suwaidi JA. Endothelial dysfunction in diabetes mellitus. *Vasc Health Risk Manag*. 2007;3(6):853–876.
35. Wadhwa C, Parker A, Wang XP. High Glucose Attenuates Protein S -Nitrosylation in. *Diabetes*. 2007;56(November):2715–2721. doi:10.2337/db06-1294
36. Virtanen KA, Iozzo P, Hällsten K, et al. Increased fat mass compensates for insulin resistance in abdominal obesity and type 2 diabetes: a positron-emitting tomography study. *Diabetes*. 2005;54(9):2720–2726. doi:10.2337/diabetes.54.9.2720
37. Rakhmat II, Putra ICS, Wibowo A, et al. Cardiometabolic risk factors in adults with normal weight obesity: a systematic review and meta-analysis. *Clin Obes*. 2022;12(4). doi:10.1111/cob.12523
38. Kamso S, Purwastyastuti P, Lubis DU, Juwita R, Robbi YK, Besral B. Prevalensi dan Determinan Sindrom Metabolik pada Kelompok Eksekutif di Jakarta dan Sekitarnya. *Kesmas Natl Public Heal J*. 2011;6(2):85. doi:10.21109/kesmas.v6i2.110
39. Zahtamal Z, Prabandari YS, Setyawati L. Prevalensi Sindrom Metabolik pada Pekerja Perusahaan. *Kesmas Natl Public Heal J*. 2014;9(2):113. doi:10.21109/kesmas.v9i2.499
40. Inayati Rakhmat I, Jusuf H, Rachmanita S. Prevalence and Characteristics of Metabolic Syndrome Patients in Cibeber Village Cimahi Indonesia. *J Heal Dent Sci*. 2021;1(Volume 1 No 1):56.
41. Saklayen MG. The Global Epidemic of the Metabolic Syndrome. *Curr Hypertens Rep*. 2018;20(2):1–8. doi:10.1007/s11906-018-0812-z
42. Jahangiry L, Khosravi-Far L, Sarbakhsh P, Kousha A, Entezarmahdi R, Ponnet K. Prevalence of metabolic syndrome and its determinants among Iranian adults: evidence of IraPEN survey on a bi-ethnic population. *Sci Rep*. 2019;9(1):1–7. doi:10.1038/s41598-019-44486-8

43. Alowfi A, Binladen S, Iqrqous S, Khashoggi A, Khan MA, Calacattawi R. Metabolic syndrome: prevalence and risk factors among adolescent female intermediate and secondary students in Saudi Arabia. *Int J Environ Res Public Health*. 2021;18(4):1–13. doi:10.3390/ijerph18042142
44. Cook S, Weitzman M, Auinger P, Nguyen M, Dietz WH. Prevalence of a Metabolic Syndrome Phenotype in Adolescents. *Arch Pediatr Adolesc Med*. 2003;157(8):821. doi:10.1001/archpedi.157.8.821
45. Elabbassi WN, Haddad HA. The epidemic of the metabolic syndrome. *Saudi Med J*. 2005;26(3):373–375.

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>