



Prevalence, awareness, and associated factors of high blood pressure among female migrant workers in Central South China

Hua Peng¹, Mei Sun², Xin Hu³, Huiwu Han⁴, Jing Su⁵, Emin Peng⁶, James Wiley⁷, Lisa Lommel⁸ and Jyu-Lin Chen⁸

¹ Department of Nursing, Xiangya Hospital of Central South University, Changsha, China

² Community Nursing Department, Xiangya Nursing School of Central South University, Hunan Women Research Association, Changsha, China

³ Community Nursing Department, Xiangya Nursing School of Central South University, Changsha, China

⁴ Department of Cardiology, Xiangya Hospital of Central South University, Changsha, China

⁵ Department of Nursing, Shantou University Medical College, Shantou, China

⁶ Outpatient Clinic, Xiangya Hospital of Central South University, Changsha, China

⁷ Institute for Health Policy Studies, University of California, San Francisco, CA, United States of America

⁸ Department of Family Health Care Nursing, School of Nursing, University of California, San Francisco, CA, United States of America

ABSTRACT

Background. Although many young Chinese women migrate to urban regions for better opportunities, little is known about the prevalence and awareness of having high blood pressure (HBP) in this population. This study investigated the prevalence, awareness, and factors associated with HBP among young female migrant workers in Central South China.

Methods. We conducted a cross-sectional study to identify HBP (2017 ACC/AHA guidelines) among female migrant workers aged 18–45 years in Central South China. Demographics, anthropometric measurements, hypertension-related lifestyle, awareness of HBP, and blood pressure were recorded. Logistic regression analysis was used to identify the factors associated with HBP (blood pressure \geq 130/80 mmHg).

Results. Overall, 232 female migrants participated in the study (mean age 34.4; standard deviation: 6.4 years). The prevalence of HBP was 27.2% (95% CI [21.6–33.2]), and 88.9% of the participants were unaware of their HBP status. Having rural medical insurance (odds ratio [OR] = 20.7; 95% confidence interval 95% CI [2.1–204.8]), awareness of having HBP (OR = 5.1; 95% CI [1.4–18.5]), physical inactivity (OR = 2.9; 95% CI [1.1–7.9]), and being overweight/obese (OR = 2.7; 95% CI [1.3–6.1]) were independently associated with HBP.

Conclusions. This study revealed a high prevalence of HBP among young Chinese female migrant workers, as well as a high frequency of being unaware of their condition and some associated factors (rural medical insurance, awareness of having HBP, physical inactivity, and overweight/obesity). The uncontrolled HBP among young Chinese female migrant workers suggested that health education needs further promotion in such a population.

Submitted 8 July 2020

Accepted 10 April 2022

Published 4 May 2022

Corresponding author

Mei Sun, smnjw2008@126.com

Academic editor

Andrew Gray

Additional Information and
Declarations can be found on
page 17

DOI 10.7717/peerj.13365

© Copyright
2022 Peng et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Cardiology, Public Health, Women's Health

Keywords Hypertension, Migrant workers, Prevalence, Awareness

INTRODUCTION

Hypertension is a leading cause of cardiovascular diseases (CVD) worldwide (*Roth et al., 2020; Wu et al., 2015*) and is also strongly associated with many cerebrovascular diseases such as stroke (*Roth et al., 2020; Wajngarten & Silva, 2019*). The Chinese hypertension guidelines are similar to the recommendations from the Eighth Joint National Committee (JNC-8) (*James et al., 2014; Wang, 2015*), defining hypertension as systolic blood pressure (SBP) ≥ 140 mmHg or diastolic blood pressure (DBP) ≥ 90 mmHg, or use of antihypertensive medication, while high blood pressure (HBP) is defined as blood pressure that is high but without necessarily qualifying for hypertension, *i.e.*, SBP ≥ 130 mmHg or DBP ≥ 80 mmHg (*Li et al., 2019*). The high prevalence of hypertension significantly aggravates the public health burden demonstrated in morbidity, mortality, and medical care expenses, particularly in low- and middle-income countries, including China (*Sarki et al., 2015; van Kleef & Spiering, 2017*). As well as having the largest proportion of the world's population, China has experienced a rapidly increased prevalence of hypertension among adults aged ≥ 18 years, from 26.6% in 1990 to 33.6% in 2016 (*Bundy & He, 2016*). According to the most recent national survey on adults aged 35–75 years in China (in 2017), the prevalence of hypertension has reached 44.7%, but the overall rates for awareness of having HBP, hypertension treatment, and control ($<140/90$ mmHg) were poor (44.7%, 30.1%, and 5.7%, respectively) (*Lu et al., 2017*).

As prevention of hypertension is key to preventing CVD, the clinical practice guidelines for detecting and treating hypertension were updated in 2017 by the American College of Cardiology/American Heart Association (ACC/AHA) (*Whelton et al., 2018*). By lowering the blood pressure level for hypertension diagnosis, the new 2017 guidelines targeted BP reduction in adults at high risk of hypertension or adults with hypertension to prevent developing HBP-related CVD, especially adults who are overweight, consume excessive dietary sodium and alcohol, or those who are physically inactive (*Whelton et al., 2018*). Considering the importance of early prevention for hypertension management of young patients, this study used the new definition of hypertension stage I recommended by the ACC/AHA as the cutoff point (SBP ≥ 130 mmHg or DBP ≥ 80 mmHg) for HBP.

Mainland China has the largest population of native migrant workers (*i.e.*, born in China and migrated to another region/province for work) globally, with most of them being rural-to-urban migrants (*National Bureau of Statistics of the People's Republic of China, 2016*). In 2016, rural-to-urban migrant workers had increased to 281.7 million, accounting for 20.4% of the total Chinese population (*National Bureau of Statistics of the People's Republic of China, 2017*). Young migrants aged 26–40 years having a greater prevalence of prehypertension (SBP 120–139 mmHg and/or DBP 80–89 mmHg) compared to older migrants (≥ 41 years) has been observed (*Guan, 2018*). *Ying et al. (2015)* reported that young migrant people aged ≥ 30 years had a higher prevalence of hypertension than migrant adults aged <30 years (12.4% vs. 5.5%, $p = 0.001$; odds ratio [OR]: 2.0, 95% CI [1.3–3.0]). Many studies have found a higher prevalence of hypertension in people aged ≥ 35 years than in those <35 years (*Donghui et al., 2015; Gao et al., 2013; Hu, Gao & Liu, 2019; Li et al., 2019; Liu et al., 2017*). Still, it remains unclear whether the epidemiology

and risk factors of HBP among women migrant workers are the same as in the general population.

Lifestyle modification could significantly reduce the prevalence of HBP and hypertension, especially in urban dwellers. The potentially modifiable factors related to lifestyle include smoking cessation, restricted alcohol intake, increased fruit and vegetable consumption, weight loss, increased physical activity, and reduced dietary salt intake (Juraschek et al., 2017; Masterson Creber et al., 2010; Zhou, 2002; Zhou et al., 2002). Bi et al. (2016) suggested that obesity might be the earliest manifestation of hypertension and CVD among Chinese migrant workers after transitioning to an urbanized lifestyle. Another study found that long working hours (>8 hours/day), higher income (>USD 490/month), and single marital status are associated with unhealthy lifestyles in female migrant workers (Yang et al., 2015a). Understanding the prevalence of hypertension and health behaviors related to hypertension, which is the leading cause of CVD, is critical in eliminating health disparities in this young and understudied population.

Therefore, this study aimed (a) to estimate the prevalence and awareness of having HBP and (b) to identify factors associated with HBP among female migrant workers from Central South China.

METHOD

Study design and setting

This cross-sectional study was conducted in Changsha, Hunan Province, Central South China, between March 2017 and June 2017. Hunan has a total population of 68.99 million, with 38.65 million (56.0%) urban and 30.34 million (44.0%) rural residents by the end of 2018. However, the number of domestic migrant workers in Hunan has reached approximately 17.58 million, accounting for approximately 25.5% of the total provincial population (Hunan Provincial Bureau of Statistics, 2018). Hunan ranked third in the total number of migrant workers in China and first in Central South China (National Bureau of Statistics of China, 2020). Among the migrant population in Hunan, nearly 9.59 million (54.6% of the total migrants in Hunan) were relatively young, born after 1980. Changsha is the biggest commercial, transportation, and manufacturing center in Hunan, with approximately 8.15 million people in 2018 (Hunan Provincial Bureau of Statistics, 2018).

According to the Chinese National Bureau of Statistics, all qualified workplaces were classified into three levels: large, $\geq 1,000$ employees; medium, 300–999 employees; and small, 10–299 employees (National Bureau of Statistics of China, 2020). Workplaces with female migrant workers accounting for 50% of the total employees or above were eligible.

The institutional review board of the University of California, San Francisco, and Central South University reviewed and approved the study protocol (ID: 2017036). Patients' personal information and anthropometric data are kept strictly confidential and only used for scientific research. The participants all signed the informed consent form.

Study population

Mandarin-speaking research assistants recruited a convenience sample of young female participants from custodians in a tertiary hospital, workers in a medical device factory,

and staff in a restaurant in Changsha, representing large-, medium-, and small-sized workplaces, respectively, considering that these migrant women are mainly engaged in the labor-intensive light industry and service sectors due to their traits of physical strength and education. The inclusion criteria were (a) Chinese female migrants from rural regions working in Changsha, (b) age between 18 and 45 years, (c) able to read and write in Mandarin Chinese, and (d) has stayed in Changsha as a migrant worker for at least 6 months. Participants were not recruited if they could not provide consent due to physical or mental illness or declined to participate.

Sample size

The sample size was estimated using an online sample size calculator for prevalence studies (<https://www.calculator.net/sample-size-calculator.html>). Assuming that the prevalence of hypertension in the female migrant workers was similar to the reported prevalence of 15.6% (Wang et al., 2018) in the population aged ≥ 18 years in Hunan province in 2012–2015, we calculated the minimum sample size by taking 5% permission error with 95% confidence interval (CI) and the population size as 8.15 million. Considering the nonresponse rate of 20%, a final sample size of 254 was calculated.

Measurements

All participants were informed about the purpose of the study and details of their enrollment by trained interviewers, and they provided written informed consent at their workplace. After completing the screening questionnaire, each subject's weight, height, waist circumference (WC), and blood pressure were measured by trained research assistants in a separate room to ensure privacy. All measurements and screenings were completed at the place of their employment. The interviewers were trained by the research principal investigator to perform the procedure in a standardized manner, collect data, and interpret the questionnaire in standard language. Meanwhile, the research assistants were trained about physical measurements by a professional cardiac clinical instructor. Each participant received a small umbrella as a token of appreciation for their participation.

Demographic survey

A questionnaire was developed to examine the sociodemographic characteristics of the sample population, including their age (≥ 35 or < 35 years), the highest level of school (higher than high school vs. high school or less), marital status (married, single, or widowed), family location (village or town), whether a mother with children (yes or no), job category (manager or worker), the length of being a migrant worker (< 1 , 1–3, 4–5, or > 5 years), average monthly income (≤ 420 or > 420 USD), and hours worked per day (≤ 8 or > 8 h).

The participants were also asked about whether they had rural medical insurance (yes or no), had ever taken HBP pills (yes or no), and received an annual health examination (yes or no). Regarding whether their parents had ever been diagnosed with CVD, the responses included yes, no, or don't know.

Survey on hypertension-related lifestyle

The questionnaire also obtained lifestyle behaviors, including dietary behaviors, physical activity level, and cigarette smoking status. Items related to dietary behaviors included whether the participants consumed five servings of fruits and vegetables per day, were told by a physician or a nurse to reduce salt intake, or were told by a physician or a nurse to reduce carbohydrate intake. Physical activity level was assessed by asking participants how many days per week they engage in regular physical exercise of ≥ 30 min per session. The responses included ≥ 3 days per week and < 3 days per week. Regarding smoking status, information was collected about whether they were current smokers (yes or no). These lifestyle behavior questions are included in the Chinese version of the Canadian Diabetes Risk Questionnaire, which has been confirmed to have good reliability and validity (*Guo et al., 2018*).

Awareness of having HBP and self-rated health

The question assessed the awareness of having HBP, “Have you ever been told by a physician or nurse that your blood pressure was high?” Overall health was self-rated by participants as poor, fair, good, very good, or excellent.

Blood pressure

Blood pressure was measured using an OMRON automatic blood pressure monitor (HBP-1100; Omron Corporation, Dalian, China) and validated using mercury sphygmomanometer measurements. The participants had to avoid caffeine, exercise, and smoking for at least 30 min before their visit. They were asked to empty their bladder before entering the test room. The participants were told to relax and rest in the seated position (feet on the floor and back supported) for 10 min, without talking, and two blood pressure measurements were obtained at 60-s intervals, also without talking. Blood pressure measurements were made on the left arm, with the transducer placed at the level of the heart and the cuff adapted to the arm circumference. The final blood pressure result was the arithmetic mean of the two measurements taken on the same arm. The precision of the automatic blood pressure monitor was one mmHg. The participants were considered to have HBP if the SBP was ≥ 130 mmHg or the DBP was ≥ 80 mmHg (including stage I and above). Hypertension was defined as having HBP or self-reported antihypertensive medication use.

Body composition

Body composition was assessed by measuring the body mass index (BMI). The weight (kg) and height (cm) of the participants were measured, and BMI was calculated as weight in kilograms divided by the square of height in meters ($\text{weight}/\text{height}^2$ (kg/m^2)). With the participants wearing light clothing and no shoes, weight was measured to the nearest 0.5 kg in increments of up to 120 kg using the Penguin TCS weighing scale (TCS-766; Wuyi Penguin Industry and Trade Co. Ltd., Zhejiang, China). Height was measured to the nearest 0.5 cm using a stadiometer, with the participant not wearing footwear. Overweight ($24 \text{ kg}/\text{m}^2 \leq \text{BMI} < 28 \text{ kg}/\text{m}^2$) and obesity ($\text{BMI} \geq 28 \text{ kg}/\text{m}^2$) were identified using standard Chinese-specific BMI cutoff values (*Jin et al., 2013; Yang et al., 2017*).

Waist circumference

The WC was measured using standard protocols. With the participant in a standing position, the abdomen was fully exposed, and measurement was conducted in a state of natural respiration. The WC was determined as the smallest measure of the waist at the level of the umbilicus by using a nonelastic plastic ribbon with a width of 0.7 cm. The reading was measured to the nearest 0.1 cm. Based on previous studies, the participants were defined to have central obesity when the WC was ≥ 80 cm (Du et al., 2017; Xu et al., 2016).

Statistical analysis

The participants' sociodemographic characteristics were presented using descriptive statistics, including frequencies, percentages, and means with standard deviations (SDs). Bivariable logistic regression analyses were performed to examine the odds ratio (OR) and 95% CI of hypertension by separate parameters. A multivariable logistic regression analysis was performed to estimate the OR and 95% CI of hypertension after adjustment for variables such as age, education, cigarette use, physical activity, fruit and vegetable intake, receiving advice to reduce salt and carbohydrates, parental history of CVD, WC, BMI, awareness of having HBP and rural medical insurance. A directed acyclic graph is a method to illustrate causal relationships between variables based on professional and epidemiological knowledge (Etmnan, Collins & Mansournia, 2020). Then, the variance inflation factor (VIF) was used to check the collinearity of the selected variables. VIF < 5 indicated that collinearity was unlikely to be a problem (Gueugnon et al., 2019; Kim, 2019; Marcoulides & Raykov, 2019). The multivariable regression model for the outcome (HBP) was described using a receiver operating characteristic (ROC) curve and the area under the curve (AUC) (Hajian-Tilaki, 2013; Zou, O'Malley & Mauri, 2007). The Hosmer-Lemeshow test and calibration curves were used to test the calibration of the model. All *P*-values were two-sided and considered statistically significant at < 0.05 . Statistical analyses were conducted using IBM SPSS Statistics v.25.0 (International Business Machines Corp., Armonk, New York, USA).

RESULTS

Participant characteristics

Of the 268 participants considered for participation, nine declined to participate in the study because of time constraints, thirteen did not complete the questionnaire, and fourteen missed at least one of the physical measurements. The overall response rate was 86.6% (232 participants), with missing data in $< 5\%$ of the participants (Fig. 1). There were 167 custodians from a tertiary hospital, 51 workers from a medical device factory, and 14 staff in a restaurant in Changsha, representing large-, medium-, and small-sized workplaces, respectively.

Of the 232 female migrant workers who participated in this study, 157 (67.7%) migrated from rural villages, while 75 (32.3%) migrated from small towns located in suburban areas where the living conditions are more similar to rural conditions. The mean age of the participants was 34.4 years (SD = 6.4, range 18–45 years). Most of the migrant women




Figure 1 Flowchart.

Full-size  DOI: [10.7717/peerj.13365/fig-1](https://doi.org/10.7717/peerj.13365/fig-1)

had a high school diploma or less education (88.8%), were married and also had children (79.7%), and made <USD 420/month (77.2%). Additionally, about one-third of these women reported working >8 hours/day, and one-half has migrated for more than 5 years (Table 1).

The mean BMI was 22.17 kg/m² (SD = 3.28), with 27.2% of the migrant women being overweight (22.4%) or obese (4.7%). The mean WC was 79.30 cm (SD = 8.20), with 44.4% of the participants meeting the cutoff for central obesity (≥ 80 cm).

Prevalence/awareness of having HBP and lifestyle behaviors

The prevalence of HBP and hypertension (with HBP or antihypertensive medication use) was 27.2% (63/232; 95% CI [21.6–33.2]) and 28.9% (67/232; 95% CI [23.3–34.9]), respectively. Of the female migrant workers with HBP, 69.8% (44/63) had stage I hypertension, while 30.2% (19/63) had stage II hypertension (Table 2).

Approximately 5.6% (13/232) of female migrants believed that they currently or formerly had HBP (Table 3). Notably, 88.9% of the female migrants with HBP reported that they were unaware of their HBP status, and only four used antihypertensive medication (Table 3). Moreover, 50.8% of these workers who had HBP self-rated their overall health status as good, very good, or excellent. Overall, 59.1% had undergone annual health physical

Table 1 Characteristics of the participants, frequencies (percentage) [n (%)].

| Characteristic | Total (n = 232) | High blood pressure ^a | |
|-----------------------------|--------------------|----------------------------------|-----------------|
| | | No (n = 169) | Yes (n = 63) |
| Age (years) | | | |
| <35 | 127 (54.7) | 100 (59.2) | 27 (42.9) |
| ≥35 | 105 (45.3) | 69 (40.8) | 36 (57.1) |
| Education | | | |
| Higher than high school | 26 (11.2) | 21 (12.4) | 5 (7.9) |
| High school or less | 206 (88.8) | 148 (87.6) | 58 (92.1) |
| Marital status | | | |
| Married | 212 (91.4) | 153 (90.5) | 59 (93.7) |
| Single/widowed | 20 (8.6) | 16 (9.5) | 4 (6.3) |
| Family location | | | |
| Village | 157 (67.7) | 114 (67.5) | 43 (68.3) |
| Town | 75 (32.3) | 55 (32.5) | 20 (31.7) |
| Mother with children | | | |
| No | 47 (20.3) | 33 (19.5) | 14 (22.2) |
| Yes | 185 (79.7) | 136 (80.5) | 49 (77.8) |
| Job category | | | |
| Manager | 33 (14.2) | 26 (15.4) | 7 (11.1) |
| Worker | 199 (85.8) | 143 (84.6) | 56 (88.9) |
| Length of migration (years) | | | |
| <1 | 25 (10.8) | 17 (10.1) | 8 (12.7) |
| 1–3 | 59 (25.4) | 47 (27.8) | 12 (19.0) |
| 4–5 | 21 (9.1) | 13 (7.7) | 8 (12.7) |
| >5 | 127 (54.7) | 92 (54.4) | 35 (55.6) |
| Monthly income | | | |
| ≤420 USD | 179 (77.2) | 131 (77.5) | 48 (76.2) |
| >420 USD | 53 (22.8) | 38 (22.5) | 15 (23.8) |
| Hours worked per day | | | |
| ≤8 | 147 (63.4) | 111 (65.7) | 36 (57.1) |
| >8 | 85 (36.6) | 58 (34.3) | 27 (42.9) |
| Rural medical insurance | | | |
| No | 33 (14.2) | 32 (18.9) | 1 (1.6) |
| Yes | 199 (85.8) | 137 (81.1) | 62 (98.4) |
| Annual health examination | | | |
| No | 95 (40.9) | 74 (43.8) | 21 (33.3) |
| Yes | 137 (59.1) | 95 (56.2) | 42 (66.7) |
| Parents with CVD | | | |
| No/do not know | 196 (84.5) | 142 (84.0) | 54 (85.7) |
| Yes | 36 (15.5) | 27 (16.0) | 9 (14.3) |

(continued on next page)

Table 1 (continued)

| Characteristic | Total (n = 232) | High blood pressure ^a | |
|---|--------------------|----------------------------------|-----------------|
| | | No (n = 169) | Yes (n = 63) |
| Consume 5 servings of fruits/vegetables per day | | | |
| No | 149 (64.2) | 108 (63.9) | 41 (65.1) |
| Yes | 83 (35.8) | 61 (36.1) | 22 (34.9) |
| Advised to reduce salt intake | | | |
| No | 152 (65.5) | 114 (67.5) | 38 (60.3) |
| Yes | 80 (34.5) | 55 (32.5) | 25 (39.7) |
| Advised to reduce carbohydrate intake | | | |
| No | 189 (81.5) | 139 (82.2) | 50 (79.4) |
| Yes | 43 (18.5) | 30 (17.8) | 13 (20.6) |
| Physical activity ≥ 3 days/week | | | |
| No | 189 (81.5) | 132 (78.1) | 57 (90.5) |
| Yes | 43 (18.5) | 37 (21.9) | 6 (9.5) |
| Current smoker | | | |
| No | 229 (98.7) | 168 (99.4) | 61 (96.8) |
| Yes | 3 (1.3) | 1 (0.6) | 2 (3.2) |
| Self-rated overall health | | | |
| Good/ very good/ excellent | 135 (58.2) | 103 (60.9) | 32 (50.8) |
| Fair/poor | 97 (41.8) | 66 (39.1) | 31 (49.2) |
| Body mass index (kg/m ²) | | | |
| <24 | 169 (72.8) | 132 (78.1) | 37 (58.7) |
| ≥ 24 | 63 (27.2) | 37 (21.9) | 26 (41.3) |
| Waist circumference (cm) | | | |
| <80 | 129 (55.6) | 103 (60.9) | 26 (41.3) |
| ≥ 80 | 103 (44.4) | 66 (39.1) | 37 (58.7) |

Notes.^aSBP ≥ 130 mmHg or DBP ≥ 80 mmHg.

CVD, cardiovascular disease; DBP, diastolic blood pressure; SBP, systolic blood pressure; USD, United States dollar.

Table 2 Distribution of blood pressure among the female migrant workers in Central South China, frequencies (percentage) [n (%)].

| BP category (SBP/DBP, mmHg) | n (%) | 95% CI ^a |
|--|------------|---------------------|
| Normal (<120/80) | 158 (68.1) | 62.1–74.1 |
| Elevated (120–129/<80) | 11 (4.7) | 2.2–7.3 |
| Hypertension stage I (130–139 /or 80–89) | 44 (19.0) | 14.2–24.1 |
| Hypertension stage II (≥ 140 /or ≥ 90) | 19 (8.2) | 4.7–12.5 |

Notes.

DBP, diastolic blood pressure; SBP, systolic blood pressure; CI, confidence interval.

^a95% CI of the percentage.

Table 3 Awareness and treatment of participants.

| | High blood pressure ^a | | P |
|---------------------------------|----------------------------------|--------------|-------|
| | No (n = 169) | Yes (n = 63) | |
| Awareness | | | 0.026 |
| Yes | 6 (3.6) | 7 (11.1) | |
| No | 163 (96.4) | 56 (88.9) | |
| Antihypertensive medication use | | | 0.218 |
| Yes | 4 (2.4) | 4 (6.3) | |
| No | 165 (97.6) | 59 (93.7) | |

Notes.

Data displayed as frequency (percentage) [n (%)].

^aSBP \geq 130 mmHg or DBP \geq 80 mmHg.

examinations, 1.3% reported smoking, 64.2% reported consuming less than five servings of fruits and vegetables daily, and 81.5% reported not engaging in physical activity at least three days per week.

Additionally, among the 67 participants with hypertension (SBP \geq 130 mmHg or DBP \geq 80 mmHg or antihypertensive medication use), 83.6% were unaware of it, 11.9% used antihypertensives, and 94.0% had uncontrolled blood pressure.

Factors associated with HBP

The bivariable analyses showed that age \geq 35 years, physical activity $<$ 3 days/week, waist circumference \geq 80 cm, BMI \geq 24 kg/m², awareness of having HBP, and having rural medical insurance were associated with higher odds of HBP (all $P <$ 0.05, Table 4).

The multivariable logistic regression analysis (with all variables entered at the same time) showed that the significant factors associated with higher odds of HBP were having rural medical insurance (OR = 20.7, 95%CI [2.1–204.8]), physical activity less than three days per week (OR = 2.9, 95%CI [1.1–7.9]), being overweight or obese (OR = 2.7, 95%CI [1.2–6.1]), having parents without CVD or uncertain (OR = 2.8, 95%CI [1.0–7.7]), and awareness of having HBP (OR = 5.1, 95%CI [1.4–18.5]) (Table 4). No collinearity was observed among these included variables (all VIF $<$ 5). The five statistically significant variables above were incorporated into a logistic regression model. The AUC of this model was 0.742 (95%CI [0.673–0.811], Fig. 2A), and Fig. 2B presents the calibration curve. The Hosmer-Lemeshow test revealed $P =$ 0.801.

Parental CVD history may directly increase the risk of HBP among migrant workers. A current awareness of having high blood pressure (being informed of HBP by a physician or nurse) may indirectly affect the prevalence of HBP by influencing physical activity, BMI, and purchase of rural medical insurance (Fig. S1).

DISCUSSION

This study investigated the prevalence, awareness, and factors associated with HBP (SBP \geq 130 mmHg or DBP \geq 80 mmHg) among young female migrant workers in Central South China. We found that approximately 27% of female migrant workers had HBP, and the majority of them were unaware of their condition. In total, 50% reported having good, very good, or excellent overall health status. Nearly one-third of the migrant women were

Table 4 Logistic regressions for high blood pressure among female migrant workers in China.

| Variables | Bivariable analysis | | | Multivariable analysis | | |
|---|---------------------|-------------|-------|------------------------|-------------|-------|
| | OR | 95% CI | P | OR | 95% CI | P |
| Age (years) | | | | | | |
| <35 | reference | | | reference | | |
| ≥35 | 1.93 | 1.08–3.47 | 0.028 | 1.33 | 0.66–2.69 | 0.426 |
| Education | | | | | | |
| Higher than high school | reference | | | reference | | |
| High school or less | 1.65 | 0.59–4.57 | 0.339 | 1.23 | 0.40–3.81 | 0.714 |
| Physical activity ≥ 3 days/week | | | | | | |
| No | 2.66 | 1.07–6.66 | 0.036 | 2.94 | 1.09–7.90 | 0.033 |
| Yes | reference | | | reference | | |
| Consume 5 servings of fruits/vegetables per day | | | | | | |
| No | 1.05 | 0.57–1.93 | 0.868 | 1.32 | 0.66–2.62 | 0.437 |
| Yes | reference | | | reference | | |
| Current smoker | | | | | | |
| No | reference | | | reference | | |
| Yes | 5.51 | 0.49–61.84 | 0.167 | 3.29 | 0.25–42.48 | 0.362 |
| Advised to reduce salt intake | | | | | | |
| No | reference | | | reference | | |
| Yes | 1.36 | 0.75–2.48 | 0.310 | 1.97 | 0.83–4.68 | 0.125 |
| Advised to reduce carbohydrate intake | | | | | | |
| No | reference | | | reference | | |
| Yes | 1.21 | 0.58–2.49 | 0.615 | 0.46 | 0.16–1.30 | 0.142 |
| Parents with CVD | | | | | | |
| No/do not know | 1.14 | 0.50–2.58 | 0.752 | 2.78 | 1.01–7.69 | 0.049 |
| Yes | reference | | | reference | | |
| Waist circumference (cm) | | | | | | |
| <80 | reference | | | reference | | |
| ≥80 | 2.22 | 1.23–4.00 | 0.008 | 1.65 | 0.78–3.47 | 0.189 |
| Body mass index (kg/m ²) | | | | | | |
| <24 | reference | | | reference | | |
| ≥24 | 2.51 | 1.35–4.66 | 0.004 | 2.74 | 1.23–6.12 | 0.014 |
| Awareness of having HBP | | | | | | |
| No | reference | | | reference | | |
| Yes | 3.40 | 1.10–10.53 | 0.034 | 5.06 | 1.39–18.49 | 0.014 |
| Rural medical insurance | | | | | | |
| No | reference | | | reference | | |
| Yes | 14.48 | 1.94–108.39 | 0.009 | 20.69 | 2.09–204.76 | 0.010 |

Notes.

CI, confidence interval; CVD, cardiovascular disease; HBP, high blood pressure; OR, odds ratio.




Figure 2 The receiver operating characteristic curve (ROC) (A) and calibration curve (B) of multivariable analysis model.

Full-size  DOI: [10.7717/peerj.13365/fig-2](https://doi.org/10.7717/peerj.13365/fig-2)

overweight or obese, and 44% had central obesity. Many of these female migrants did not adhere to the recommended level of fruit or vegetable intake and physical activity. We also found that having rural medical insurance, physical activity less than three days per week, being overweight or obese, and being aware of having HBP were associated with the higher odds of HBP. The validation tests showed that the model's results had good goodness of fit. The results have implications for the public health management of migrant women. The uncontrolled HBP among young Chinese female migrant workers suggested that this specific population needs health education promotion and specific programs.

In our study, the prevalence of HBP (27.2%) and hypertension (28.9%) according to the 2017 ACC/AHA guidelines were higher than both the recent 2017 prevalence report of 15.6% in Hunan province of Central South China (Wang et al., 2018) and the prevalence of 13.5% among Chinese women at reproductive age (Wang et al., 2017) based on the JNC-8 guidelines (SBP \geq 140 mmHg or DBP \geq 90 mmHg or use of antihypertensive medication) (James et al., 2014). Since an SBP of 130–139 mmHg or a DBP of 80–89 mmHg was included in the stage I diagnosis for hypertension according to the new 2017 ACC/AHA criteria, an additional 44 women (19.0% of the migrant workers) were classified as hypertensive. Therefore, the proportion of Chinese female migrant workers aged between 18 and 45 years classified as having hypertension would increase from 8.2% based on the JNC-8 guidelines to 27.2% based on the 2017 ACC/AHA guidelines, representing a remarkably higher prevalence of hypertension compared with the general population. Of note, comparison of prevalence studies should be interpreted with caution because of the differences in blood pressure thresholds over the years and across different guidelines from different countries.

Whether the new ACC/AHA standard is applicable for assessing the prevalence of hypertension in China is highly controversial. Several cardiovascular experts argued that the strategy of hypertension management must combine scientific evidence with the current socioeconomic capacity because of its high incidence and differences in each country or

region based on blood pressure target achievement. Meanwhile, some researchers advocated for implementing early management of hypertension in China to address the increasing prevalence of hypertension in younger adults. *Li et al. (2019)* found that Chinese young adults (25–34 years) accounted for the highest proportion of newly diagnosed individuals with HBP (130–40 mmHg/80–90 mmHg according to the 2017 ACC/AHA hypertension guidelines). To reduce the national health burden of HBP-related diseases in the long term, the significantly high prevalence among young migrant female workers in our study should be considered a public health issue that should be further investigated.

Our results also showed that only a minority (11.1%) of the female migrant workers with current HBP were aware of their condition. The awareness level of currently or formerly having HBP in our subjects was far lower than among migrant workers reported by *Li et al. (2017)* (5.6% vs. 44.3%). This difference might be due to the general traits of the participants in the current study, who were younger and had lower levels of education (secondary school and above). Other studies showed that a disadvantaged socioeconomic status, including younger age, low level of education, having an agricultural or rural migrant occupation, and low income, were significantly associated with poor health literacy (*Wang et al., 2015; Xie et al., 2019*). Poor health literacy might influence an individual's ability to understand hypertension and lead to disparities in access to hypertension care (*Ahuja et al., 2018*). Another explanation for this high unawareness of having HBP might be the disparity in utilization of public health resources because of economic or insurance limitations. A low socioeconomic status was more associated with HBP in women than men (*Leng et al., 2015*). In our study, most of the migrant female workers had a monthly income of \leq USD 420, and nearly 14.2% had no insurance; however, 98.4% of those migrant women diagnosed with hypertension also had rural insurance. The rural forms of health insurance are distinct from those in the urban systems in China and are not transferable to the urban insurance system and/or may not be reimbursed in the same amount in the rural region. Thus, the possibility of willingness to receive medical treatment for these migrant workers in their new living environment is limited by their economic status and because their health insurance has higher rates of reimbursement for care received in rural and town hospitals and lower rates of reimbursement in city hospitals. Some migrants have to go back to rural areas for cheaper medical services. It is probably why having rural insurance had the largest OR for HBP. Theoretically, the direct effect of the medical insurance system is to reduce the medical burden of the insured so that they can get timely treatment for illnesses. Therefore, once people with medical insurance have health-related problems, they are less likely to be affected by economic factors. On the other hand, people without medical insurance tend not to seek medical treatment even if they know they have health conditions because their economic situation is not good enough to cope with the follow-up and treatment costs. But some migrant workers choose to buy medicines by themselves, considering the low cost of medical treatment and the low reimbursement rate of the whole, and the extra cost of transportation and loss of work when they return to the countryside. According to a Chinese survey, about 24.3% of migrant workers self-medicated, and 29.0% did nothing when they developed symptoms in the past month, which might be because the rural and urban insurance systems are not fully connected (*Shao et al., 2018*). Therefore,

eliminating the regional disparities in rural and urban health insurance for female migrants and enhancing the health service availability will probably benefit the prevention and management of hypertension. But it also means increasing out-of-pocket costs for migrant workers.

Our study found that general obesity (determined by BMI) and a low level of physical activity but not central obesity (determined by WC) were independently associated with HBP. Epidemiological studies have demonstrated that obesity is one of the most important factors associated with hypertension (*Nurdiantami et al., 2018; Susic & Varagic, 2017*). Moreover, women with obesity are seven times more likely to develop hypertension than their normal-weight counterparts (*Aucott et al., 2005*). The high prevalence of obesity might result from limited adherence to the recommended dietary practices and physical activity levels. Rapid urbanization in China has contributed to lifestyle changes, including unhealthy dietary habits (intake of more refined foods and sugary drinks) and reduced physical activity, which have led to the increased prevalence of general and central obesity and the development of hypertension (*Schwingshackl et al., 2017; Su, Sun & Xu, 2018; Yan et al., 2016*). Observational data indicate that maintaining a minimum of 150 min of moderate physical activity per week for adults aged ≥ 18 years would potentially prevent 46% of inactivity-related deaths (*Mok et al., 2019*). The proportion of female migrant workers who met the recommended level of physical activity was only 18.5%, which was lower than the average national rate of 22.8% (*Tian et al., 2016*). According to a 20-year cohort study in China, physical activity among adults has decreased by half from 1991 to 2011, with much steeper declines in women than in men. Each new generation had lower activity levels than the previous one (*Zang & Ng, 2016*). Strong evidence has demonstrated that physical activity could reduce blood pressure among adults with prehypertension and hypertension (*Pescatello et al., 2019*). Additionally, physical inactivity independently contributes to the development of CVD in women (*Li et al., 2006*). These findings highlight the importance of developing a healthy lifestyle program to increase awareness and reduce obesity and hypertension among female migrant workers.

Many studies explored the increases in hypertension prevalence by age (*Bundy & He, 2016; Gao et al., 2013; Hu, Gao & Liu, 2019; Zhou, 2002*). The prevalence of HBP has been recently reported to be high in young Chinese people. The prevalence of hypertension in younger adults (20–44 years) doubled from 2002 to 2010 (12% vs. 23% in men, 6% vs. 14% in women), which was much greater than the growth rate in older adults (29% vs. 46% in aged 45 years or above) (*Bundy & He, 2016*). The mean age of Chinese migrant workers was 39 years, and 53.9% of them were aged 40 and below. Furthermore, women accounted for 34.5% of Chinese migrant workers (*Department of Service and Management of Migrant Population NHaFPCoC, 2017*). The study by Guan in 2009 showed that about 25.0% of Chinese migrants aged at least 18 years had prehypertension (19.9%) or hypertension (5.1%) (*Guan, 2018*). In addition, the prevalence of hypertension rapidly increased to 16.3% in 2012 among Chinese migrant workers aged 18–59 years (*Bi et al., 2016*). *Li et al. (2017)* found that in 2015, the prevalence of hypertension was similar between migrants and urban residents (17.3% vs. 18.1%). Although many female migrant workers are more likely to be mainly engaged in manual labor, have a low level of education, and

lack access to healthcare services (Zheng *et al.*, 2018), the studies on key health issues, including hypertension and CVD, are limited in this population (Fang *et al.*, 2017; Yang *et al.*, 2015b; Zhang *et al.*, 2018). We also found that age ≥ 35 years was associated with HBP in the bivariable analysis but not in the multivariable one. Jin *et al.* reported that the prevalence of HTN in the migrant population aged <35 years was lower than that in the older population aged ≥ 35 years (5.2% vs. 28.4%) in Hunan in 2012 (Donghui *et al.*, 2015). Similar conclusions were also reported in other studies in China (Liu *et al.*, 2017; Zhang *et al.*, 2013). With increasing age, the stiffness of the aorta and arterial walls increases, which contributes to the high prevalence of hypertension in the older group (Sabbahi *et al.*, 2016; Yoo *et al.*, 2018). Although aging cannot be reversed, increasing the awareness of hypertension and promoting a healthy lifestyle among female migrant workers may be an effective way to delay the development of this condition, especially for those aged older than 35 years.

The bivariable analysis suggested that the difference was not significant in terms of the odds of HBP in female migrant workers whose parents did not have (or were uncertain of having) a history of CVD and those whose parents had CVD, but the association was significant in the multivariable analysis. Indeed, increased awareness of family history might increase an individual's awareness of the disease. Future studies should explore other potential mechanisms for hypertension among this population. Of note, it is common in medical research to include in a multivariable analysis only the variables that were significant in the bivariable analyses. Still, this practice can be misleading because some of the association of specific variables with the outcome might become significant only when considering the other variables. Indeed, such a circumstance can occur in the presence of groups with different sizes, missing data, an extreme within-group variation, and the presence of interaction (Lo *et al.*, 1995). Although the present study was not designed to determine causality, this association was significant in the multivariable analysis but not in the bivariable one. Findings regarding this association are hypothesis-generating and should be explored in future studies.

Although being aware of having HBP, blood pressure control was unsatisfactory among young female migrants, with a control rate of 46.2% (6/13). Public health workers should pay more attention to young female migrants, educate them about the increasing prevalence of HBP and the optimal goal for blood pressure control, and implement measures to reduce their risk factors for active prevention of hypertension. Improving knowledge and promoting a healthy lifestyle are crucial to alleviating the growing hypertension burden in the female migrant population. There were many female workers with obesity and HBP. This phenomenon reveals that prevention strategies are still needed for female workers. The awareness of whether one and their parents have HBP needs to be improved, which has a certain relationship with the health education level of female workers. It is necessary to strengthen health literacy interventions related to the prevention and treatment of chronic diseases.

Limitations

This is one of the first studies to examine the prevalence, awareness, and factors associated with HBP in young female migrant workers in Central South China. However, the study also has some limitations. First, a causal relationship could not be established due to the cross-sectional design. Second, our study included only female migrant workers aged between 18 and 45 years, thus limiting the generalizability of our results to other populations. In addition, the present study did not conduct surveys on female workers in every category of occupation. Still, it adopted the principle of convenient sampling and volunteers from large, medium, and small enterprises with $\geq 50\%$ of women, which certainly introduced selection biases. Third, family and personal health history were based on self-reports, introducing recall biases. HBP medications were also based on the participant's response (yes or no) to a question, without asking the types of medications or confirming using the participants' pill bottles from home. Fourth, for accurate measurement of blood pressure, the 2017 AHA guideline suggested using an average of ≥ 2 readings obtained on ≥ 2 occasions to estimate the individual's level of BP (*Whelton et al., 2018*). The failure to measure blood pressure on another occasion was a limitation in our study. Fifth, this study was not designed to take into account white coat HBP (which could overestimate the prevalence of HBP) or masked hypertension (which could underestimate the prevalence of HBP). Finally, and most importantly, the patients were divided based on having HBP or not. Therefore, some patients who self-reported hypertension history with controlled blood pressure were grouped as non-HBP. Nevertheless, the results indicated that only 6% of the hypertensive patients had controlled BP.

CONCLUSIONS AND IMPLICATIONS FOR PRACTICE

This study revealed a high prevalence of HBP among young Chinese female migrant workers, as well as a high frequency of being unaware of their condition and some associated factors (rural medical insurance, awareness of having HBP, physical inactivity, and overweight/obesity). The uncontrolled HBP among young Chinese female migrant workers suggested that health education needs further promotion in such a population.

Abbreviations

| | |
|--------------|---------------------------------|
| CVD | cardiovascular disease |
| JNC-8 | Eighth Joint National Committee |
| SBP | systolic blood pressure |
| DBP | diastolic blood pressure |
| HBP | having high blood pressure |
| ACC | American College of Cardiology |
| AHA | American Heart Association |
| CI | confidence interval |
| WC | waist circumference |
| HTN | hypertension |
| SDs | standard deviations |
| OR | odds ratio |

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by the grant from TT and WF Chao Presidential Chair in Global Health Nursing Research Grant, Hunan Social Science Research Fund 18YBA440 and Hunan Key Nursing Laboratory Program 2017TP1004. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

TT and WF Chao Presidential Chair in Global Health Nursing Research Grant.

Hunan Social Science Research Fund: 18YBA440.

Hunan Key Nursing Laboratory Program: 2017TP1004.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Hua Peng analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, contributed reagents/materials/analysis tools, and approved the final draft.
- Mei Sun and Lisa Lommel conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Xin Hu, Huiwu Han and Emin Peng performed the experiments, prepared figures and/or tables, and approved the final draft.
- Jing Su performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- James Wiley analyzed the data, authored or reviewed drafts of the paper, contributed reagents/materials/analysis tools, and approved the final draft.
- Jyu-Lin Chen conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Human Ethics

The following information was supplied relating to ethical approvals (*i.e.*, approving body and any reference numbers):

The Institutional Review Board of the University of California, San Francisco, and Central South University reviewed and approved the study protocol (ID: 2017036).

Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplementary File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.13365#supplemental-information>.

REFERENCES

- Ahuja R, Ayala C, Tong X, Wall HK, Fang J. 2018.** Public awareness of health-related risks from uncontrolled hypertension. *Preventing Chronic Disease* 15:E40 DOI 10.5888/pcd15.170362.
- Aucott L, Poobalan A, Smith WC, Avenell A, Jung R, Broom J. 2005.** Effects of weight loss in overweight/obese individuals and long-term hypertension outcomes: a systematic review. *Hypertension* 45:1035–1041 DOI 10.1161/01.HYP.0000165680.59733.d4.
- Bi Y, Wang L, Xu Y, Jiang Y, He J, Zhang M, Zhou M, Li Y, Xu M, Lu J, Wang T, Ding L, Dai M, Zhang D, Chen CS, Lai S, Wang W, Wang L, Ning G. 2016.** Diabetes-related metabolic risk factors in internal migrant workers in China: a national surveillance study. *Lancet Diabetes Endocrinology* 4:125–135 DOI 10.1016/s2213-8587(15)00366-6.
- Bundy JD, He J. 2016.** Hypertension and related cardiovascular disease burden in China. *Annals of Global Health* 82:227–233 DOI 10.1016/j.aogh.2016.02.002.
- Department of Service and Management of Migrant Population NHaFPCoC. 2017.** Report on China's migrant population development 2016. Beijing: China Population Publishing House (in Chinese).
- Donghui J, Qiaohua X, Huilin L, Jiawu L, Biyun C. 2015.** Management of hypertension and diabetes mellitus among floating population in Hunan. *Practical Preventive Medicine* 22:1211–1212.
- Du P, Wang HJ, Zhang B, Qi SF, Mi YJ, Liu DW, Tian QB. 2017.** Prevalence of abdominal obesity among Chinese adults in 2011. *Journal of Epidemiology* 27:282–286 DOI 10.1016/j.je.2017.03.001.
- Etminan M, Collins GS, Mansournia MA. 2020.** Using causal diagrams to improve the design and interpretation of medical research. *Chest* 158:S21–S28 DOI 10.1016/j.chest.2020.03.011.
- Fang H, Jin Y, Zhao M, Zhang H, AR J, Zhang D, Hou Z. 2017.** Does migration limit the effect of health insurance on hypertension management in China? *International Journal of Environmental Research and Public Health* 14:1256 DOI 10.3390/ijerph14101256.
- Gao Y, Chen G, Tian H, Lin L, Lu J, Weng J, Jia W, Ji L, Xiao J, Zhou Z, Ran X, Ren Y, Chen T, Yang W. 2013.** Prevalence of hypertension in china: a cross-sectional study. *PLOS ONE* 8:e65938 DOI 10.1371/journal.pone.0065938.
- Guan M. 2018.** Epidemiology of hypertensive state among Chinese migrants: effects of unaffordable medical care. *International Journal of Hypertension* 2018:5231048 DOI 10.1155/2018/5231048.
- Gueugnon M, Stapley PJ, Gouteron A, Lecland C, Morisset C, Casillas JM, Ornetti P, Laroche D. 2019.** Age-related adaptations of lower limb intersegmental coordination during walking. *Frontiers in Bioengineering and Biotechnology* 7:173 DOI 10.3389/fbioe.2019.00173.

- Guo J, Shi Z, Chen JL, Dixon JK, Wiley J, Parry M. 2018.** Translation and validation of the Canadian diabetes risk assessment questionnaire in China. *Public Health Nursing* 35:18–28 DOI [10.1111/phn.12380](https://doi.org/10.1111/phn.12380).
- Hajian-Tilaki K. 2013.** Receiver Operating Characteristic (ROC) curve analysis for medical diagnostic test evaluation. *Caspian Journal of Internal Medicine* 4:627–635.
- Hunan Provincial Bureau of Statistics. 2018.** *Hunan province 2018 national economic and social development statistical bulletin*. Changsha: Hunan Provincial Bureau of Statistics Available at <http://tjj.hunan.gov.cn/hntj/English/hs/index.html>.
- Hu S, Gao R, Liu L. 2019.** Summary of the 2018 report on cardiovascular diseases in China. *Chinese Circulation Journal* 34:209–220.
- James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Oggedegbe O, Smith Jr SC, Svetkey LP, Taler SJ, Townsend RR, Wright Jr JT, Narva AS, Ortiz E. 2014.** 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 311:507–520 DOI [10.1001/jama.2013.284427](https://doi.org/10.1001/jama.2013.284427).
- Jin MJ, Chen BB, Mao YY, Zhu YM, Yu YX, Wu YY, Zhang MW, Zhu SK, Chen K. 2013.** Prevalence of overweight and obesity and their associations with socioeconomic status in a rural Han Chinese adult population. *PLOS ONE* 8:e79946 DOI [10.1371/journal.pone.0079946](https://doi.org/10.1371/journal.pone.0079946).
- Juraschek SP, Miller 3rd ER, Weaver CM, Appel LJ. 2017.** Effects of sodium reduction and the DASH Diet in relation to baseline blood pressure. *Journal of the American College of Cardiology* 70:2841–2848 DOI [10.1016/j.jacc.2017.10.011](https://doi.org/10.1016/j.jacc.2017.10.011).
- Kim JH. 2019.** Multicollinearity and misleading statistical results. *Korean Journal of Anesthesiology* 72:558–569 DOI [10.4097/kja.19087](https://doi.org/10.4097/kja.19087).
- Leng B, Jin Y, Li G, Chen L, Jin N. 2015.** Socioeconomic status and hypertension: a meta-analysis. *Journal of Hypertension* 33:221–229 DOI [10.1097/hjh.0000000000000428](https://doi.org/10.1097/hjh.0000000000000428).
- Li D, Zeng X, Huang Y, Lei H, Li G, Zhang N, Huang W. 2019.** Increased risk of hypertension in young adults in Southwest China: impact of the 2017 ACC/AHA high blood pressure guideline. *Current Hypertension Reports* 21:21 DOI [10.1007/s11906-019-0926-y](https://doi.org/10.1007/s11906-019-0926-y).
- Li H, Yan X, Deng X, Yang L, Zhao S, Zou J, Luo Y, Cao S. 2017.** A cross-sectional study of the prevalence, awareness, treatment and control of hypertension in Shenzhen, China. *BMJ Open* 7:e015206 DOI [10.1136/bmjopen-2016-015206](https://doi.org/10.1136/bmjopen-2016-015206).
- Li TY, Rana JS, Manson JE, Willett WC, Stampfer MJ, Colditz GA, Rexrode KM, Hu FB. 2006.** Obesity as compared with physical activity in predicting risk of coronary heart disease in women. *Circulation* 113:499–506 DOI [10.1161/circulationaha.105.574087](https://doi.org/10.1161/circulationaha.105.574087).
- Liu X, Gu W, Li Z, Lei H, Li G, Huang W. 2017.** Hypertension prevalence, awareness, treatment, control, and associated factors in Southwest China: an update. *Journal of Hypertension* 35:637–644 DOI [10.1097/hjh.0000000000001203](https://doi.org/10.1097/hjh.0000000000001203).
- Lo SK, Li IT, Tsou TS, See L. 1995.** Non-significant in univariate but significant in multivariate analysis: a discussion with examples. *Changgeng Yi Xue Za Zhi* 18:95–101.

- Lu J, Lu Y, Wang X, Li X, Linderman GC, Wu C, Cheng X, Mu L, Zhang H, Liu J, Su M, Zhao H, Spatz ES, Spertus JA, Masoudi FA, Krumholz HM, Jiang L. 2017. Prevalence, awareness, treatment, and control of hypertension in China: data from 1.7 million adults in a population-based screening study (China PEACE Million Persons Project). *Lancet* 390:2549–2558 DOI 10.1016/s0140-6736(17)32478-9.
- Marcoulides KM, Raykov T. 2019. Evaluation of variance inflation factors in regression models using latent variable modeling methods. *Educational and Psychological Measurement* 79:874–882 DOI 10.1177/0013164418817803.
- Masterson Creber RM, Smeeth L, Gilman RH, Miranda JJ. 2010. Physical activity and cardiovascular risk factors among rural and urban groups and rural-to-urban migrants in Peru: a cross-sectional study. *Revista Panamericana de Salud Pública* 28:1–8 DOI 10.1590/s1020-49892010000700001.
- Mok A, Khaw KT, Luben R, Wareham N, Brage S. 2019. Physical activity trajectories and mortality: population based cohort study. *BMJ* 365:l2323 DOI 10.1136/bmj.l2323.
- National Bureau of Statistics of the People's Republic of China. 2016. The 2015 report on the surveillance of China's rural-to-urban migrant workers. Available at http://www.stats.gov.cn/tjsj/zxfb/201604/t20160428_1349713.html (accessed on 8 October 2016).
- National Bureau of Statistics of the People's Republic of China. 2017. 2016 Survey report of migrant workers. Beijing, China: National Bureau of Statistics of the People's Republic of China (in Chinese).
- National Bureau of Statistics of China. 2020. *China Statistical Yearbook 2019*. Beijing: National Bureau of Statistics of China. Available at <http://www.stats.gov.cn/tjsj/ndsj/2019/indexeh.htm>.
- Nurdiantami Y, Watanabe K, Tanaka E, Pradono J, Anme T. 2018. Association of general and central obesity with hypertension. *Clinical Nutrition* 37:1259–1263 DOI 10.1016/j.clnu.2017.05.012.
- Pescatello LS, Buchner DM, Jakicic JM, Powell KE, Kraus WE, Bloodgood B, Campbell WW, Dietz S, Dipietro L, George SM, Macko RF, McTiernan A, Pate RR, Piercy KL. 2019. Physical activity to prevent and treat hypertension: a systematic review. *Medicine & Science in Sports & Exercise* 51:1314–1323 DOI 10.1249/mss.0000000000001943.
- Roth GA, Mensah GA, Johnson CO, Addolorato G, Ammirati E, Baddour LM, Barengo NC, Beaton AZ, Benjamin EJ, Benziger CP, Bonny A, Brauer M, Brodmann M, Cahill TJ, Carapetis J, Catapano AL, Chugh SS, Cooper LT, Coresh J, Criqui M, De Cleene N, Eagle KA, Emmons-Bell S, Feigin VL, Fernandez-Sola J, Fowkes G, Gakidou E, Grundy SM, He FJ, Howard G, Hu F, Inker L, Karthikeyan G, Kassebaum N, Koroshetz W, Lavie C, Lloyd-Jones D, Lu HS, Mirijello A, Temesgen AM, Mokdad A, Moran AE, Muntner P, Narula J, Neal B, Ntsekhe M, Moraes de Oliveira G, Otto C, Owolabi M, Pratt M, Rajagopalan S, Reitsma M, Ribeiro ALP, Rigotti N, Rodgers A, Sable C, Shakil S, Sliwa-Hahnle K, Stark B, Sundstrom J, Timpel P, Tleyjeh IM, Valgimigli M, Vos T, Whelton PK, Yacoub M, Zuhlke L, Murray C, Fuster V, Group G-N-JGBoCDW. 2020. Global burden of cardiovascular

- diseases and risk factors, 1990–2019: update From the GBD. *Journal of the American College of Cardiology* 76:2982–3021 DOI 10.1016/j.jacc.2020.11.010.
- Sabbahi A, Arena R, Elokda A, Phillips SA. 2016. Exercise and hypertension: uncovering the mechanisms of vascular control. *Progress in Cardiovascular Diseases* 59:226–234 DOI 10.1016/j.pcad.2016.09.006.
- Sarki AM, Nduka CU, Stranges S, Kandala NB, Uthman OA. 2015. Prevalence of hypertension in low- and middle-income countries: a systematic review and meta-analysis. *Medicine* 94:e1959 DOI 10.1097/md.0000000000001959.
- Schwingshackl L, Schwedhelm C, Hoffmann G, Knüppel S, Iqbal K, Andriolo V, Bechthold A, Schlesinger S, Boeing H. 2017. Food groups and risk of hypertension: a systematic review and dose-response meta-analysis of prospective studies. *Advances in Nutrition* 8:793–803 DOI 10.3945/an.117.017178.
- Shao S, Wang M, Jin G, Zhao Y, Lu X, Du J. 2018. Analysis of health service utilization of migrants in Beijing using Anderson health service utilization model. *BMC Health Services Research* 18:462 DOI 10.1186/s12913-018-3271-y.
- Su L, Sun L, Xu L. 2018. Review on the prevalence, risk factors and disease management of hypertension among floating population in China during 1990–2016. *Global Health Research and Policy* 3:24 DOI 10.1186/s41256-018-0076-9.
- Susic D, Varagic J. 2017. Obesity: a perspective from hypertension. *Medical Clinics of North America* 101:139–157 DOI 10.1016/j.mcna.2016.08.008.
- Tian Y, Jiang C, Wang M, Cai R, Zhang Y, He Z, Wang H, Wu D, Wang F, Liu X, He Z, An P, Wang M, Tang Q, Yang Y, Zhao J, Lv S, Zhou W, Yu B, Lan J, Yang X, Zhang L, Tian H, Gu Z, Song Y, Huang T, McNaughton LR. 2016. BMI, leisure-time physical activity, and physical fitness in adults in China: results from a series of national surveys, 2000–14. *Lancet Diabetes Endocrinology* 4:487–497 DOI 10.1016/s2213-8587(16)00081-4.
- Van Kleef ME, Spiering W. 2017. Hypertension: overly important but under-controlled. *European Journal of Preventive Cardiology* 24:36–43 DOI 10.1177/2047487317709116.
- Wajngarten M, Silva GS. 2019. Hypertension and stroke: update on treatment. *European Cardiology* 14:111–115 DOI 10.15420/ecr.2019.11.1.
- Wang JG. 2015. Chinese hypertension guidelines. *Pulse* 3:14–20 DOI 10.1159/000382025.
- Wang Z, Chen Z, Zhang L, Wang X, Hao G, Zhang Z, Shao L, Tian Y, Dong Y, Zheng C, Wang J, Zhu M, Weintraub WS, Gao R. 2018. Status of hypertension in China: results from the china hypertension survey, 2012–2015. *Circulation* 137:2344–2356 DOI 10.1161/circulationaha.117.032380.
- Wang X, Guo H, Wang L, Li X, Huang M, Liu Z, Liu X, Wang K, Alamian A, Anderson JL. 2015. Investigation of residents' health literacy status and its risk factors in Jiangsu Province of China. *Asia-Pacific Journal of Public Health* 27:Np:2764–2772 DOI 10.1177/1010539513487012.
- Wang ZQ, Zhao YF, Yang J, Wang LM, Zhao ZP, Zeng XY, Wang LH. 2017. Rate of prevalence, awareness, treatment and control of hypertension among women at reproductive age in China in 2013. *Zhonghua Yu Fang Yi Xue Za Zhi* 51:1086–1090 DOI 10.3760/cma.j.issn.0253-9624.2017.12.007.

- Whelton PK, Carey RM, Aronow WS, Casey Jr DE, Collins KJ, Himmelfarb CD, Nelson, Palma SM, De, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbigele B, Smith Jr SC, Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Sr KA, Williams, Williamson JD, Wright Jr JT. 2018. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension* 71:e13-e115 DOI 10.1161/hyp.0000000000000065.
- Wu CY, Hu HY, Chou YJ, Huang N, Chou YC, Li CP. 2015. High blood pressure and all-cause and cardiovascular disease mortalities in community-dwelling older adults. *Medicine* 94:e2160 DOI 10.1097/MD.00000000000002160.
- Xie Y, Ma M, Zhang Y, Tan X. 2019. Factors associated with health literacy in rural areas of Central China: structural equation model. *BMC Health Services Research* 19:300 DOI 10.1186/s12913-019-4094-1.
- Xu W, Zhang H, Paillard-Borg S, Zhu H, Qi X, Rizzuto D. 2016. Prevalence of overweight and obesity among Chinese adults: role of adiposity indicators and age. *Obesity Facts* 9:17–28 DOI 10.1159/000443003.
- Yan S, Hong X, Yu H, Yang Z, Liu S, Quan W, Xu J, Zhu L, Cheng W, Xiao H, Kitzman-Ulrich H, De Haven MJ. 2016. Prevalence of diabetes and health-related quality of life among rural-to-urban Nong Zhuan Fei migrants in an urban area of Northern China, 2013. *Public Health Reports* 131:167–176 DOI 10.1177/003335491613100124.
- Yang H, He F, Wang T, Liu Y, Shen Y, Gong J, Dai W, Zhou J, Gu J, Tu Y, Wang T, Shen L, Wu Y, Xia X, Xu D, Pan Z, Zhu S. 2015a. Health-related lifestyle behaviors among male and female rural-to-urban migrant workers in Shanghai, China. *PLOS ONE* 10:e0117946 DOI 10.1371/journal.pone.0117946.
- Yang W, Li H, Fu X, Lu J, Xue Z, Wu C. 2015b. Inequalities in cardiovascular health between local and migrant residents: a cross-sectional study of 6934 participants in Longhua District, Shenzhen. *Medicine* 94:e2103 DOI 10.1097/md.00000000000002103.
- Yang Y, Ma CW, Yang Y, Wang X, Lin X, Fu L, Wang S, Yang Z, Wang Z, Meng X, Ma D, Ma R, Ma J. 2017. Effects of integrated health management intervention on overweight and obesity. *Evidence-Based Complementary and Alternative Medicine* 2017:1239404 DOI 10.1155/2017/1239404.
- Ying D, Zeng J, Ji K, Liu X, Wu X. 2015. Analysis of hypertension morbidity among floating population of Sichuan in 2012. *Chinese Journal of Prevention and Control of Chronic Diseases* 23:27–30.
- Yoo JK, Okada Y, Best SA, Parker RS, Hieda M, Levine BD, Fu Q. 2018. Left ventricular remodeling and arterial afterload in older women with uncontrolled and controlled hypertension. *Menopause* 25:554–562 DOI 10.1097/gme.0000000000001046.
- Zang J, Ng SW. 2016. Age, period and cohort effects on adult physical activity levels from 1991 to 2011 in China. *International Journal of Behavioral Nutrition and Physical Activity* 13:40 DOI 10.1186/s12966-016-0364-z.

- Zhang YR, Ge PF, Liao YJ, Guo LL, Zhang CJ, Dong CX, Fan HQ, Zhang XR. 2013.** The tendency of prevalence of hypertension in two countries in Gansu province from 1990 to 2012. *Zhonghua Yu Fang Yi Xue Za Zhi* **47**:306–311.
- Zhang J, Wan S, Zhang B, Dong F, Pan L, Yihuo W, Gong H, Yang F, Xu G, Li Z, Li G, Li Y, Wang X, Shan G. 2018.** Twenty-year time trends in hypertension prevalence in Yi people of China: three successive cross-sectional studies, 1996-2015. *BMJ Open* **8**:e022714 DOI [10.1136/bmjopen-2018-022714](https://doi.org/10.1136/bmjopen-2018-022714).
- Zheng L, Hu R, Dong Z, Hao Y. 2018.** Comparing the needs and utilization of health services between urban residents and rural-to-urban migrants in China from 2012 to 2016. *BMC Health Services Research* **18**:717 DOI [10.1186/s12913-018-3522-y](https://doi.org/10.1186/s12913-018-3522-y).
- Zhou B. 2002.** Prospective study for cut-off points of body mass index in Chinese adults. *Zhonghua Liu Xing Bing Xue Za Zhi* **23**:431–434.
- Zhou B, Wu Y, Yang J, Li Y, Zhang H, Zhao L. 2002.** Overweight is an independent risk factor for cardiovascular disease in Chinese populations. *Obesity Reviews* **3**:147–156 DOI [10.1046/j.1467-789x.2002.00068.x](https://doi.org/10.1046/j.1467-789x.2002.00068.x).
- Zou KH, O'Malley AJ, Mauri L. 2007.** Receiver-operating characteristic analysis for evaluating diagnostic tests and predictive models. *Circulation* **115**:654–657 DOI [10.1161/CIRCULATIONAHA.105.594929](https://doi.org/10.1161/CIRCULATIONAHA.105.594929).