

A Profile of Avelumab Plus Axitinib in the Treatment of Renal Cell Carcinoma

Manuela Tiako Meyo^{1,2}, Jeanne Chen^{1,2}, Francois Goldwasser^{1,2}, Laure Hirsch^{1,2}, Olivier Huillard^{1,2} 

¹Department of Medical Oncology, Institut du Cancer Paris CARPEM, AP-HP, APHP.Centre, Hôpital Cochin, Paris, France; ²Immunomodulatory Therapies Multidisciplinary Study Group (CERTIM), AP-HP, APHP.Centre, Hôpital Cochin, Paris, France

Correspondence: Olivier Huillard, Department of Medical Oncology, Institut du Cancer Paris CARPEM, AP-HP, APHP.Centre, Hôpital Cochin, 123 Bd de Port Royal, Paris, 75014, France, Tel +33 | 58 41 14 39, Fax +33 | 58 41 14 34, Email olivier.huillard@aphp.fr

Abstract: Until recently, the approved first-line treatment for metastatic RCC (mRCC) consisted of tyrosine kinase inhibitors (TKI) targeting the vascular endothelial growth factor receptors (VEGFR) monotherapy. The landscape of first-line treatment has been transformed in the last few years with the advent of immune checkpoint inhibitors (ICI) or VEGFR TKI plus ICI combinations. This article focuses on the profile of one of these ICI plus VEGFR TKI combination, avelumab plus axitinib. We detail the characteristics of each drug separately, and then we explore the rationale for their association, its efficacy and the resulting toxicity. Finally, we examine the factors associated with avelumab plus axitinib outcomes, and their impact on therapeutic strategy.

Keywords: renal cell carcinoma, vascular endothelial growth factor, immune checkpoint inhibitor, axitinib, avelumab, pharmacology

Introduction

Renal cell carcinoma (RCC) is a frequent cancer representing 5% of the estimated new cases of cancer in males and 3% of the estimated new cases of cancer in females and is estimated to be responsible for 13,920 deaths in 2022 in the United States (US).¹ When diagnosed at a local stage, RCC can be treated with curative intent surgery. When diagnosed at an advanced or metastatic stage, active surveillance can be an option in the case of indolent growth of metastases, but disease will eventually progress in most cases requiring systemic therapy.^{2,3} Until recently, the approved first-line treatment for metastatic RCC (mRCC) consisted of tyrosine kinase inhibitors (TKI) targeting the vascular endothelial growth factor receptors (VEGFR)^{4,5} but the landscape of first-line treatment has been transformed in the last few years with the advent of immune checkpoint inhibitors (ICI) or VEGFR TKI plus ICI combinations.^{6–10} This article focuses on one of these associations, avelumab plus axitinib, detailing each drug separately, then exploring the rationale for their association, its efficacy and the resulting toxicity, and finally examining the factors associated with avelumab plus axitinib outcomes and their impact on therapeutic strategy.

Axitinib

We previously reviewed the design, development, and place in therapy of axitinib monotherapy in the treatment of renal cell carcinoma,¹¹ but some aspects must be emphasized in the context of its association with ICI.

Pharmacodynamics

Axitinib (Inlyta[®], Pfizer) is an indazole derivative produced by chemical synthesis, whose molecular weight is 386,47 Da. It is an oral second-generation TKI with an inhibitory concentration for the VEGF family receptors 10-fold lower than other TKI, but contrary to multiple TKIs, axitinib is weakly active in other receptors like KIT or PDGFR. Axitinib is therefore a potent and selective VEGFR inhibitor.¹²

Pharmacokinetics

Orally taken, axitinib is rapidly absorbed and achieves maximal plasma concentration after 4 hours. There is no clinically significant difference between the fasted and fed states,¹³ with a mean bioavailability of 58%.¹⁴ Axitinib absorption is pH-dependent, with higher absorption at acidic pH, but anti-acid agents have a limited impact on axitinib area under the curve and therefore are not contraindicated during axitinib treatment. At therapeutic dose axitinib has a high protein-binding rate exceeding 99%, and preferentially binds to albumin. Particular attention must therefore be paid to patient with hypoalbuminemia. At steady state, axitinib pharmacokinetic is approximately linear over a dosing range of 1 to 20 mg.¹⁵ Axitinib is mainly metabolized in the liver by CYP3A4/5 and to a lesser extent by CYP1A2, 2C19 and UGT1A1, producing pharmacologically inactive metabolites. Attention should therefore be paid when axitinib is prescribed with known CYP3A4/5 inducer or inhibitor. However, axitinib is not considered as a cytochrome inducer or inhibitor but is a substrate for the efflux transporter P-glycoprotein (P-gp) and for the hepatic organic anion-transporting polypeptides (OATP1B1). Axitinib has been shown to inhibit the efflux transporter P-gp in vitro but not at therapeutic plasma concentration. Axitinib is mainly eliminated in feces due to hepatobiliary excretion, whereas axitinib renal excretion accounts for less than 20%. Unchanged axitinib is detected only in feces and not in urine.¹⁶ A significant increase in axitinib plasma concentration has been observed in patients with moderate hepatic impairment (Child-Pugh B), but not for mild impairment (Child-Pugh A). Axitinib elimination remains constant during chronic dosing, without argument for auto-induction or auto-inhibition.¹⁵ Axitinib half-life is shorter than other TKI and varies from 2.5 to 6.1 hours, justifying its twice daily schedule of prescription. Due to this short half-life, a steady state is expected in less than 3 days, and the plasma concentration quickly decreases after treatment interruption. Axitinib, as most TKI, shows an important pharmacokinetic variability. Inter-individual variability for AUC at the 5 mg bid standard dose is estimated around 80%. This inter-individual variability is quite the same after intravenous or oral administration, suggesting a greater role of metabolism than absorption.¹⁵ Intra-individual variability is estimated around 20–22%.¹⁴

Pharmacokinetics-Pharmacodynamics Analysis (PK/PD)

An increase in blood pressure (BP) is a common on-target effect of VEGFR-TKIs. Thus, diastolic BP (dBP) as a biomarker of axitinib efficacy in solid tumors was investigated and increased dBP > 90 mm Hg was associated with axitinib efficacy.¹⁷ A PK-PD model was later developed, showing that dBP increased with increasing drug exposure.¹⁸ This relationship was not proportional, suggesting that an increase in dBP was not entirely explained by axitinib exposure, and thus that dBP should not be used exclusively to guide axitinib dosing. In a retrospective study, higher exposure and dBP were independently associated with longer PFS and overall survival (OS) and higher probability of partial response.¹⁹ Another analysis reported that patients receiving axitinib dose titration for mRCC had an increased axitinib exposure and a significantly higher objective response rate (ORR) than those with placebo titration (54% versus 34%; $P = 0.019$).²⁰ Another study demonstrated that axitinib exposure could be increased with individual dose titration, leading to a greater response rate.²¹ However, there was no improvement in PFS when axitinib plasma exposure increased, possibly due to toxicity from titration and the necessity of dose reduction after initial dose titration.

Axitinib Monotherapy Prescription

Axitinib is available as 1, 3, 5 and 7 mg coated tablets. The recommended starting dose is 5 mg twice daily administered with continuous daily dosing. Dose adjustments are recommended based on individual tolerability. For patients with good tolerance after two weeks (no adverse event > grade 2, and no increase in BP > 150/90 mm Hg or introduction of antihypertensive treatment), the dose should be increased to 7 mg twice daily. With the same criteria, a good tolerance of axitinib 7 mg bid may lead to an increase in the dose to 10 mg twice daily. Conversely, adverse reaction could require treatment interruption and reintroduction after dose reduction. No dose adjustment is recommended based on race,²² gender, weight,¹⁵ renal function²³ or drug-metabolizing enzymes genotype.²⁴ Axitinib clearance decreases modestly in subjects older than 60 years but these changes are not considered clinically significant and no dose adjustment is recommended according to age.¹⁵ No association has been identified between genetic polymorphisms in drug-metabolizing enzymes or transporters and axitinib pharmacokinetic variability.²⁴ As it has been described with the use of other VEGFR-TKI in mRCC patients, sarcopenia associated with a body mass index of less than 25 kg.m⁻² could help

identify patients at high risk of severe toxicity for whom particular attention may be needed at treatment initiation.^{25,26} For patients with moderate hepatic impairment (Child-Pugh B), a two-fold higher axitinib exposure has been observed. A decrease of half the dose is recommended for these patients.²⁷ Chronic kidney disease (CKD) and end-stage renal disease (ESRD) are not unusual in mRCC patients because of comorbidities and previous local treatments. However, there are few available data regarding the use of axitinib in patients with renal insufficiency, especially for patients with creatinine clearance of <15 mL/min. A population pharmacokinetic model found no difference according to basal renal function, in favor of no dose adjustment in patients with renal impairment.²³

Drug Interaction

Acids suppressing agents have a slight effect on axitinib overall exposure and thus are not contraindicated during axitinib treatment.¹⁶ CYP3A4/5 inhibitors²⁸ may increase axitinib plasma concentration²⁹ resulting in clinically significant effects.^{30,31} If an alternative treatment is not possible, a dose decrease of axitinib is recommended (approximately half the dose). CYP3A4/5 inducers²⁸ may decrease axitinib exposure.³² When CYP3A4/5 inducer cannot be discontinued, axitinib dose should be increased progressively with enhanced toxicity and therapeutic drug monitoring.

Clinical Development and Place in Therapy

Axitinib was approved by both American and European Agencies in 2012 for the treatment of advanced RCC after failure of one prior systemic therapy.³³ Table 1 summarizes the randomized Phase III clinical trials of axitinib alone or in combination with ICI in mRCC. The demonstration of clinical benefit for axitinib was based on a Phase III, randomized, open-label, multicenter study of axitinib compared with sorafenib in patients with advanced RCC after failure of a prior systemic first-line regimen.³⁴ In this study, 723 patients were assigned to receive axitinib (n = 361) or sorafenib (n = 362). The median PFS was 6.7 months with axitinib compared to 4.7 months with sorafenib (hazard ratio (HR) 0.665; 95% confidence interval (CI) 0.544–0.812; one-sided p < 0.0001). However, the updated median OS was 20.1 months (95% CI 16.7–23.4) with axitinib and 19.2 months (17.5–22.3) with sorafenib (HR 0.969, 95% CI 0.800–1.174; one-sided p = 0.3744). Importantly, axitinib dose increase to 7 mg and then to 10 mg, twice daily, was allowed for patients without hypertension or adverse reactions above grade 2 but dose increase was not allowed for patients receiving sorafenib. This difference could explain the difference in efficacy between axitinib and sorafenib.³⁵

Table 1 Randomized Phase III Trials of Avelumab and/or Axitinib in Renal Cell Carcinoma

Clinical Trials Number	Purpose	Patients Included	Results	Year First Published
NCT00678392	Axitinib versus sorafenib as second-line treatment for advanced renal cell carcinoma (AXIS)	723	Axitinib resulted in significantly longer progression-free survival compared with sorafenib.	2011
NCT00920816	Axitinib versus sorafenib as first-line therapy for advanced renal-cell carcinoma	288	Axitinib did not significantly increase progression-free survival compared to sorafenib.	2013
NCT01599754	Axitinib versus placebo as an adjuvant treatment of renal cell carcinoma (ATLAS)	724	The trial was stopped due to futility. There was no significant difference in disease-free survival.	2018
NCT02684006	Avelumab plus axitinib versus sunitinib in advanced renal cell carcinoma (Javelin Renal 101)	886	Treatment with avelumab plus axitinib resulted in a statistically significant improvement in progression-free survival versus sunitinib.	2019
NCT02853331	Pembrolizumab plus axitinib versus sunitinib in advanced renal cell carcinoma (KEYNOTE-426)	861	Treatment with pembrolizumab plus axitinib resulted in significantly longer overall survival and progression-free survival, as well as a higher objective response rate, than treatment with sunitinib.	2019

Plasma monitoring can help identify disease progression linked to sorafenib underexposure and lead to dose optimization to restore efficacy.^{36–39} Therefore sorafenib with plasma monitoring⁴⁰ may represent an alternative to axitinib.

Avelumab

Pharmacodynamics

Avelumab (Bavencio[®], Merck) is a fully human immunoglobulin G1 (IgG1) monoclonal antibody directed against programmed death ligand 1 (PD-L1). Avelumab binds PD-L1 and blocks the interaction between PD-L1 and the programmed death 1 (PD-1) and B7.1 receptors removing the suppressive effects of PD-L1 on cytotoxic CD8+ T-cells, resulting in the restoration of anti-tumor T-cell responses. Avelumab has also been shown to induce natural killer (NK) cell-mediated direct tumor cell lysis via antibody-dependent cell-mediated cytotoxicity (ADCC).⁴¹

Pharmacokinetics

Avelumab is expected to be distributed in the systemic circulation and to a lesser extent in the extracellular space. Consistent with a limited extravascular distribution, the volume of distribution of avelumab at steady state is small. As expected for an antibody, avelumab does not bind to plasma proteins in a specific manner.⁴¹

Steady-state concentrations of avelumab were reached after approximately 4 to 6 weeks of repeated dosing at 10 mg/kg every 2 weeks, and systemic accumulation was approximately 1.25-fold. The elimination half-life at the recommended dose is 6.1 days based on the population PK analysis.⁴¹

The exposure of avelumab increased dose-proportionally in the dose range from 10 mg/kg to 20 mg/kg every 2 weeks. The avelumab dose of 10 mg/kg every 2 weeks was selected based on adequate safety and tolerability, as well as PK and target occupancy data.⁴² No maximal tolerated dose was reached with doses up to 20 mg/kg. The mean target occupancy was 90% or higher at doses of 3 and 10 mg/kg. However, several patients treated at 3 mg/kg (but none treated at 10 mg/kg) had trough levels below 1 µg/mL, which in vitro studies suggested was the serum level required to ensure 90% target occupancy.⁴³ Population PK, exposure-efficacy, and exposure-safety models and simulations in Merkel cell carcinoma and urothelial carcinoma^{44,45} patient populations provided the basis for the subsequent approval of an avelumab 800-mg flat dose every two weeks.

A population PK analysis suggested no difference in the total systemic clearance of avelumab based on age, gender, race, PD-L1 status, tumour burden, renal impairment and mild or moderate hepatic impairment.⁴¹

Pharmacokinetics-Pharmacodynamics Analysis

Logistic regression analysis of data from 88 patients with Merkel cell carcinoma treated with avelumab 10 mg/kg every 2 weeks identified an apparent relationship between trough concentration at steady state and best objective response, with higher exposure associated with improved response contrasting with findings concerning other PD-(L)1 inhibitors.⁴³ Concerning exposure–safety relationship there was a weak association between increased exposure and increased incidence of immune-related adverse events, but none with the risk of treatment emergent adverse events or infusion-related reactions.⁴³

Avelumab Prescription

Avelumab was developed with infusions of 10 mg/kg every two weeks but was later modified to a flat dose of 800 mg infusions every two weeks.

No clinically important difference in the clearance of avelumab was found between patients with mild (glomerular filtration rate (GFR) 60 to 89 mL/min, Cockcroft-Gault Creatinine Clearance (CrCL); n = 623), moderate (GFR 30 to 59 mL/min, n = 320) and patients with normal (GFR ≥ 90 mL/min, n = 671) renal function.⁴¹

No clinically important difference in the clearance of avelumab was found between patients with mild hepatic impairment (bilirubin ≤ ULN and AST > ULN or bilirubin between 1 and 1.5 times ULN, n = 217) and normal hepatic function (bilirubin and AST ≤ ULN, n = 1388) in a population PK analysis.⁴¹

Drug Interaction

No formal drug–drug interaction studies were conducted for avelumab. However, because monoclonal antibodies are large molecules that are not commonly metabolized by CYP450 enzymes, the potential for co-administered small-molecule drugs (most of which are metabolized by CYP450) to alter the exposure of monoclonal antibodies by the route of metabolism and transport is low.⁴³ In particular, avelumab did not induce cytokines in humans at concentrations needed to affect either transporters involved in the distribution or CYP450 metabolism of small-molecule drugs.⁴³

Complementary Mechanisms of Action of Immune Checkpoint Inhibitors and Vascular Endothelial Growth Factor Pathway Inhibitors

The combination of ICI plus VEGFR inhibitor is a new first-line option for patients with advanced RCC.⁴⁶ This combination downregulates two major hallmarks of cancer: avoiding immune destruction and inducing angiogenesis.⁴⁷ The rationale for this combination relies on the interplay between the immune and angiogenic systems (Figure 1). Avelumab plus axitinib could have a synergistic effect similar to the other ICI plus VEGFR inhibitor combinations. Avelumab inhibits interactions between PD-L1 and programmed cell death protein (PD-1), leading to immunosuppression signal elimination and the restoration of anti-tumor T-cell functions in the tumor microenvironment. It is important to note that avelumab could also mediate antibody-dependent cell-mediated cytotoxicity (ADCC) lysis of tumor cells, which could be an additional method of action compared to other anti-PD-(L)1 antibody.^{42,48} Axitinib was selected for the association with avelumab because of a lower incidence of hepatic toxic effects compared to other VEGFR inhibitors previously tested.⁴⁹ The hypothesis of synergy over additive effects is based on two main concepts: the immunomodulatory roles of VEGF pathway inhibitors and the vascular-remodeling of immunotherapy.

VEGF pathway inhibitors regulate immune cells and their interaction with tumor microenvironment in two different ways: direct effects on immune cells and indirect effects through endothelial cells and hypoxia. Proangiogenic factors such as VEGF-A can have direct immunosuppressive effects on innate and adaptive immune cells. VEGF-A/VEGFR-1 and VEGF-A/VEGFR-2 signaling downregulate dendritic cells maturation and differentiation, respectively.⁵⁰ Mature dendritic cells play a critical role for the antigenic presentation and the induction of anti-tumor T-cell functions. Proangiogenic factors recruit tumor-associated macrophages, which can contribute to tumor immunosuppression.⁵¹ VEGF signaling also impacts adaptive immune cells, such as anti-tumor effector T cells and pro-tumor regulatory T cells. VEGF-A reduces proliferation and function and promotes exhaustion of anti-tumor effector T cells.^{52,53} Conversely, VEGF-A can favor regulatory T cells tumor infiltration and accumulation.⁵⁴ VEGF pathway inhibitors, including axitinib, could reverse direct immunosuppressive effects of VEGF. VEGF pathway inhibitors reduce tumor immunosuppressive cells, such as myeloid-derived suppressor cells (MDSC) and regulatory T cells and promote mature dendritic cells and effector T cells.⁵⁵ Some studies focused particularly on axitinib effects. In murine RCC xenografts model, axitinib treatment decreases MDSC in the spleens and tumor beds of animals and promotes anti-tumor CD8+




Figure 1 Complementary mechanisms of action of immune checkpoint inhibitors and Vascular endothelial growth factor pathway inhibitors.

Abbreviations: ADCC, antibody-dependent cell-mediated cytotoxicity; ICI, Immune checkpoint inhibitor; VEGFR, Vascular endothelial growth factor receptor.

T-cell functions.⁵⁶ Axitinib treatment also promotes natural killer cell recognition and degranulation in RCC cell lines model.⁵⁷ Natural killer cells play an important role in anti-tumor innate response and their degranulation triggers tumor cells destruction. In murine models, axitinib treatment reduces the number of immune-suppressive cells such as tumor-associated macrophages and improves T cell response.⁵⁸ VEGF pathway inhibitors also reverse immunosuppressive tumor microenvironment through an indirect effect on endothelial cells. VEGF pathway inhibitors can increase anti-tumor effector lymphocytes infiltration through normalization of vascularization.⁵⁹

Tumor angiogenesis favors immunosuppression through tumor endothelial cells and hypoxia. Proangiogenic factors affect several steps involved in tumor infiltration by circulating immune cells: homing of cells, adhesion to endothelium, and extravasation through endothelial intercellular junctions.⁵⁵ For example, VEGF-A in association with interleukin 10 and prostaglandin E2 induces Fas ligand expression on endothelial cells, which acquire the capacity to kill effector lymphocytes T cells, but not regulatory T cells.⁶⁰ Anarchic proliferation of abnormal tumor vessel can lead to hypoxic areas creation and disrupted immune cell infiltration. Hypoxia promotes pro-tumor macrophages, which exhibit pro-angiogenic activity and could suppress effector lymphocytes T cells proliferation.⁶¹ Tumor hypoxia also promotes the recruitment of regulatory T cell via the expression of the chemokine CC-chemokine ligand 28.⁶² VEGF pathway inhibitors could modulate tumor immunosuppression mediated by indirect effects through endothelial cells and hypoxia. Sunitinib, a VEGFR TKI such as axitinib, increases anti-tumor lymphocytes T cells infiltration through endothelial activation and up-regulation of T lymphocyte-attracting chemokines, CXCL10 and CXCL11.⁶³ Normalization of tumor vasculature related to VEGF pathway inhibitor effects can also promote immune-cell infiltration into tumors, but more studies should be conducted to determine the optimal window of tumor vascular normalization.⁶³ Indeed, VEGF pathway inhibitors can also damage tumor vessel and reduce immune cells and therapeutics infiltration in some cases. The dosing of antiangiogenic seems particularly important for the tumor vascular normalization and immune cells infiltration. For example, lower doses of anti-VEGFR2 antibody seems superior to the high doses in polarizing tumor macrophage from pro- toward anti-tumor phenotype and promoting infiltration of effector T cells.⁶⁴ Overall, VEGF pathway inhibitors, including axitinib, help to reverse the immunosuppressive tumor microenvironment, increase antitumor immunity and are thus good candidates for combination with ICI.

ICI efficacy relies mainly on the restoration of anti-tumor T-cell functions in the tumor microenvironment, but more and more studies show that immunotherapy might also improve the efficacy of VEGF pathway inhibitors through changes to the tumor vasculature. Interferon- γ activated endothelial cells can inhibit lymphocytes T cells activation via the expression of immune checkpoint molecules, such as PD-L1 or PD-L2.⁶⁵ Treatment with anti-PDL-1 inhibits immune checkpoint expressed on endothelial cell and promotes lymphocytes T cells activation but could also directly destroy tumor vasculature through ADCC, particularly in the case of avelumab.⁴⁸ Effector T cells activated by ICI secrete interferon- γ which has direct and indirect anti-vascular effects. Interferon- γ inhibits the expression of delta-like protein 4 on endothelial cells, which is necessary to Notch signaling and tumor angiogenesis.⁶⁶ Furthermore, interferon- γ down-regulates the mRNA of VEGF-A production.⁶⁶ Overall, synergy of ICI plus VEGF pathway inhibitor combination could be based on an immunostimulatory vascular-modulating cycle.⁶⁷ An important study illustrated this mutual regulation and showed an impact of vascular normalization disruption on effector T lymphocytes infiltration and an impact of T lymphocytes depletion on vascular normalization in tumor microenvironment.⁶⁸ Avelumab plus axitinib falls within the framework of this win–win relation.

Avelumab Plus Axitinib Combination in mRCC: The JAVELIN 101 Trial

In the phase III trial comparing avelumab plus axitinib to sunitinib in 886 advanced, treatment naïve, RCC patients, after a minimum follow-up of 13 months, PFS was significantly longer in the avelumab plus axitinib arm than in the sunitinib arm (PD-L1+ population: hazard ratio (HR) 0.62 [95% confidence interval (CI) 0.490–0.777]; one-sided $P < 0.0001$; median 13.8 (95% CI 10.1–20.7) versus 7.0 months (95% CI 5.7–9.6); overall population: HR 0.69 (95% CI 0.574–0.825); one-sided $P < 0.0001$; median 13.3 (95% CI 11.1–15.3) versus 8.0 months (95% CI 6.7–9.8)) but no overall survival benefit was demonstrated at the last data cut off, possibly reflecting immature data.^{10,69} In this phase III trial, avelumab 10 mg/kg was administered in combination with axitinib 5 mg and the respective exposures of avelumab and axitinib were unchanged compared to the single agents. There was no evidence to suggest a clinically relevant change in

avelumab clearance over time in patients with advanced RCC.⁴¹ Furthermore, analyses using PK and E-R modeling and simulation provide support for the labeling of an 800-mg flat dose of avelumab in combination with axitinib in patients with treatment-naïve RCC.⁷⁰ In the setting of the treatment combination, no titration of axitinib is recommended in the current label.

These results lead to the approval of axitinib plus avelumab by the FDA in the US. However, in the absence of a demonstrated benefit to overall survival, the combination is not currently recommended in the main international guidelines. Its place is all the more compromised as such approval was also obtained for the alternative combinations that were all compared to sunitinib: ipilimumab plus nivolumab, pembrolizumab plus axitinib, pembrolizumab plus lenvatinib and nivolumab plus cabozantinib (Table 2). To date, only indirect comparison in meta-analyses exists between the different treatment combinations.^{71–73} These meta-analyses, which vary in their methodology, tend to report better efficacy outcomes with nivolumab plus cabozantinib or pembrolizumab plus lenvatinib. Ipilimumab plus nivolumab is consistently associated with the highest likelihood of complete responses. Finally, in one of these reports, avelumab plus axitinib had the highest likelihood of an OS benefit for patients with favorable risk of advanced RCC.⁷¹

Toxicity Profile of Avelumab Axitinib Combination

The frequency and severity of adverse events observed with the combination of avelumab plus axitinib seem consistent with the known safety profiles of avelumab and axitinib when used as monotherapy.

Table 2 Randomized Phase III Trials of First-Line Immune Checkpoint Inhibitors and Vascular Endothelial Growth Factor Inhibitors Combinations in Advanced Renal Cell Carcinoma

Clinical Trials Number	Purpose	Patients Included	Results	Year First Published
NCT02231749	Nivolumab plus Ipilimumab versus sunitinib in patients with previously untreated advanced renal-cell carcinoma (CheckMate 214)	1096	Overall survival and objective response rates were significantly higher with nivolumab plus ipilimumab than with sunitinib among intermediate- and poor-risk patients.	2018
NCT02420821	Atezolizumab plus bevacizumab versus sunitinib in patients with previously untreated advanced renal-cell carcinoma (IMmotion151)	915	Atezolizumab plus bevacizumab prolonged progression-free survival versus sunitinib, although no improvement in overall survival was observed with atezolizumab plus bevacizumab at the final analysis.	2019
NCT02684006	Avelumab plus axitinib versus sunitinib in patients with previously untreated advanced renal-cell carcinoma (Javelin Renal 101)	886	Treatment with avelumab plus axitinib resulted in a statistically significant improvement in progression-free survival versus sunitinib.	2019
NCT02853331	Pembrolizumab plus axitinib versus sunitinib in patients with previously untreated advanced renal-cell carcinoma (KEYNOTE-426)	861	Treatment with pembrolizumab plus axitinib resulted in significantly longer overall survival and progression-free survival, as well as a higher objective response rate, than treatment with sunitinib.	2019
NCT03141177	Nivolumab plus cabozantinib versus sunitinib in patients with previously untreated advanced renal-cell carcinoma (CheckMate 9ER)	651	Nivolumab plus cabozantinib had significant benefits over sunitinib with respect to progression-free survival, overall survival, and likelihood of response.	2021
NCT02811861	Lenvatinib plus pembrolizumab or everolimus in patients with previously untreated advanced renal-cell carcinoma (CLEAR)	1069	Lenvatinib plus pembrolizumab was associated with significantly longer progression-free survival and overall survival than sunitinib.	2021

Avelumab and Axitinib Monotherapy Safety Profile

Avelumab is generally well tolerated and demonstrates a manageable safety profile in a large population of patients with advanced solid tumors. Among 1738 patients in the phase IB Javelin solid tumor trial,⁷⁴ the incidence of grade 3 treatment related adverse events (TRAEs) or immune related adverse events (IRAEs) of any grade was low. The most common TRAEs reported with avelumab monotherapy were fatigue, pruritus, nausea, diarrhea, and increased serum biomarkers. Infusion-related reactions (IRRs) and related symptoms occurred in 439 patients (25.3%) and mostly at first infusion. IRAEs occurred mainly during the first 2 infusions and were generally of low grade, easily manageable, and reversible. Most common were thyroid disorders and immune-related rashes.⁷⁵ Treatment discontinuation was rarely required. This is consistent with TRAEs reported in other trials of anti-PD-L1/PD-1 antibodies, although avelumab might appear with most favorable safety profile according to some meta-analyses.^{76,77} It has been suggested that this might be linked to avelumab shorter half-life, its ADCC activity or its specificity towards PD-L1, leaving PD-L2/PD-1 interaction free.⁷⁷⁻⁷⁹ Axitinib-related toxicity is globally similar to that reported with other VEGFR inhibitors. Dose limiting toxicities occurring in first in human study of axitinib were hypertension and stomatitis.¹⁶ In Phase 3 trials, the most common grade 3 TRAEs reported with axitinib were hypertension, diarrhea, decreased appetite, dysphonia and hypothyroidism.^{19,80,81} These have been reported with other VEGFR inhibitors, but other AEs often reported with currently approved VEGFR inhibitors are less commonly seen with axitinib: palmar-plantar erythrodysesthesia (PPE) syndrome, cutaneous toxicities, and myelosuppression. This might constitute a potential advantage of using more specific VEGFR inhibitors.

Toxicity Profile of Avelumab Plus Axitinib Combination

The combination of avelumab plus axitinib was initially evaluated in the phase Ib JAVELIN Renal 100 trial.⁸² In the dose-finding stage, patients received axitinib monotherapy twice daily for 1 week, followed by intravenous avelumab every 2 weeks plus axitinib twice daily. Only grade 3 proteinuria was dose-limiting. The maximum tolerated dose was defined as avelumab 10 mg/kg every 2 weeks and axitinib 5 mg twice daily. In the dose-expansion stage, 58% of grade 3-5 treatment-related adverse events were reported, the most frequent being hypertension, increased alanine aminotransferase, amylase, and lipase, and PPE syndrome. In the phase 3 JAVELIN 101 study,¹⁰ TRAEs occurred in 95.4% of patients who received avelumab plus axitinib versus 96.4% receiving sunitinib. Adverse events of grade 3 or higher occurred in 309 patients (56.7%) in the combination group versus 314 patients (55.4%) in the sunitinib group. Overall, 183 (42.2%) had at least one dose reduction of axitinib in the combination group compared to 187 (42.6%) in the sunitinib group. TRAEs leading to discontinuation of both avelumab and axitinib occurred in 33 patients (7.6%), which was lower than in the sunitinib group (13.4%). In both groups, the most frequent TRAEs were diarrhea, hypertension, PPE syndrome, and alanine aminotransferase increase. Adverse events categorized as IRAEs occurred in 166 out of 434 patients (38.2%), 9% of which were grade 3 or higher. The most frequent was immune-related thyroid disorders (24.7%). High-dose glucocorticoids were administered to 48 patients (11.1%). The JAVELIN 101 safety results were consistent with those of other phase 3 studies comparing TKI plus ICI combination strategies to sunitinib, in which TRAEs were similar between the study arms for all grade adverse events, grade 3 TRAEs being hypertension, diarrhea, PPE syndrome and ALT increase.⁷⁻⁹

Toxicity Management, a New Issue: Managing Overlapping Toxicities

A particular concern when combining VEGFR TKI plus ICI is overlapping toxicities. As with other approved combinations of VEGFR inhibitors plus ICI, axitinib plus ICI therapies have shown toxicity, likely due to overlapping toxicities, including fatigue, diarrhea, and hepatic toxicity.^{83,84} Therefore, a new issue to be addressed is the early identification of the underlying etiology of the toxicity: axitinib or immune related, which is crucial to set up appropriate management strategies. IRAEs are related to a non-specific activation of the immune system that can impact multiple organs. ICI-related toxicity management relies on the administration of immunosuppressors such as systemic corticosteroids.⁸⁵⁻⁸⁷ Conversely, antiangiogenic agents related to toxicity result from various underlying mechanisms due to on-target effects and are mainly managed by dose reductions and treatment interruption. The use of ICI plus TKI combination thus leads

to a more complex management and recognition of treatment-related toxicity, as there is an overlap between some ICI related and VEGFR TKI related adverse events.

As the number of patients treated with axitinib-based combination has increased, it appears necessary to get new guidance on how to distinguish IRAEs from axitinib-related adverse events and how to manage them specifically. An evidence and expert-based class IV consensus recommendation for treatment optimization and management of related adverse events in the management of axitinib plus ICI combination was published in 2020.⁸⁸ In this international consensus, a systematic review of the literature with avelumab plus axitinib and pembrolizumab plus axitinib combinations was performed, and adverse events associated with each combination were compared with the monotherapy treatments in similar indication. The most common (>10%) treatment-related adverse events associated with each combination that may be immune-related or associated with axitinib monotherapy included endocrinopathies, dermatologic, gastrointestinal, and musculoskeletal disorders. TRAEs that required additional guidance based on overlapping toxicities were identified, and recommendations were set up focusing on adverse events requiring differential diagnosis of etiology, for which no published guidelines were yet available.^{84,85,87,89,90} These events were identified as diarrhea, hepatitis, cardiovascular events, and fatigue and lead to specific recommendations.⁸⁸

As a general rule for low-grade adverse events, it is recommended to first assess the absence of severity symptoms that would require immediate intervention. In the absence of severe clinical signs of immune-related toxicity, the recommended first step is the interruption of axitinib and observation for resolution or improvement. In case of quick improvement or resolution, the adverse event is most likely due to axitinib, and intermittent treatment or dose reduction is recommended. If persistence, immune-related etiology must be considered and may require treatment with steroids.

Time to Treatment Resolution

Considering this, the time to resolution (TTR) of VEGFR TKI toxicity might be helpful in clinical practice. A shorter toxicity resolution decreases the risk of cumulative toxicity, and in combination regimens, TTR may help in determining the etiology of the toxicity. In a post hoc analysis of data from 5 studies of axitinib monotherapy or in combination with immunotherapy, Rini et al described the TTR of axitinib-related toxicity. Data from this analysis⁹¹ suggested that TTR for axitinib-related adverse events seems shorter after treatment interruption for axitinib monotherapy than for other VEGFR inhibitors. This was the same for axitinib monotherapy compared with combined axitinib and immunotherapy. In addition to fatigue, the majority of TTRs associated with combined axitinib plus ICI were longer than for axitinib monotherapy. The most common grade 3 events were fatigue, nausea, hypertension, and diarrhea. In the axitinib monotherapy cohort, the median TTR was 3 days for diarrhea, 8 days for fatigue, 2 days for hypertension, 4 days for nausea, and 3 days for PPE. In the axitinib plus ICI cohort, the median TTR was 4 to 11 days for all TRAEs. Although fully descriptive and extracted from clinical trials and not real-world data, these data suggest that axitinib-based combination toxicity may be easier to manage and the etiology easier to determine than with other combination regimens. The fast resolution of the adverse events (1–3 days except for fatigue) for axitinib monotherapy is likely related to its half-life (2.5 to 6.1 hours) which is notably shorter than that of other VEGFR TKI (approximately 28 hours for lenvatinib, 99 hours for cabozantinib, 50 hours for sunitinib). This may advantage axitinib in the management of TKI plus ICI combination toxicity and particularly in determining the underlying etiology.

Real Life Data

Phase 3 trials reporting the efficacy of ICI plus TKI, or ICI plus ICI combination have dramatically changed the standard of care for front-line therapy in mRCC. Physicians are now faced with larger therapeutic options with different toxicity profiles, but real-world evidence is lacking to accurately describe TKI plus ICI combination toxicity profile in non-trial selected populations, and to help better address patient selection. In a review, Allison et al reported their real-world experience of axitinib plus avelumab in 44 patients.⁹² The combination was well tolerated and the rate of grade 3 or higher adverse events was 36%, most common being hypertension (11%) and hepatitis (11%). Overall, 22% of patients had limiting toxicity and 20% of the patients received corticosteroids for suspicion of IRAEs, which was higher than data shown in the JAVELIN 101. As physicians are facing a growing number of patients eligible for this combination, more real-life data will be needed in the future to assess its true toxicity profile and set up optimal management strategies.

Factors Associated with Avelumab – Axitinib Outcomes and Their Impact on Therapeutic Strategy

Avelumab plus axitinib significantly improves PFS among treatment-naïve mRCC patients.⁶⁹ However, data suggest that almost half of patients will not benefit from this regimen. Thus, identifying factors associated with response remains a major challenge.

Tumor-Related Factors

The effectiveness of a treatment depends on its sufficient delivery to the tumor site, and the bone and brain are two potential sanctuaries for mRCC therapies. In the Phase II trial NIVOREN, nivolumab monotherapy failed to demonstrate efficacy in brain metastases from mRCC.⁹³ However, nivolumab plus ipilimumab combination showed encouraging results in this same situation, with a 32% to 44% ORR.^{94,95} Among the VEGFR TKIs, cabozantinib has shown significant activity in brain and bone metastases, while data concerning axitinib and other TKIs in the same setting lacks.⁹⁶ Therefore, sites of cancer extension are key factors in the treatment decision process, and avelumab plus axitinib may not be the preferred regimen for brain and bone involvement. Of all pathological subtypes, sarcomatoid mRCC have the poorest prognosis.⁹⁷ VEGFR TKIs have shown limited efficacy in this population, whereas nivolumab plus ipilimumab combination has become a standard.^{98,99} Interestingly, in a subgroup analysis of the sarcomatoid mRCC population of JAVELIN 101 trial, the objective response rate was twice as high with avelumab plus axitinib combination as with sunitinib.¹⁰⁰ Thus, this combination could be considered as an eligible option for these patients. As RCC patients represent a heterogeneous cancer population, IMDC risk criteria are critical in determining patient prognosis and can help predict patient response to systemic therapies. In the CHECKMATE 214 trial, nivolumab plus ipilimumab combination has shown limited benefit in favorable IMDC risk group in comparison to sunitinib monotherapy. Progression-free survival and objective response rates were even higher with sunitinib in this population (25.1 months [20.9 – NE] vs 15.3 months [9.7–20.3] HR: 0.45 [0.27–0.78], $p < 0.001$; 52% vs 29%, $p < 0.001$), suggesting pathological and molecular features associated with immune-resistance or VEGFR TKI high sensibility in this subgroup.⁶ Similarly, the benefit of pembrolizumab plus axitinib and nivolumab plus cabozantinib combinations in the favorable-risk subpopulations of KEYNOTE-426 and CHECKMATE-9ER trials failed to reach significance.^{7,8} However, in the JAVELIN 101 trial, avelumab plus axitinib showed consistent efficacy in all IMDC risk groups. This questions the impact of PD-L1 tumor expression, as the population is enriched in PD-L1-positive tumors. PD-L1 expression by tumor cells or tumor microenvironment has been widely investigated as a predictive biomarker for ICI⁶⁹(p1),¹⁰¹(p1). In lung cancer, the tumor PD-L1 expression determines the choice of treatment for first line of therapy.¹⁰² In RCC, this biomarker seems to be associated with a poor prognosis,¹⁰³ but to this day its predictive value remains controversial^{104,105} and the interpretation of the predictive value of PD-L1 tumor expression in mRCC is all the more complex in the era of ICI plus TKI association. Recently, Damotte et al have demonstrated that an 18-gene expression assay measuring the level of tumor microenvironment inflammation (TIS score) is independently predictive of tumor response to anti-PD1 therapies in a cohort of 58 solid cancer patients including 10 RCC.¹⁰⁶ Given the limited sample size, further trials are warranted to implement these results in clinical practice. Other biomarkers such as tumor mutational burden, soluble biomarkers and molecular characterization have shown encouraging yet preliminary results.^{107–112}

Host-Related Factors

The factors associated with immune response appear to be in a large part host-related and independent of the tumor. The ELY study investigated the measured resting energy expenditure (REE) as a predictive clinical marker for 6-month progression-free survival among 110 metastatic non-small cell lung cancer patients treated with ICI.¹¹³ Hypermetabolism, defined as a measured/theoretical REE ratio of 110% or above, was associated with a 35% decrease of 6-months PFS rates (22% versus 55%; odds ratio: 4.76; IC95 [1.87–12.89], $p < 0.001$). Hypermetabolism is also correlated with sarcopenia and cancer cachexia and is associated with a shorter survival in metastatic cancer patients.¹¹⁴ A previous study has also highlighted sarcopenia as a predictive factor for anti-PD1 toxicity.¹¹⁵ Taken together, these results point to hypermetabolism not only as a major prognostic marker but also as a potential target for improving ICI response. The gut microbiota is another host-related marker, which may reflect the immune system efficiency. The impact

of antibiotic consumption on ICI efficacy is now thoroughly documented.^{116,117} Its deleterious effect results from excessive bacterial depletion, as specific commensal germs such as *Akkermansia muciniphila*, *Alistipes indistinctus* and *Enterococcus hirae* seem to be critical for T-cell activation.^{118,119} Among the various corrective strategies under investigation, such as probiotic use or antibiotic sparing, fecal microbiota transplantation (FMT) is one of the most promising. A preliminary clinical study has investigated FMT obtained from long-responding melanoma patient as a therapy for anti-PD1 refractory melanoma patients and demonstrated that a single transplantation could help overcome resistance in a subset of patients.¹²⁰ A very recent Phase 1 trial showed that a bifidogenic live bacterial product appears to enhance the clinical outcome in patients with mRCC treated with nivolumab-ipilimumab.¹²¹ Interestingly, although diarrhea is a common adverse event of anti-VEGF therapies, very few studies have investigated the clinical impact of TKI-induced dysbiosis. In 2015, Pal et al have conducted stool bacteriomic profiling of patients with mRCC receiving VEGF TKIs, and found that high rates of *Bacteroides spp.* and low rates of *Prevotella spp.* were associated with higher risks of diarrhea.¹²² These findings are supported by recent studies suggesting that microbioma modulating therapies such as probiotics and FMT from healthy donors may also be effective treatments for VEGFR TKIs induced diarrhea in mRCC patients.^{123,124} In summary, the understanding of the host, its metabolism and its immunity could ultimately lead to therapeutic strategies based on the optimization of the immune response and limit the stakes of the choice of combination.

Treatment-Related Factors

Therapeutic drug monitoring is a powerful instrument for dose adjustment of targeted therapies.^{125,126} A strong PK/PD relationship has been demonstrated for cabozantinib and sunitinib.^{127,128} However, the mechanism of inter-individual variability in exposure remains unclear. For axitinib, several data suggest that dose titration is insufficient to overcome underexposure.^{21,129} Various factors, such as body composition, drug–drug interactions and self-medication, are likely to influence exposure as well.¹³⁰ Conversely, there is no demonstrated exposure–response relationship with anti-PD(L)1 therapies, which confirms that addressing host-related factors associated with response is crucial to derive the highest benefit of these therapies.^{44,131} Therefore, a tailored pre-therapeutic pharmacological risk assessment, taking into account individual patient characteristics at the same level as the plasma concentration of the drug, should be the first step into decision-making.

Conclusion

Avelumab plus axitinib combination was developed in times of dramatically increasing therapeutic opportunities for mRCC patients. Although ICI plus TKI and ICI plus ICI combinations are standard of care, the added value of each combination in comparison to the others is debatable. Axitinib has some advantages over other TKIs for a combination with ICI: it displays few risks of drug–drug interaction, its dose can be easily optimized with up-titration based on hypertension (a non-overlapping toxicity), and it has a short half-life with therefore a short time to toxicity resolution. For avelumab, the induced ADCC and a favorable safety profile may be advantageous over other ICI. Nevertheless, better understanding of the complex interaction between immunomodulation and angiogenesis and improving our ability to restore the immune system may be more discriminating than the choice of systemic combination.

Disclosure

Pr. Goldwasser reports grants and personal fees from BAXTER, and personal fees from FRESENIUS, KABI, and NUTRICIA, outside the submitted work. Dr. Huillard reports personal fees from BAYER, SANOFI, MSD, BMS, IPSEN, PFIZER, EISAI, JANSSEN, and ASTRA ZENECA, outside the submitted work. The authors report no other conflicts of interest in this work.

References

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin.* 2022;72(1):7–33. doi:10.3322/caac.21708
2. Rini BI, Dorff TB, Elson P, et al. Active surveillance in metastatic renal-cell carcinoma: a prospective, Phase 2 trial. *Lancet Oncol.* 2016;17(9):1317–1324. doi:10.1016/S1470-2045(16)30196-6

3. Marchioni M, Bandini M, Pompe RS, et al. Survival of metastatic renal cell carcinoma patients continues to improve over time, even in targeted therapy era. *Int Urol Nephrol*. 2017;49(12):2143–2149. doi:10.1007/s11255-017-1703-y
4. Motzer RJ, Hutson TE, Cella D, et al. Pazopanib versus sunitinib in metastatic renal-cell carcinoma. *N Engl J Med*. 2013;369(8):722–731. doi:10.1056/NEJMoa1303989
5. Motzer RJ, Hutson TE, Tomczak P, et al. Overall survival and updated results for sunitinib compared with interferon alfa in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2009;27:3584–3590. doi:10.1200/JCO.2008.20.1293
6. Motzer RJ, Tannir NM, McDermott DF, et al. Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma. *N Engl J Med*. 2018;2018:14.
7. Choueiri TK, Powles T, Burotto M, et al. Nivolumab plus cabozantinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med*. 2021;384(9):829–841. doi:10.1056/NEJMoa2026982
8. Rini BI, Plimack ER, Stus V, et al. Pembrolizumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med*. 2019;380(12):1116–1127. doi:10.1056/NEJMoa1816714
9. Motzer R, Alekseev B, Rha SY, et al. Lenvatinib plus pembrolizumab or everolimus for advanced renal cell carcinoma. *N Engl J Med*. 2021;384(14):1289–1300. doi:10.1056/NEJMoa2035716
10. Motzer RJ, Penkov K, Haanen J, et al. Avelumab plus axitinib versus sunitinib for advanced renal-cell carcinoma. *N Engl J Med*. 2019;380(12):1103–1115. doi:10.1056/NEJMoa1816047
11. Bellesoeur A, Carton E, Alexandre J, Goldwasser F, Huillard O. Axitinib in the treatment of renal cell carcinoma: design, development, and place in therapy. *Drug Des Devel Ther*. 2017;11:2801–2811. doi:10.2147/DDDT.S109640
12. Escudier B, Gore M. Axitinib for the management of metastatic renal cell carcinoma. *Drugs RD*. 2011;11(2):113–126. doi:10.2165/11591240-000000000-00000
13. Pithavala YK, Chen Y, Toh M, et al. Evaluation of the effect of food on the pharmacokinetics of axitinib in healthy volunteers. *Cancer Chemother Pharmacol*. 2012;70(1):103–112. doi:10.1007/s00280-012-1888-9
14. European Medicines Agency. Inlyta (Axitinib): EU EPAR public assessment report; 2012.
15. Chen Y, Tortorici MA, Garrett M, Hee B, Klammer KJ, Pithavala YK. Clinical pharmacology of axitinib. *Clin Pharmacokinet*. 2013;52(9):713–725. doi:10.1007/s40262-013-0068-3
16. Rugo HS, Herbst RS, Liu G, et al. Phase I trial of the oral antiangiogenesis agent AG-013736 in patients with advanced solid tumors: pharmacokinetic and clinical results. *J Clin Oncol off J Am Soc Clin Oncol*. 2005;23(24):5474–5483. doi:10.1200/JCO.2005.04.192
17. Rini BI, Schiller JH, Fruehauf JP, et al. Diastolic blood pressure as a biomarker of axitinib efficacy in solid tumors. *Clin Cancer Res off J Am Assoc Cancer Res*. 2011;17(11):3841–3849. doi:10.1158/1078-0432.CCR-10-2806
18. Chen Y, Rini BI, Bair AH, Mugundu GM, Pithavala YK. Population pharmacokinetic-pharmacodynamic modelling of 24-h diastolic ambulatory blood pressure changes mediated by axitinib in patients with metastatic renal cell carcinoma. *Clin Pharmacokinet*. 2015;54(4):397–407. doi:10.1007/s40262-014-0207-5
19. Rini BI, Garrett M, Poland B, et al. Axitinib in metastatic renal cell carcinoma: results of a pharmacokinetic and pharmacodynamic analysis. *J Clin Pharmacol*. 2013;53(5):491–504. doi:10.1002/jcph.73
20. Rini BI, Melichar B, Ueda T, et al. Axitinib with or without dose titration for first-line metastatic renal-cell carcinoma: a randomised double-blind phase 2 trial. *Lancet Oncol*. 2013;14(12):1233–1242. doi:10.1016/S1470-2045(13)70464-9
21. Rini BI, Melichar B, Fishman MN, et al. Axitinib dose titration: analyses of exposure, blood pressure and clinical response from a randomized phase II study in metastatic renal cell carcinoma. *Ann Oncol off J Eur Soc Med Oncol*. 2015;26(7):1372–1377. doi:10.1093/annonc/mdv103
22. Chen Y, Suzuki A, Tortorici MA, et al. Axitinib plasma pharmacokinetics and ethnic differences. *Invest New Drugs*. 2015;33(2):521–532. doi:10.1007/s10637-015-0214-x
23. Chen Y, Rini BI, Motzer RJ, et al. Effect of renal impairment on the pharmacokinetics and safety of axitinib. *Target Oncol*. 2016;11(2):229–234. doi:10.1007/s11523-015-0389-2
24. Brennan M, Williams JA, Chen Y, Tortorici M, Pithavala Y, Liu YC. Meta-analysis of contribution of genetic polymorphisms in drug-metabolizing enzymes or transporters to axitinib pharmacokinetics. *Eur J Clin Pharmacol*. 2012;68(5):645–655. doi:10.1007/s00228-011-1171-8
25. Huillard O, Mir O, Peyromaure M, et al. Sarcopenia and body mass index predict sunitinib-induced early dose-limiting toxicities in renal cancer patients. *Br J Cancer*. 2013;108:1034–1041. doi:10.1038/bjc.2013.58.bjc201358
26. Antoun S, Baracos VE, Birdsell L, Escudier B, Sawyer MB. Low body mass index and sarcopenia associated with dose-limiting toxicity of sorafenib in patients with renal cell carcinoma. *Ann Oncol off J Eur Soc Med Oncol ESMO*. 2010;21(8):1594–1598. doi:10.1093/annonc/mdp605
27. Tortorici MA, Toh M, Rahavendran SV, et al. Influence of mild and moderate hepatic impairment on axitinib pharmacokinetics. *Invest New Drugs*. 2011;29(6):1370–1380. doi:10.1007/s10637-010-9477-4
28. Thomas-Schoemann A, Blanchet B, Bardin C, et al. Drug interactions with solid tumour-targeted therapies. *Crit Rev Oncol Hematol*. 2014;89:179–196. doi:10.1016/j.critrevonc.2013.08.007S1040-8428(13)00184-4[pii]
29. Pithavala YK, Tong W, Mount J, et al. Effect of ketoconazole on the pharmacokinetics of axitinib in healthy volunteers. *Invest New Drugs*. 2012;30(1):273–281. doi:10.1007/s10637-010-9511-6
30. Gomo C, Coriat R, Faivre L, et al. Pharmacokinetic interaction involving sorafenib and the calcium-channel blocker felodipine in a patient with hepatocellular carcinoma. *Invest New Drugs*. 2011;29(6):1511–1514. doi:10.1007/s10637-010-9514-3
31. Da Silva F, Thomas-Schoemann A, Huillard O, Goldwasser F, Blanchet B. Benefit of therapeutic drug monitoring to disclose pharmacokinetic interaction between sunitinib and calcium channel blocker. *Ann Oncol off J Eur Soc Med Oncol*. 2016;27(8):1651–1652. doi:10.1093/annonc/mdw182
32. Pithavala YK, Tortorici M, Toh M, et al. Effect of rifampin on the pharmacokinetics of Axitinib (AG-013736) in Japanese and Caucasian healthy volunteers. *Cancer Chemother Pharmacol*. 2010;65(3):563–570. doi:10.1007/s00280-009-1065-y

33. Tzoganis K, Skibeli V, Westgaard I, et al. The European Medicines Agency approval of axitinib (Inlyta) for the treatment of advanced renal cell carcinoma after failure of prior treatment with sunitinib or a cytokine: summary of the scientific assessment of the committee for medicinal products for human use. *Oncologist*. 2015;20(2):196–201. doi:10.1634/theoncologist.2014-0177
34. Motzer RJ, Escudier B, Tomczak P, et al. Axitinib versus sorafenib as second-line treatment for advanced renal cell carcinoma: overall survival analysis and updated results from a randomised phase 3 trial. *Lancet Oncol*. 2013;14(6):552–562. doi:10.1016/S1470-2045(13)70093-7
35. Huillard O, Boissier E, Blanchet B, et al. Drug safety evaluation of sorafenib for treatment of solid tumors: consequences for the risk assessment and management of cancer patients. *Expert Opin Drug Saf*. 2014;13(5):663–673. doi:10.1517/14740338.2014.907270
36. Arrondeau J, Mir O, Boudou-Rouquette P, et al. Sorafenib exposure decreases over time in patients with hepatocellular carcinoma. *Invest New Drugs*. 2012;30(5):2046–2049. doi:10.1007/s10637-011-9764-8
37. Boudou-Rouquette P, Ropert S, Mir O, et al. Variability of sorafenib toxicity and exposure over time: a pharmacokinetic/pharmacodynamic analysis. *Oncologist*. 2012;17(9):1204–1212. doi:10.1634/theoncologist.2011-0439
38. Bellesoeur A, Carton E, Mir O, et al. Critical role of sorafenib exposure over time for its antitumor activity in thyroid cancer. *Invest New Drugs*. 2014;32(3):569–572. doi:10.1007/s10637-013-0052-7
39. Boudou-Rouquette P, Thomas-Schoemann A, Bellesoeur A, Goldwasser F. Sorafenib for patients with differentiated thyroid cancer. *Lancet*. 2015;385(9964):227–228. doi:10.1016/S0140-6736(15)60054-X
40. Blanchet B, Billemonet B, Cramard J, et al. Validation of an HPLC-UV method for sorafenib determination in human plasma and application to cancer patients in routine clinical practice. *J Pharm Biomed Anal*. 2009;49:1109–1114. doi:10.1016/j.jpba.2009.02.008.S0731-7085(09)00112-5
41. European Medicines Agency. Bavencio (Avelumab): EU EPAR public assessment report; 2022.
42. Heery CR, O’Sullivan-Coyne G, Madan RA, et al. Avelumab for metastatic or locally advanced previously treated solid tumours (JAVELIN Solid Tumor): a phase 1a, multicohort, dose-escalation trial. *Lancet Oncol*. 2017;18(5):587–598. doi:10.1016/S1470-2045(17)30239-5
43. Sheng J, Srivastava S, Sanghavi K, et al. Clinical pharmacology considerations for the development of immune checkpoint inhibitors. *J Clin Pharmacol*. 2017;57(Suppl 10):S26–S42. doi:10.1002/jcph.990
44. Novakovic AM, Wilkins JJ, Dai H, et al. Changing body weight-based dosing to a flat dose for avelumab in metastatic Merkel cell and advanced urothelial carcinoma. *Clin Pharmacol Ther*. 2020;107(3):588–596. doi:10.1002/cpt.1645
45. Wilkins JJ, Brockhaus B, Dai H, et al. Time-varying clearance and impact of disease state on the pharmacokinetics of avelumab in Merkel cell carcinoma and urothelial carcinoma. *CPT Pharmacomet Syst Pharmacol*. 2019;8(6):415–427. doi:10.1002/psp4.12406
46. Powles T, Albiges L, Bex A, et al. ESMO Clinical Practice Guideline update on the use of immunotherapy in early stage and advanced renal cell carcinoma. *Ann Oncol*. 2021;32(12):1511–1519. doi:10.1016/j.annonc.2021.09.014
47. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646–674. doi:10.1016/j.cell.2011.02.013
48. Boyerinas B, Jochems C, Fantini M, et al. Antibody-dependent cellular cytotoxicity activity of a novel anti-PD-L1 antibody avelumab (MSB0010718C) on human tumor cells. *Cancer Immunol Res*. 2015;3(10):1148–1157. doi:10.1158/2326-6066.CIR-15-0059
49. Amin A, Plimack ER, Infante JR, et al. Nivolumab (anti-PD-1; BMS-936558, ONO-4538) in combination with sunitinib or pazopanib in patients (pts) with metastatic renal cell carcinoma (mRCC). *J Clin Oncol*. 2014;32(15_suppl):5010. doi:10.1200/jco.2014.32.15_suppl.5010
50. Dikov MM, Ohm JE, Ray N, et al. Differential roles of vascular endothelial growth factor receptors 1 and 2 in dendritic cell differentiation. *J Immunol Baltim Md*. 2005;174(1):215–222. doi:10.4049/jimmunol.174.1.215
51. Sidibe A, Ropraz P, Jemelin S, et al. Angiogenic factor-driven inflammation promotes extravasation of human proangiogenic monocytes to tumours. *Nat Commun*. 2018;9(1):355. doi:10.1038/s41467-017-02610-0
52. Ohm JE, Gabrilovich DI, Sempowski GD. VEGF inhibits T-cell development and may contribute to tumor-induced immune suppression. *Blood*. 2003;101(12):4878–4886. doi:10.1182/blood-2002-07-1956
53. Voron T, Colussi O, Marcheteau E, et al. VEGF-A modulates expression of inhibitory checkpoints on CD8+ T cells in tumors. *J Exp Med*. 2015;212(2):139–148. doi:10.1084/jem.20140559
54. Terme M, Pernet S, Marcheteau E, et al. VEGFA-VEGFR pathway blockade inhibits tumor-induced regulatory T-cell proliferation in colorectal cancer. *Cancer Res*. 2013;73(2):539–549. doi:10.1158/0008-5472.CAN-12-2325
55. Hirsch L, Flippot R, Escudier B, Albiges L. Immunomodulatory roles of VEGF pathway inhibitors in renal cell carcinoma. *Drugs*. 2020;80(12):1169–1181. doi:10.1007/s40265-020-01327-7
56. Yuan H, Cai P, Li Q, et al. Axitinib augments antitumor activity in renal cell carcinoma via STAT3-dependent reversal of myeloid-derived suppressor cell accumulation. *Biomed Pharmacother*. 2014;68(6):751–756. doi:10.1016/j.biopha.2014.07.002
57. Morelli MB, Amantini C, Santoni M, et al. Axitinib induces DNA damage response leading to senescence, mitotic catastrophe, and increased NK cell recognition in human renal carcinoma cells. *Oncotarget*. 2015;6(34):36245–36259. doi:10.18632/oncotarget.5768
58. Läubli H, Müller P, D’Amico L, Buchi M, Kashyap AS, Zippelius A. The multi-receptor inhibitor axitinib reverses tumor-induced immunosuppression and potentiates treatment with immune-modulatory antibodies in preclinical murine models. *Cancer Immunol Immunother CII*. 2018;67(5):815–824. doi:10.1007/s00262-018-2136-x
59. Shrimali RK, Yu Z, Theoret MR, Chinnasamy D, Restifo NP, Rosenberg SA. Antiangiogenic agents can increase lymphocyte infiltration into tumor and enhance the effectiveness of adoptive immunotherapy of cancer. *Cancer Res*. 2010;70(15):6171–6180. doi:10.1158/0008-5472.CAN-10-0153
60. Motz GT, Santoro SP, Wang LP, et al. Tumor endothelium FasL establishes a selective immune barrier promoting tolerance in tumors. *Nat Med*. 2014;20(6):607–615. doi:10.1038/nm.3541
61. Movahedi K, Laoui D, Gysemans C, et al. Different tumor microenvironments contain functionally distinct subsets of macrophages derived from Ly6C (high) monocytes. *Cancer Res*. 2010;70(14):5728–5739. doi:10.1158/0008-5472.CAN-09-4672
62. Facciabene A, Peng X, Hagemann IS, et al. Tumour hypoxia promotes tolerance and angiogenesis via CCL28 and Treg cells. *Nature*. 2011;475(7355):226–230. doi:10.1038/nature10169
63. Huang Y, Goel S, Duda DG, Fukumura D, Jain RK. Vascular normalization as an emerging strategy to enhance cancer immunotherapy. *Cancer Res*. 2013;73(10):2943–2948. doi:10.1158/0008-5472.CAN-12-4354
64. Huang Y, Yuan J, Righi E, et al. Vascular normalizing doses of antiangiogenic treatment reprogram the immunosuppressive tumor microenvironment and enhance immunotherapy. *Proc Natl Acad Sci U S A*. 2012;109(43):17561–17566. doi:10.1073/pnas.1215397109

65. Rodig N, Ryan T, Allen JA, et al. Endothelial expression of PD-L1 and PD-L2 down-regulates CD8+ T cell activation and cytotoxicity. *Eur J Immunol*. 2003;33(11):3117–3126. doi:10.1002/eji.200324270
66. Lu Y, Yang W, Qin C, et al. Responsiveness of stromal fibroblasts to IFN-gamma blocks tumor growth via angiostasis. *J Immunol Baltim Md*. 2009;183(10):6413–6421. doi:10.4049/jimmunol.0901073
67. Khan KA, Kerbel RS. Improving immunotherapy outcomes with anti-angiogenic treatments and vice versa. *Nat Rev Clin Oncol*. 2018;15(5):310–324. doi:10.1038/nrclinonc.2018.9
68. Tian L, Goldstein A, Wang H, et al. Mutual regulation of tumour vessel normalization and immunostimulatory reprogramming. *Nature*. 2017;544(7649):250–254. doi:10.1038/nature21724
69. Choueiri TK, Motzer RJ, Rini BI, et al. Updated efficacy results from the JAVELIN Renal 101 trial: first-line avelumab plus axitinib versus sunitinib in patients with advanced renal cell carcinoma. *Ann Oncol*. 2021;31:1030–1039. doi:10.1016/j.annonc.2020.04.010
70. Masters JC, Khandelwal A, Di Pietro A, Dai H, Brar S. Model-informed drug development supporting the approval of the avelumab flat-dose regimen in patients with advanced renal cell carcinoma (aRCC). *CPT Pharmacomet Syst Pharmacol*. 2022;11(4):458–468. doi:10.1002/psp4.12771
71. Bosma NA, Warkentin MT, Gan CL, et al. Efficacy and safety of first-line systemic therapy for metastatic renal cell carcinoma: a systematic review and network meta-analysis. *Eur Urol Open Sci*. 2022;37:14–26. doi:10.1016/j.euro.2021.12.007
72. Quhal F, Mori K, Bruchbacher A, et al. First-line immunotherapy-based combinations for metastatic renal cell carcinoma: a systematic review and network meta-analysis. *Eur Urol Oncol*. 2021;4(5):755–765. doi:10.1016/j.euo.2021.03.001
73. Riaz IB, He H, Ryu AJ, et al. A living, interactive systematic review and network meta-analysis of first-line treatment of metastatic renal cell carcinoma. *Eur Urol*. 2021;80(6):712–723. doi:10.1016/j.eururo.2021.03.016
74. Vaishampayan U, Schöffski P, Ravaud A, et al. Avelumab monotherapy as first-line or second-line treatment in patients with metastatic renal cell carcinoma: phase Ib results from the JAVELIN Solid Tumor trial. *J Immunother Cancer*. 2019;7(1):275. doi:10.1186/s40425-019-0746-2
75. Kelly K, Infante JR, Taylor MH, et al. Safety profile of avelumab in patients with advanced solid tumors: a pooled analysis of data from the phase 1 JAVELIN solid tumor and phase 2 JAVELIN Merkel 200 clinical trials. *Cancer*. 2018;124(9):2010–2017. doi:10.1002/cncr.31293
76. Al-Showbaki L, Nadler MB, Desnoyers A, Almugbel FA, Cescon DW, Amir E. Network meta-analysis comparing efficacy, safety and tolerability of anti-PD-1/PD-L1 antibodies in solid cancers. *J Cancer*. 2021;12(14):4372–4378. doi:10.7150/jca.57413
77. Sonpavde GP, Grivas P, Lin Y, Hennessy D, Hunt JD. Immune-related adverse events with PD-1 versus PD-L1 inhibitors: a meta-analysis of 8730 patients from clinical trials. *Future Oncol Lond Engl*. 2021;17(19):2545–2558. doi:10.2217/fon-2020-1222
78. Spain L, Diem S, Larkin J. Management of toxicities of immune checkpoint inhibitors. *Cancer Treat Rev*. 2016;44:51–60. doi:10.1016/j.ctrv.2016.02.001
79. de Velasco G, Je Y, Bossé D, et al. Comprehensive meta-analysis of key immune-related adverse events from CTLA-4 and PD-1/PD-L1 inhibitors in cancer patients. *Cancer Immunol Res*. 2017;5(4):312–318. doi:10.1158/2326-6066.CIR-16-0237
80. Hutson TE, Lesovoy V, Al-Shukri S, et al. Axitinib versus sorafenib as first-line therapy in patients with metastatic renal-cell carcinoma: a randomised open-label phase 3 trial. *Lancet Oncol*. 2013;14(13):1287–1294. doi:10.1016/S1470-2045(13)70465-0
81. Rini BI, Escudier B, Tomczak P, et al. Comparative effectiveness of axitinib versus sorafenib in advanced renal cell carcinoma (AXIS): a randomised phase 3 trial. *Lancet Lond Engl*. 2011;378(9807):1931–1939. doi:10.1016/S0140-6736(11)61613-9
82. Choueiri TK, Larkin J, Oya M, et al. Preliminary results for avelumab plus axitinib as first-line therapy in patients with advanced clear-cell renal-cell carcinoma (JAVELIN Renal 100): an open-label, dose-finding and dose-expansion, phase 1b trial. *Lancet Oncol*. 2018;19(4):451–460. doi:10.1016/S1470-2045(18)30107-4
83. Amin A, Plimack ER, Ernstoff MS, et al. Safety and efficacy of nivolumab in combination with sunitinib or pazopanib in advanced or metastatic renal cell carcinoma: the CheckMate 016 study. *J Immunother Cancer*. 2018;6(1):109. doi:10.1186/s40425-018-0420-0
84. Rini BI, Battle D, Figlin RA, et al. The Society for Immunotherapy of Cancer consensus statement on immunotherapy for the treatment of advanced renal cell carcinoma (RCC). *J Immunother Cancer*. 2019;7(1):354. doi:10.1186/s40425-019-0813-8
85. Haanen JB, Carbone G, Robert C F, et al. Management of toxicities from immunotherapy: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol off J Eur Soc Med Oncol*. 2017;28(suppl_4):iv119–iv142. doi:10.1093/annonc/mdx225
86. Brahmer JR, Lacchetti C, Schneider BJ, et al. Management of immune-related adverse events in patients treated with immune checkpoint inhibitor therapy: American Society of clinical oncology clinical practice guideline. *J Clin Oncol off J Am Soc Clin Oncol*. 2018;36(17):1714–1768. doi:10.1200/JCO.2017.77.6385
87. Puzanov I, Diab A, Abdallah K, et al. Managing toxicities associated with immune checkpoint inhibitors: consensus recommendations from the Society for Immunotherapy of Cancer (SITC) Toxicity Management Working Group. *J Immunother Cancer*. 2017;5(1):95. doi:10.1186/s40425-017-0300-z
88. Grünwald V, Voss MH, Rini BI, et al. Axitinib plus immune checkpoint inhibitor: evidence- and expert-based consensus recommendation for treatment optimisation and management of related adverse events. *Br J Cancer*. 2020;123(6):898–904. doi:10.1038/s41416-020-0949-9
89. Brahmer JR, Tykodi SS, Chow LQM, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *N Engl J Med*. 2012;366(26):2455–2465. doi:10.1056/NEJMoa1200694
90. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009;339:b2700. doi:10.1136/bmj.b2700
91. Rini BI, Atkins MB, Choueiri TK, et al. Time to resolution of axitinib-related adverse events after treatment interruption in patients with advanced renal cell carcinoma. *Clin Genitourin Cancer*. 2021;19(5):e306–e312. doi:10.1016/j.clgc.2021.03.019
92. Allison J, Charnley N, Stevenson R, Waddell T, Pillai MR. Axitinib and avelumab (AA) as first-line treatment of metastatic renal cell carcinoma (mRCC): a real-world outcome review in the Northwest of England, United Kingdom. *J Clin Oncol*. 2021;39(6_suppl):294. doi:10.1200/JCO.2021.39.6_suppl.294
93. Flippot R, Dalban C, Laguerre B, et al. Safety and efficacy of nivolumab in brain metastases from renal cell carcinoma: results of the GETUG-AFU 26 NIVOREN Multicenter Phase II Study. *J Clin Oncol off J Am Soc Clin Oncol*. 2019;37(23):2008–2016. doi:10.1200/JCO.18.02218

94. Enamekhoo H, Olsen MR, Carthon BC, et al. Safety and efficacy of nivolumab plus ipilimumab in patients with advanced renal cell carcinoma with brain metastases: checkMate 920. *Cancer*. 2022;128(5):966–974. doi:10.1002/cncr.34016
95. Brown LC, Desai K, Wei W, et al. Clinical outcomes in patients with metastatic renal cell carcinoma and brain metastasis treated with ipilimumab and nivolumab. *J Immunother Cancer*. 2021;9(9):e003281. doi:10.1136/jitc-2021-003281
96. Escudier B, Powles T, Motzer RJ, et al. Cabozantinib, a new standard of care for patients with advanced renal cell carcinoma and bone metastases? Subgroup analysis of the METEOR trial. *J Clin Oncol off J Am Soc Clin Oncol*. 2018;36(8):765–772. doi:10.1200/JCO.2017.74.7352
97. Blum KA, Gupta S, Tickoo SK, et al. Sarcomatoid renal cell carcinoma: biology, natural history and management. *Nat Rev Urol*. 2020;17(12):659–678. doi:10.1038/s41585-020-00382-9
98. Golshayan AR, George S, Heng DY, et al. Metastatic sarcomatoid renal cell carcinoma treated with vascular endothelial growth factor-targeted therapy. *J Clin Oncol off J Am Soc Clin Oncol*. 2009;27(2):235–241. doi:10.1200/JCO.2008.18.0000
99. Tannir NM, Signoretti S, Choueiri TK, et al. Efficacy and safety of nivolumab plus ipilimumab versus sunitinib in first-line treatment of patients with advanced sarcomatoid renal cell carcinoma. *Clin Cancer Res off J Am Assoc Cancer Res*. 2021;27(1):78–86. doi:10.1158/1078-0432.CCR-20-2063
100. Choueiri TK. Efficacy and correlative analyses of avelumab plus axitinib versus sunitinib in sarcomatoid renal cell carcinoma: post hoc analysis of a randomized clinical trial. *ESMO Open*. 2021;6(3):12.
101. Carbognin L, Pilotto S, Milella M, et al. Differential activity of nivolumab, pembrolizumab and MPDL3280A according to the tumor expression of programmed death-ligand-1 (PD-L1): sensitivity analysis of trials in melanoma, lung and genitourinary cancers. *PLoS One*. 2015;10(6):e0130142. doi:10.1371/journal.pone.0130142
102. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med*. 2016;375(19):1823–1833. doi:10.1056/NEJMoa1606774
103. Iacovelli R, Nolè F, Verri E, et al. Prognostic role of PD-L1 expression in renal cell carcinoma. a systematic review and meta-analysis. *Target Oncol*. 2016;11(2):143–148. doi:10.1007/s11523-015-0392-7
104. Mori K, Abufaraj M, Mostafaei H, et al. The predictive value of programmed death ligand 1 in patients with metastatic renal cell carcinoma treated with immune-checkpoint inhibitors: a systematic review and meta-analysis. *Eur Urol*. 2020. doi:10.1016/j.eururo.2020.10.006
105. Noel J, Huillard O, Goldwasser F, et al. The predictive value of programmed death ligand 1 in patients with metastatic renal cell carcinoma treated with immune-checkpoint inhibitors: a systematic review and meta-analysis. *Eur Urol*. 2021;79(4):e112. doi:10.1016/j.eururo.2021.01.013
106. Damotte D, Warren S, Arrondeau J, et al. The tumor inflammation signature (TIS) is associated with anti-PD-1 treatment benefit in the CERTIM pan-cancer cohort. *J Transl Med*. 2019;17(1):357. doi:10.1186/s12967-019-2100-3
107. Zhu J, Armstrong AJ, Friedlander TW, et al. Biomarkers of immunotherapy in urothelial and renal cell carcinoma: PD-L1, tumor mutational burden, and beyond. *J Immunother Cancer*. 2018;6(1):4. doi:10.1186/s40425-018-0314-1
108. Larrinaga G, Solano-Iturri JD, Errarte P, et al. Soluble PD-L1 is an independent prognostic factor in clear cell renal cell carcinoma. *Cancers*. 2021;13(4):667. doi:10.3390/cancers13040667
109. Zizzari IG, Napoletano C, Di Filippo A, et al. Exploratory pilot study of circulating biomarkers in metastatic renal cell carcinoma. *Cancers*. 2020;12(9):E2620. doi:10.3390/cancers12092620
110. Tiako Meyo M, Jouinot A, Fabre E, et al. Predictive value of soluble PD-1, PD-L1, VEGFA, CD40 ligand and CD44 for nivolumab therapy in advanced non-small cell lung cancer: a case-control study. *Cancers*. 2020;12(2):473. doi:10.3390/cancers12020473
111. Epaillard N, Simonaggio A, Elaidi R, et al. BIONIKK: a phase 2 biomarker driven trial with nivolumab and ipilimumab or VEGFR tyrosine kinase inhibitor (TKI) in naïve metastatic kidney cancer. *Bull Cancer*. 2020;107(5S):eS22–eS27. doi:10.1016/S0007-4551(20)30283-6
112. Motzer RJ, Robbins PB, Powles T, et al. Avelumab plus axitinib versus sunitinib in advanced renal cell carcinoma: biomarker analysis of the phase 3 JAVELIN Renal 101 trial. *Nat Med*. 2020;26(11):1733–1741. doi:10.1038/s41591-020-1044-8
113. Boudou-Rouquette P, Arrondeau J, Gervais C, et al. Development and validation of a host-dependent, PDL1-independent, biomarker to predict 6-month progression-free survival in metastatic non-small cell lung cancer (mNSCLC) patients treated with anti-PD1 immune checkpoint inhibitors (ICI) in the CERTIM Cohort: the ELY study. *EBioMedicine*. 2021;73:103630. doi:10.1016/j.ebiom.2021.103630
114. Vazeille C, Jouinot A, Durand JP, et al. Relation between hypermetabolism, cachexia, and survival in cancer patients: a prospective study in 390 cancer patients before initiation of anticancer therapy. *Am J Clin Nutr*. 2017;105(5):1139–1147. doi:10.3945/ajcn.116.140434
115. Hirsch L, Bellesoeur A, Boudou-Rouquette P, et al. The impact of body composition parameters on severe toxicity of nivolumab. *Eur J Cancer Oxf Engl*. 2020;124:170–177. doi:10.1016/j.ejca.2019.11.003
116. Derosa L, Hellmann MD, Spaziano M, et al. Negative association of antibiotics on clinical activity of immune checkpoint inhibitors in patients with advanced renal cell and non-small-cell lung cancer. *Ann Oncol*. 2018;29(6):1437–1444. doi:10.1093/annonc/mdy103
117. Pinato DJ, Howlett S, Ottaviani D, et al. Association of prior antibiotic treatment with survival and response to immune checkpoint inhibitor therapy in patients with cancer. *JAMA Oncol*. 2019;5(12):1774. doi:10.1001/jamaoncol.2019.2785
118. Elkrief A, Derosa L, Zitvogel L, Kroemer G, Routy B. The intimate relationship between gut microbiota and cancer immunotherapy. *Gut Microbes*. 2019;10(3):424–428. doi:10.1080/19490976.2018.1527167
119. Matson V, Chervin CS, Gajewski TF. Cancer and the microbiome—influence of the commensal microbiota on cancer, immune responses, and immunotherapy. *Gastroenterology*. 2021;160(2):600–613. doi:10.1053/j.gastro.2020.11.041
120. Davar D, Dzutsev AK, McCulloch JA, et al. Fecal microbiota transplant overcomes resistance to anti-PD-1 therapy in melanoma patients. *Science*. 2021;371(6529):595–602. doi:10.1126/science.abf3363
121. Dizman N, Meza L, Bergerot P, et al. Nivolumab plus ipilimumab with or without live bacterial supplementation in metastatic renal cell carcinoma: a randomized phase 1 trial. *Nat Med*. 2022;28(4):704–712. doi:10.1038/s41591-022-01694-6
122. Pal SK, Li SM, Wu X, et al. Stool bacteriomic profiling in patients with metastatic renal cell carcinoma receiving vascular endothelial growth factor-tyrosine kinase inhibitors. *Clin Cancer Res off J Am Assoc Cancer Res*. 2015;21(23):5286–5293. doi:10.1158/1078-0432.CCR-15-0724
123. Ianiro G, Rossi E, Thomas AM, et al. Faecal microbiota transplantation for the treatment of diarrhoea induced by tyrosine-kinase inhibitors in patients with metastatic renal cell carcinoma. *Nat Commun*. 2020;11(1):4333. doi:10.1038/s41467-020-18127-y

124. Dizman N, Hsu J, Bergerot PG, et al. Randomized trial assessing impact of probiotic supplementation on gut microbiome and clinical outcome from targeted therapy in metastatic renal cell carcinoma. *Cancer Med.* 2021;10(1):79–86. doi:10.1002/cam4.3569
125. Mueller-Schoell A, Groenland SL, Scherf-Clavel O, et al. Therapeutic drug monitoring of oral targeted antineoplastic drugs. *Eur J Clin Pharmacol.* 2021;77(4):441–464. doi:10.1007/s00228-020-03014-8
126. Bardin C, Veal G, Paci A, et al. Therapeutic drug monitoring in cancer—are we missing a trick? *Eur J Cancer Oxf Engl.* 2014;50(12):2005–2009. doi:10.1016/j.ejca.2014.04.013
127. Cerbone L, Combarel D, Geraud A, et al. Association of cabozantinib pharmacokinetics, progression and toxicity in metastatic renal cell carcinoma patients: results from a pharmacokinetics/pharmacodynamics study. *ESMO Open.* 2021;6(6):100312. doi:10.1016/j.esmoop.2021.100312
128. Cabel L, Blanchet B, Thomas-Schoemann A, et al. Drug monitoring of sunitinib in patients with advanced solid tumors: a monocentric observational French study. *Fundam Clin Pharmacol.* 2018;32(1):98–107. doi:10.1111/fcp.12327
129. Beinse G, Hulin A, Rousseau B. Axitinib pharmacologic therapeutic monitoring reveals severe under-exposure despite titration in patients with metastatic renal cell carcinoma. *Invest New Drugs.* 2019;37(6):1289–1291. doi:10.1007/s10637-019-00743-1
130. Bellesoeur A, Nagai H, Matsui D. Prevalence of drug–drug interactions in sarcoma patients: key role of the pharmacist integration for toxicity risk management. *Cancer Chemother Pharmacol.* 2021;89(1):11. doi:10.1007/s00280-021-04359-2
131. Bellesoeur A, Ollier E, Allard M, et al. Is there an exposure–response relationship for nivolumab in real-world NSCLC patients?. *Cancers.* 2019;16.

Therapeutics and Clinical Risk Management

Dovepress

Publish your work in this journal

Therapeutics and Clinical Risk Management is an international, peer-reviewed journal of clinical therapeutics and risk management, focusing on concise rapid reporting of clinical studies in all therapeutic areas, outcomes, safety, and programs for the effective, safe, and sustained use of medicines. This journal is indexed on PubMed Central, CAS, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/therapeutics-and-clinical-risk-management-journal>