

Increasing arbovirus risk in Chile and neighboring countries in the Southern Cone of South America

Elizabeth L. Estallo,^{a,b,*} Rachel Sippy,^c Michael A. Robert,^d Salvador Ayala,^e Carlos J. Barboza Pizarra,^f Pastor E. Pérez-Estigarribia,^g and Anna M. Stewart-Ibarra^h

^aUniversidad Nacional de Córdoba, Facultad de Ciencias Exactas, Físicas y Naturales, Centro de Investigaciones Entomológicas de Córdoba, Córdoba, Argentina

^bInstituto de Investigaciones Biológicas y Tecnológicas, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de Córdoba, Córdoba, Argentina

^cDepartment of Psychiatry, University of Cambridge, Cambridge, England, United Kingdom

^dDepartment of Mathematics and Center for Emerging, Zoonotic, and Arthropod-borne Pathogens (CeZAP), Virginia Tech, Blacksburg, VA, USA

^eInstituto de Salud Pública de Chile, Santiago, Chile

^fMinisterio de Salud Pública de Uruguay, Montevideo, Uruguay

^gPolytechnic School, National University of Asunción, San Lorenzo, Paraguay

^hInter-American Institute for Global Change Research, Montevideo, Uruguay

On April 26, 2023, Chile declared a public health emergency in its northern and central regions in response to the presence of *Aedes aegypti* and the epidemiological situation of neighboring countries.¹ *Aedes aegypti* is a highly efficient vector of dengue, chikungunya, Zika, and other arboviruses. The mosquito recently expanded within Parinacota (northern continental Chile) and was detected in Los Andes, Valparaíso Region (32°50'S, 70°32'W, elevation ~800 m), a location with no previous history of the vector. The emergency declaration also noted the proliferation of *Anopheles pseudopuntipenis*, a malaria vector, in Pica, Tarapacá region (20°28'S, 69°18'W, elevation 1320 m).

Continental Chile faces an impending risk of vector-borne disease emergence. *Aedes aegypti* was first detected on Easter Island in 2000 and in continental Chile in 2015, Arica region, 18°27'S, 70°21'W, elevation 200 m). In March 2002, the first dengue case was confirmed in an Easter Island resident with no travel history.² In the past decade, the frequency and magnitude of dengue outbreaks have increased in neighboring countries, and cases are regularly imported to continental Chile (28 cases to date, 2023), increasing the risk of local transmission. Continental Chile remains the only South American country without autochthonous dengue transmission.

Since 2022, dengue and chikungunya cases have escalated across the Americas,³ including the temperate southern cone, the range limit of transmission. Concurrent dengue and chikungunya outbreaks are ongoing in Argentina. Due to heightened arbovirus activity,

Argentina's Ministry of Health declared a public health emergency on April 18, 2023. Between December 2022 and May 2023 (EW 20/2023), 106,672 dengue cases were reported, with 99,456 autochthonous cases and 59 deaths.⁴

In 2023 Argentina and Uruguay reported the first instances of local chikungunya transmission.³ In Argentina, 1993 chikungunya cases were detected (EW 52/2022 to 20/2023), with 1336 autochthonous cases, while Uruguay reported 73 chikungunya cases (52 autochthonous).

Paraguay has also experienced recent large-scale arboviral epidemics,⁵ with a major dengue epidemic in late 2019–early 2020. The country is currently experiencing an explosive chikungunya epidemic. This outbreak, driven by the East-Central-South-Africa (ECSA) genotype, has alarmed all countries in the region as it is among the largest reported in the Americas.

Climate change is a critical driver of arboviral transmission. Rising temperatures,⁶ shifting rainfall and humidity,⁶ and the genetic plasticity of *Ae. aegypti* facilitate the ongoing expansion of the mosquito's range.⁷ Global forecasting models predict an expansion of vector distribution and disease transmission under all climate change scenarios. Arbovirus transmission responds to ambient temperature due to the thermal biology of mosquitoes.⁸ Climate change is projected to increase net and new exposures to *Aedes*-borne viruses, with the most extreme increase in *Ae. albopictus*-borne transmission under intermediate climate change scenarios; more severe scenarios are expected to produce poleward shifts in arboviruses and broader exposure of humans to *Ae. aegypti*.⁸ Shifting distributions are aggravated by insecticide resistance in vectors. Several countries are exploring new biotechnologies for control.⁹

Regional cooperation—including sharing data, surveillance and control tools, expertise, and supplies—is urgently needed to respond to arboviruses. South



The Lancet Regional Health - Americas 2023;23: 100542

Published Online xxx
<https://doi.org/10.1016/j.lana.2023.100542>

*Corresponding author. Universidad Nacional de Córdoba, Facultad de Ciencias Exactas, Físicas y Naturales, Centro de Investigaciones Entomológicas de Córdoba, Córdoba, Argentina.

E-mail address: elizabeth.estallo@mi.unc.edu.ar (E.L. Estallo).

© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

American countries have a history of transnational cooperation in response to health emergencies. In the mid-20th century, a unanimously-supported campaign led by national vector control services, the Pan American Health Organization (PAHO), and ministries of health led to the eradication of *Ae. aegypti* throughout most of South America.¹⁰ Another example is the Southern Cone Initiative formed in 1991 by Argentina, Bolivia, Brazil, Chile, Paraguay, and Uruguay to control Chagas disease (*Triatoma* vectors), reducing the disease.⁹ National collaboration between malaria control programs along the coastal border of Ecuador–Peru during the mid-1990s–2010s led to local malaria elimination. In 2003, PAHO introduced an integrated management plan for *Ae. aegypti* across sectors with coordination among sub-regions.⁹ The resurgence and emergence of *Aedes* vectors and viral epidemics over the last 20 years is a clear call to action. Bi-national and multinational collaboration is needed to respond to this public health emergency.

As globalization, urbanization, and climate change alter our communities, arbovirus importation and subsequent local transmission will continue. Addressing these challenges requires a transdisciplinary systems-thinking approach, uniting scientists from diverse disciplines, public health practitioners, urban planners, civil society, and the private sector to co-develop solutions. Equitable collaborations can facilitate the sharing of resources across nations with similar challenges, including the Southern Cone. These partnerships have the potential to generate the tools and evidence that are urgently needed to guide policies that protect the health of populations worldwide.

“Eradication campaigns cannot be restricted by the boundaries of any one country.”

(Camargo, 1967).¹⁰

Contributors

ELE, RS, MAR and AMSE: conceptualisation, writing - original draft, writing - review & editing; SA, CJPB and PEPE: writing - review & editing. All authors have read and agreed with the final manuscript submitted.

Declaration of interests

The authors state that no conflicts of interest exist.

Acknowledgments

ELE is a researcher at the National Scientific and Technical Research Council–Argentina (CONICET). ELE and MAR are supported in part by a Burroughs Wellcome Fund Climate and Health Interdisciplinary Award (BWF #1022621).

References

- 1 Decreto 12. Ministerio de Salud, Gobierno de Chile; 2023. <https://senapred.cl/se-declara-alerta-temprana-preventiva-para-la-region-de-atacama-por-alteracion-sanitaria/>. Accessed May 10, 2023.
- 2 Perret C, Abarca K, Ovalle J, et al. Dengue-1 virus isolation during first dengue fever outbreak on Easter Island, Chile. *Emerg Infect Dis*. 2003;9:1465–1467.
- 3 Expansión geográfica de los casos de dengue y chikungunya más allá de las áreas históricas de transmisión en la Región de las Américas. WHO; 2023. <https://www.who.int/es/emergencias/disease-outbreak-news/item/2023-DON448>. Accessed May 31, 2023.
- 4 Boletín epidemiológico 654, SE 20. Ministerio de Salud de la Nación; 2023. <https://bancos.salud.gob.ar/recurso/boletin-epidemiologico-nacional-654-se-20>. Accessed May 29, 2023.
- 5 Sequera G. ¿Por qué esta gran epidemia de Chikungunya? ¿Qué paso del Dengue? *An Fac Cienc Méd (Asunción)*. 2023;56:19–24.
- 6 Robert MA, Stewart-Ibarra AM, Estallo EL. Climate change and viral emergence: evidence from *Aedes*-borne arboviruses. *Curr Opin Virol*. 2020;40:41–47.
- 7 Obholz G, San Blas G, Fischer S, Diaz A. Winter survival of *Aedes aegypti* (Diptera: Culicidae) eggs at its southern limit distribution. *Acta Trop*. 2022;231:106471.
- 8 Colón-González FJ, Sewe MO, Tompkins AM, et al. Projecting the risk of mosquito-borne diseases in a warmer and more populated world: a multi-model, multi-scenario intercomparison modelling study. *Lancet Planet Health*. 2021;5:e404–e414.
- 9 Wilson AL, Courtenay O, Kelly-Hope LA, et al. The Importance of vector control for the control and elimination of vector-borne diseases. *PLoS Neglected Trop Dis*. 2020;14(1):e0007831.
- 10 Camargo S. History of *Aedes aegypti* eradication in the Americas. *Bull World Health Organ*. 1967;36:602–603.