

# Disseminated gonorrhoea with laryngeal involvement in a 25-year-old man

François Voruz MD, Igor Leuchter MD

■ Cite as: *CMAJ* 2021 May 3;193:E646. doi: 10.1503/cmaj.202183


**A** healthy immunocompetent 25-year-old married man presented to the emergency department with a 5-day history of severe odynophagia, fever (peak 40°C), and a scattered pustular skin eruption. He denied extramarital sexual relations. The patient's physical examination was remarkable for scattered fibrinous lesions in his pharynx and larynx (Figure 1; Appendix 1 video, available at [www.cmaj.ca/lookup/doi/10.1503/cmaj.202183/tab-related-content](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.202183/tab-related-content)) and pustular lesions on the trunk, limbs and scrotum. He did not have urethritis.

Polymerase chain reaction of a skin pustule identified *Neisseria gonorrhoeae*, consistent with a diagnosis of disseminated gonococcal infection with laryngeal involvement. Investigations for HIV, syphilis, chlamydia and herpes simplex infection were negative. The patient was admitted to hospital and received intravenous fluids, nasogastric tube feeding, fentanyl and intravenous ceftriaxone, after consultation with our infectious disease team. The lesions resolved in 9 days. His spouse tested negative and was not treated for gonorrhoea. We were unable to determine the source of his infection but concluded that asymptomatic gonococcal carriage was a plausible explanation.

Gonococcal infection is a sexually transmitted disease contracted during vaginal, anal or oral sex that largely affects mucosal surfaces and manifests mostly as a localized genitourinary tract infection.<sup>1</sup> In 2017, the incidence of gonorrhoea in Canada was 79.5/100 000, which was more than double the 2013 incidence.<sup>2</sup>

Disseminated infection occurs through hematogenous spreading of *N. gonorrhoeae*, and usually manifests as tenosynovitis, arthritis, skin lesions and, very rarely, meningitis, endocarditis and osteomyelitis.<sup>3</sup> Pharyngeal involvement occurs in about 3% of infections and is often asymptomatic, with oral sex being the main risk factor.<sup>4,5</sup> The differential diagnoses include infectious and reactive arthritis, herpes simplex, HIV and syphilis infections.

Because of penicillin and fluoroquinolone resistance, third-generation cephalosporins, such as ceftriaxone, are recommended, although resistant strains are emerging and becoming problematic.<sup>1</sup> When assessing cutaneous lesions, consider sexually transmitted infections, take a thorough sexual history and examine for atypical systemic manifestations, such as pharyngolaryngitis.



**Figure 1:** (A) Pharynx of a 25-year-old man with scattered whitish macules with perilesional erythema on the palate, uvula and tonsils. (B) Larynx with the fibrinous scattered lesions involving the surrounding structures, without vocal folds edema. Note: Yellow arrows = arytenoids, green arrow = left vocal fold, black arrow = epiglottis.

## References

- Hill SA, Masters TL, Wachter J. Gonorrhoea — an evolving disease of the new millennium. *Microb Cell* 2016;3:371-89.
- Report on sexually transmitted infections in Canada, 2017. Gonorrhoea. Ottawa: Public Health Agency of Canada; modified 2020 Jan. 27. Available: [www.canada.ca/en/public-health/services/publications/diseases-conditions/report-sexually-transmitted-infections-canada-2017.html#a5](http://www.canada.ca/en/public-health/services/publications/diseases-conditions/report-sexually-transmitted-infections-canada-2017.html#a5) (accessed 2020 Oct. 29).
- Bleich AT, Sheffield JS, Wendel GD Jr., et al. Disseminated gonococcal infection in women. *Obstet Gynecol* 2012;119:597-602.
- Chan PA, Robinette A, Montgomery M, et al. Extragenital infections caused by *Chlamydia trachomatis* and *Neisseria gonorrhoeae*: a review of the literature. *Infect Dis Obstet Gynecol* 2016;2016:5758387.
- Balmelli C, Gunthard HF. Gonococcal tonsillar infection — a case report and literature review. *Infection* 2003;31:362-5.

**Competing interests:** None declared.

This article has been peer reviewed.

The authors have obtained patient consent.

**Affiliation:** Otorhinolaryngology and Head and Neck Surgery, Geneva University Hospitals, Switzerland

**Content licence:** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NCND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is non-commercial (i.e., research or educational use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Correspondence to:** François Voruz, [francois.voruz@hcuge.ch](mailto:francois.voruz@hcuge.ch)



Please see the accompanying video online, “Disseminated gonorrhoea: scattered fibrinous lesions in the pharynx and larynx,” available at [www.cmaj.ca/lookup/doi/10.1503/cmaj.202183/tab-related-content](http://www.cmaj.ca/lookup/doi/10.1503/cmaj.202183/tab-related-content)