


Article

Virus-like Vesicles Expressing Multiple Antigens for Immunotherapy of Chronic Hepatitis B



Timur O.
Yarovinsky,
Stephen W.
Mason, Manisha
Menon, ..., John K.
Rose, Michael D.
Robek, Valerian
Nakaar

tyarovinsky@carogencorp.com
(T.O.)
vnakaar@carogencorp.com
(V.N.)

HIGHLIGHTS
Virus-like vesicles (VLV)
designed to express three
HBV antigens from a
single vector

Expression of HBV
polymerase, core, and
middle S antigen-induced
specific CD8⁺ T cells

A single immunization
protects mice from acute
HBV infection

VLV prime and DNA boost
decreased HBV
replication in a model of
chronic hepatitis B

Yarovinsky et al., iScience 21,
391–402
November 22, 2019 © 2019
The Authors.
<https://doi.org/10.1016/j.isci.2019.10.040>



Article

Virus-like Vesicles Expressing Multiple Antigens for Immunotherapy of Chronic Hepatitis B

Timur O. Yarovinsky,^{1,2,4,5,*} Stephen W. Mason,^{1,4} Manisha Menon,¹ Marie M. Krady,¹ Maria Haslip,¹ Bhaskara R. Madina,¹ Xianyong Ma,¹ Safiehkhatoon Moshkani,³ Carolina Chiale,³ Anasuya Chattopadhyay Pal,² Bijan Almassian,¹ John K. Rose,² Michael D. Robek,³ and Valerian Nakaar^{1,*}

SUMMARY

Infections with hepatitis B virus (HBV) can initiate chronic hepatitis and liver injury, causing more than 600,000 deaths each year worldwide. Current treatments for chronic hepatitis B are inadequate and leave an unmet need for immunotherapeutic approaches. We designed virus-like vesicles (VLV) as self-amplifying RNA replicons expressing three HBV antigens (polymerase, core, and middle surface) from a single vector (HBV-VLV) to break immune exhaustion despite persistent HBV replication. The HBV-VLV induces HBV-specific T cells in naive mice and renders them resistant to acute challenge with HBV. Using a chronic model of HBV infection, we demonstrate efficacy of HBV-VLV priming in combination with DNA booster immunization, as 40% of treated mice showed a decline of serum HBV surface antigen below the detection limit and marked reduction in liver HBV RNA accompanied by induction of HBsAg-specific CD8 T cells. These results warrant further evaluation of HBV-VLV for immunotherapy of chronic hepatitis B.

INTRODUCTION

Human hepatitis B virus (HBV), a partially double-stranded DNA virus of the genus Orthohepadnavirus, replicates in hepatocytes and causes liver inflammation and injury (Ganem and Prince, 2004; Seeger and Mason, 2015; Tang et al., 2018). Although most adults recover from the acute infection and become immune to subsequent exposures, 2%–6% of adults and 90% of infants develop chronic hepatitis B (CHB) and are at increased risk of developing liver cirrhosis or liver cancer (Ganem and Prince, 2004; Tang et al., 2018). Multiple factors, including vertical transmission to newborns before the immune system is fully formed and active suppression of the immune system by hepatitis B surface antigen (HBsAg) and hepatitis B e antigen (HBeAg), contribute to HBV persistence (Tsai et al., 2018). Worldwide, 240–350 million people are chronically infected with HBV and more than 600,000 die each year from HBV-associated liver diseases. In the United States, CHB prevalence is estimated at 850,000 and the rate of new infections remains about 1 per 100,000 population despite routine use of a prophylactic HBV vaccine (Schillie et al., 2018). Currently approved therapeutic approaches for CHB, including pegylated interferon- α (IFN α) and nucleoside/nucleotide analogues, may reduce and control HBV replication in a subset of patients (Tang et al., 2018). However, adverse effects of IFN α treatment, lack of complete HBV clearance, and development of drug resistance after prolonged treatment with nucleosides/nucleotides emphasize the unmet need for alternative therapeutic approaches that can lead to functional cure (Block et al., 2018; Schinazi et al., 2018; Tang et al., 2018).

Spontaneous clearance of HBV through immune-mediated mechanisms, particularly T cell-mediated immune responses, in acute and in some chronically infected patients provides a rationale for CHB immunotherapy (Kosinska et al., 2017). Ongoing efforts at preclinical and early clinical stages include activation of innate immunity through TLR ligands, treatment with immune checkpoint inhibitors, and/or immunization with HBV antigens delivered by plasmid DNA or viral vectors (Kosinska et al., 2017; Schinazi et al., 2018). DNA immunization is a safe and reliable method for induction of antibody responses, but this approach requires special delivery such as electroporation, and remains poorly immunogenic for T cells (Cova, 2017). On the other hand, viral vectors are highly immunogenic and efficient at mounting T cell responses, but their safety remains a concern even for attenuated and replication-deficient viral vectors (Kosinska et al., 2017).

Virus-like vesicles (VLVs), a form of self-amplifying RNA replicons, are a non-pathogenic and immunogenic alternative to viral vectors. VLVs have an excellent safety profile and induce T cell and antibody responses

¹CaroGen Corporation, Farmington, CT 06032, USA

²Department of Pathology, Yale University School of Medicine, New Haven, CT 06510, USA

³Department of Immunology and Microbial Disease, Albany Medical College, Albany, NY 12208, USA

⁴These authors contributed equally

⁵Lead Contact

*Correspondence:
tyarovinsky@carogenCorp.com (T.O.Y.),
vnaaar@carogenCorp.com (V.N.)

<https://doi.org/10.1016/j.isci.2019.10.040>



(Rose et al., 2014, 2008; Schell et al., 2011). The RNA-dependent RNA polymerase from Semliki Forest virus and glycoprotein from vesicular stomatitis virus (VSV-G) expressed from the same vector are essential components of the VLV platform that enables rapid and efficient production of VLVs in BHK-21 cells and expression of the desired antigens in the immunized host (Rose et al., 2014). A previous study showed that immunization with VLVs expressing HBV middle surface antigen (MHBs) induced HBV-specific CD8⁺ T cells and protected mice against experimental acute HBV infection (Reynolds et al., 2015). Furthermore, HBV-transgenic mice with low HBeAg produced HBV-specific CD8⁺ T cells after priming with plasmid DNA and boosting with the VLV expressing MHBs, suggesting that VLV immunization may break immune tolerance, a characteristic feature of CHB (Reynolds et al., 2015). Although that study demonstrated promise for VLV-based immunotherapy, it is evident from other studies that immunization with multiple HBV antigens to enhance and/or broaden T cell responses is necessary for therapeutic effect and HBV elimination (King et al., 2014; Martin et al., 2014; Obeng-Adjei et al., 2013).

In this report, we describe the design, production, immunogenicity, and immunotherapeutic potential of a polycistronic VLV vector that expresses three HBV antigens: polymerase (Pol), core (HBcAg), and MHBs. Our goal was to target multiple HBV antigens simultaneously and to elicit the broad and vigorous multi-specific immune responses needed for the resolution of CHB infection in humans. To that end, we co-expressed structural (MHBs and HBcAg) and non-structural (Pol) HBV proteins as a single polycistronic open reading frame. This approach enabled the induction of broad T cell responses to multiple antigens that were sufficient to protect mice from experimental acute HBV replication. In a murine model of experimental CHB driven by AAV-HBV transduction, prime immunization with the multiantigen VLV vector followed by two booster DNA immunizations induced HBsAg-specific CD8⁺ T cell responses and reduced serum HBsAg and liver HBV RNA.

RESULTS

VLV Design and Production

Because HBV Pol and HBcAg are important targets for CD8⁺ T cells (Akbar et al., 2012; Mizukoshi et al., 2004; Zhuo et al., 2015), we aimed to incorporate these antigens into the new VLV vectors for immunization. The previously reported VLV vector (referred to as MT2A and depicted in Figure 1A) relied on the ribosome-skipping 2A peptide from *Thosaea asigna* virus (T2A) to express both MHBs and VSV-G from the same subgenomic RNA (Reynolds et al., 2015). Using that vector as a backbone, we designed and constructed VLVs for expression of HBV polymerase alone (Polt2A) or as a polycistronic vector for expression of Pol, HBcAg, and MHBs by interspersing the HBV antigens with the ribosome skipping peptides (3xT2A) as depicted in Figure 1A. We replaced several key amino acids in the terminal protein domain of Pol (Y63A, W74A, Y147A, and Y173A) to abolish its DNA binding (primase) activity (Lanford et al., 1999). In addition, we deleted amino acids 538–544 encompassing the active site (residues YMDD) of the reverse transcriptase domain to remove its RNA-dependent DNA polymerase and DNA-dependent DNA polymerase activities. Furthermore, we introduced mutations into the RNase H domain (amino acids 680–832) in order to abolish the ribonuclease H activity, as previously described (Kim et al., 1999). We altered the HBcAg sequence to improve processing and peptide presentation by substituting the disulfide bond-forming cysteine residues (C48S, C61S, C107S and C183S) and adding an N-terminal fusion with the 76-amino acid ubiquitin (Ub) peptide (Nassal et al., 1992; Zhuo et al., 2015). To produce VLVs for evaluation of antigen expression and immunogenicity, we transfected BHK-21 cells with the plasmid DNA expressing 3xT2A VLV RNA under the control of CMV promoter, collected the master VLV stocks, and used them for propagation and concentration of VLVs by ultrafiltration. The resulting VLV titers routinely exceeded 10⁹ pfu/mL.

To mitigate potential risks of homologous recombination of the repeating T2A peptide sequences in the 3xT2A construct during VLV replication, we designed and evaluated an alternative construct, Mix2A, in which we replaced the two T2A peptides with the porcine teschovirus-1 (P2A) and equine rhinitis A virus (E2A) peptides (Liu et al., 2017) (Figure 1A). These peptides are structurally similar, but encoded by different nucleotide sequences. When produced in BHK-21 cells under identical conditions, the resulting titers were similar for Mix2A VLVs (2.2*10⁹ pfu/mL) and 3xT2A VLVs (1.8*10⁹ pfu/mL).

Validation of Antigen Expression

We evaluated expression of Pol and HBcAg in BHK-21 cells infected with PolT2A and 3xT2A VLVs by immunofluorescence (Figure 1B). Using HBV Pol specific mAb clone 2C8 (zu Putlitz et al., 1999), we observed a characteristic fine granular staining in the cytoplasm at 24 h after infection with either of the VLVs (Figure 1B,




Figure 1. Virus-like Vesicle Platform for Therapeutic HBV Vaccine Expressing Polymerase (Pol), Ubiquitinylated Core (Ub-HBc), and Middle Surface (MHBs) Antigens

(A) Design of replicating VLV for expression of MHBs (MT2A), HBV Pol (PoIT2A) single antigens, and for polycistronic expression of the three HBV antigens using different 2A self-cleaving peptides (3xT2A and Mix2A).

(B) Validation of antigen expression in BHK-21 cells after infection with PoIT2A or 3xT2A VLV (MOI = 1) by immunofluorescence at 20 h post infection using the antibodies specific for HBV Pol and HBcAg.

(C) Validation of antigen expression in BHK-21 cells after infection with VLV expressing MHBs only (MT2A), Pol only (PoIT2A), or the three antigens (3xT2A) by flow cytometry using antibodies specific for Pol, MHBs, HBcAg, and VSV-G. Geometric mean fluorescence intensity is shown for each antibody and VLV.

(D) Evaluation of antigen expression and VLV replication in BHK-21 cells transfected with plasmid DNA for 3xT2A or Mix2A VLV using 2A-peptide specific antibody in western blots. Anti-VSV-G and anti-actin were also used as controls. The data are representative of three independent experiments.

top row). Staining with the HBcAg-specific polyclonal antibody showed variable levels of HBcAg expression in most of the BHK-21 cells infected with 3xT2A, but not with the PoIT2A VLV (Figure 1B, second row).

To compare expression levels of HBV antigens in BHK-21 cells infected with polycistronic VLVs 3xT2A versus VLVs expressing Pol or MHBs alone, all at MOI = 1 at 24 h post infection, we used intracellular staining with antibodies for Pol, HBcAg, and MHBs, followed by flow cytometry analyses (Figure 1C). Cells infected with 3xT2A VLVs showed slightly higher levels of Pol expression, compared with PoIT2A VLVs. On the contrary, cells infected with 3xT2A VLVs showed slightly lower levels of MHBs expression compared with MT2A VLVs. As expected, no Pol staining was detected in cells infected with the MT2A VLVs and no MHBs staining was detected in cells infected with PoIT2A VLVs. Staining for HBcAg was detectable only in cells infected with 3xT2A, albeit at relatively low levels. It is important to note that VSV-G, the only structural component of VLVs, was detectable in most BHK-21 cells after infection with all three VLV vectors above (Figure 1C).

We also compared antigen expression from 3xT2A or Mix2A VLVs by immunoblotting of BHK-21 cell lysates at 24 h post-infection at MOI = 1 (Figure 1D). Using a polyclonal anti-2A peptide antibody, which recognizes the GDVESNPG epitope shared by all 2A peptides, we detected bands corresponding to the predicted molecular weights of HBV polymerase (92 kDa), middle S antigen (32 kDa), and modified core (29 kDa),

which is consistent with the expectation that each of the antigens is tagged with the 2A-peptide sequences at the C-terminus. We also observed additional bands at ~16, 18, and 40 kDa in the lysates from Mix2A VLV-infected cells, which may be due to incomplete recognition of the 2A peptide sequences during translation or subsequent proteolytic degradation. In addition, we noted a high molecular weight (>250 kDa) and a 37-kDa non-specific bands present in lysates from non-infected and VLV-infected cells, which most likely are the result of the antibody cross-reactivity with the endogenous proteins in BHK-21 cells. VSV-G was detected by polyclonal anti-VSV-G antibody only in the VLV-infected cells with comparable intensity between 3xT2A and Mix2A.

Immunogenicity of 3xT2A and Mix2A VLV Vaccine Candidates in Naive Mice

Administration of 3xT2A or Mix2A VLVs at 10^8 pfu/mouse resulted in an apparent increase in frequency (Figure 2A) and absolute numbers (Figure 2B) of CD44^{hi}CD62L^{lo} effector CD4⁺ and CD8⁺ T cells in spleens at day 7 post-immunization. Furthermore, we quantified the frequencies (Figure 2C) and absolute numbers (Figure 2D) of CD8⁺ T cells that produce IFN γ after ex vivo restimulation with the peptides derived from the epitopes from within HBV Pol, HBcAg, and MHBs. The cumulative frequencies of the IFN γ ⁺ (i.e., HBV-specific) CD8⁺ T cells exceeded by 2% among those the CD8⁺ T cells that recognized the HBS-371 peptide were most prevalent. Furthermore, substantial numbers of CD8⁺ T cells responded to HBV peptide stimulation by producing IFN γ and TNF (Figures 2E and 2F), indicating that VLV may induce double-cytokine producing CD8⁺ T cells, which are efficient in non-cytolytic elimination of HBV from the infected hepatocytes (Guidotti et al., 1996). The 3xT2A vector showed a trend toward higher frequency and absolute numbers of MHBs-specific CD8⁺ T cells when compared with Mix2A, albeit with no statistically significant difference. Since the use of different 2A peptides in the Mix2A VLV construct did not provide any apparent advantage for HBV antigen immunogenicity in naive mice, we proceeded to use the simpler of the two constructs, 3xT2A, for efficacy studies.

Immunization with 3xT2A VLV Protects Mice from Experimental Acute Challenge with AAV-HBV

To determine whether VLV-mediated expression of HBV Pol, HBcAg, and MHBs generates protective immunity against HBV, we immunized CB6F1/J mice with 3xT2A or empty vector VLV and administered an equivalent volume of PBS to a control group of mice. Six weeks later, we challenged these mice with HBV delivered to the liver via recombinant adeno-associated virus (AAV-HBV) encoding a 1.3-overlength HBV genome and monitored serum HBsAg and HBeAg as a measure of HBV replication for additional four weeks. HBsAg levels peaked in PBS and empty vector VLV control groups of mice 14 days after infection, but remained low in 3xT2A VLV-immunized mice throughout the experiment (Figure 3A). HBeAg levels steadily increased during infection in the PBS and empty vector VLV groups (Figure 3B) but barely rose above background at days 14 and 21 and declined by day 28 in 3xT2A VLV-immunized mice. Quantitative analyses of HBV RNA in the livers of mice euthanized on day 28 showed a statistically significant decrease in the 3xT2A VLV immunized group compared with empty vector VLV group (Figure 3C). 3xT2A VLV-immunized mice, but not mice immunized with the empty vector VLV, showed a statistically significant increase in HBsAg-specific cells in the spleens that produce IFN γ after stimulation with the dominant peptide epitopes (Figure 3D). Thus, immunization with the 3xT2A VLV expressing HBV Pol, HBcAg, and MHBs protects mice from experimental acute HBV replication.

Evaluation of 3xT2A VLV in an Experimental Model of Chronic HBV Infection

We explored the therapeutic potential of 3xT2A in the setting of the AAV-HBV mouse model of chronic HBV infection (Dion et al., 2013). To initiate this model, we transduced C57BL/6J male mice with AAV-HBV-1.2 (2×10^{11} genome copies of AAV-HBV particles per mouse) and monitored serum HBsAg levels every week. As expected, we observed a steady rise of serum HBsAg from 11.52 ± 2.40 ng/mL (mean \pm SEM, n = 20) at day 7 to 70.04 ± 6.25 ng/mL (mean \pm SEM, n = 20) at day 21 post-AAV-HBV transduction. Based on the measurements at day 21, we formed 2 groups of 10 mice each with equivalent levels of serum HBsAg prior to immunization with 3xT2A or empty vector VLV on day 28 post-AAV-HBV transduction. Figure 4A shows responses to the treatment through the full course of the experiment as the fold change of HBsAg levels (OD₄₅₀) relative to day 28, whereas Figure 4B shows the concentrations of serum HBsAg in ng/ml after VLV prime. Both VLV vectors caused an apparent, but transient, decline in HBsAg levels at day 35, which may reflect host interferon response to VLVs. HBsAg levels rebounded in both groups by day 42 and remained essentially steady for the empty vector VLV group. By day 49, the average HBsAg levels of the 3xT2A VLV-immunized groups started to decline and become significantly lower at days 56 and 63. Importantly, by day 63, the concentration of HBsAg dropped below the limit of detection (3.12 ng/mL) in four out




Figure 2. Comparison of Immunogenicity of the 3xT2A and Mix2A VLV in Naive C57BL/6J Mice

Accumulation of effector CD4⁺ and CD8⁺ T cells at day 7 after immunization with 3xT2A or Mix2A VLV.

(A) Representative plots for CD62L and CD44 staining after gating on CD4⁺ or CD8⁺ T cells.


(B) Absolute numbers of effector (CD44^{hi}CD62L^{lo}) T cells.

(C) Frequency of HBV-specific IFN γ -producing CD8⁺ T cells identified after ex vivo stimulation with the indicated peptides from Pol (HBP-44, HBP-396, HBP-419), HBcAg (HBC-93), and surface (HBS-353 and HBS-371) quantified by intracellular staining and flow cytometry.

(D) Numbers of IFN γ ⁺ CD8⁺ T cells per spleen.

(E) Frequency of HBV-specific double cytokine (TNF $^+$ IFN γ ⁺)-producing CD8⁺ T cells, respectively.

(F) Numbers of TNF $^+$ IFN γ ⁺-double positive CD8⁺ T cells. Values in (C and E) are mean of 3–5 mice per group.

**Figure 3. Efficacy of 3xT2A VLV as Prophylactic Vaccine in Acute HBV Infection**

CB6F1 mice were immunized with 3xT2A VLV or empty vector VLV for 6 weeks prior to challenge with AAV-HBV.

(A) Levels of serum HBsAg (OD450) measured by specific ELISA from blood samples taken weekly.

(B) Weekly levels of HBeAg. The values are mean \pm SEM ($n = 5$). Asterisks indicate significant difference (*, $p < 0.05$, **, $p < 0.01$) in 3xT2A-immunized mice compared with PBS or empty vector VLV.

Figure 3. Continued

(C) Abundance of HBV RNA in liver samples from mice euthanized on day 28 after challenge with AAV-HBV. TaqMan-based qRT-PCR was used to quantify HBV RNA and mouse GAPDH mRNA. The HBV RNA data were normalized to GAPDH and expressed as relative to the mean in the PBS group. Individual values and mean \pm SD are shown.

(D) Frequency of HBV-specific IFN γ ⁺ producing cells was measured by ELISPOT on day 28 after challenge with AAV-HBV after ex vivo stimulation of splenocytes with the indicated peptides. Individual values and mean \pm SD are shown.

of ten 3xT2A VLV-immunized mice. These results indicate that 40% of the chronically infected animals responded to the treatment with 3xT2A VLV.

Because only 40% of the animals showed responses to 3xT2A VLV and considering that a single VLV immunization would be unlikely to overcome HBV tolerance in CHB, we added DNA immunization as a boost after blood collection on days 65 and 84 (Figure 4A). We administered a pool of three DNA plasmids expressing MHBs, HBcAg, and HBV Pol in pVAC1 vector (further referred as HBVpDNA) to the group of mice immunized with 3xT2A and empty vector plasmid to the group of mice immunized with empty vector VLV. HBsAg levels continued to decline in the group of mice treated with 3xT2A VLV and HBVpDNA and, with the exception of a single time point (day 98), were significantly lower ($p < 0.05$) than in the group of mice treated with empty vector VLV and DNA (Figure 4C).

To determine whether 3xT2A VLV-induced decreases in serum HBsAg were accompanied by reduction of HBV RNA in liver, we euthanized five animals in each group to collect liver samples on day 70 and the remaining five animals at the termination of the experiment on day 119. Quantitative analyses of HBV RNA showed no difference between the groups at day 70 (Figure 5A). However, at day 119 three out of five mice primed with 3xT2A VLV and boosted twice with HBVpDNA showed more than 100-fold reduction of liver HBV RNA. Remarkably, this group showed a very strong correlation between serum HBsAg and liver HBV RNA ($R^2 = 0.905$, $p = 0.013$ for 3xT2A VLV plus HBVpDNA vs. $R^2 = 0.179$, $p = 0.478$ for empty vector VLV plus control DNA), suggesting that the observed decrease of serum HBsAg is due to a reduction of HBV replication.

Finally, we analyzed HBV-specific CD8⁺ T cell responses in the immunized groups. At day 70, there was no significant increase over background in the numbers of IFN γ -producing CD8⁺ T cells (Figure 5B). There was also no apparent increase in the numbers of IFN γ -producing CD8⁺ T cells after restimulation with HBV Pol or HBcAg peptides at day 119. However, restimulation with HBsAg peptides at the same time point resulted in a statistically significant increase over background in the 3xT2A VLV-primed/HBVpDNA-boosted group of mice (Figure 5C).

DISCUSSION

Our results demonstrate feasibility and potential of using the VLV platform for polycistronic expression of HBV antigens and immunotherapy for CHB. The HBV antigens expressed by 3xT2A and Mix2A VLV constitute three of the four major proteins expressed by HBV: the entire polymerase protein (HBV pol) with mutations to inactivate reverse transcriptase activity, ubiquitin-tagged core (UbHBc) with mutations to prevent the formation of stable disulfide-mediated dimers, and the middle surface antigen (MHBs). The combination of three different 2A peptides in Mix2A VLVs did not provide any obvious advantage over repeating T2A peptides in 3xT2A VLVs. The selected 3xT2A VLVs replicate to high titers ($>10^9$ pfu/mL), express the individual HBV proteins of interest (Figure 1), and induce slightly higher specific CD8⁺ T cell responses to each antigen than Mix2A VLV (Figure 2). Taking advantage of mouse strain-specific differences in persistence of AAV-mediated transduction of HBV (Huang et al., 2006), we demonstrated that a single immunization with 3xT2A VLVs provides protection of CB6F1 mice from an acute challenge (Figure 3). Using the AAV-HBV-transduced mice as a model of persistent HBV replication, we evaluated the utility of the 3xT2A VLV for CHB immunotherapy. Since persistent HBV replication in liver leads to tolerance to HBV antigens (Dion et al., 2013), we did not expect that a single immunization with 3xT2A VLV would be sufficient to clear the virus or induce high numbers of HBV-specific T cells. Thus, we used 3xT2A VLV for prime immunization followed by 2 booster immunizations with DNA plasmids expressing the same antigens. This immunization schedule resulted in sustained decline of serum HBsAg and liver HBV RNA in a subset of mice with established chronic HBV replication (Figures 4 and 5A). These data indicate that 3xT2A VLVs function not only as a prophylactic vaccine but could also be used as a component of comprehensive immunotherapy.

Immunotherapeutic approaches for CHB, including DNA vaccination and therapeutic vaccines based on viral vectors, promise to provide functional cure by reducing or eliminating the virus from infected hepatocytes (Ko-sinska et al., 2017; Schinazi et al., 2018). However, none of the experimental immunotherapies tested to date




Figure 4. Effects of 3xT2A VLV on HBsAg levels in Chronic AAV-HBV Model

Persistent HBV replication was established in C57BL/6 mice by delivery of HBV genome with AAV. The groups were balanced for HBsAg prior to immunization with empty vector VLV or 3xT2A at day 28 after AAV-HBV transduction. Booster DNA immunizations were carried out on days 65 and 84, as indicated by arrows, using a pool of 3 plasmids expressing MHBs, HBcAg, and HBV Pol for 3xT2A group or empty plasmid vectors for the empty vector VLV. (A–C) (A) Fold change in HBsAg levels (OD450). Values are mean \pm SEM ($n = 10$ for days 28–70 and $n = 5$ thereafter). Quantification of HBsAg (ng/mL) after VLV prime at day 28 (B) and DNA boost at day 65 (C). Individual values and median are shown by symbols and bars, respectively. Asterisks in A indicate significant difference between 3xT2A and empty vector VLV-immunized animals (*, $p < 0.05$; **, $p < 0.01$).

have shown sufficient efficacy in clinical trials (Kosinska et al., 2017). This suggests that these vaccines lacked sufficient potency, were not targeting the correct antigen(s), were targeting too few antigens or, perhaps, all the above. It appears that success depends on the induction of CD8⁺ T cells that are both cytolytic and non-cytolytic in action rather than induction of humoral responses by the present prophylactic vaccines (Kosinska et al., 2017; Webster et al., 2004). Although therapeutic vaccines using recombinant VSV Moshkani et al., 2019 or modified vaccinia virus have a potential to induce more robust T cell-mediated immune responses to HBV, concerns about their safety profile may represent an obstacle for clinical applications (Kosinska et al., 2017).

Immunotherapeutic vaccines expressing HBV Pol, HBcAg, and MHBs using VLV vectors address the safety concerns and provide sufficient efficacy by expressing multiple HBV antigens that are optimized for induction of T cell responses. Restriction of VLV replication by type I interferons enables 100% survival rate of mice even after intracranial administration of VLV (van den Pol et al., 2017). Inclusion of HBV Pol provides a number of highly immunogenic and conserved epitopes for CTL that can detect and eliminate HBV-infected cells and limit viral spread (Mizukoshi et al., 2004). Because expression of intact (wild-type) HBcAg from a single antigen-expressing VLV did not induce sufficient T cell responses to control HBV replication (Reynolds et al., 2015), the polycistronic constructs described in this report included ubiquitinated HBcAg with mutated cysteine residues. These changes may have improved peptide presentation to T cells through prevention of HBcAg dimerization and enhancement of proteolytic degradation (Zhuo et al., 2015). Even though MHBs are expressed in the third position in the 3xT2A and Mix2A vectors, it appeared as the dominant antigen, as demonstrated by the frequency of IFN γ ⁺ and IFN γ ⁺TNF⁺ CD8⁺ T cells after immunization (Figures 2C and 2D). Thus, the polycistronic design of the VLV vector induces broad T cell responses in naive mice with a potential to eliminate HBV through cytolytic and non-cytolytic mechanisms.

To evaluate prophylactic and therapeutic efficacy of 3xT2A VLV, we employed both acute and chronic models of HBV infection established by delivery of HBV genome DNA via liver-tropic AAV (Dion et al., 2013). This approach overcomes the species-specific restrictions in early steps in the HBV replication cycle and enables homogeneous transduction of mouse hepatocytes, HBV replication, and secretion of HBsAg (Dion et al., 2013). Delivery of HBV via AAV also provides more reproducible results than hydrodynamic delivery of HBV genome in plasmid DNA for acute challenge. Although the HBV-specific T cell responses are muted as the chronicity of infection causes T cell exhaustion and tolerance to HBV antigens, the mechanism and level of tolerance are different from those in the HBV transgenic mouse model (Dion et al., 2013; Huang et al., 2006; Sette et al., 2001). Thus, despite some limitations, including highly variable levels of viral load and HBsAg levels, the AAV-HBV model is more suitable for evaluation of immune-mediated clearance of HBV.

After determining that 3xT2A VLV can protect mice from acute AAV-HBV challenge (Figure 3), we evaluated the immunotherapeutic potency of 3xT2A in a mouse model of CHB. As a precaution that a single immunization with 3xT2A VLV might not be sufficient to break tolerance in the chronically infected mice, we added DNA booster immunization to the treatment protocol. We observed that priming of chronically infected mice with 3xT2A VLV, followed by booster DNA immunization, results in a statistically significant decline of serum HBsAg (Figure 4), with 40% of the mice dropping their HBsAg levels to the limit of detection and showing marked reduction of liver HBV RNA (Figure 5A). Such a partial response is not unheard of with immunotherapeutic approaches and may be due to variable levels of HBV replication or differences in the TCR repertoire of T cells responding to the viral antigens. Following this immunization schedule, a subset of mice also showed a statistically significant accumulation of HBsAg-specific IFN γ -producing CD8⁺ T cells in the spleens at later time points (Figures 5B and 5C). Although the design of the study described here did not allow us to be certain that the booster DNA immunizations would be necessary, our data clearly indicate that the multiantigen HBV-VLV shows promise as an immunotherapeutic tool for treatment of CHB.

Among other therapeutic vaccines for CHB evaluated in the HBV-AAV model, the adenovirus serotype 5 (Ad5)-based therapeutic vaccine (TG1050) is comparable to our VLV-based vector in approach and goals




Figure 5. Effects of 3xT2A VLV on Liver HBV RNA in Chronic AAV-HBV Model

Liver samples were collected at day 70 or day 119 from the animals immunized as shown in Figure 4 for qRT-PCR analyses. (A) Abundance of HBV RNA was determined after normalization to mouse GAPDH mRNA and expressed relative to the mean of the empty vector VLV group ($n = 5$). Numbers of IFN γ $^{+}$ CD8 $^{+}$ T cells per spleen at day 70 (B) and day 119 (C) were determined by intracellular flow cytometry after stimulation with HBV antigen peptides.

of inducing HBV-specific T cell responses (Martin et al., 2014). TG1050 expresses a fusion protein consisting of a deleted and mutated HBVpol, truncated HBcAg, and two regions of HBsAg (Martin et al., 2014). Similar to our observations with 3xT2A VLVs, the TG1050-treated group of mice showed a three-fold reduction in

HBsAg levels compared to those in the control group. While the effects of TG1050 on individual animals were not reported, we observed that 40% of the animals in our study responded by decline of HBsAg to the limit of detection. TG1050 presently is in Phase I/II clinical trials (NCT02428400) in HBV-infected patients with the results of this trial expected in 2019.

In order to improve immunogenicity and efficacy of therapeutic vaccine candidates for chronic HBV infection, others have used a combination of approaches, such as adjuvants, addition of checkpoint inhibitors, and prime/boost regimens (Kosinska et al., 2017; Schinazi et al., 2018). Recent studies showed that an optimized, heterologous prime/boost strategy produced strong T and B responses to HBV in various preclinical models of CHB (Backes et al., 2016; Chuai et al., 2017; Liu et al., 2014). A combination of an HBsAg/HBcAg protein prime with TLR9 agonists followed by a boost with non-replicating modified vaccinia Ankara viruses, optimized for higher level expression of the same antigens, was able to overcome the high-level tolerance observed in HBV-transgenic mice (Backes et al., 2016).

Given the evidence of high efficacy in at least 40% of the chronically infected mice, we plan to develop the VLV-based immunotherapy further and initiate clinical trials to demonstrate safety and efficacy in CHB patients. Further improvement of VLV-based immunotherapy may include establishing a prime-boost strategy and/or combination with nucleoside/nucleotide treatment to lower HBV replication.

Limitations of the Study

Although the AAV-HBV model of persistent HBV replication is widely accepted as the state-of-the-art system to study immune-mediated clearance of HBV, its limitations include lack of reinfection of the host with circulating HBV and differences in mouse and human immune systems.

METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2019.10.040>.

ACKNOWLEDGMENTS

We thank Lorraine Apuzzo, Mehdi Pouresmail, and Seyedehfatemeh Fazlhashemi at CaroGen Corporation for assistance with DNA cloning, production of VLV and characterization of the VLVs. We are also grateful to Drs. Raj Kalker, Kevin Walters, Krista Salley, Elizabeth Janci Wonderlich, and Jennifer Carl at Southern Research for their efforts on the chronic AAV-HBV model and to Shuguang Huang at Stat4ward for assistance with statistical analyses. This study was funded by Connecticut Innovations, Connecticut Bioscience Innovation Fund, Gerpang Healthcare, and NIH grants R43DK113858 (V.N.) and R01AI124006 (M.D.R.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

Conceptualization, T.O.Y., S.W.M., B.A., J.K.R., M.D.R., and V.N.; Methodology, T.O.Y., S.W.M., M.D.R., and V.N.; Investigation, T.O.Y., M.M., M.M.K., M.H., B.R.M., X.M., S.M., C.C., and A.C.P.; Formal Analysis, T.O.Y., and S.W.M.; Writing – Original Draft, T.O.Y., and S.W.M.; Writing – Review & Editing, T.O.Y., S.W.M., B.A., J.K.R., M.D.R., and V.N.; Funding Acquisition, B.A., M.D.R., and V.N.; Supervision, S.W.M., B.A., J.K.R., M.D.R., and V.N.

DECLARATION OF INTERESTS

T.O.Y., S.W.M., B.A., J.K.R., M.D.R., and V.N. have financial relationships with CaroGen Corporation.

Received: June 28, 2019

Revised: September 24, 2019

Accepted: October 21, 2019

Published: November 22, 2019

REFERENCES

- Akbar, S.M.F., Chen, S., Al-Mahtab, M., Abe, M., Hiasa, Y., and Onji, M. (2012). Strong and multi-antigen specific immunity by hepatitis B core antigen (HBcAg)-based vaccines in a murine model of chronic hepatitis B: HBcAg is a candidate for a therapeutic vaccine against hepatitis B virus. *Antivir. Res.* 96, 59–64.
- Backes, S., Jäger, C., Dembek, C.J., Kosinska, A.D., Bauer, T., Stephan, A.-S., Dißlers, A., Mutwiri, G., Busch, D.H., Babiuk, L.A., et al. (2016). Protein-prime/modified vaccinia virus Ankara vector-boost vaccination overcomes tolerance in high-antigenemic HBV-transgenic mice. *Vaccine* 34, 923–932.
- Block, T.M., Alter, H., Brown, N., Brownstein, A., Brosgart, C., Chang, K.-M., Chen, P.-J., Cohen, C., El-Serag, H., Feld, J., et al. (2018). Research priorities for the discovery of a cure for chronic hepatitis B: report of a workshop. *Antivir. Res.* 150, 93–100.
- Chuai, X., Chen, P., Chen, H., Wang, W., Deng, Y., Ruan, L., Li, W., and Tan, W. (2017). Protective efficacy and hepatitis B virus clearance in mice enhanced by cell-mediated immunity with novel prime-boost regimens. *J. Viral Hepat.* 24, 337–345.
- Cova, L. (2017). Present and future DNA vaccines for chronic hepatitis B treatment. *Expert Opin. Biol. Ther.* 17, 185–195.
- Dion, S., Bourgine, M., Godon, O., Levillayer, F., and Michel, M.-L. (2013). Adeno-associated virus-mediated gene transfer leads to persistent hepatitis B virus replication in mice expressing HLA-A2 and HLA-DR1 molecules. *J. Virol.* 87, 5554–5563.
- Ganem, D., and Prince, A.M. (2004). Hepatitis B virus infection — natural history and clinical consequences. *N. Engl. J. Med.* 350, 1118–1129.
- Guidotti, L.G., Ishikawa, T., Hobbs, M.V., Matzke, B., Schreiber, R., and Chisari, F.V. (1996). Intracellular inactivation of the hepatitis B virus by cytotoxic T lymphocytes. *Immunity* 4, 25–36.
- Huang, L.-R., Wu, H.-L., Chen, P.-J., and Chen, D.-S. (2006). An immunocompetent mouse model for the tolerance of human chronic hepatitis B virus infection. *Proc. Natl. Acad. Sci. U S A* 103, 17862–17867.
- Kim, Y., Hong, Y.B., and Jung, G. (1999). Hepatitis B virus: DNA polymerase activity of deletion mutants. *Biochem. Mol. Biol. Int.* 47, 301–308.
- King, T.H., Kemmler, C.B., Guo, Z., Mann, D., Lu, Y., Coeshott, C., Gehring, A.J., Bertolletti, A., Ho, Z.Z., Delaney, W., et al. (2014). A whole recombinant yeast-based therapeutic vaccine elicits HBV X, S and Core specific T cells in mice and activates human T cells recognizing epitopes linked to viral clearance. *PLoS One* 9, e101904.
- Kosinska, A.D., Bauer, T., and Protzer, U. (2017). Therapeutic vaccination for chronic hepatitis B. *Curr. Opin. Virol.* 23, 75–81.
- Lanford, R.E., Kim, Y.H., Lee, H., Notvall, L., and Beames, B. (1999). Mapping of the hepatitis B virus reverse transcriptase TP and RT domains by transcomplementation for nucleotide priming and by protein-protein interaction. *J. Virol.* 73, 1885–1893.
- Liu, J., Kosinska, A., Lu, M., and Roggendorf, M. (2014). New therapeutic vaccination strategies for the treatment of chronic hepatitis B. *Virol. Sin.* 29, 10–16.
- Liu, Z., Chen, O., Wall, J.B.J., Zheng, M., Zhou, Y., Wang, L., Ruth Vaseghi, H., Qian, L., and Liu, J. (2017). Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector. *Sci. Rep.* 7, <https://doi.org/10.1038/s41598-017-02460-2>.
- Martin, P., Dubois, C., Jacquier, E., Dion, S., Mancini-Bourgine, M., Godon, O., Kratzer, R., Lelu-Santolaria, K., Evlachev, A., Meritet, J.-F., et al. (2014). TG1050, an immunotherapeutic to treat chronic hepatitis B, induces robust T cells and exerts an antiviral effect in HBV-persistent mice. *Gut* 64, 1961–1971.
- Mizukoshi, E., Sidney, J., Livingston, B., Ghany, M., Hoofnagle, J.H., Sette, A., and Rehermann, B. (2004). Cellular immune responses to the hepatitis B virus polymerase. *J. Immunol.* 173, 5863–5871.
- Moshkani, S., Chiale, C., Lang, S.M., Rose, J.K., and Robek, M.D. (2019). A highly attenuated vesicular stomatitis virus-based vaccine platform controls hepatitis B virus replication in mouse models of hepatitis B. *J. Virol.* 93, e01586–18.
- Nassal, M., Rieger, A., and Steinau, O. (1992). Topological analysis of the hepatitis B virus core particle by cysteine-cysteine cross-linking. *J. Mol. Biol.* 225, 1013–1025.
- Obeng-Adjei, N., Hutnick, N.A., Yan, J., Chu, J.S., Myles, D.J.F., Morrow, M.P., Sardesai, N.Y., and Weiner, D.B. (2013). DNA vaccine cocktail expressing genotype A and C HBV surface and consensus core antigens generates robust cytotoxic and antibody responses in mice and Rhesus macaques. *Cancer Gene Ther.* 20, 652–662.
- Reynolds, T.D., Buonocore, L., Rose, N.F., Rose, J.K., and Robek, M.D. (2015). Virus-like vesicle-based therapeutic vaccine vectors for chronic hepatitis B virus infection. *J. Virol.* 89, 10407–10415.
- Rose, N.F., Buonocore, L., Schell, J.B., Chattopadhyay, A., and Rose, J.K. (2014). In vitro evolution of high-titer, virus-like vesicles containing a single structural protein. *Proc. Natl. Acad. Sci. U S A* 111, 16866–16871.
- Rose, N.F., Publicover, J., Chattopadhyay, A., and Rose, J.K. (2008). Hybrid alphavirus-rhabdovirus propagating replicon particles are versatile and potent vaccine vectors. *Proc. Natl. Acad. Sci. U S A* 105, 5839–5843.
- Schell, J.B., Rose, N.F., Bahl, K., Diller, K., Buonocore, L., Hunter, M., Marx, P.A., Gambhira, R., Tang, H., Montefiori, D.C., et al. (2011). Significant protection against high-dose simian immunodeficiency virus challenge conferred by a new prime-boost vaccine regimen. *J. Virol.* 85, 5764–5772.
- Schillie, S., Vellozzi, C., Reingold, A., Harris, A., Haber, P., Ward, J.W., and Nelson, N.P. (2018). Prevention of hepatitis B virus infection in the United States: recommendations of the advisory committee on immunization practices. *MMWR Recomm. Rep.* 67, 1–31.
- Schinazi, R.F., Ehteshami, M., Bassit, L., and Asselah, T. (2018). Towards HBV curative therapies. *Liver Int.* 38, 102–114.
- Seeger, C., and Mason, W.S. (2015). Molecular biology of hepatitis B virus infection. *Virology* 479–480, 672–686.
- Sette, A.D., Oseroff, C., Sidney, J., Alexander, J., Chesnut, R.W., Kakimi, K., Guidotti, L.G., and Chisari, F.V. (2001). Overcoming T cell tolerance to the hepatitis B virus surface antigen in hepatitis B virus-transgenic mice. *J. Immunol.* 166, 1389–1397.
- Tang, L.S.Y., Covert, E., Wilson, E., and Kottilil, S. (2018). Chronic hepatitis B infection. *JAMA* 319, 1802–1813.
- Tsai, K.-N., Kuo, C.-F., and Ou, J.-H.J. (2018). Mechanisms of hepatitis B virus persistence. *Trends Microbiol.* 26, 33–42.
- van den Pol, A.N., Mao, G., Chattopadhyay, A., Rose, J.K., and Davis, J.N. (2017). Chikungunya, influenza, nipah, and Semliki forest virus chimeric viruses with vesicular stomatitis virus: actions in the brain. *J. Virol.* 91, e02154–16.
- Webster, G.J.M., Reignat, S., Brown, D., Ogg, G.S., Jones, L., Seneviratne, S.L., Williams, R., Dusheiko, G., and Bertolletti, A. (2004). Longitudinal analysis of CD8+ T cells specific for structural and nonstructural hepatitis B virus proteins in patients with chronic hepatitis B: implications for immunotherapy. *J. Virol.* 78, 5707–5719.
- Zhuo, M., Song, L., Tang, Y., Dai, S., Chen, X., Yu, Y., Zang, G., and Tang, Z. (2015). Vaccination with ubiquitin-hepatitis B core antigen-cytoplasmic transduction peptide enhances the hepatitis B virus-specific cytotoxic T-lymphocyte immune response and inhibits hepatitis B virus replication in transgenic mice. *Mol. Med. Rep.* 12, 3591–3598.
- zu Putlitz, J., Lanford, R.E., Carlson, R.I., Notvall, L., de la Monte, S.M., and Wands, J.R. (1999). Properties of monoclonal antibodies directed against hepatitis B virus polymerase protein. *J. Virol.* 73, 4188–4196.

Supplemental Information

**Virus-like Vesicles Expressing Multiple Antigens
for Immunotherapy of Chronic Hepatitis B**

Timur O. Yarovinsky, Stephen W. Mason, Manisha Menon, Marie M. Krady, Maria Haslip, Bhaskara R. Madina, Xianyong Ma, Safiehkhatoon Moshkani, Carolina Chiale, Anasuya Chattopadhyay Pal, Bijan Almassian, John K. Rose, Michael D. Robek, and Valerian Nakaar

Supplemental Information

Transparent Methods

Plasmids and cells for VLV production. Plasmids for generation of empty vector VLV and the VLV expressing MHBs (MT2A) have been previously described (Reynolds et al., 2015; Rose et al., 2014). We custom synthesized DNA fragments encoding HBV Pol and HBcAg with 2A peptides and cloned them into the MT2A plasmid using conventional cloning and/or Gibson assembly at SbfI and Pael restriction sites. We used BHK-21 cells for generation of VLV master stocks by Lipofectamine-mediated transfection and production of VLV working stocks by infection. We maintained BHK-21 cells in Dulbecco's modified Eagle's medium (DMEM) supplemented with 5% fetal bovine serum (FBS), 50 U/ml of penicillin, and 2 mM L-glutamine and switched them to Opti-MEM I reduced serum medium for VLV production for 48 to 72 h after infection. After preclearing of the conditioned medium by centrifugation at 600g for 10 min, we concentrated working stocks by ultrafiltration using MacroSep® Advance 100 MWCO (Pall Laboratory) or Amicon Ultra 100K MWCO (EMD Millipore) centrifugal filter units. To quantify VLV titers, we used plaque assay or indirect immunofluorescence for VSV-G in BHK-21 as previously described (Reynolds et al., 2015; Rose et al., 2014).

Indirect immunofluorescence microscopy and flow cytometry. We seeded BHK-21 cells on coverslips one day prior to infection with VLVs at MOI=1 plaque forming unit (PFU)/cell. We fixed cells with 3% paraformaldehyde at 20 h post infection, washed with PBS containing 10 mM glycine (PBS-glycine), permeabilized with 0.1% Triton X-100 in PBS-glycine, and incubated with a 1:200 dilution of mouse monoclonal antibodies for MHBS (preS2, clone S 26),

HBV Pol (clone 2C8) from Santa Cruz Biotechnology, VSV-G (clones I1 and I1-4), or rabbit polyclonal for HBcAg (DAKO). After extensive washes, we stained the cells with 1:500 dilution of goat anti-mouse AlexaFluor® 488 IgG or goat anti-rabbit AlexaFluor® 594 (ThermoFisher). We mounted washed coverslips on slides using ProLong Gold antifade reagent with DAPI (ThermoFisher) and imaged with Leica DMIRB microscope with 20×/0.75 objective, DC350FX camera (Leica) and ImagePro software or Nikon Eclipse 80i epifluorescence microscope with Retiga 2000R camera and NIS Elements software.

To validate expression of antigens by flow cytometry, we seeded BHK-21 cells in 12-well or 6-well tissue culture plates one day prior to infection with VLVs at MOI=1 plaque forming unit (PFU)/cell. We harvested cells at 20 h post infection with TrypLE Express (ThermoFisher), washed in FACS buffer (3% FBS in PBS supplemented with 0.1% sodium azide), treated with fixation and permeabilization buffer (ThermoFisher), and incubated with the PreS2, Pol, HBcAg and VSV-G antibodies listed above followed by rat monoclonal anti-mouse IgG1 conjugated to PE-Cy7 (BioLegend), goat anti-mouse IgG2a conjugated to AlexaFluor® 647 and goat anti-rabbit conjugated to AlexaFluor® 488 (ThermoFisher) antibodies. We acquired a minimum of 10,000 cells satisfying live cell forward and side scatter parameters using LSR II flow cytometer at the Cell Sorter Core Facility (Yale University School of Medicine) and analyzed the data with FlowJo software (version 7.6.5).

Immunoblotting. We seeded BHK-21 cells in 100-mm tissue culture plates one day prior to infection with VLVs at MOI=1 plaque forming unit (PFU)/cell. We prepared protein lysates by scraping the cells in cold lysis buffer containing 1% Igepal (Sigma) and Complete™ protease inhibitor cocktail and preclearing by centrifugation at 14,000 g for 10 min at 4°C. We separated

proteins by SDS-PAGE in 4%–15% precast gradient gels (Biorad), transferred them onto nitrocellulose membranes, blocked, and incubated with rabbit polyclonal anti-2A-peptide (EMD Millipore), anti-VSV-G or anti-actin (SantaCruz) antibodies and HRP-conjugated or near infrared fluorescently labeled secondary antibodies. We scanned and analyzed immunoblots using ChemiDoc (Bio-Rad) or Odyssey imaging system (LICOR).

In vivo studies (immunogenicity studies and models of HBV challenge and chronic infection). All studies with animals were designed and carried out according to the recommendations in the Guide for the Care and Use of Laboratory Animals of the U. S. National Institutes of Health, Eighth Edition. IACUCs of University of Connecticut Health Center, Yale University, Albany Medical Center and Southern Research approved the studies carried out at the corresponding institutions. For immunogenicity studies, we used 6-8 week-old C57BL/6J male mice (Jackson Laboratory) at Yale University and CaroGen laboratory at UCHC TIP facility. We immunized animals with 10^8 PFU/mouse in 1 ml of PBS via i.p. route and collected spleens from euthanized mice at day 7 post immunization. Control mice received equal volume of PBS at the time of immunization. We prepared single cell suspension of splenocytes by ACK lysis buffer, counted cells and stained with directly conjugated antibodies for T cell markers (CD4-Pacific Blue clone GK1.5, CD8-AlexaFluor® 488 clone 53-6.7, CD62L-PE clone MEL-14, CD44-AlexaFluor® 647 clone IM7) in the presence of FcBlock (BD Biosciences). We also stimulated 10^6 splenocytes per well with custom synthesized HBV peptides (Genscript) in the presence of 1X Brefeldin A (BioLegend) for 6 to 12 h prior to fixable viability staining eFluor-780 (ThermoFisher) and surface staining with CD4-Pacific Blue, CD8-AlexaFluor® 488 (BioLegend). After extensive washes and treatment with fixation and permeabilization buffer

(ThermoFisher), we stained cells with PE-conjugated anti-mouse IFN γ (clone XMG1.2) and AlexaFluor® 647-conjugated anti-mouse TNF (clone MP6-XT22). We acquired a minimum of 50,000 cells satisfying live cell forward and side scatter parameters using LSR II flow cytometer at the Cell Sorter Core Facility (Yale University School of Medicine) and analyzed the data with FlowJo software (version 7.6.5). The H-2^b peptides used for CD8⁺ T cell stimulation were as follows: HBP-44 (NLNVSI PWTHK VGNF), HBP-396 (FAVPNLQSL), HBP-419 (VSAAFYHLPL), HBC-93 (MGLKFRQL), HBS-353 (VWL SVI WM), HBS-371 (ILSPFLPL). As a control peptide, we used H-2^d restricted peptide from HBV Pol (HYFQTRHYL). The threshold for cytokine positive CD8⁺ T cells was manually set using similarly gated and stimulated CD8⁺ T cells from spleens of non-immunized mice.

For acute challenge with HBV-AAV, we used 6-8 week-old CB6F1/J male mice (Jackson Laboratory) at Albany Medical College. We chose the CB6F1/J strain (H2^{d/b} haplotype) over C57BL/6 strain (H2^b haplotype) for the HBV challenge because they generate a response to a dominant protective epitope of the surface antigen (HBS-191) restricted to H2^d haplotype that facilitates HBV clearance. We immunized animals with 10⁸ PFU/mouse via the i.p. route six weeks prior to delivery of 10¹¹ genome copies of the AAV2/8 vector carrying 1.3 copies of the HBV genome (genotype D) via the i.v. route. Frequency of IFN γ -producing cells was measured by ELISPOT assay as previously described (Reynolds et al., 2017).

We established experimental chronic HBV replication by transduction of 6-8 week old C57BL/6J male mice in the facility of Southern Research (Birmingham, AL) following description of the model (Dion et al., 2013). After the HBV infection was confirmed by rising levels of serum HBsAg, we formed experimental groups with 10 animals per group based on normalization of

HBsAg levels at day 21 post infection such that each group had the same mean and standard-deviation of 70 ± 29 ng/mL HBsAg. Therefore, each group had a very similar range of HBsAg levels just prior to administration of VLVs. We immunized mice with 3xT2A or empty vector VLV (10^8 PFU/mouse) via i.p. route at day 28 post infection. We continued monitoring HBsAg levels for additional 13 weeks. On days 65 and 84, we administered plasmid DNA adjuvanted with 20 μ g/mouse monophosphoryl lipid A from *Salmonella minnesota* (Invivogen) via the i.m. route as follows: the group of mice immunized with 3XT2A received three separate plasmids expressing HBsAg, HBV Pol and HBcAg under the promoter region of the rhesus monkey EF1 α gene in pVAC1 vector (Invivogen, 50 μ g/mouse ea.); the group of mice immunized with empty vector VLV received empty pVAC1 vector (150 μ g/mouse). We cloned HBsAg, HBcAg and Pol into the pVAC1 vector (Invivogen) and used EndoFree plasmid maxi kit (QIAGEN) to isolate DNA for the booster immunizations. To measure liver HBV RNA, we euthanized 5 mice per group to collect liver samples on day 70 and at the conclusion of the study on day 119. The numbers of IFN γ -producing CD8 $^+$ T cells were determined similarly to the analyses of immunogenicity in the naïve mice, except that HBV Pol and HBsAg antigen peptides were pooled for each antigen separately and no peptide control was used for background determination.

Analytical assays for serum HBV antigens and liver HBV RNA. We collected blood samples from anesthetized mice weekly to measure serum HBsAg and HBeAg in the acute challenge experiment using ELISA kits from International Immunodiagnostics as previously described (Reynolds et al., 2015). We used HBsAg ELISA kit from XpressBio and the HBsAg standard from CellBioLabs to measure serum HBsAg in the chronic HBV infection model.

We collected liver samples from euthanized mice, snap-froze them in liquid nitrogen, and stored at -80° C or on dry ice until processing for RNA isolation. We isolated total RNA using the RNeasy Mini kit (QIAGEN) and used High Capacity cDNA Reverse Transcription kit (Applied Biosystems) for cDNA synthesis. We used quantitative TaqMan™ Fast Advanced Master Mix (Applied Biosystems) and TaqMan™ Assay Mix containing previously described probe and specific primers for detection of HBV: probe, 6FAM- 5' CCT CTT CAT CCT GCT GCT ATG CCT CAT C 3'-MGBNFQ, forward primer 5'-GTG TCT GCG GCG TTT TAT CA-3', and reverse primer 5'-GAC AAA CGG GCA ACA TAC CTT-3', and mouse GAPDH as an endogenous loading control on a StepOnePlus real time PCR system (Applied Biosystems) using StepOne software v2.3 (Garson et al., 2005). We used the comparative ΔCT method to determine HBV RNA abundance relative to mouse GAPDH mRNA.

Statistical analyses. To determine the difference between the experimental groups, we used 1-way ANOVA for endpoint analyses (HBV RNA) or 2-way ANOVA for analyses over the time course (HBsAg) with Sidak's multiple comparison test. To perform all calculations we used GraphPad Prism software, version 7 (GraphPad Software, San Diego, CA).

References for methods:

- Dion, S., Bourgine, M., Godon, O., Levillayer, F., Michel, M.-L., 2013. Adeno-associated virus-mediated gene transfer leads to persistent hepatitis B virus replication in mice expressing HLA-A2 and HLA-DR1 molecules. *J. Virol.* 87, 5554–63. <https://doi.org/10.1128/JVI.03134-12>
- Garson, J.A., Grant, P.R., Ayliffe, U., Ferns, R.B., Tedder, R.S., 2005. Real-time PCR quantitation of hepatitis B virus DNA using automated sample preparation and murine cytomegalovirus internal control. *J. Virol. Methods* 126, 207–213. <https://doi.org/10.1016/j.jviromet.2005.03.001>
- Reynolds, T.D., Buonocore, L., Rose, N.F., Rose, J.K., Robek, M.D., 2015. Virus-Like Vesicle-Based Therapeutic Vaccine Vectors for Chronic Hepatitis B Virus Infection. *J. Virol.* 89, 10407–10415. <https://doi.org/10.1128/jvi.01184-15>

Reynolds, T.D., Moshkani, S., Robek, M.D., 2017. An ELISPOT-based assay to measure HBV-specific CD8+T cell responses in immunocompetent mice, in: Methods in Molecular Biology. pp. 237–247. https://doi.org/10.1007/978-1-4939-6700-1_20

Rose, N.F., Buonocore, L., Schell, J.B., Chattopadhyay, A., Bahl, K., Liu, X., Rose, J.K., 2014. In vitro evolution of high-titer, virus-like vesicles containing a single structural protein. Proc. Natl. Acad. Sci. U. S. A. 111, 16866–71. <https://doi.org/10.1073/pnas.1414991111>