

# Beneficial Antioxidative and Antiperoxidative Effect of Cinnamaldehyde Protect Streptozotocin-Induced Pancreatic $\beta$ -Cells Damage in Wistar Rats

P. Subash-Babu<sup>1,\*</sup>, Ali A. Alshatwi<sup>1</sup> and S. Ignacimuthu<sup>2</sup>

<sup>1</sup>Department of Food Sciences and Nutrition, College of Food Sciences and Agriculture, King Saud University, P.O. Box 2460 Riyadh 11451, Kingdom of Saudi Arabia, <sup>2</sup>Division of Ethnopharmacology, Entomology Research Institute, Loyola College, Chennai 600 034, Tamil Nadu, India

## Abstract

The present study was aimed to evaluate the antioxidant defense system of cinnamaldehyde in normal, diabetic rats and its possible protection of pancreatic  $\beta$ -cells against its gradual loss under diabetic conditions. *In vitro* free radical scavenging effect of cinnamaldehyde was determined using DPPH (1,1-diphenyl-2-picrylhydrazyl), superoxide radical, and nitric oxide radical. Streptozotocin (STZ) diabetic rats were orally administered with cinnamaldehyde at concentrations of 5, 10 and 20 mg/kg body weight for 45 days. At the end of the experiment, the levels of plasma lipid peroxides and antioxidants such as vitamin C, vitamin E, ceruloplasmin, catalase, superoxide dismutase, reduced glutathione and glutathione peroxidase were determined. A significant increase in the levels of plasma glucose, vitamin E, ceruloplasmin, and lipid peroxides and significant decrease in the levels of plasma insulin and reduced glutathione were observed in the diabetic rats. Also the activities of pancreatic antioxidant enzymes were altered in the STZ-induced diabetic rats. The altered enzyme activities were reverted to near-normal levels after treatment with cinnamaldehyde and glibenclamide. Histopathological studies also revealed a protective effect of cinnamaldehyde on pancreatic  $\beta$ -cells. Cinnamaldehyde enhances the antioxidant defense against reactive oxygen species produced under hyperglycemic conditions and thus protects pancreatic  $\beta$ -cells against their loss and exhibits antidiabetic properties.

**Key Words:** Cinnamaldehyde, Diabetes,  $\beta$ -Islets, Streptozotocin, *Cinnamomum zeylanicum*

## INTRODUCTION

Diabetes have been estimated to affect 180 million people worldwide in 2000, and this number is expected to double by 2030 (WHO, 2008). Diabetes is associated with an increased generation of reactive oxygen species (ROS) or impaired antioxidant defense systems which results in oxidative damage leading to ROS-mediated diabetic pathogenesis (Pitozzi *et al.*, 2003). Moreover, it has been reported that an enhancement of lipid peroxidation, an alteration in antioxidant enzymes, and impaired glutathione metabolism may cause disturbances to the antioxidant systems in diabetes (Bagri *et al.*, 2009). Chronic hyperglycemia is the key clinical manifestation of diabetes and leads to a series of biochemical events those results in

the production of high levels of ROS and eventual oxidative stress (Rajarajeswari and Pari, 2011). The increased generation of free radicals and the subsequent impairment of antioxidant defense capabilities indicate that these processes play a central role in the onset, progression and complications of diabetes (Rolo and Palmeira, 2006). Hyperglycemia increases oxidative stress via the overproduction of ROS (Nogueira *et al.*, 2005). The concentrations of ROS have been modulated by nonenzymatic and enzymic antioxidants (Saxena *et al.*, 1993). Alterations in the activities of antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPx) which are the three primary antioxidant enzymes have been demonstrated in different tissues of diabetic animals (Kakkar *et al.*, 1995).

**Open Access** <http://dx.doi.org/10.4062/biomolther.2013.100>

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Nov 22, 2013 Revised Dec 18, 2013 Accepted Jan 10, 2014

### \*Corresponding Author

E-mail: subashbabu80@hotmail.com  
Tel: +996 1 467 7122, Fax: +996 1 467 8394

Free radicals are continuously produced in the body as a result of normal metabolic processes and interactions with environmental stimuli. Oxidative stress results from an imbalance between radical-generating and radical-scavenging systems due to increased free radical production, reduced antioxidant defense activity or both. The implication of oxidative stress in the pathogenesis of diabetes is suggested not only by the generation of oxygen free radicals but also by non-enzymatic protein glycation, the auto-oxidation of glucose (Mullarkey *et al.*, 1990), impaired glutathione metabolism, alterations in antioxidant enzymes, the formation of lipid peroxides and a decrease in ascorbic acid levels (Ravi *et al.*, 2004). In addition to reduced glutathione (GSH), there are other defense mechanisms that may play a major role in protecting against free radicals by eliminating superoxide anion radicals ( $O_2^{\cdot-}$ ), hydrogen peroxide ( $H_2O_2$ ), hydroxyl radicals ( $\cdot OH$ ) and singlet oxygen ( $^1O_2$ ) (Soto *et al.*, 2003).

Recent investigations demonstrated that the antioxidant properties of plants can be correlated with the defense mechanism against oxidative stress and different human diseases including cancer, and arteriosclerosis and aging processes (Rolo and Palmeira, 2006). Antioxidants can interfere with the oxidative process by reacting with free radicals, chelating free catalytic metals and acting as oxygen scavengers. Phenolic antioxidants function as free radical terminators and sometimes as metal chelators (Sanchez-Moreno *et al.*, 1999). Thus, antioxidant defense systems have been co-evolved with aerobic metabolism to counteract oxidative damage from ROS.

We have isolated the potential antidiabetic compound cinnamaldehyde from the bark of *Cinnamomum zeylanicum* (Subash-Babu *et al.*, 2007). It has been well established that cinnamaldehyde possesses a wide variety of bioactive properties (Lee, 2002; Lee *et al.*, 2003; Qin *et al.*, 2003). However, no information is available regarding the determination of the antioxidant defense system using cinnamaldehyde treatment in streptozotocin (STZ) diabetic rats. We investigated the antioxidant properties of cinnamaldehyde to evaluate its medicinal value and to characterize its natural antioxidant effect which could be used as a food supplement to the diabetic people. In this study, we evaluated the antioxidant defense system of cinnamaldehyde (20 mg/kg body weight [bw]) in normal and diabetic rats and determined its possible protection of pancreatic  $\beta$ -cell against their gradual loss under diabetic conditions.

## MATERIALS AND METHODS

### Chemicals

STZ was purchased from Sigma Chemicals Co., St. Louis, MO, USA. All other chemicals used were of analytical grade.

### Plant material

*Cinnamomum zeylanicum* Blume. (Lauraceae) bark was collected from the Kanyakumari district, Tamil Nadu, India, and dried in the shade. The species was identified and authenticated by Dr. D. Narasimhan, Taxonomist, Department of Botany, Madras Christian College, Chennai, and a voucher specimen (MPC-301) was deposited at the Ethnopharmacology Research Unit, Loyola College, Chennai, Tamil nadu, India.

### Isolation and identification of the active compounds

Based on a bioassay-guided fractionation, the active com-

pound was isolated and identified; this compound was shown to decrease plasma glucose levels. The active isolate was purified by repeat of column chromatography, and the structure of cinnamaldehyde was determined on the basis of a spectral analysis (Subash-Babu *et al.*, 2007).

### In vitro free radical scavenging assay

**In vitro DPPH assay:** The hydrogen-donating ability of cinnamaldehyde was examined in the presence of a DPPH stable radical using the method of Blois with a slight modification (Blois, 1958). The reaction mixture contained 1.0 ml of 0.1 mM DPPH-ethanol solution, 1.0 ml of ethanol, 0.95 ml of 0.05 M Tris-HCl buffer (pH 7.4), and 50  $\mu$ l of cinnamaldehyde in different concentrations (2.5, 5, 10, 20 and 40  $\mu$ g). Reduction of the DPPH free radical was measured by reading the absorbance at 517 nm at exactly 30 sec after adding different concentrations of cinnamaldehyde. Vitamin C was used as a positive control. The inhibition percentage (%) of the radical scavenging activity was calculated using the following equation:

Inhibition (%) =  $(A_c - A_s/A_c) \times 100$ , where  $A_c$  is the absorbance of the control, and  $A_s$  is the absorbance of the sample at 515 nm.

From the inhibition (%), the amount of the sample ( $\mu$ g) reducing the absorbance by 50% was determined ( $IC_{50}$ ).

### Superoxide anion scavenging activity

The superoxide anion scavenging activity of cinnamaldehyde was determined according to the method of Nishimiki *et al.* (1972) with slight modifications. Briefly, generation of superoxide anion was measured in a reaction mixture containing 160 mM NADH, 40 mM Nitro blue tetrazolium (NBT) and 8 mM phenazine methosulphate (PMS) in phosphate buffered saline (PBS) at pH 7.4. The reduction of NBT was followed by measuring the change in absorbance at 560 nm for 2 min, a period in which the absorbance increased linearly. Different doses of cinnamaldehyde were prepared in dimethylsulfoxide (DMSO) and added to the reaction mixture to give a final concentration of 0.7% DMSO. The activity of the test compound was determined in comparison with the control sample.

### Scavenging of NO radical

Nitric oxide was generated from sodium nitropruside and measured by Griess' reaction as described by Green *et al.* (1982). Sodium nitropruside (5 mM) in standard phosphate buffer saline solution (0.025 M, pH: 7.4) was incubated with different concentrations (2-200  $\mu$ g/ml) of cinnamaldehyde dissolved in phosphate buffer saline (0.025 M, pH: 7.4) and the tubes were incubated at 27°C for 5 hr. A control without the test compound but with an equivalent amount of methanol was taken. After incubation 5 ml of incubation mixture was removed and diluted with 0.5 ml of Griess' reagent (1% sulphanilamide, 2% O-phosphoric acid and 0.1% naphthylethylenediamine dihydrochloride). The absence of chromophore formed during diazotization of nitrite with sulphanilamide and its subsequent coupling with naphthyl ethylene diamine was read at 546 nm against blank. The experiment was repeated in triplicate.

### In vivo biochemical assays

**Estimation of thiobarbituric acid reactive substances (TBARS) and hydroperoxides (HP):** Plasma thiobarbituric acid reactive substances (TBARS) and hydroperoxides (HP) were determined by the methods of Yagi (1976) and Jiang *et al.*

(1992) respectively. Briefly, 0.1 ml of Plasma/ tissue homogenate was treated with 2 ml of (1:1:1 ratio) TBA-TCA-HCL (TBA 0.37%, 0.25 N HCL and 15% TCA) reagent and placed in water bath for 15 mins, cooled and centrifuged and then clear supernatant was measured at 535 nm against reference blank.

Hydroperoxides was expressed as mM/dl. 0.1 ml of Plasma/ tissue homogenate was treated with 0.9 ml of Fox reagent (88 mg BHT, 7.6 mg xylene orange and 0.8 mg ammonium iron sulphate) were added to 90 ml of methanol and 10 ml of 250 mM sulphuric acid) and incubated at 37°C for 30 min. The colour developed was read at 560 nm.

#### Determination of non-enzymatic antioxidants

Reduced glutathione (GSH) was determined by the method of Ellman (1959). 1 ml of plasma/ tissue homogenate was taken and 0.5 ml of Ellman's reagent (0.0198% DTNB in 1% sodium citrate) and 3 ml of phosphate buffer (pH-8.0) were added. The colour developed was read at 412 nm.

Ascorbic acid (vitamin C) concentration was measured by Omaye *et al.* (1979) method. To 0.5 ml of plasma/ tissue homogenate 1.5 ml of 6% TCA was added and centrifuged (3,500 g, 20 min). To 0.5 ml of supernatant, 0.5 ml of DNPH reagent (2% DNPH and 4% thiourea in 9N sulphuric acid) was added and incubated for 3 hours at room temperature. After incubation 2.5 ml of 85% sulphuric acid was added and colour developed was read at 530 nm after 30 min.

Vitamin E was estimated by the method of Desai (1984). Vitamin E was extracted from plasma/ tissue homogenate by addition of 1.6 ml ethanol and 2.0 ml petroleum ether to 0.5 ml plasma and centrifuged. The supernatant was separated and evaporated. To the residue, 0.2 ml of 0.2% 2,2'-dipyridyl, 0.2 ml of 0.5% Ferric chloride was added and kept in dark for 5 min. An intense red colored layer obtained on addition of 4 ml butanol was read at 520 nm. And plasma ceruloplasmin was determined according to the methods of Ravin (1961).

#### Determination of enzymic antioxidants

Superoxide dismutase (SOD) activity was determined by the modified method of NADH-Phenazinemethosulphate-nitroblue tetrazolium formazan inhibition reaction spectrophotometrically at 560 nm (Misra and Fridovich, 1972). A single unit of enzyme was expressed as 50% inhibition of NBT (Nitroblue tetrazolium) reduction/min/mg protein. Catalase (CAT) was assayed colorimetrically as described by Takahara *et al.* (1960) using dichromate-acetic acid reagent (5% potassium dichromate and glacial acetic acid were mixed in 1:3 ratio). The intensity was measured at 620 nm and the amount of hydrogen peroxide hydrolyzed was calculated for the catalase activity.


Glutathione peroxidase (GPx) activity was measured by the method described by Rotruck *et al.* (1973). The activity was expressed based on inhibition of GSH. Protein was determined by the method of Lowry *et al.* (1951) using Bovine Serum Albumin (BSA) as standard, at 660 nm.

#### Histopathological study


A portion of pancreatic tissue was fixed in 10% buffered neutral formal saline for histological studies. After fixation, tissues were embedded in paraffin; solid sections were cut at 5  $\mu$ m and stained with aldehyde. The sections were examined under light microscope and photomicrographs were taken.

#### Statistical Analysis


Statistical analysis was performed using the SPSS software package, version 6.0. The values were analyzed by one-way analysis of variance (ANOVA) followed by Duncan's multiple range test (DMRT) (Duncan, 1957). All the results were expressed as the mean  $\pm$  SD for six rats in each group. *p* values less than 0.05 were considered significant.



**Fig. 1.** Comparison of abilities to scavenge DPPH (A), superoxide (B) and nitric oxide (C) radicals *in vitro* between cinnamaldehyde and ascorbic acid.



**Fig. 2.** Effect of cinnamaldehyde on plasma glucose levels (mg/dl) in streptozotocin-induced diabetic male Wistar rats. Group 1: Normal; Group 2: Normal+Cinnamaldehyde (20 mg/Kg bw); Group 3: Diabetic control; Group 4: Diabetic+Cinnamaldehyde (5 mg/kg bw); Group 5: Diabetic+Cinnamaldehyde (10 mg/kg bw); Group 6: Diabetic+Cinnamaldehyde (20 mg/kg bw); Group 7: Diabetic+Glibenclamide (0.6 mg/kg bw). Each value is mean  $\pm$  SD for 6 rats in each group. Values not sharing a common superscript differ significantly at  $p < 0.05$  (DMRT).



**Fig. 3.** Effect of cinnamaldehyde on plasma insulin levels ( $\mu$ U/ml) in streptozotocin-induced diabetic male Wistar rats. Group 1: Normal; Group 2: Normal+Cinnamaldehyde (20 mg/Kg bw); Group 3: Diabetic control; Group 4: Diabetic+Cinnamaldehyde (20 mg/kg bw); Group 5: Diabetic+Glibenclamide (0.6 mg/kg bw). Each value is mean  $\pm$  SD for 6 rats in each group. Values not sharing a common superscript differ significantly at  $p < 0.05$  (DMRT).

**Table 1.** Effect of cinnamaldehyde on plasma levels of TBARS, hydroperoxides, and non-enzymatic antioxidants in male Wistar rats with and without diabetes

Groups	Normal	Normal + cinnamaldehyde (20 mg/kg bw)	Diabetic control	Diabetic + cinnamaldehyde (20 mg/kg bw)	Diabetic + glibenclamide (0.6 mg/kg bw)
TBARS (nMoles/ml)	2.02 $\pm$ 0.12 <sup>a</sup>	2.18 $\pm$ 0.05 <sup>a,b</sup>	4.98 $\pm$ 0.15 <sup>d</sup>	2.24 $\pm$ 0.08 <sup>b</sup>	2.65 $\pm$ 0.09 <sup>c</sup>
Hydroperoxides (value $\times 10^{-5}$ mMoles/dl)	7.23 $\pm$ 0.28 <sup>a</sup>	7.84 $\pm$ 0.61 <sup>a,b</sup>	14.97 $\pm$ 0.74 <sup>d</sup>	8.10 $\pm$ 0.25 <sup>b</sup>	9.18 $\pm$ 0.29 <sup>c</sup>
GSH (mg/dl)	28.81 $\pm$ 1.19 <sup>a</sup>	30.10 $\pm$ 1.27 <sup>a,b</sup>	17.92 $\pm$ 0.77 <sup>c</sup>	26.37 $\pm$ 0.82 <sup>a</sup>	24.16 $\pm$ 0.57 <sup>b</sup>
Vitamin C (mg/dl)	1.71 $\pm$ 0.14 <sup>a</sup>	1.83 $\pm$ 0.10 <sup>a</sup>	0.85 $\pm$ 0.04 <sup>c</sup>	1.67 $\pm$ 0.03 <sup>a</sup>	1.59 $\pm$ 0.08 <sup>a,b</sup>
Vitamin E (mg/dl)	1.82 $\pm$ 0.25 <sup>a</sup>	1.78 $\pm$ 0.42 <sup>a</sup>	3.42 $\pm$ 0.37 <sup>b</sup>	1.68 $\pm$ 0.40 <sup>a</sup>	1.70 $\pm$ 0.39 <sup>a</sup>
Ceruloplasmin (mg/dl)	18.23 $\pm$ 1.75 <sup>a</sup>	22.94 $\pm$ 1.19 <sup>a,b</sup>	32.18 $\pm$ 2.01 <sup>c</sup>	20.63 $\pm$ 1.41 <sup>a,b</sup>	22.11 $\pm$ 1.26 <sup>b</sup>

Each value is mean  $\pm$  SD for 6 rats in each group. Values not sharing a common superscript differ significantly at  $p < 0.05$  (DMRT).

## RESULTS

### In vitro free radical scavenging effect of cinnamaldehyde

The free radical scavenging effects of cinnamaldehyde on DPPH, superoxide radical, and NO radical were determined. The results are shown in Fig. 1. On a comparative basis, cinnamaldehyde exhibited better free radical scavenging activity in quenching DPPH, with an  $IC_{50}$  value of 8.2  $\mu$ g/ml, and the superoxide radical, with an  $IC_{50}$  value of 13.3  $\mu$ g/ml. Cinnamaldehyde also exhibited a better response in quenching NO radicals, with an  $IC_{50}$  value of 12.1  $\mu$ g/ml. The results were compared with ascorbic acid as a positive control, indicating the potential antioxidant properties of cinnamaldehyde.

### Effect of cinnamaldehyde on plasma glucose levels

Fig. 2 shows the plasma glucose levels in the normal and experimental rats. The STZ-diabetic rats showed a significant increase in the levels of plasma glucose compared with normal rats. Oral administration of cinnamaldehyde at a dose of 20 mg/kg bw resulted in a highly significant ( $p < 0.05$ ) effect compared with 5 and 10 mg/kg bw doses. Therefore, cinnamaldehyde was selected for further biochemical studies.

aldehyde was selected for further biochemical studies.

### Effect of cinnamaldehyde on plasma insulin levels

Fig. 3 shows the plasma insulin levels of normal and experimental rats. Plasma insulin level was significantly ( $p < 0.05$ ) decreased in diabetic rats when compared to the normal levels. Oral administration of cinnamaldehyde significantly ( $p < 0.05$ ) increased the plasma insulin levels compared with the levels observed in the untreated diabetic rats. The results were compared with glibenclamide, a standard reference drug.

### In vivo antioxidant effect of cinnamaldehyde

Table 1 shows the levels of TBARS and HP and nonenzymatic antioxidants such as GSH, vitamin C, vitamin E and ceruloplasmin in the plasma of normal and STZ-diabetic rats. The rats treated with cinnamaldehyde alone showed increased levels of TBARS, HP and ceruloplasmin levels when compared with normal rats, but the differences were not significant. The diabetic rats showed a significant ( $p < 0.05$ ) increase in plasma TBARS, hydroperoxides, vitamin E and ceruloplasmin levels, whereas the diabetic rats showed a significantly ( $p < 0.05$ ) de-


**Table 2.** Effect of cinnamaldehyde on pancreatic levels of TBARS, hydroperoxides, GSH, and enzymatic antioxidants in male Wistar rats with and without diabetes

Groups	Normal	Normal + cinnamaldehyde (20 mg/kg bw)	Diabetic control	Diabetic + cinnamaldehyde (20 mg/kg bw)	Diabetic + glibenclamide (0.6 mg/kg bw)
TBARS	32.40 $\pm$ 2.20 <sup>a</sup>	31.75 $\pm$ 2.30 <sup>a</sup>	67.28 $\pm$ 3.90 <sup>c</sup>	38.91 $\pm$ 3.10 <sup>b</sup>	36.27 $\pm$ 2.90 <sup>a,b</sup>
Hydroperoxides	12.65 $\pm$ 0.56 <sup>a</sup>	12.66 $\pm$ 1.18 <sup>a</sup>	22.16 $\pm$ 0.87 <sup>b</sup>	12.24 $\pm$ 0.51 <sup>a</sup>	13.03 $\pm$ 0.35 <sup>a</sup>
GSH	19.34 $\pm$ 0.62 <sup>b</sup>	20.46 $\pm$ 1.14 <sup>b,c</sup>	11.67 $\pm$ 1.18 <sup>a</sup>	21.43 $\pm$ 0.80 <sup>c</sup>	20.94 $\pm$ 0.61 <sup>c</sup>
GPx	28.31 $\pm$ 1.08 <sup>c</sup>	25.08 1.65 <sup>b</sup>	17.13 $\pm$ 0.56 <sup>a</sup>	26.44 $\pm$ 1.54 <sup>b,c</sup>	25.95 $\pm$ 1.44 <sup>b,c</sup>
SOD	4.05 $\pm$ 0.17 <sup>b</sup>	4.93 0.22 <sup>c</sup>	2.53 $\pm$ 0.34 <sup>a</sup>	3.96 $\pm$ 0.23 <sup>b</sup>	3.67 $\pm$ 0.22 <sup>b</sup>
CAT	17.90 $\pm$ 0.54 <sup>e</sup>	16.09 0.47 <sup>d</sup>	7.81 $\pm$ 0.29 <sup>a</sup>	15.17 $\pm$ 0.58 <sup>c</sup>	14.40 $\pm$ 0.42 <sup>b</sup>

Units: TBARS- nmols/100 g tissue; HP- nmols/100 g tissue; GSH- m moles/ g wet tissue; GP<sub>x</sub> -  $\mu$ g GSH consumed/ min/ mg protein; SOD- Enzyme concentration required to inhibit the O.D at 560nm of chromogen production by 50% in one min / mg protein; CAT-  $\mu$ moles of H<sub>2</sub>O<sub>2</sub> consumed/ min/ protein.

Each value is mean  $\pm$  SD for 6 rats in each group.

Values not sharing a common superscript differ significantly at  $p < 0.05$  (DMRT).



**Fig. 4.** Histopathological studies of normal and experimental diabetic rat pancreas (A-E). (A) Normal rat pancreas. Pancreas showing  $\beta$ -islets. (B) Normal+Cinnamaldehyde treated rat pancreas. Normal acini with islet cells. (C) Diabetic rat pancreas. Pancreatic acini with destroyed islet cells. (D) Diabetic+Cinnamaldehyde treated rats pancreas. Regenerating islet cells. (E) Diabetic+glibenclamide treated rat pancreas. Normal islets cells with mild fatty change.

creased levels of vitamin C compared with normal rats. The altered levels of lipid peroxides and non-enzymic antioxidants in diabetic rats have been significantly ( $p < 0.05$ ) restored to near control levels by the administration of cinnamaldehyde and glibenclamide.

Table 2 shows the concentration of TBARS, HP and GSH

and the activities of enzymic antioxidants such as GPx, SOD and CAT in the pancreas of normal and diabetic rats. In diabetic rats, we observed significantly ( $p < 0.05$ ) increased levels of TBARS and HP and diminished levels of GSH, SOD, CAT and GPx compared with the normal rats. We also observed decreased activities of GPx and CAT in the rats treated with

cinnamaldehyde alone compared with the normal rats. The oral administration of cinnamaldehyde and glibenclamide significantly ( $p < 0.05$ ) decreased the levels of lipid peroxides and increased the enzymic antioxidant activities and the GSH levels in the pancreas of diabetic rats compared with the diabetic control rats.

### Histopathological observation of the pancreas

The histopathological examination revealed a preventive effect of cinnamaldehyde administration on extensive alteration in the pancreas of STZ-induced diabetic rats (Fig. 4). Fig. 4A depicts the pancreas of a normal rat, showing a normal architecture of the islets. Fig. 4B depicts the pancreas of a normal rat treated with cinnamaldehyde. The pancreas appeared to have normal islets. Fig. 4C depicts the pancreas of a diabetic rat, showing pancreatic acini and shrunken islet cells. Fig. 4D depicts the pancreas of a diabetic rat treated with cinnamaldehyde, showing regenerated pancreatic islets. Fig. 4E depicts the pancreas of a diabetic rat treated with glibenclamide, showing the regeneration of the pancreatic islets.

## DISCUSSION

Diabetes is one of the pathological processes known to be related to an unbalanced production of ROS, such as hydroxyl radicals ( $\cdot\text{OH}$ ), superoxide anions ( $\text{O}_2\cdot^-$ ), and  $\text{H}_2\text{O}_2$ . Therefore, cells must be protected from this oxidative injury by antioxidant enzymes (Mullarkey *et al.*, 1990). Antioxidants may offer resistance against oxidative stress by scavenging free radicals, inhibiting lipid peroxidation by many other mechanisms and preventing disease (Qujeq and Rezvani, 2007). DPPH is a stable free radical that is often used to evaluate the antioxidant activity of several natural compounds (Miller and Rice-Evans, 1997). Antioxidants interact with DPPH and thus neutralize its free-radical character.

SOD catalyzes the dismutation of the highly reactive superoxide anion to form oxygen and hydrogen peroxide (Rajarajeswari and Pari, 2011). The superoxide anion is the first reduction product of oxygen, which is measured in terms of the inhibition of generation of  $\text{O}_2\cdot^-$ . NO is a free radical produced in mammalian cells and is involved in the regulation of various physiological processes. However, an excessive production of NO is associated with several diseases (Yokozawa *et al.*, 1998). NO is a very unstable species under aerobic conditions. NO reacts with  $\text{O}_2$  to produce the products nitrate and nitrite via the intermediates  $\text{NO}_2$ ,  $\text{N}_2\text{O}_4$  and  $\text{N}_3\text{O}_4$ . The level of NO is estimated using Griess' reagent. NO produced by a solution of sodium nitroprusside was scavenged by the presence of cinnamaldehyde, and the production of nitrous acid has been scavenged by the presence of cinnamaldehyde. This result may be due to the antioxidant principle whereby cinnamaldehyde competes with oxygen to react with nitric oxide (Ray and Husain, 2002).

STZ-diabetes resulted an increase in lipid peroxidation, which occurs indirectly by the intensified free radical production (Ialenti *et al.*, 1993). Most of the tissue damage is considered to be mediated by free radicals, which attack the membrane via the peroxidation of unsaturated fatty acids (Maritim *et al.*, 2003). Increased levels of peroxides are observed in the plasma due to the consequence of the increased free radical production, and the lipid peroxides are liberated into circula-

tion due to pathological changes. A marked increase in the levels of TBARS may be due to an increased susceptibility of pancreas to lipid peroxidation (Mullarkey *et al.*, 1990). The oral administration of cinnamaldehyde significantly protects against the formation of lipid peroxides. This result suggests that the protective role of cinnamaldehyde could be due to the antioxidative effect of cinnamaldehyde. In addition, oral administration of glibenclamide also significantly decreased TBARS and HP levels in the pancreas of diabetic rats. The alterations in lipid peroxide and antioxidant levels in the rats treated with cinnamaldehyde alone were not significant, which may be due to the counter-defense mechanism between the native and developed antioxidant defense systems.

The plasma protein ceruloplasmin is a powerful free radical scavenger that inhibits lipid peroxidation by binding to copper (Cao *et al.*, 1997). Ceruloplasmin levels increased under conditions leading to the generation of oxygen products such as the superoxide radical and hydrogen peroxide. The observed rise in the plasma ceruloplasmin levels in the diabetic rats may be due to the increased levels of lipid peroxides (Halliwell and Gutteridge, 1990).

GSH, the most important biomolecule protecting against chemically induced toxicity, can participate in the elimination of reactive intermediates via reduction of hydroperoxides in the presence of GPx (Rajarajeswari and Pari, 2011). The observed decrease in the GSH level in diabetic rats represents an increased utilization of GSH due to oxidative stress (Dormandy, 1980). It appears that generation of oxygen radicals by increased levels of glucose causes utilization of GSH and thus lowers GSH levels in plasma and pancreas. Oral administration of cinnamaldehyde and glibenclamide increased the glutathione content in the diabetic animals.

Vitamin C, or L-ascorbic acid, is considered to be the most important antioxidant in extracellular fluids, and it has many cellular activities of an antioxidant nature (Stocker and Frei, 1991). Vitamin C functions as a free radical scavenger of active and stable oxy radicals. The marked decrease in the plasma levels of ascorbic acid levels might be due to its increased utilization in antioxidant defense against increased levels of ROS or to a decrease in the GSH levels because GSH is required for the recycling of vitamin C (Moser and Bendich, 1991). Ascorbic acid can also protect membranes against peroxidation by enhancing the activity of a tocopherol, the chief lipid-soluble, chain breaking antioxidant (Wefers and Sies, 1988).  $\alpha$ -Tocopherol interrupts the chain reaction of lipid peroxidation by reacting with lipid peroxy radicals and protects cell membranes against damage by lipid hydroperoxides (Bhatia and Jain, 2003). The elevated levels of  $\alpha$ -tocopherol observed in diabetic rats play a protective role against increased lipid peroxidation in diabetes mellitus (Kinalski *et al.*, 2000). In our study, oral administration of cinnamaldehyde and glibenclamide gradually decreased the  $\alpha$ -tocopherol level in a manner that was directly proportional to the production of free radicals by STZ.

Pancreatic islet cells possess very low levels of free radical scavenging enzymes, including SOD, CAT and GPx and are vulnerable to free radical toxicity (Subash Babu and Prince, 2004). Pancreatic  $\beta$ -cells are considered to be exceptionally vulnerable to the cytotoxic actions of oxygen free radicals because of their low levels of antioxidant enzymes (Simmons, 2006). STZ-induced cytotoxicity of islets is reduced by antioxidant enzymes. SOD in particular is considered to be the

first cellular defense against superoxide radical toxicity (Bhattacharya *et al.*, 1997). The antioxidant activity of cinnamaldehyde may be due to the inhibition of the glycation of the antioxidant enzymes SOD, CAT and GPx. Glucose, which forms a Schiff base with proteins, has been reported to have a high affinity for proteins, especially those containing transition metal ions. Increased glycation of Cu-Zn-SOD has been reported in diabetes. Many reports have shown that curcumin inhibits the formation of advanced glycation end products in STZ-induced diabetic rats (Jia *et al.*, 2009). In our previous study, we observed the decreased glycosylation of hemoglobin (HbA<sub>1c</sub>) in cinnamaldehyde-treated diabetic rats (Subash-Babu *et al.*, 2007). In the present study, we concluded that the increased activities of CAT and GPx in the cinnamaldehyde-treated STZ-diabetic rats might be due to the prevention of glycation.

An increase in the activities of the enzymes SOD and CAT may protect  $\beta$ -cells against damage from ROS. In this context, treatment with the antioxidant N-acetyl cysteine in diabetic mice was shown to suppress apoptosis in  $\beta$ -cells and decrease blood glucose levels (Subash-Babu *et al.*, 2009). A histopathological observation also revealed that an alteration occurred in the architecture of the pancreatic islets in STZ-diabetic rats.

Reduction of  $\beta$ -cell mass is critical in the pathogenesis of diabetes mellitus. The discovery of agents that induce regeneration of pancreatic  $\beta$ -cells would be useful to develop a new therapeutic approach to treat diabetes. After 45 days of oral administration of cinnamaldehyde, blood glucose returned to normal levels, and the islets of Langerhans showed an improvement in the  $\beta$ -cell granulation. The plasma glucose lowering activity was compared with glibenclamide a standard hypoglycemic drug. Glibenclamide has been used for many years to treat diabetes by stimulating insulin secretion from pancreatic  $\beta$ -cells. Sharma *et al.* (2003) have also reported an improvement of the islet after one month of oral feeding of *E. jambolana* seed powder and some alkaloids. Our results are consistent with previous reports suggesting that the regeneration of islet cells by dietary components and the stimulation of insulin secretion by different plant extracts have been achieved (Sharma *et al.*, 2003).

In conclusion, the present study demonstrates that cinnamaldehyde decreased the levels of lipid peroxidation products and increased the activities of antioxidant enzymes. The results of the present study indicated the glucose lowering effect of cinnamaldehyde due to the insulin release from pancreatic  $\beta$ -cells of the islets of Langerhans. Cinnamaldehyde augmented the release of insulin many folds probably by protecting  $\beta$ -cells from free radicals through its antioxidant activity and stimulation of  $\beta$ -cells resembling direct insulin secretagogue effect. Histological assessment also showed that the damage caused by STZ in pancreatic  $\beta$ -cells was markedly reduced, and that the damaged pancreatic  $\beta$ -cells were regenerated by the administration of cinnamaldehyde. Thus, it is suggested that cinnamaldehyde plays an antioxidant role in addition to its antidiabetic activity.

## ACKNOWLEDGMENTS

The authors would like to extend their sincere thanks to the Deanship of Scientific Research, King Saud University for its funding of this research through the Research Group Project

no: RGP- VPP-276.

## REFERENCES

- Bagri, P., Ali, M., Aeri, V., Bhowmik, M. and Sultana, S. (2009) Antidiabetic effect of *Punica granatum* flowers: effect on hyperlipidemia, pancreatic cells lipid peroxidation and antioxidant enzymes in experimental diabetes. *Food Chem. Toxicol.* **47**, 50-54.
- Bhatia, A. L. and Jain, M. (2003) *Amaranthus paniculatus* (Linn.) improves learning after radiation stress. *J. Ethnopharmacol.* **85**, 73-79.
- Bhattacharya, S. K., Satyan, K. S. and Chakrabarti, A. (1997) Effect of trasina, an ayurvedic herbal formulation on pancreatic islet superoxide dismutase activity in hyperglycaemic rats. *Indian J. Exp. Biol.* **35**, 297-299.
- Blois, M. S. (1958) Antioxidant determinations by the use of stable free radical. *Nature* **181**, 1199-1200.
- Cao, G., Sofic, E. and Prior, R. L. (1997) Antioxidant and prooxidant behaviour of flavonoids: structure-activity relationships. *Free Radic. Bio. Med.* **22**, 749-760.
- Desai, I. D. (1984) Vitamin E analysis methods for animal tissues. *Methods Enzymol.* **105**, 138-147.
- Dormandy, T. L. (1980) Free radical reactions in biological systems. *Ann. R. Coll. Surg. Engl.* **62**, 188-194.
- Duncan, B. D. (1957) Multiple range tests for correlated and heteroscedastic means. *Biometrics* **13**, 164-176.
- Ellman, G. C. (1959) Tissue Sulfhydryl groups. *Arch. Biochem. Biophys.* **82**, 70-77.
- Green, L. C., Wanger, D. A., Glogowski, J., Skipper, P. L., Wishnok, J. S. and Tannenbaum, S. R. (1982) Analysis of nitrate and [<sup>15</sup>N] nitrate in biological fluids. *Anal. Biochem.* **126**, 131-138.
- Halliwell, B. and Gutteridge, J. M. (1990) The antioxidants of human extracellular fluids. *Arch. Biochem. Biophys.* **280**, 1-8.
- Ialenti, S., Moncada, M. and Rosa, D. (1993) Modulation of adjuvant arthritis by endogenous nitric oxide. *Brit. J. Pharmacol.* **110**, 701-706.
- Jia, J., Zhang, X., Hu, Y. S., Wu, Y., Wang, Q. Z., Li, N. N., Guo, Q. C. and Dong, X. C. (2009) Evaluation of in vivo antioxidant activities of *Ganoderma lucidum* polysaccharides in STZ-diabetic rats. *Food Chem.* **115**, 32-36.
- Jiang, Z. Y., Hunt, J. V. and Wolff, S. P. (1992) Ferrous ion oxidation in the presence of Xylenol orange for detection of lipid hydroperoxide in low density lipoprotein. *Anal. Biochem.* **202**, 384-389.
- Kakkar, R., Kalra, J., Mantha, S. V. and Prasad, K. (1995) Lipid peroxidation and activity of antioxidant enzymes in diabetic rats. *Mol. Cell. Biochem.* **151**, 113-119.
- Kinalska, M., Sledziowski, A., Telejko, B., Zarzycki, W. and Kinalska, I. (2000) Lipid peroxidation and scavenging enzyme activity in streptozotocin-induced diabetes. *Acta Diabetol.* **37**, 179-183.
- Lee, H. S. (2002) Inhibitory activity of Cinnamomum cassia bark derived component against rat lens aldolase reductase. *J. Pharm. Pharm. Sci.* **5**, 226-230.
- Lee, J. S., Jeon, S. M., Park, E. M., Huk, T. L., Kwon, O. S., Lee, M. K. and Cois, M. S. (2003) Cinnamate supplementation enhances hepatic lipid metabolism and antioxidant defense systems in high cholesterol-fed rats. *J. Med. Food* **6**, 183-191.
- Lowry, O. H., Rosenbrough, N. J., Farr, A. L. and Randall, R. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
- Maritim, A. C., Sanders, R. A. and Watkins, J. B. (2003) Effects of alpha-lipoic acid on biomarkers of oxidative stress in streptozotocin-induced diabetic rats. *J. Nutr. Biochem.* **14**, 288-94.
- Miller, N. J. and Rice-Evans, C. (1997) Factors influencing the antioxidant activity determined by the ABTS<sup>•</sup> radical cation assay. *Free Radic. Res.* **26**, 195-199.
- Misra, H.P. and Fridovich, I. (1972) The role of superoxide anion in the auto oxidation of epinephrine and a simple assay of superoxide dismutase. *J. Biol. Chem.* **247**, 3170-3175.
- Moser, U. and Bendich, A. (1991) Vitamin C. In Handbook of vitamins, 2nd Ed, (Machlin, L. J., Ed), pp195-232. Marcel Dekker, New York.

- Mullarkey, C. J., Edelstein, D. and Brownlee, L. (1990) Free radical generation by early glycation products: a mechanism for accelerated atherogenesis in diabetes. *Biochem. Biophys. Res. Commun.* **173**, 932-939.
- Nishimiki, M., Rao, N. A. and Yagi, K. (1972) The occurrence of superoxide anion in the reaction of reduced phenazine methosulphate and molecular oxygen. *Biochem. Biophys. Res. Commun.* **46**, 849-854.
- Nogueira, F. N., Carvalho, A. M., Yamaguti, P. M. and Nicolau, J. (2005) Antioxidants parameters and lipid peroxidation in salivary glands of streptozotocin-induced diabetic rats. *Clin. Chim. Acta* **353**, 133-139.
- Omaye, S. T., Turnabull, J. C. and Sanberlick, H. E. (1979) Selected methods for the determination of ascorbic acid in animal cells, tissues and fluids. *Methods Enzymol.* **62**, 3-11.
- Pitozzi, V., Giovannelli, L., Bardini, G., Rotella, C. M. and Dolara, P. (2003) Oxidative DNA damage in peripheral blood cells in type 2 diabetes mellitus: higher vulnerability of polymorphonuclear leukocytes. *Mutat. Res.* **529**, 129-133.
- Qin, B., Nagasaki, M., Ren, M., Bajotto, G., Oshida, Y. and Sato, Y. (2003) Cinnamon extract (traditional herb) potentiates in vivo insulin regulated glucose utilization via enhancing insulin signaling in rats. *Diabetes Res. Clin. Pract.* **62**, 139-148.
- Qujeq, D. and Rezvani, T. (2007) Catalase (antioxidant enzyme) activity in streptozotocin-induced diabetic rats. *Int. J. Diabetes Metab.* **15**, 22-24.
- Rajarajeswari, N. and Pari, L. (2011) Antioxidant role of coumarin on streptozotocin-nicotinamide-induced type 2 diabetic rats. *J. Biochem. Mol. Toxicol.* **25**, 355-361.
- Ravi, K., Ramachandran, B. and Subramaniyan, S. (2004) Effect of *Eugenia jambolana* seed kernel on antioxidant defense system in streptozotocin-induced diabetes in rats. *Life Sci.* **75**, 2717-2731.
- Ravin, H. A. (1961) An improved colorimetric enzymatic assay of ceruloplasmin. *J. Lab. Clin. Med.* **58**, 161-168.
- Ray, G. and Husain, S. A. (2002) Oxidants, antioxidants and carcinogenesis. *Indian J. Exp. Biol.* **40**, 1213-1232.
- Rolo, A. P. and Palmeira, C. M. (2006) Diabetes and mitochondrial function: role of hyperglycemia and oxidative stress. *Toxicol. Appl. Pharmacol.* **212**, 167-178.
- Rotruck, J. T., Pope, A. L., Ganther, H. E., Swanson, A. B., Hafeman, D. G. and Hoekstra, W. G. (1973) Selenium; Biochemical role as a component of glutathione peroxidase. *Science* **179**, 588-590.
- Sanchez-Moreno, C., Larrauri, J. A. and Saura-Calixto, F. (1999) Free radical scavenging capacity and inhibition of lipid peroxidation of wines, grape juices and related polyphenolic constituents. *Food Res. Int.* **32**, 407-412.
- Saxena, A. K., Srivastava, P., Kale, R. K. and Baquer, N. Z. (1993) Impaired antioxidant status in diabetic rat liver: effect of vanadate. *Biochem. Pharmacol.* **45**, 539-542.
- Sharma, S. B., Nasir, A., Prabhu, K. M., Murthy, P. S. and Dev, G. (2003) Hypoglycemic and hypolipidemic effect of ethanolic extract of seeds of *Eugenia jambolana* in alloxan induced diabetic rabbits. *J. Ethnopharmacol.* **85**, 201-206.
- Simmons, R. A. (2006) Developmental origins of diabetes: the role of oxidative stress. *Free Radic. Biol. Med.* **40**, 917-922.
- Soto, C., Recoba, R., Barron, H., Alvarez, C. and Favari, L. (2003) Silymarin increases antioxidant enzymes in alloxan induced diabetes in rat pancreas. *Comp. Biochem. Physiol. C. Toxicol. Pharmacol.* **136**, 205-212.
- Stocker, R. and Frei, B. (1991) Endogenous antioxidant defense in human blood plasma. In *Oxidative Stress: Oxidants and antioxidants* (H. Sies, Ed), pp213-243. Academic Press, London.
- Subash Babu, P. and Prince, P. S. M. (2004) Antihyperglycemic and antioxidant effect of Hyponid, an ayurvedic herbomineral formulation in streptozotocin induced diabetic rats. *J. Pharm. Pharmacol.* **56**, 1435-1442.
- Subash-Babu, P., Prabuseenivasan, S. and Ignacimuthu, S. (2007) Cinnamaldehyde- A potential antidiabetic agent. *Phytomedicine* **14**, 15-22.
- Subash-Babu, P., Ignacimuthu, S., Agastian, P. and Varghese, B. (2009) Partial regeneration of beta-cells in the islets of Langerhans by Nymphayol a sterol isolated from *Nymphaea stellata* (Willd.) flowers. *Bioorg. Med. Chem.* **17**, 2864-2870.
- Takahara, S., Hamilton, B. H., Nell, J. V., Kobara, T. Y., Ogura, Y. and Nishimura, E. T. (1960) Hypocatalasemia, a new genetic carrier state. *J. Clin. Inv.* **39**, 610-619.
- Wefers, H. and Sies, H. (1988) The protection by ascorbate and glutathione against microsomal lipid peroxidation is dependent on vitamin E. *Eur. J. Biochem.* **174**, 353-357.
- World Health Organization. (2008) Prevalence data of diabetes worldwide. Available at <http://www.who.int/media centre/fact sheets/fs312/en/index.html>, Accessed 24.04.2008.
- Yagi, K. (1976) A simple fluorometric assay for lipid peroxide in blood plasma. *Biochem. Med.* **15**, 212-216.
- Yokozawa, T., Chen, C. P., Dong, E., Tanaka, T., Nonaka, G. I. and Nishioka, I. (1998) Study on the inhibitory effect of tannins and flavonoids against the 1,1-diphenyl-2-picrylhydrazyl radical. *Biochem. Pharmacol.* **56**, 213-222.