

SCIENTIFIC REPORTS



OPEN

Nutrient enrichment alters impacts of *Hydrocotyle vulgaris* invasion on native plant communities

Lin Liu¹, Han Quan¹, Bi-Cheng Dong¹, Xiang-Qi Bu¹, Lin Li¹, Fu-De Liu², Guang-Chun Lei¹ & Hong-Li Li¹

Received: 25 April 2016
Accepted: 23 November 2016
Published: 20 December 2016

Nutrients may affect the invasiveness of alien plants and the invasibility of native plant communities. We performed a greenhouse experiment to investigate the interactive effect of invasion by a clonal herb *Hydrocotyle vulgaris* and nutrient enrichment on biomass and evenness of native plant communities. We established three types of plant communities (*H. vulgaris* alone, native plant communities without or with *H. vulgaris*) under low and high levels of nutrients. Native communities consisted of eight native, terrestrial species of three functional groups, i.e. four grasses, two legumes, and two forbs. Invasion of *H. vulgaris* had no effect on biomass of the native community, the functional groups, or the individual species. High nutrients increased biomass of grasses, but reduced evenness of the community. High nutrients also decreased the competitive effect, and the relative dominance index of *H. vulgaris*. Therefore, high nutrients reduced the competitive ability of *H. vulgaris* and enhanced the resistance of the native community to invasion. The results provide a basis for management strategies to control the invasion and spread of *H. vulgaris* by manipulating resource availability to support native communities.

Biological invasion has become a serious ecological problem¹. Invasion of plant species can decrease native biodiversity^{1,2} and alter community structure³, biogeochemical cycles⁴ and ecosystem services¹. In the past few decades, the number of introduced invasive plant species has increased dramatically⁵, and many of them are clonal plants^{6–8}. Nutrient enrichment as a result of anthropogenic landscape modifications has also become widely recognized as a serious threat to biodiversity maintenance and ecosystem functioning^{9,10}. Although many studies have investigated responses of invasive plants and native plant communities to nutrient enrichment^{11,12}, impacts of nutrient enrichment on the interaction between invasive plants and native plant communities remains an unresolved issue in invasion biology^{13–15}.

The invasibility of native communities depends on biotic factors^{16,17}. For example, native plants can create spatial heterogeneity in soil nutrients¹⁸ or act as physical obstacles, blocking the spread of the belowground rhizomes and tubers of invasive clonal plants^{19,20}. The invasibility of native communities also depends on the functional similarity between the invasive species and the dominant species of the native communities²¹. Native species with ecological characteristics that are similar to those of invasive species tend to resist invasions more strongly than those exhibit distinct characteristics because they have a higher niche overlap and thus a higher demand for the same resources^{22,23}.

Abiotic factors, such as resource enrichment, can confer invasive species with advantages over native species²⁴. Nutrients are an important resource and may therefore affect the invasibility of native plant communities and the invasiveness of exotic species^{25,26}. Many studies have shown that the invasion success of exotic plants can be enhanced by enrichment of nutrients that are limiting²⁷. For example, nitrogen addition enhanced richness and abundance of invasive annual herbs^{10,11}, promoted the spread of introduced plants into terrestrial habitats¹², but increased the resistance of resident communities to the invasion by *Bromus tectorum*²⁸. Thus, different habitat-dependent pathways mediate interactions between invasion and nutrient enrichment to drive community change²⁹.

Nutrients are correlated with primary production of ecosystems at large scales because nitrogen and phosphorus are limiting in most freshwater, marine, and terrestrial ecosystems^{26,30}. Post-industrial anthropogenic activities have amplified nitrogen and phosphorus cycles by 100% and 400%, respectively³⁰, and rates of nutrient

¹School of Nature Conservation, Beijing Forestry University, Beijing 100083, China. ²School of Environmental Science and Safety Engineering, Tianjin University of Technology, Tianjin 300384, China. Correspondence and requests for materials should be addressed to H.-L.L. (email: lihongli327@163.com)

deposition have increased dramatically³¹. Higher rates of atmospheric nutrient deposition can enhance plant invasion³². Native species have different nutrient-acquisition strategies and are key functional components of vegetation. Changes in functional group representation can disrupt key ecosystem function such as productivity. Nutrient enrichment may favor invasive species over native species and thus change the structure and function of communities. However, the complex associations among nutrient enrichment, plant species invasion, and specific invasion patterns are unclear^{33,34}.

We conducted a greenhouse experiment to test whether high nutrient availability would promote the invasion of a clonal exotic plant species *Hydrocotyle vulgaris* into a terrestrial plant community. We simulated the invasion of *H. vulgaris* into a community consisting of eight native terrestrial plant species of three functional groups (grasses, legumes, and forbs) under two levels of nutrient availability. Specifically, we addressed four questions. (1) Does nutrient availability affect the invasiveness of *H. vulgaris*? (2) Does increasing nutrient availability decrease the invasibility of the native communities? (3) Does nutrient availability affect performance of the functional groups of the plant community? (4) Does the interaction between the invasive plant species and the native community rely on the nutrient availability?

Results

The growth of *H. vulgaris*. The presence of the native plant community significantly decreased in all traits of *H. vulgaris* measured, except stem mass and petiole length. Nutrient enrichment also increased all traits, except total mass and root mass (Fig. 1, Table 1). There were interactive effects of native plant communities and nutrient enrichment on leaf mass, number of nodes, and stem length of *H. vulgaris*. Nutrient enrichment significantly increased these three traits when *H. vulgaris* grew alone, but had little effect when it grew with the native plant community (Fig. 1D,E and G, Table 1).

Responses of communities, functional groups, and species. Invasion by *H. vulgaris* significantly affected none of the indexes of the native community, but nutrient enrichment significantly increased its biomass and decreased its evenness (Fig. 2A–D, Table 2A). Grasses were the dominant functional group of the native communities and their mass constituted over 90% mass of the native communities (Fig. 3). Invasion by *H. vulgaris* affected none of the parameters of any of the functional groups (Table 3). Nutrient enrichment significantly increased biomass (total, aboveground, and underground) of grasses, but had no effect on that of legumes or forbs.

At the species level, *S. viridis* exhibited the highest mass (Fig. 4A). Invasion by *H. vulgaris* had no effect on biomass of any native species, while nutrient enrichment significantly increased biomass of *S. viridis* (Fig. 4, Table 2B). Neither invasion nor nutrient enrichment had a significant effect on seedling establishment of any species, except for *A. sinicus* whose seedling establishment increased due to the presence of *H. vulgaris* (Supplementary Table S1 and Figure S1).

Interactions between *H. vulgaris* and the native plant community. Nutrient enrichment significantly decreased the community-level competitive effect of *H. vulgaris* ($F_{1,10} = 14.27, P = 0.005$) and relative dominance index of *H. vulgaris* ($F_{1,10} = 6.61, P = 0.027$; Fig. 5). Nutrient enrichment can even invert the competitive effect from positive to negative (Fig. 5).

Discussion

The presence of the terrestrial plant community and nutrient enrichment had opposite effects on the growth and establishment of *H. vulgaris*. There were also significant interaction effects of these two factors on leaf mass, number of nodes, and stem length of *H. vulgaris*. In particular, high nutrient availability enhanced the growth of *H. vulgaris* when growing without a native plant community, but had no effect when growing with the native plant community. Our results thus contradict with those of previous studies, which have typically demonstrated that the invasion success of exotic plants into native plant communities can be enhanced by the enrichment of growth-limiting nutrients^{27,35,36}. There are several potential explanations for this phenomenon. First, biological interactions among species may limit the establishment and spread of introduced species^{17,37}. The introduced and native species may affect each other by directly competing for soil nutrients, light, water, and physical space³⁸. As such, in the high nutrient level treatment, the native species *S. viridis* can capture more light before interspecific competition can suppress its growth, thus developing a higher and denser canopy than *H. vulgaris* and shading *H. vulgaris* and thereby preventing its spread³⁹. Alternatively, the spread of stolons and root development at nodes of *H. vulgaris* may have been limited to vacant spaces between native plants, which were diminished by the vigorous growth of the resident root systems under high nutrient conditions²⁹. Previous studies have suggested that root phenotypic plasticity within native communities enables native plants to persist in the context of plant invasion or changing resource levels⁴⁰. Furthermore, *S. viridis* can interfere with and suppress the normal root growth of forbs by releasing nonspecific allelochemicals into the rhizosphere²⁰. A combination of these factors may explain the observed results.

Overall, *H. vulgaris* invasion had no effect on the recipient native plant community in terms of biomass or seeding establishment for all species except for *A. sinicus*. This result indicates that the resident vegetation community could resist invasion. Nutrient enrichment promoted biomass of the native communities and decreased also their evenness (Fig. 2). The decreased community evenness was because nutrient enrichment mainly increased biomass of *S. viridis* and had no significant effect on biomass of other species. We found no interaction effect of *H. vulgaris* invasion and nutrient availability on biomass of native communities, agreeing with previous findings²⁹. However, most previous studies have shown that plant invasion reduces community biomass^{41–43}. This discrepancy may reflect differences in characteristics of invasive species or composition of native communities.




Figure 1. Effects of plant community and nutrient level on the growth of *H. vulgaris* (mean \pm SE): (A) total mass; (B) root mass; (C) stem mass; (D) leaf mass; (E) number of nodes; (F) petiole length; (G) stem length; and (H) leaf area. See Table 1 for ANOVA summaries.

We found that nutrient enrichment increased the productivity of the grasses, mainly *S. viridis*, while had little impact on that of the other two functional groups (legumes and forbs). Grasses, especially *S. viridis*, produced the majority of total biomass of the community, suggesting that grasses were the dominant functional group and *S. viridis* was the dominant species in this community. Grasses were more sensitive to nutrient addition and able to obtain more nutrients under the enrichment treatment, as observed in other studies^{9,10,44,45}. As such, the resistance of the community to invasion may be attributed to the functional identity of resident competitors. It is likely that the fast-growing, native grasses and the dense canopy formed by them, especially that of *S. viridis*³⁹, may have

Traits	Plant community (P)		Nutrient (N)		P × N	
	<i>F</i> _{1,20}	<i>P</i>	<i>F</i> _{1,20}	<i>P</i>	<i>F</i> _{1,20}	<i>P</i>
Total mass	10.72	0.004	3.60	0.072	3.25	0.087
Root mass	8.77	0.008	1.34	0.261	3.37	0.081
Stem mass	3.28	0.085	10.28	0.004	2.93	0.103
Leaf mass ^a	14.14	0.001	7.36	<i>0.013</i>	4.72	<i>0.042</i>
No. of nodes	7.73	<i>0.012</i>	8.49	0.009	5.05	<i>0.036</i>
Petiole length ^b	0.65	0.429	11.19	0.003	1.61	0.692
Stem length ^a	9.42	0.006	7.66	<i>0.012</i>	4.69	<i>0.043</i>
Leaf area ^a	7.66	<i>0.012</i>	9.55	0.006	3.44	0.079

Table 1. Summary of ANOVAs for the effects of plant community and nutrient level on the growth of *Hydrocotyle vulgaris*. Values are in bold if $P < 0.01$ and in italics if $P < 0.05$. See Fig. 1 for graphical representation of data. ^aIndicates log-transformed data. ^bIndicates square root-transformed data.




Figure 2. Effects of *H. vulgaris* invasion and nutrient level on measure of the native plant communities (mean \pm SE): (A) total mass; (B) aboveground mass; (C) belowground mass; and (D) evenness of mass. See Table 2A for ANOVA summaries.

helped the community resist competition from *H. vulgaris*. The absence of interactive effects of these two factors suggests that such differential functional responses to invasion are not dependent on resource availability⁴⁶.

The interaction between the native plant community and *H. vulgaris* was mediated by the nutrient availability. Specifically, nutrient enrichment inverted the competitive effect of *H. vulgaris* from positive to negative, such that *H. vulgaris* only had a detrimental effect on the growth of the plant community when nutrient levels were low. This suggests that changes in nutrient availability altered the competition pattern between the native plant community and *H. vulgaris*, although the influence of *H. vulgaris* invasion was limited. This positive or negative competitive effect suggests that *H. vulgaris* can decrease or increase biomass of the plant community according to nutrient availability. One possible explanation for this facilitation of the community by *H. vulgaris* under high nutrient availability is that exorbitantly high nutrient availability might inhibit plant growth by changing physical and chemical soil properties, causing a nutrient imbalance and reducing photosynthesis by hindering the absorption of Ca^{2+} and Mg^{2+} ^{47,48}. The addition of *H. vulgaris* may have helped to reduce these disruptive nutrient resources and therefore reduce the possible negative effects of high nutrients so that the invaded plant community tended to gain more biomass than the uninvaded community. As shown in a previous study, *S. viridis* grown in close proximity to

Traits	Invasion (I)		Nutrient (N)		I × N	
	<i>F</i> _{1,20}	<i>P</i>	<i>F</i> _{1,20}	<i>P</i>	<i>F</i> _{1,20}	<i>P</i>
(A) Native plant communities						
Total mass ^b	0.06	0.808	17.84	0.001	3.05	0.100
Aboveground mass ^b	0.01	0.942	20.41	<0.001	3.17	0.094
Underground mass	0.55	0.467	7.35	<i>0.015</i>	2.58	0.128
Evenness of mass	1.49	0.240	9.39	0.007	1.53	0.233
(B) Native species						
<i>Setaria viridis</i>	0.05	0.819	19.21	<0.001	2.25	0.152
<i>Festuca arundinacea</i>	0.01	0.930	0.48	0.495	0.76	0.397
<i>Poa pratensis</i>	0.03	0.859	2.46	0.136	0.75	0.398
<i>Bromus inermis</i>	0.31	0.588	0.25	0.623	1.60	0.224
<i>Trifolium repens</i>	1.17	0.293	0.02	0.889	0.36	0.556
<i>Astragalus sinicus</i>	4.78	0.049	0.10	0.756	0.20	0.661
<i>Plantago asiatica</i>	1.89	0.188	0.84	0.374	0.14	0.711
<i>Oxalis corniculata</i>	1.12	0.305	0.93	0.350	0.00	0.986

Table 2. Summary of ANOVAs for the effects of *Hydrocotyle vulgaris* invasion and nutrient level on biomass and evenness of the native communities (A) and biomass of each native species (B). Values are in bold if $P < 0.01$ and in italics if $P < 0.05$. See Figs 2 and 5 for data. ^bIndicates square root-transformed data.

Eupatorium adenophorum accumulates more biomass than *S. viridis* grown in the native control soil⁴⁹. Such impacts may explain the facilitation of the native community by *H. vulgaris* under high nutrient availability.

Moreover, high nutrient availability reduced the relative dominance index of *H. vulgaris* grown in the native plant community. This result suggests that the interspecific competitive ability of *H. vulgaris* was relatively high under low nutrient availability and the ability of the native plant community to resist invasion was relative high under high nutrient availability. Many studies have shown that environmental conditions can change the intensity of interspecific interactions^{50,51}. Moreover, the interaction between resources and competition may increase the capacity of native plants to resist exotic invasive species by reducing the availability of other resources²⁸. Thus, it may be difficult for *H. vulgaris* to spread in a nutrient-rich and species-rich native plant community.

Our results indicate that intrinsic community attributes and nutrient availability can affect the impacts of invasion. Specifically, native species, especially the grass *S. viridis*, may enhance community resistance to invasion by *H. vulgaris*. Accordingly, increases in nutrient deposition due to future global change may not promote *H. vulgaris* invasion into such plant communities, especially those dominated by *S. viridis*. Because our experiment did not test the physiological and biochemical parameters of *H. vulgaris* and the plant community, the potential mechanisms mediating these interactions between *H. vulgaris* and the plant community remain to be explored. Therefore, further studies should be designed to examine the mechanism underlying the interaction between *H. vulgaris* and nutrients available in the rhizosphere to fully understand how nutrient availability affects the invasion process of *H. vulgaris* or other similar clonal plants. In the present study, the effects of *H. vulgaris* invasion on species interactions could not be verified for lack of a monoculture community treatment for each species. To separate the effects on the growth of individual species from effects on competitive interactions among species, it is essential to further construct monocultures containing a single species.

Methods

Study species. *Hydrocotyle vulgaris* L. (Apiaceae) is a perennial clonal herb⁵². It commonly occurs in bogs, valleys, and dune grasslands. It was introduced to China as an ornamental plant in the 1990s and is now considered to be a species with high potential invasiveness⁵³. Each ramet, which consists of a leaf and adventitious roots, may be formed by a node along stolons^{52,54}. In the field, *H. vulgaris* can produce extensive shoot systems and experience heterogeneous micro-environments created by either resource availability or aggregations of neighboring plants⁵⁴. *H. vulgaris* plants used in this experiment were collected from a wetland in the suburbs of Hangzhou, Zhejiang Province, China and were propagated vegetatively in a greenhouse at Forest Science Co. Ltd. of Beijing Forestry University.

The constructed plant communities consisted of a suite of species commonly found in the steppe of northern China⁵⁵. The plant communities consisted of eight species drawn from a random pool. They were classified into three functional groups: four grasses (*Setaria viridis*, *Festuca arundinacea*, *Poa pratensis*, and *Bromus inermis*), two legumes (*Trifolium repens* and *Astragalus sinicus*), and two forbs (*Plantago asiatica* and *Oxalis corniculata*).

Experimental design. We set up three plant arrangement treatments, i.e. *H. vulgaris* alone, a native plant community without *H. vulgaris*, and a native plant community with *H. vulgaris*, crossed with two nutrient addition treatments (i.e., a low or high nutrient level). Each of the six treatment combinations was replicated six times and distributed randomly amongst 36 plastic containers (length, 40 cm; width, 40 cm; depth, 40 cm) filled with 24 L of a 1:1:1 (v/v/v) mixture of quartz sand, vermiculite, and peat. On July 4, 2014, 80 seeds of each of the eight native species were sown uniformly into each container in the treatments with the native plant community with or without *H. vulgaris*. One week later, we planted nine ramets of *H. vulgaris* in the treatment with both the native




Figure 3. Effects of *H. vulgaris* invasion and nutrient level on biomass of functional groups (mean \pm SE). (A,B and C) total mass; (D,E and F) aboveground mass; and (G,H and I) belowground mass. See Table 3 for ANOVA summaries.

Traits	Invasion (I)		Nutrient (N)		I \times N	
	$F_{1,20}$	<i>P</i>	$F_{1,20}$	<i>P</i>	$F_{1,20}$	<i>P</i>
(A) Grasses						
Total mass	0.54	0.470	22.82	<0.001	3.01	0.100
Aboveground mass	0.40	0.533	25.61	<0.001	2.69	0.117
Underground mass	1.10	0.307	9.63	0.006	3.51	0.077
(B) Legumes						
Total mass	1.13	0.302	0.27	0.609	0.75	0.397
Aboveground mass	0.87	0.364	0.30	0.590	0.78	0.388
Underground mass	0.69	0.416	0.00	0.991	0.12	0.732
(C) Forbs						
Total mass	0.54	0.473	0.05	0.832	0.03	0.856
Aboveground mass	0.52	0.479	0.11	0.741	0.10	0.753
Underground mass	0.86	0.366	0.00	0.953	0.21	0.655

Table 3. Summary of ANOVAs for the effects of *Hydrocotyle vulgaris* invasion and nutrient level on biomass of each functional group. Values are in bold if $P < 0.01$ and in italics if $P < 0.05$. See Fig. 3 for data.

plant community and *H. vulgaris*, and the treatment with *H. vulgaris* only (no native plant community). Each *H. vulgaris* ramet was composed of a node and a leaf, and was about 10 cm in height.

For the low-nutrient treatment, 100 mL of 0.3 g L⁻¹ water-soluble fertilizer (20:20:20, N:P:K; Peters Professional; Everris, Geldermalsen, The Netherlands) was added once every two weeks. The low-nutrient




Figure 4. Effects of *H. vulgaris* invasion and nutrient level on total mass of each of eight species (mean \pm SE). See Table 2B for ANOVA summaries.

treatment consisted of the application of 1.5 g nitrogen (N) and 1.5 g phosphorous (P) $\text{m}^{-2} \text{year}^{-1}$. The low-nutrient treatment was designed to simulate field conditions⁵⁶. In the high-nutrient treatment nutrient concentrations were five times as high as those in the low-nutrient treatment (i.e., 7.5 g N and 7.5 g P $\text{m}^{-2} \text{year}^{-1}$). The nutrient concentrations in the high nutrient treatment were based on previous research in grasslands⁵⁷.

Measurements. On September 6, 2014, we harvested the surviving plants of *H. vulgaris* in each container and counted the number of seedlings of each native species. We counted the total number of stem nodes and




Figure 5. Effects of nutrition level on (A) competitive effect and (B) relative dominance index of *H. vulgaris* (mean \pm SE).

measured both petiole length and total stem length of *H. vulgaris*. Leaf area of *H. vulgaris* was measured using WinFOLIA (Pro2004a, Regent Instruments, Québec, Canada). All *H. vulgaris* were separated into three parts: leaves, stems, and roots. All plant parts were separately oven-dried at 70 °C for at least 48 hours and then weighed. The surviving plants of each of the forbs in each container were separated into shoots and roots, similarly dried, and weighed. Then, we calculated the sum of the mass, aboveground mass, and belowground mass of all native species as biomass measures of the native plant community. We also calculated biomass of each functional group (grasses, legumes and forbs).

Data analysis. We calculated the Pielou evenness index as $H/\ln S$, where H is the Shannon–Wiener diversity index based on the proportional final dry mass and S is the number of species. H was calculated as: $H = -\sum P_i \ln(P_i)$ ($i = 1, 2, \dots, S$), where S is the number of plant community species and P_i is biomass of species i divided by the sum of biomass for all eight species in the community⁵⁸.

The competitive effect of *H. vulgaris* was calculated as: competitive effect = \ln (biomass of the native community not invaded by *H. vulgaris*/biomass of the native community invaded by *H. vulgaris*)^{32,59}. A positive value suggests competition between *H. vulgaris* and the native plant community, while a negative value indicates that *H. vulgaris* invasion promotes the growth of the community⁵⁹. We also calculated the dominance of *H. vulgaris* as: the relative dominance index = biomass of *H. vulgaris*/total biomass of all plants in the invaded community⁶⁰.

We performed two-way ANOVAs to test effects of plant community (*H. vulgaris* alone vs. the native community with *H. vulgaris*) and nutrient level (low or high) on total mass, root mass, stem mass, leaf mass, number of node, petiole length, stem length, and leaf area of *H. vulgaris*. We also used two-way ANOVAs to examine effects of invasion (native plant communities without vs. with *H. vulgaris*) and nutrient level (low or high) on total biomass, above- or belowground biomass, evenness of the native plant community, biomass of each functional group, and biomass of each plant species. A one-way ANOVA was performed to test the effect of nutrient level (low or high) on the competitive effect and relative dominance index of *H. vulgaris*.

Data were transformed to natural log or square root before analysis when necessary to remove heteroscedasticity. Leaf mass, stem length, and leaf area of *H. vulgaris* were transformed using the natural log transformation, and petiole length of *H. vulgaris* as well as total mass and aboveground mass of plant communities were transformed using the square root transformation. All data analyses were conducted with SPSS 19.0 (IBM Corp., Armonk, NY, USA).

References

- Mack, R. N. *et al.* Biotic invasions: causes, epidemiology, global consequences, and control. *Ecol. Appl.* **10**, 689–710 (2000).
- Yu, H., Yu, F. H., Miao, S. L. & Dong, M. Holoparasitic *Cuscuta campestris* suppresses invasive *Mikania micrantha* and contributes to native community recovery. *Biol. Conserv.* **141**, 2653–2661 (2008).
- Vorsino, A. E. *et al.* Modeling Hawaiian ecosystem degradation due to invasive plants under current and future climates. *Plos One* **9**, e95427 (2014).
- Dassonville, N., Vanderhoeven, S., Gruber, W. & Meerts, P. Invasion by *Fallopia japonica* increases topsoil mineral nutrient concentrations. *Ecoscience* **14**, 230–240 (2007).
- Colomer-Ventura, F. *et al.* Contemporary evolution of an invasive plant is associated with climate but not with herbivory. *Funct. Ecol.* **29**, 1475–1485 (2015).
- Yu, F. H., Wang, N., Alpert, P., He, W. M. & Dong, M. Physiological integration in an introduced, invasive plant increases its spread into experimental communities and modifies their structure. *Am. J. Bot.* **96**, 1983–1989 (2009).
- Wang, N. *et al.* Clonal integration affects growth, photosynthetic efficiency and biomass allocation, but not the competitive ability, of the alien invasive *Alternanthera philoxeroides*. *Ann. Bot.* **101**, 671–678 (2008).
- Song, Y. B. *et al.* United we stand, divided we fall: a meta-analysis of experiments on clonal integration and its relationship to invasiveness. *Oecologia* **171**, 317–327 (2013).
- Song, M. H. *et al.* Different responses to availability and form of nitrogen in space and time explain species coexistence in an alpine meadow community after release from grazing. *Global Change Biol.* **18**, 3100–3111 (2012).
- Song, M. H. & Yu, F. H. Reduced compensatory effects explain the nitrogen-mediated reduction in stability of an alpine meadow on the Tibetan Plateau. *New Phytol.* **207**, 70–77 (2015).

11. Siemann, E. & Rogers, W. E. The role of soil resources in an exotic tree invasion in Texas coastal prairie. *J. Ecol.* **95**, 689–697 (2007).
12. Li, H. L. *et al.* Nitrogen level changes the interactions between a native (*Scirpus triquetus*) and an exotic species (*Spartina anglica*) in coastal China. *Plos One* **6**, e25629. (2011).
13. Heberling, J. M. & Fridley, J. D. Resource-use strategies of native and invasive plants in Eastern North American forests. *New Phytol.* **200**, 523–533 (2013).
14. Daniel, S. *et al.* Unraveling the life history of successful invaders. *Science* **337**, 580–583 (2012).
15. Davidson, A. M., Jennions, M. & Nicotra, A. B. Do invasive species show higher phenotypic plasticity than native species and, if so, is it adaptive? A meta-analysis. *Ecol. Lett.* **14**, 419–431 (2011).
16. Lonsdale, W. M. Global patterns of plant invasions and the concept of invasibility. *Ecology* **80**, 1522–1536 (1999).
17. Derivera, C. E., Ruiz, G. M., Hines, A. H. & Jivoff, P. Biotic resistance to invasion: native predator limits abundance and distribution of an introduced crab. *Ecology* **86**, 3364–3376 (2005).
18. Richards, C. L. & Pigliucci, M. Plasticity in salt tolerance traits allows for invasion of novel habitat by Japanese knotweed *S. L.* (*Fallopia japonica* and *F. xbohemica*, Polygonaceae). *Am. J. Bot.* **95**, 931–942 (2008).
19. Huber-Sannwald, E. & Durham, S. Effects of nutrient patches and root systems on the clonal plasticity of a rhizomatous grass. *Ecology* **79**, 2267–2280 (1998).
20. Bais, H. P., Weir, T. L., Perry, L. G., Gilroy, S. & Vivanco, J. M. The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Plant Biol.* **57**, 233–266 (2006).
21. Vitousek, P. M. & Hooper, D. U. The effects of plant composition and diversity on ecosystem processes. *Science* **277**, 1302–1305 (1997).
22. Young, S. L., Barney, J. N., Kyser, G. B., Jones, T. S. & Ditomaso, J. M. Functionally similar species confer greater resistance to invasion: implications for grassland restoration. *Restor. Ecol.* **17**, 884–892 (2009).
23. Blumenthal, D. Interrelated causes of plant invasion. *Science* **310**, 243–244 (2005).
24. Davis, M. A., Philip, G. J. & Ken, T. Fluctuating resources in plant communities: a general theory of invasibility. *J. Ecol.* **88**, 528–534 (2000).
25. Fetene, M. Intra- and inter-specific competition between seedlings of *Acacia etbaica* and a perennial grass (*Hyparrhenia hirta*). *J. Arid. Environ.* **55**, 441–451 (2003).
26. Kiehl, K., Esselink, P. & Bakker, J. P. Nutrient limitation and plant species composition in temperate salt marshes. *Oecologia* **111**, 325–330 (1997).
27. Leishman, M. R. & Thomson, V. P. Experimental evidence for the effects of additional water, nutrients and physical disturbance on invasive plants in low fertility Hawkesbury Sandstone soils, Sydney, Australia. *J. Ecol.* **93**, 38–49 (2005).
28. Going, B. M., Hillerislambers, J. & Levine, J. M. Abiotic and biotic resistance to grass invasion in serpentine annual plant communities. *Oecologia* **159**, 839–847 (2009).
29. Gooden, B. & French, K. Impacts of alien plant invasion on native plant communities are mediated by functional identity of resident species, not resource availability. *Oikos* **124**, 298–306 (2014).
30. Elser, J. J. *et al.* Global analysis of nitrogen and phosphorus limitation of primary producers in freshwater, marine and terrestrial ecosystems. *Ecol. Lett.* **10**, 1135–1142 (2008).
31. Huang, J. *et al.* Urbanization in China changes the composition and main sources of wet inorganic nitrogen deposition. *Environ. Sci. Pollut. R.* **22**, 6526–6534 (2015).
32. He, W. M., Yu, G. L. & Sun, Z. K. Nitrogen deposition enhances *Bromus tectorum* invasion: biogeographic differences in growth and competitive ability between China and North America. *Ecography* **34**, 1059–1066 (2011).
33. Bradley, B. A., Blumenthal, D. M., Wilcove, D. S. & Ziska, L. H. Predicting plant invasions in an era of global change. *Trends. Ecol. Evol.* **25**, 310–318 (2010).
34. Seastedt, T. R. & Suding, K. N. Biotic constraints on the invasion of diffuse knapweed (*Centaurea diffusa*) in North American grasslands. *Oecologia* **151**, 626–636 (2007).
35. Barger, N. N., D'Antonio, C. M., Ghneim, T. & Cuevas, E. Constraints to colonization and growth of the African grass, *Melinis minutiflora*, in a Venezuelan savanna. *Plant Ecol.* **167**, 31–43 (2003).
36. Sorrell, B. K., Brix, H., Fitridge, I., Konnerup, D. & Lambertini, C. Gas exchange and growth responses to nutrient enrichment in invasive *Glyceria maxima* and native New Zealand *Carex* species. *Aquat. Bot.* **103**, 37–47 (2012).
37. Gestoso, I., Arenas, F. & Olabarria, C. Feeding behaviour of an intertidal snail: Does past environmental stress affect predator choices and prey vulnerability? *J. Sea. Res.* **97**, 66–74 (2015).
38. Leger, E. A. & Espeland, E. K. Coevolution between native and invasive plant competitors: implications for invasive species management. *Evol. Appl.* **3**, 169–178 (2010).
39. Perry, L. G., Cronin, S. A. & Paschke, M. W. Native cover crops suppress exotic annuals and favor native perennials in a greenhouse competition experiment. *Plant Ecol.* **204**, 247–259 (2009).
40. Phillips, A. J. & Leger, E. A. Plastic responses of native plant root systems to the presence of an invasive annual grass. *Am. J. Bot.* **102**, 73–84 (2015).
41. Maron, J. L. & Marler, M. Field-based competitive impacts between invaders and natives at varying resource supply. *J. Ecol.* **96**, 1187–1197 (2008).
42. Flory, S. L. & Bauer, J. T. Experimental evidence for indirect facilitation among invasive plants. *J. Ecol.* **102**, 12–18 (2014).
43. Michelan, T. S., Thomaz, S. M., Mormul, R. P. & Carvalho, P. Effects of an exotic invasive macrophyte (tropical signalgrass) on native plant community composition, species richness and functional diversity. *Freshwater Biol.* **55**, 1315–1326 (2009).
44. Robledo, E. A., Kmiecik, K., Oplinger, E. S., Nienhuis, J. & Triplett, E. W. Trifolotoxin production increases nodulation competitiveness of *Rhizobium etli* CE3 under agricultural conditions. *Appl. Environ. Microb.* **64**, 2630–2633 (1998).
45. Song, L., Bao, X. M., Liu, X. J. & Zhang, F. S. Impact of nitrogen addition on plant community in a semi-arid temperate steppe in China. *J. Arid. Land.* **4**, 3–10 (2012).
46. Shea, K. & Chesson, P. Community ecology theory as a framework for biological invasions. *Trends. Ecol. Evol.* **17**, 170–176 (2002).
47. McNulty, S. G., Aber, J. D. & Newman, S. D. Nitrogen saturation in a high elevation New England spruce-fir stand. *Forest Ecol. Manag.* **84**, 109–121 (1996).
48. Persson, H., Ahlström, K. & Clemensson-Lindell, A. Nitrogen addition and removal at Gårdsjön — effects on fine-root growth and fine-root chemistry. *Forest Ecol. Manag.* **101**, 199–205 (1998).
49. Xiao, B., Zhou, W., Liu, W. X., Jiang, Z. L. & Wan, F. H. Feedback of *Ageratina adenophora* soil microbe on *A. adenophora* and native plants. *J. Agr. Sci. Tech.* **16**, 151–158 (2014).
50. Liu, L., Dong, B. C., Aplert, P. & Yu, F. H. 2016. Effects of soil substrate heterogeneity and moisture on interspecific competition between *Alternanthera philoxeroides* and four native species. *J. Plant Ecol.*, doi: 10.1093/jpe/rtw052 (2016).
51. Zhang, J. Y., Cheng, G. W., Yu, F. H., Krauchi, N. & Li, M. H. Intensity and importance of competition for a grass (*Festuca rubra* L.) and a legume (*Trifolium pretense* L.) vary with environmental changes. *J. Integr. Plant Biol.* **50**, 1570–1579 (2008).
52. Dong, B. C. *et al.* Soil heterogeneity affects ramet placement of *Hydrocotyle vulgaris*. *J. Plant Ecol.* **8**, 91–100 (2015).
53. Miao, L. H., Ji, M. C., Wang, Y. Y., Qiao, D. D. & Chen, Y. C. Study on invasion risk of *Hydrocotyle vulgaris* as an alien species in wetlands. *J. Zhejiang Uni.* **37**, 425–431 (2011).
54. Dong, B. C., Wang, J. Z., Liu, R. H., Zhang, M. X. & Yu, F. H. Effects of heterogeneous competitor distribution and ramet aggregation on the growth and size structure of a clonal plant. *Plos One* **8**, 68557 (2013).

55. Hou, X. *Vegetation atlas of China*. (Scientific Press, Beijing, 2001).
56. Ying, Z. *et al.* Spatial and temporal variation of atmospheric nitrogen deposition in North China Plain. *Acta Ecol. Sin.* **26**, 1633–1638 (2006).
57. Kolb, A., Alpert, P., Enters, D. & Holzappel, C. Patterns of invasion within a grassland community. *J. Eco.* **90**, 871–881 (2002).
58. Kent, M. & Coker, P. Vegetation description and analysis. *Area* **12**, 76–77 (1992).
59. Wang, P., Lei, J. P., Li, M. H. & Yu, F. H. Spatial heterogeneity in light supply affects intraspecific competition of a stoloniferous clonal plant. *Plos One* **7**, e39105 (2012).
60. Lei, Y. B., Wang, W. B., Feng, Y. L., Zheng, Y. L. & Gong, H. D. Synergistic interactions of CO₂ enrichment and nitrogen deposition promote growth and ecophysiological advantages of invading *Eupatorium adenophorum* in Southwest China. *Planta* **236**, 1205–1213 (2012).

Acknowledgements

This research was supported by the Fundamental Research Funds for the Central Universities (2015ZCQ-BH-01), the National Natural Science Foundation of China (31470475 and 41303057), and the National Key Research and Development Program (2016YFC1201100).

Author Contributions

H.Q., B.-C.D., and H.-L.L. designed the experiment, H.Q., X.-Q.B., and L.L. (Lin Li) conducted the experiment. L.L. (Lin Liu), H.Q. and H.-L.L. performed the statistical analysis, L.L. (Lin Liu), H.Q., F.-D. L., G.-C.L., and H.-L.L. wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Liu, L. *et al.* Nutrient enrichment alters impacts of *Hydrocotyle vulgaris* invasion on native plant communities. *Sci. Rep.* **6**, 39468; doi: 10.1038/srep39468 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016