


Potential of a Pharmacogenetic-Guided Algorithm to Predict Optimal Warfarin Dosing in a High-Risk Hispanic Patient: Role of a Novel *NQO1**2 Polymorphism

Journal of Investigative Medicine High Impact Case Reports
 October-December 2016: 1–6
 © 2016 American Federation for Medical Research
 DOI: 10.1177/2324709616682049
journals.sagepub.com/home/hic


Dagmar F. Hernandez-Suarez, MD¹, Karla Claudio-Campos, BSc¹,
 Javier E. Mirabal-Arroyo, MD¹, Bianca A. Torres-Hernández, PhD^{1,2},
 Angel López-Candales, MD, FACC, FASE¹, Kyle Melin, PharmD¹,
 and Jorge Duconge, PhD, BScPharm¹

Abstract

Deep abdominal vein thrombosis is extremely rare among thrombotic events secondary to the use of contraceptives. A case to illustrate the clinical utility of ethno-specific pharmacogenetic testing in warfarin management of a Hispanic patient is reported. A 37-year-old Hispanic Puerto Rican, non-gravid female with past medical history of abnormal uterine bleeding on hormonal contraceptive therapy was evaluated for abdominal pain. Physical exam was remarkable for unspecific diffuse abdominal tenderness, and general initial laboratory results—including coagulation parameters—were unremarkable. A contrast-enhanced computed tomography showed a massive thrombosis of the main portal, splenic, and superior mesenteric veins. On admission the patient was started on oral anticoagulation therapy with warfarin at 5 mg/day and low-molecular-weight heparin. The prediction of an effective warfarin dose of 7.5 mg/day, estimated by using a recently developed pharmacogenetic-guided algorithm for Caribbean Hispanics, coincided with the actual patient's warfarin dose to reach the international normalized ratio target. We speculate that the slow rise in patient's international normalized ratio observed on the initiation of warfarin therapy, the resulting high risk for thromboembolic events, and the required warfarin dose of 7.5 mg/day are attributable in some part to the presence of the *NQO1**2 (g.559C>T, p.P187S) polymorphism, which seems to be significantly associated with resistance to warfarin in Hispanics. By adding genotyping results of this novel variant, the predictive model can inform clinicians better about the optimal warfarin dose in Caribbean Hispanics. The results highlight the potential for pharmacogenetic testing of warfarin to improve patient care.

Keywords

warfarin, Caribbean Hispanics, pharmacogenetics, *NQO1*, thrombosis, dosing algorithm

Introduction

Warfarin (Coumadin) is an oral anticoagulant drug that inhibits the vitamin K–dependent clotting factors. To date, it has demonstrated to be highly effective in the treatment and prevention of several thromboembolic disorders, including atrial fibrillation, artificial heart valves, pulmonary embolism, antiphospholipid syndrome, and deep vein thrombosis.^{1,2} Despite the advent of the new direct oral anticoagulants,³ warfarin continues to be the mainstay in oral anticoagulation for many patients. According to IMS Health, about 20 million of warfarin prescriptions were dispensed annually in recent years within the United States.^{4,5} However, warfarin ranks as one of the top 10 prescription drugs by reports of adverse episodes in outpatients.^{6,7} In spite of its proven clinical utility and the fact that this drug has been on the market for a long time, optimal warfarin dosing continues to be a

challenge due to the wide variability among patients, narrow therapeutic index, high potential for drug interactions, high potential for food-based interactions (vitamin K–containing products), and variability in pharmacogenomics.^{1,8}

Pharmacogenetic algorithms to predict effective warfarin dose have been developed in several populations worldwide.¹

¹University of Puerto Rico Medical Sciences Campus, San Juan, PR, USA

²University of Puerto Rico Humacao Campus, Humacao, PR, USA

Received September 30, 2016. Revised November 5, 2016. Accepted November 8, 2016.

Corresponding Author:

Dagmar F. Hernandez-Suarez, MD, University of Puerto Rico School of Medicine, Medical Sciences Building, PO Box 365067, San Juan, PR 00936-5067, USA.

Email: dagmar.hernandez@upr.edu



Creative Commons CC-BY: This article is distributed under the terms of the Creative Commons Attribution 3.0 License

(<http://www.creativecommons.org/licenses/by/3.0/>) which permits any use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Nonetheless, individuals of Hispanic heritage have been largely excluded from derivation cohorts.⁹⁻¹¹ This seems to exacerbate existing health care disparities due to the omission of certain ethno-specific genetic polymorphisms on relevant pharmacogenes (eg, *NQO1*2*, *CYP4F2*3*, *CYP2C9*8*) of clinical significance among individuals from this minority group.

To our knowledge, there are no reported data regarding the application of pharmacogenetic-guided algorithm in Hispanics with massive abdominal vein thrombosis to determine the optimal warfarin dose. In this report, a recently developed admixture-adjusted pharmacogenetic algorithm for Caribbean Hispanics is applied to a patient carrying a loss of function *NQO1*2* (g.559C>T, p.P187S) polymorphism, which seems to be significantly associated with resistance to warfarin in this population.

Case Presentation

A case is described of a 37-year-old white Hispanic Puerto Rican non-gravid female (147 lbs) with past medical history of abnormal uterine bleeding for 4 months who visited the emergency room due to abdominal pain. The patient mentioned having been evaluated by gynecology service a month ago, and she was started on medroxyprogesterone acetate (Provera, Pfizer, New York, NY), although she just used the medication for 10 days. Moreover, she underwent endometrial curettage procedure and biopsy with no further complications a week before starting with symptoms. She had no history of tobacco, alcohol, or illegal drugs use. On initial evaluation, the patient stated having started with abdominal bloating and upper mild abdominal pain, constant, no radiating, dull, not related with meals or bowel movements, about 2 weeks prior to her visit. She denied nausea, vomiting, fever, anorexia, diarrhea, and constipation. Due to no improvement of symptoms—despite being treated with antacid medications for a suspected gastritis—she went to seek medical attention to the emergency room. Physical examination was just remarkable for unspecific diffuse abdominal tenderness. Additionally, general initial laboratory results were unremarkable with a prothrombin time and partial thromboplastin time of 10.0 and 26.6 seconds, respectively, and international normalized ratio (INR) of 0.9. No electrolyte abnormalities or anemia were observed. As part of an initial imaging workup, a contrast-enhanced abdominal computed tomography was ordered and an acute on chronic massive thrombosis of the main portal, splenic, and superior mesenteric veins was diagnosed (Figure 1). The patient was then admitted under oral anticoagulation therapy with warfarin at 5 mg/day and low-molecular-weight heparin (ie, Lovenox [enoxaparin] injections) 70 mg every 12 hours, the second discontinued at day 7 of therapy. The need to use a short course of low-molecular-weight heparin was due to the initial temporal effect of Coumadin in promoting clot formation as a consequence to the secondary increase of protein C and protein S factors, also dependent of vitamin K. Follow-up




Figure 1. Contrast-enhanced computed tomography of the abdomen and pelvis showing a massive thrombosis of the main portal, splenic, and superior mesenteric veins.

laboratories showed normal results in antithrombin III, protein C and protein S activity, antiphospholipid antibodies, factor V Leiden mutation, JAK2 mutation, prothrombin gene mutation, flow cytometry for CD55 and CD59, and direct Coombs test.

On initial evaluation, the patient was approached by our team and agreed to participate as a volunteer in an Institutional Review Board–approved research study aiming to investigate the clinical utility of a recently developed pharmacogenetic-guided algorithm for Caribbean Hispanics.¹² Consequently, we collected a buccal swap sample and genotype testing was performed after obtaining her written consent. The sample was interrogated for 6 relevant genetic variants on 4 candidate pharmacogenes (ie, *CYP2C9*, *CYP4F2*, *NQO1*, and *VKORC1*) using the corresponding probes from TaqMan SNP assay technology (Applied Biosystems, Foster City, CA). Consequently, she was found to be single carrier of the *NQO1*2* (g.559C>T, p.P187S) polymorphism and noncarrier of the the remaining analyzed variants.

Based on the diagnosis of deep vein thrombosis, the recommended INR therapeutic range is between 2.0 and 3.0. The patient was initiated on oral warfarin at 5 mg/day following dose recommendations in clinical charts (nomograms) and current American College of Chest Physicians guidelines.^{13,14} The baseline INR value on admission was 0.9 (Table 1). Consecutive INR value of 1.1 was measured on day 3 of therapy and the warfarin dose was increased to

Table 1. Consecutive International Normalized Ratio (INR) Measurements and Daily Warfarin Dose Over the Entire Follow-up Period (ie, 12 weeks). Therapeutic Range Was Set at 2.0 to 3.0.

No.	Day/Week	Test Date	INR Value	Dosing (mg/day)
1	Baseline	March 26, 2016	0.9 ^a	5 mg/day
2	Day 3/week 1	March 28, 2016	1.1 ^a	Rise to 10 mg/day
3	Day 7/week 1	April 1, 2016	4.0 ^b	10 mg/day and then stop warfarin
4	Day 8/week 1	April 2, 2016	4.7 ^b	One day off
5	Day 9/week 2	April 3, 2016	2.8 ^c	Resume warfarin at 7.5 mg/day
6	Day 10/week 2	April 4, 2016	1.3 ^a	7.5 mg/day
7	Day 11/week 2	April 5, 2016	1.6 ^a	7.5 mg/day
8	Day 12/week 2	April 6, 2016	2.2 ^c	7.5 mg/day
9	Day 13/week 2	April 7, 2016	2.6 ^c	7.5 mg/day and discharged
10	Day 17/week 3	April 11, 2016	5.09 ^b	Stop warfarin for 1 day
11	Day 18/week 3	April 12, 2016	3.86 ^b	Resume warfarin at 5 mg/day
12	Day 21/week 3	April 15, 2016	2.93 ^c	5 mg Monday to Friday; 7.5 mg Saturday and Sunday
13	Day 26/week 4	April 20, 2016	3.30 ^b	5 mg Monday to Friday; 7.5 mg Saturday and Sunday
14	Week 7	May 8, 2016	3.35 ^b	5 mg Monday to Friday; 7.5 mg Saturday and Sunday
15	Week 9	May 26, 2016	3.17 ^b	5 mg Monday to Friday; 7.5 mg Saturday and Sunday
16	Week 11	June 4, 2016	2.15 ^c	5 mg Monday to Friday; 7.5 mg Saturday and Sunday
17	Week 11	June 8, 2016	2.07 ^c	7.5 mg/day
18	Week 12	June 11, 2016	2.18 ^c	7.5 mg/day
19	Week 12	June 16, 2016	1.81 ^a	5 mg Monday to Friday; 7.5 mg Saturday and Sunday

^aBelow therapeutic range.

^bAbove therapeutic range.

^cINR levels on target.

10 mg/day (the highest dose given). Following the pharmacogenetic-driven algorithm recommendations, the patient was prescribed warfarin at the dose of 7.5 mg/day for further prevention of thrombus formation. She was kept on this dose until discharge when she reported 2 continue INR values on therapeutic range. Notably, the dose prescribed at the time of achieving the first target INR was 7.5 mg/day (Figure 2). It is important to mention that during the follow-up period, warfarin dosing was stopped on 2 occasions. First, on day 7 due to an INR value of 4.0 (patient was on 10 mg/day) and on day 17 after a reported INR of 5.09.

As per protocol, a 3-month follow-up period was undertaken for identifying warfarin-related adverse episodes as well as other relevant on-treatment events in the patient. Warfarin dosing was changed 6 times in this patient during the course of 90 days. There were 19 INR assessments, with 4 tests made before the patient reached first INR on target (Table 1). Thirty-seven percent of these INRs exceeded target range, peaking at 5.09. Last INR measured on week 12 was 1.81, suggesting a mild sub-therapeutic level of warfarin dosage, and re-exposing her to a slightly higher risk of thrombosis. Previous reports suggest a 17-fold increase in stroke as INR fell below 2.0, whereas the risk for major and life-threatening bleeding events increases 42% for each unit of INR increment above therapeutic range.¹⁵ Fortunately, in this case neither strokes nor thromboembolic events or major life-threatening bleeding episodes were reported. Over the follow-up




Figure 2. Serial INR values and daily warfarin doses over the follow-up period. The optimal warfarin dose of 7.5 mg/day was predicted (green line) by a pharmacogenetic-guided algorithm. Therapeutic range (2.0-3.0) is represented as a gray shaded area. Routinely prescribed “standard” average daily dose of warfarin (ie, 5 mg/day) is depicted as a dashed line in red.

period, the patient reported 2 bruises (one in the calves and the other in the thighs) on weeks 2 to 4, moderate palpitations and dark stools on week 3, chest pain on weeks 3 and 4 and weeks 11 and 12, as well as recurrent fatigue, weakness, anxiety, and discomfort.

Discussion

Deep abdominal vein thrombosis is extremely rare among thrombotic events secondary to the use of contraceptives.¹⁶ Patients with this condition are at high risk of mortality due to the resulting extensive intestinal ischemia secondary to the blood flow obstruction. Thus, it represents a medical emergency that needs to be treated immediately and yet the mortality rate remains significant.¹⁷ Furthermore, the unspecific abdominal symptoms—often referred as the main complaint—and findings on physical examination make difficult the diagnosis, which results in frequent delays in therapy. To date, prompt anticoagulation represents the mainstay therapy. Nevertheless, bowel surgical resection might become the most appropriate choice when associated surgical complications are present. In the present case, we introduced for the first time genetic information to guide the anticoagulation therapy in a Hispanic female with an atypical thrombosis.

Pharmacogenetic Analysis

An admixture-adjusted pharmacogenetic algorithm that explained more than two thirds of the observed variance in stable warfarin dose within Caribbean Hispanics was recently developed by our team.¹² Among predictors of warfarin dose variability in this model, the effect on dose of all *CYP2C9* polymorphisms combined was -18%, whereas the effect of both *VKORC1*-AA and *VKORC1*-GA status combined was -13%. *CYP4F2**3 18000G>A (V433M, rs2108622) and *NQO1**2 c.559C>T (P187S, rs1800566) variants were independently associated with a 17% and 10% increase of the dose per variant allele, respectively, whereas the admixture index decreases the dose by 7%. In the proposed model, the effect (%) on the estimates of the effective dose is calculated per number of variant alleles (ie, *CYP2C9*, *CYP4F2*, *NQO1*, and *VKORC1*), per decades (Age) and per 0.25-unit increase in the dose-adjusted INR response at the third day.

The observed frequency of the *NQO1**2 (g.559C>T, p.P187S, rs1800566) “resistant” polymorphism among the cardiovascular patients from the Puerto Rican population used as the “derivation cohort” was 0.10 (95% confidence interval [CI] = 0.07-0.13).¹² There are 15.7% carriers of the C/T heterozygous genotype for the *NQO1**2 SNP (95% CI 0.12-0.21) and 2% were double carriers (T/T) of this variant (95% CI = 0.007-0.05).¹² The NAD(P)H dehydrogenase, quinone 1 (*NQO1*) gene encodes for a FAD-binding cytoplasmic 2-electron reductase enzyme that catalyzes the reduction of vitamin K to its active hydroquinone form, and it is therefore involved in the vitamin K-dependent prothrombin synthesis.^{18,19} Notably, the combination of *CYP4F2**3 and *NQO1**2 “resistant” variants, identified in as many as 20.4% and 17.7% of individuals from the Puerto Rican study cohort, respectively, had only been previously described to be associated with higher warfarin dose

requirements in another US Hispanic population of Mexican descents.¹⁰

The prediction of 7.5 mg/day of maintenance warfarin by the developed pharmacogenetic-guided algorithm coincided with the patient’s actual effective warfarin dose to reach the INR therapeutic target. The delayed increase in the INR levels of this patient during the initial phase of therapy—in addition to the life cycle of existing clotting factors—may be, in part, a consequence of the *NQO1**2 (g.559C>T, p.P187S) polymorphism found in this subject. Interestingly, this genetic variant seems to be significantly associated in Hispanics with resistance to warfarin.^{10,12}

Although American College of Chest Physicians guidelines recommends against routine use of pharmacogenetic testing for guiding doses of warfarin in patients initiating therapy (Grade 1B),¹⁴ the new and revised 2014 American Heart Association/American Stroke Association guidelines for the primary prevention of stroke recommend that pharmacogenetic dosing of vitamin K antagonists may be considered when therapy is initiated (level of evidence C).²⁰ According to the Clinical Pharmacogenomics Implementation Consortium (CPIC) guidelines for warfarin maintenance dosing, the recommendations based on genetic information are rated as level A or strong, and are derived from numerous observational studies and some prospective trials that suggest the ability to more accurately identify stable therapeutic warfarin dose requirements through the use of both individual genotypes and clinically relevant information.¹ The CPIC guideline also indicates that the use of pharmacogenetic algorithm-based dosing is a better predictive tool and, therefore, is recommended over other methods as long as clinical decision support systems or electronic means are available. This CPIC guideline is currently under revision to update recommendations and incorporate new evidences. Nevertheless, at the moment, there should be some level of consideration for the feasibility of pharmacogenetic testing in the context of warfarin treatment across various health care settings, as widespread pharmacogenetic testing in the outpatient setting may not be feasible or cost-effective.

Most of the currently available pharmacogenetic-driven warfarin dosing algorithms overlook the effect of polymorphisms occurring on pharmacogenetic loci not found to be strongly associated with warfarin dose requirements among White Europeans,²¹⁻²⁵ as opposed to the common *CYP2C9**2, *CYP2C9**3, and *VKORC1*-1639G>A variants that are indeed systematically incorporated into the developed models. It is becoming increasingly clear that the utility of existing models is limited in patients with mixed ancestry like Caribbean Hispanics or African Americans. A previously reported clinical trial (COAG, NCT00839657) raised concerns about real benefits from genotyping patients on warfarin to optimal dose predictions.²⁵ Nonetheless, we strongly believe the COAG trial failed to account for certain ethno-specific genetic polymorphisms of clinical relevance, which led to significant errors in dosing predictions among individuals of

African heritage. Accordingly, it is fairly reasonable to expect the same poor predictability by ignoring admixture and novel pharmacogenetic markers (eg, *NQO1*2*) of potential clinical interest in patients from underrepresented populations like Caribbean Hispanics.

Conclusions

The case reported supports our perception about the predictability of our pharmacogenetic model, after adding the genotyping result of this novel variant, in order to better inform clinicians on optimal warfarin dosing in the Hispanic population. Additionally, it highlights the potential for pharmacogenetic testing of warfarin to improve patient care and reduced health care costs. Further studies are warranted to determine whether the applied algorithm in this case could decrease adverse outcomes, hospital stay, and contribute as an important decision support tool on initiation of warfarin therapy.

Acknowledgments

The material presented herein is the result of work supported with resources from and the use of facilities at the UPR-MSRCMI Center for Genomics in Health Disparities and Rare Disorders for performing genetic assays. The authors also want to thank the patient for her participation in this study.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This investigation was supported in part by grants from the National Heart, Lung, and Blood Institute (Grant Number SC1 HL123911); the Research Center in Minority Institutions from the National Institute on Minority Health and Health Disparities (Award Number 8G12 MD007600); and the Minority Biomedical Research Support-Research Initiative for Scientific Enhancement at the University of Puerto Rico Medical Sciences Campus (Grant Number R25 GM061838).

References

1. Johnson JA, Gong L, Whirl-Carrillo M, et al. Clinical Pharmacogenetics Implementation Consortium Guidelines for CYP2C9 and VKORC1 genotypes and warfarin dosing. *J Clin Pharmacol Ther.* 2011;90:625-629.
2. Bristol-Myers Squibb. Coumadin (warfarin sodium) [Package insert]. Princeton, NJ: Bristol-Myers Squibb. http://packageinserts.bms.com/pi/pi_coumadin.pdf. Accessed November 21, 2016.
3. Ruff CT, Giugliano RP, Braunwald E, et al. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomized trials. *Lancet.* 2014;383:955-962.
4. Bartholow M. Top 200 drugs of 2012. *Pharmacy Times.* 2013;79(7):42-44.
5. IMS Institute for Healthcare Informatics. Declining medicine use and costs: for better or worse? A review of the use of medicines in the United States in 2012. <http://www.imshealth.com/en/thought-leadership/webinar-library/declining-medicine-use-and-costs-for-better-or-worse-u.s-2012>. Accessed November 21, 2016.
6. Wysowski DK, Nourjah P, Swartz L. Bleeding complications with warfarin use: a prevalent adverse effect resulting in regulatory action. *Arch Intern Med.* 2007;167:1414-1419.
7. Budnitz DS, Pollock DA, Weidenbach KN, Mendelsohn AB, Schroeder TJ, Annest JL. National surveillance of emergency department visits for outpatient adverse drug events. *JAMA.* 2006;296:1858-1866.
8. Gulseth MP, Grice GR, Dager WE. Pharmacogenomics of warfarin: uncovering a piece of the warfarin mystery. *Am J Health Syst Pharm.* 2009;66:123-133.
9. Duconge J, Cadilla CL, Seip RL, Ruaño G. Why admixture matters in genetically-guided therapy: missed targets in the COAG and EU-PACT trials. *PR Health Sci J.* 2015;34:175-177.
10. Bress A, Patel SR, Perera MA, Campbell RT, Kittles RA, Cavallari LH. Effect of NQO1 and CYP4F2 genotypes on warfarin dose requirements in Hispanic-Americans and African-Americans. *Pharmacogenomics.* 2012;13:1925-1935.
11. Duconge J. Population heterogeneity and genomic admixture: relevance for global pharmacogenetics. *Pharmacogenomics Pharmacoproteomics.* 2014;5:5. doi:10.4172/2153-0645.1000e141.
12. Duconge J, Ramos AS, Claudio-Campos K, et al. A novel admixture-based pharmacogenetic approach to refine warfarin dosing in Caribbean Hispanics. *PLoS One.* 2016;11:e0145480.
13. Ageno W, Gallus AS, Wittkowsky A, Crowther M, Hylek EM, Palareti G. Oral Anticoagulant Therapy Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians (ACCP) Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 suppl):e44S-e88S.
14. Holbrook A, Schulman S, Witt DM, et al; American College of Chest Physicians. Evidence-based management of anticoagulant therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians (ACCP) Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141(2 suppl):e152S-e184S.
15. Ansell JE. Optimizing the efficacy and safety of oral anticoagulant therapy: high-quality dose management, anticoagulation clinics, and patient self-management. *Semin Vasc Med.* 2003;3:261-270.
16. Lidegaard Ø, Løkkegaard E, Svendsen AL, Agger C. Hormonal contraception and risk of venous thromboembolism: national follow-up study. *BMJ.* 2009;339:b2890.
17. Brunaud L, Antunes L, Collinet-Adler S, et al. Acute mesenteric venous thrombosis: case for nonoperative management. *J Vasc Surg.* 2001;34:673-679.
18. Jaiswal AK. Human NAD(P)H: quinone oxidoreductase2. Gene structure, activity, and tissue-specific expression. *J Biol Chem.* 1994;269:14502-14508.
19. Jaiswal AK, Bell DW, Radjendirane V, Testa JR. Localization of human NQO1 gene to chromosome 16q22 and NQO2-6p25

- and associated polymorphisms. *Pharmacogenetics*. 1999;9: 413-418.
20. Meschia JF, Bushnell C, Boden-Albala B, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Functional Genomics and Translational Biology; Council on Hypertension. Guidelines for the primary prevention of stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45:3754-3832.
 21. Higashi MK, Veenstra DL, Kondo LM, et al. Association between CYP2C9 genetic variants and anticoagulation-related outcomes during warfarin therapy *JAMA*. 2002;287: 1690-1698.
 22. Sanderson S, Emery J, Higgins J. CYP2C9 gene variants, drug dose, and bleeding risk in warfarin treated patients: a HuGENet systemic review and meta-analysis. *Genet Med*. 2005;7:97-104.
 23. Takahashi H, Wilkinson GR, Nutescu EA, et al. Different contributions of polymorphisms in VKORC1 and CYP2C9 to intra- and inter-population differences in maintenance doses of warfarin in Japanese, Caucasians and African Americans. *Pharmacogenet Genomics*. 2006;16:101-110.
 24. Caldwell MD, Berg RL, Zhang KQ, et al. Evaluation of genetic factors for warfarin dose prediction. *Clin Med Res*. 2007;5: 8-16.
 25. Kimmel SE, French B, Kasner SE, et al. A pharmacogenetic versus a clinical algorithm for warfarin dosing. *N Engl J Med*. 2013;369:2283-2293.