

Identification and Characterization of a Sphere Organelle Protein

Rabiya S. Tuma,* John A. Stolk, and Mark B. Roth

Division of Basic Sciences, Fred Hutchinson Cancer Research Center, 1124 Columbia Street, Seattle, Washington 98104; and *Department of Biochemistry, University of Washington, Seattle, Washington 98104

Abstract. Sphere organelles are nuclear structures in amphibian oocytes that are easily visible by light microscopy. These structures are up to 10 μm in diameter and have been described morphologically for decades, yet their function remains obscure. The present study defines a protein component of the sphere organelle, named SPH-1, which is recognized by a mAb raised against purified *Xenopus laevis* oocyte nucleoplasm. SPH-1 is an 80-kD protein which is localized specifically to spheres and is undetectable elsewhere on lampbrush chromosomes or in nucleoli. We show using confocal microscopy that SPH-1 is localized to the cortex of sphere organelles. Furthermore, we have isolated a cDNA that can encode

SPH-1. When epitope-tagged forms of SPH-1 are expressed in *X. laevis* oocytes the protein specifically localizes to spheres, demonstrating that the cloned cDNA encodes the sphere antigen. Comparison of the predicted amino acid sequence with sequence databases shows SPH-1 is related to p80-coilin, a protein associated with coiled bodies; coiled bodies are nuclear structures found in plant and animal cells. The sphere-specific mAb stains *X. laevis* tissue culture cells in a punctate nuclear pattern, showing that spheres or sphere antigens are present in somatic cells as well as germ cells and suggesting a general and essential function for spheres in all nuclei.

THE nuclei of amphibian oocytes contain several large, morphologically distinct structures, including lampbrush chromosomes, nucleoli and "spheres." Sphere organelles were first identified as "knobs" attached to lampbrush chromosomes of the newt *Notophthalmus viridescens* (Gall, 1954; reviewed in Callan, 1986). The term organelle is used to describe spheres because they are distinct structures and larger than some whole cells, yet they are not membrane bound. Structures with similar morphology have since been observed on lampbrush chromosomes of many different amphibian species and in oocyte nuclei of several insects, suggesting that spheres may be common to all animal oocyte nuclei where they perform an unknown yet essential function (Gall and Callan, 1989). In amphibians some spheres appear attached to specific loci on lampbrush chromosomes while others are free in the nucleoplasm. In *Xenopus laevis* there are three chromosomal sphere loci, located on chromosomes VIII, IX, and XVI (Callan et al., 1987), as well as 40 to 60 free spheres (Fig. 1). Free and attached spheres are morphologically indistinguishable. They contain a central electron dense core and a surrounding cortex that is less dense (Kezer et al., 1980). Larger spheres have often been observed with 2 to 3 μm hemispherical masses on their surfaces. These hemispherical masses, which are morphologically distinct from spheres, were recently named "B snurposomes" when antibody staining experiments showed they contain a variety

of proteins and small nuclear RNAs involved in pre-mRNA splicing (Wu et al., 1991). Similar studies have also shown that spheres can be stained with two antibodies that bind snRNPs and have, as a result, been referred to as "C snurposomes" (Gall and Callan, 1989; Wu et al., 1991). These common splicing components are associated with nearly all active sites of transcription on lampbrush chromosomes (Gall and Callan, 1989; Wu et al., 1991).

We report here the generation of a mAb that binds exclusively to spheres. We have used this antibody to isolate a cDNA which encodes a sphere-specific protein, SPH-1. The primary amino acid sequence of this sphere protein is related to p80-coilin; p80-coilin is localized to a functionally undefined structure called the "coiled body" found in the nuclei of many different plant and animal cells (Andrade et al., 1991). The similarity between SPH-1 and p80-coilin suggests that spheres may be related to coiled bodies.

Materials and Methods

Isolation of Oocyte Nuclei

Ovary (30 g) was removed from albino frogs and minced with scissors into pieces containing ~ 30 mature oocytes. The ovary was then divided into two T-150 tissue culture flasks each containing 0.1% collagenase w/v/15 grams ovary in 50 mls of Barth's medium (10 mM Hepes, pH 7.4, 88 mM NaCl, 1 mM KCl, 2.3 mM NaHCO₃, 0.82 mM MgSO₄, 0.66 mM NaNO₃, 0.41 mM CaCl₂), and rotated at 75 rpm for 90–120 min at room temperature. Oocytes were sieved from the collagenase using 100 μm nitex. The large oocytes were separated from the smaller oocytes by sequential sieving on

Address correspondence to Mark B. Roth, Fred Hutchinson Cancer Research Center, 1124 Columbia Street, Seattle, WA 98104.




Figure 1. Phase contrast micrograph of nuclear contents from a *Xenopus laevis* oocyte. A nucleus was dissected from a late stage oocyte (V or VI), its nuclear envelope removed, and the nuclear contents transferred to a chambered slide. The major structures in the oocyte nucleus are clearly visible, including the bivalent lampbrush chromosomes. The many free spheres appear as phase-dark, distinctly round objects. The most abundant structures are the nucleoli, which appear phase bright. The light and dark appearance of the spheres and nucleoli is dependent on the refractive index of the buffer; in subsequent images both spheres and nucleoli appear phase dark because the preparations are mounted in a different buffer. Bar, 40 μm .

250, 500, 750, and 1,000 μm nitex. Oocytes retained on the 750 μm nitex but which passed through the 1,000 μm nitex were placed in 10 mM NaCl and hypotonically swollen for 60–70 min. Several hundred oocytes were transferred using a large bore pasteur pipette to a 150-mm petri dish containing 50 ml of buffer (10 mM Hepes, pH 7.4, 5 mM MgCl_2 , 60 mM KCl, 1 mM DTT, 5% glycerol) with a piece of 750 μm nitex in the bottom; the 150-mm dish was placed in a Tupperware dish containing a small amount of wet ice. A 100-mm petri dish was then used to break the oocytes by gently crushing them in the larger dish. Using a dissecting microscope the nuclei were observed as they emerged from the oocytes. The lysate was stored for 10–20 min in a refrigerator to allow the yolk to settle, after which time the dishes were gently swirled to bring the nuclei or germinal vesicles to the center of the dish. Swirling involved picking up the dish and rocking it in a circular motion so that “one edge” was always in contact with the ice-containing dish. Some practice was required to know how vigorously to swirl the dish because yolk obscures the ability to see the nuclei. Once the nuclei are in the center of the dish they are removed to a second dish and reswirled; the swirling was repeated five to six times until the nuclei were separated from the cytoplasmic contents. Nuclei were then transferred to a new dish. The nuclear envelopes adhered to the clean plastic and the dish was turned quickly to create a shear force sufficient to separate the congealed nucleoplasm from the nuclear envelopes. The percentage recovery of nuclei from oocytes was determined to be at least 80% and the contents of as many as 25,000 nuclei have been recovered in one day.

Monoclonal Antibody Isolation and Characterization

Several mice (RBF/Dn, Jackson Laboratories, Bar Harbor, MN) were each immunized with $\sim 10,000$ nuclei over a 6 mo period. Polyclonal antisera were tested by indirect immunofluorescent staining of chromosome preparations (Roth and Gall, 1987; Callan et al., 1987). Hybridoma cell lines were generated by fusing the spleen cells of one of these mice to Fox-NY cells (Taggart and Samloff, 1983), and the supernatants were used to test for indirect immunofluorescent staining of spheres. Hybridoma cell lines that produced sphere binding antibodies were isolated as mAb producing cell lines by limiting dilution cloning (Harlow and Lane, 1988). Immunoblotting and indirect immunofluorescent staining were done according to previously published methods (Roth and Gall, 1987; Roth et al. 1991). *X. laevis* K2 tissue culture cells were grown according to McStay and Reeder




Figure 2. Photograph of purified *Xenopus laevis* oocyte nuclei. Approximately 3,000 oocyte nuclei are shown before removal of nuclear envelopes. The inset is a dark field light micrograph of a single isolated germinal vesicle after removal of its nuclear envelope. The bright objects on the perimeter are nucleoli; because of their position in the congealed nucleoplasm many nucleoli are lost during the purification. The aggregation of material in the center of the nucleoplasm contains the chromosomes and spheres. The diameter of the single nucleoplasm is 350 μm . Bar, 4 mm.

(1990). For immunoblotting, the cells were washed once in X-PBS (68 mM NaCl, 1.3 mM KCl, 4.0 mM Na_2HPO_4 , 0.7 mM KH_2PO_4 , 0.35 mM CaCl_2 , 0.25 mM MgCl_2) and resuspended in SDS-PAGE sample buffer.

Identification and Characterization of SPH-1 cDNAs

A *X. laevis* young ovary polyA⁺ cDNA expression library made in the vector Lambda Zap (Stratagene Corp., Burlingame, CA) was screened using mAbH1 (Short et al., 1988). Two independent classes of inserts (700 and 1,900 bp) were isolated from the expression library and subcloned into Bluescript plasmids (Stratagene Corp.). Sequencing of the cDNAs revealed an open reading frame that could encode a 43-kD protein and that the 700-bp insert is identical to an internal region of the 1,900-bp clone. Because the 1,900-bp insert appeared to be lacking the 5' end of the open reading frame (ORF), the 700 bp cDNA was used as a hybridization probe to screen a lambda gt10 library of *Xenopus* ovary cDNAs to obtain a full-length cDNA (Rebagliati et al., 1985; Maniatis et al., 1982). 15 positive phage were isolated from $\sim 1.2 \times 10^5$ phage screened. The largest insert, referred to here as cSPH-1, is 2.3 kb. cDNA clones were sequenced on both strands according to published methods (Henikoff, 1987). Sequence databases were searched using BLAST (Altschul et al., 1990) at the National Center for Biotechnology Information and BLOCKS (Henikoff and Henikoff, 1991).




Figure 3. Indirect immunofluorescent staining of a *Xenopus laevis* lampbrush chromosome preparation with mAbH1. *A* and *B* are phase contrast and fluorescent micrographs, respectively, of the same preparation after staining with the mAbH1. Staining is specific for spheres; the fluorescent images of spheres are somewhat larger than the phase contrast images because of the intensity of fluorescence. Bar, 18 μm .

Expression and Detection of SPH-1 Protein

Epitope tagged SPH-1 protein was expressed in *X. laevis* oocytes according to previously published methods with minor modifications. The reiterated hexamer epitope tag, MT6 (Roth et al., 1991), was fused to the cSPH-1 sequence encoding either the second methionine (amino acid 25; construct A) or to amino acid 111 (construct B) as follows. The lambda cDNA insert was cloned into a bluescript plasmid (Stratagene Corp.) containing the MT6 sequence with the tag ORF preceding, and in the same direction as, the SPH-1 ORF. The A construct was then generated by fusing the tag ORF to the cSPH-1 ORF by site directed mutagenesis using a 31 base oligonucleotide. Construct B was made by cloning the original 1,900-bp Lambda Zap cDNA into the MT6 Bluescript plasmid; the 1,900-bp insert encodes the SPH-1 protein from amino acid 111 to the termination codon after amino acid 536. Capped runoff in vitro synthesized transcripts encoding fusion proteins were prepared (Roth and Gall, 1989), and injected into *Xenopus* oocytes according to methods described by Gurdon et al. (1971). Immunoblotting of pools of 8–20 hand isolated oocyte nuclei 48 h after injection of RNA was done according to Roth et al. (1991). *Xenopus* lampbrush chromosomes were prepared according to Callan et al. (1987); indirect immunofluorescent staining was done according to Roth and Gall (1987). mAb9E10 was isolated by Evan et al. (1985).

Results

Identification and Characterization of a Sphere-specific Monoclonal Antibody

To begin a biochemical characterization of spheres, we developed a method for isolating large numbers of oocyte nuclei (Fig. 2; see Materials and Methods). We immunized mice with *X. laevis* oocyte nucleoplasm that had been purified away from cytoplasm and nuclear envelopes (Fig. 2, inset). Immunostaining of lampbrush chromosomes with the resulting polyclonal antisera indicated high titers of antibodies against spheres as well as nucleoli, chromosomes, and B snurposomes (data not shown). The spleen cells of one of these mice were fused to a mouse myeloma cell line and the resulting hybridoma cell lines tested for production of antibodies that bind to spheres by indirect immunofluorescent staining of lampbrush chromosome preparations. Of a total of 384 cell lines tested, four produce antibodies that preferentially bind to spheres. After subcloning, four independent




Figure 4. Confocal microscopy of spheres stained with mAbH1 or mAbID7. *A* and *B* show confocal pseudo-phase contrast and fluorescent images, respectively, of a sphere stained with mAbH1; the antigen is localized to the cortex of the sphere. Note that the nucleoli and B snurposomes are not stained. *C* and *D* show mAbID7 staining is localized to B snurposomes and the internal core of the sphere. mAbID7 stains B snurposomes both when they are, and when they are not, attached to the surface of spheres. This antibody also stains most of the lateral loops of lampbrush chromosomes and is known to recognize a family of essential splicing factors called SR proteins. The staining patterns of mAbH1 and mAbID7 reflect the bipartite structure observed in electron micrographs of spheres (Kezer et al., 1980). Bar, 5 μm .

hybridoma cell lines were recovered which produce antibodies that bind spheres. While three of the antibodies show some staining of other nuclear structures, one of the antibodies, mAbH1, shows selective binding to spheres with no detectable staining of other nuclear structures (Fig. 3).

Electron microscopic observations of sections through spheres have shown that spheres have morphologically distinct domains including an electron dense core and a cortex that is less dense (Kezer et al., 1980). Confocal microscopy of spheres stained with mAbH1 shows that the antigen is concentrated in the sphere's outer cortex with negligible staining in the inner core (Fig. 4, *a* and *b*). Conversely, the inner core is preferentially stained by mAbID7 (Fig. 4, *c* and *d*); this antibody binds a conserved epitope on a class of alternative pre-mRNA splicing factors called SR proteins (Tuma and Roth, unpublished observation; Zahler et al., 1992). Core staining with mAbID7 is consistent with a previous report which shows that another anti-SR protein antibody, mAb SC35, stains the core of spheres (Wu et al., 1991; Fu and Maniatis, 1990). The spatial segregation of specific antigens within the sphere is consistent with the view that the core and cortex represent distinct functional domains.

We were also interested to determine whether spheres are present in somatic cells. We stained *X. laevis* K2 tissue culture cells with mAbH1 and found that it stains nuclei in a punctate pattern (Fig. 5). Although some of the nuclei show a few larger aggregations of staining, most show a fine granular pattern. This observation indicates the mAbH1 antigen is expressed in somatic cells. If spheres are present in somatic

ported by the fact that both spheres and coiled bodies are nuclear structures and by the observation that SPH-1 is present in oocyte and somatic cell nuclei. Furthermore, coiled bodies and spheres are thought to contain snRNPs and snRNAs (Eliceiri et al., 1984; Fakan et al., 1984; Wu et al., 1991; Carmo-Fonseca et al., 1992). However, there are apparent differences between spheres and coiled bodies. For example, coiled bodies contain fibrillarin (Andrade et al., 1991) which has been detected in nucleoli (Lischwe et al., 1985) but has not been shown to be in spheres. Similarly, mAbSC35, which stains spheres (Wu et al., 1991), does not stain coiled bodies (Carmo-Fonseca et al., 1991). Hence, the similarity of SPH-1 to p80-coilin is interesting, but further studies are necessary to determine if spheres and coiled bodies are functionally similar.

The identification of SPH-1, together with other studies, more clearly defines the structure of spheres and sheds light on how this organelle is generated. Our working hypothesis is that the formation of spheres is analogous to the formation of nucleoli. Nucleoli arise as a result of rDNA transcription and accumulation of variously assembled pre-ribosomal subunits around the site of transcription (Sheer and Benavente, 1990; Fisher et al., 1991). We propose that spheres arise by a similar mechanism; that is, DNA at the sphere loci is transcribed and a sphere-specific ribonucleoprotein complex (RNP) is formed as a result of specific proteins binding to these nascent transcripts. The accumulation of this RNP gives spheres their distinctive morphology. This model is supported by common features shared by nucleoli and spheres. Each has been mapped to distinct genetic loci on newt lampbrush chromosomes (reviewed in Callan, 1986) and each has an electron dense core and a cortex that is less dense (Kezer et al., 1980; Goessens, 1984). Another feature common to spheres and nucleoli is that they both exist as chromosomally attached and extrachromosomal "free" structures in amphibian oocyte nuclei. Extrachromosomal nucleoli are formed as a result of amplification of rDNA during oogenesis, nucleating the formation of hundreds of free nucleoli (Brown and Dawid, 1968; Gall, 1968). It is possible that sphere DNA is similarly amplified, and that this amplified DNA gives rise to the free spheres. Alternatively, free spheres may arise via "shedding" of accumulated sphere RNP from the chromosomal sphere loci (Callan, 1986). If spheres do arise in the same way as nucleoli, then spheres may be sites of assembly of a distinct RNP. The identification of rDNA as the nucleolar organizer played a fundamental role in understanding the function of nucleoli; we presume that identification of the sphere organizer sequence will be of equal importance for understanding sphere function.

In addition to providing a tool for the direct characterization of spheres, the work presented here provides some useful insight into the possible function of spheres. First, the observation that SPH-1 is present in somatic cells, as well as oocytes, suggests that the function carried out by spheres is not restricted to oocytes, and is therefore likely to be general in nature. Second, preferential staining of the sphere core by mAbID7, which binds a family of pre-mRNA splicing factors called SR proteins (Zahler et al., 1992), suggests that RNA processing may occur in the sphere core. Consistent with this, mAbSC35 (Fu and Maniatis, 1990) which binds the

same set of SR proteins as mAbID7 (Neugebauer and Roth, unpublished observations), also stains the core of spheres (Wu et al., 1991). Finally, the observation that the antibody mAbH1 binds the cortex of spheres preferentially suggests that the sphere protein SPH-1 does not directly interact with SR proteins and is therefore unlikely to be involved in pre-mRNA splicing. The binding of the cortex by mAbH1 does suggest, however, that SPH-1 may be a component of an RNP produced by spheres.

We thank Judith Roan for her extensive help in screening hybridoma cell lines and for the *Xenopus* K2 tissue culture cell line. We would also like to thank Saeko Takano and Ed Chan and Zheng'an Wu and Joe Gall for providing their unpublished sequences of the full-length p80-coilin cDNA. We thank Ali Hammati Brivanlou and Doug Melton for the use of their *Xenopus* ovary cDNA library. We thank Karla Neugebauer, Janet Partridge, Laurie B. Roth, and Alan Zahler for their comments on the manuscript, as well as Tim Knight and Paul Goodwin for help with the production of figures.

R. S. Tuma was supported by National Research Service Award T32 GM07270. This work was supported by grant 42786-02 from the National Institutes of General Medical Sciences to M. B. Roth.

Received for publication 21 April 1993 and in revised form 3 June 1993.

References

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403-410.
- Andrade, L. E. C., E. K. L. Chan, I. Raska, C. L. Peebles, G. Roos, and E. M. Tan. 1991. Human autoantibody to a novel protein of the nuclear coiled body: immunological characterization and cDNA cloning of p80-coilin. *J. Exp. Med.* 173:1407-1419.
- Brown, D. D., and I. B. Dawid. 1968. Specific gene amplification in oocytes. *Science (Wash. DC)*. 160:272-280.
- Callan, H. G. 1986. Lampbrush chromosomes. In *Molecular Biology, Biochemistry and Biophysics*. M. Solioz, editor. Springer Verlag, Berlin, Heidelberg. 1-254.
- Callan, H. G., J. G. Gall, and C. A. Berg. 1987. The lampbrush chromosomes of *Xenopus laevis*: preparation, identification, and distribution of 5S DNA sequences. *Chromosoma*. 95:236-250.
- Carmo-Fonseca, M., R. Pepperkok, M. T. Cravalho, and Lamond, A. I. 1992. Transcription-dependent colocalization of the U1, U2, U4/U6, and U5 snRNPs in coiled bodies. *J. Cell Biol.* 117:1-14.
- Carmo-Fonseca, M., R. Pepperkok, B. S. Sproat, W. Ansorge, S. M. Swanson, and A. I. Lamond. 1991. In vivo detection of snRNP-rich organelles in the nuclei of mammalian cells. *EMBO (Eur. Mol. Biol. Organ.) J.* 10:1863-1873.
- Eliceiri, G. L., and J. S. Ryerse. 1984. Detection of intranuclear clusters of Sm antigens with monoclonal anti-Sm antibodies by immunoelectron microscopy. *J. Cell. Physiol.* 121:449-451.
- Evan, G. I., G. K. Lewis, G. Ramsay, and J. M. Bishop. 1985. Isolation of monoclonal antibodies specific for human c-myc proto-oncogene product. *Mol. Cell. Biol.* 5:3610-3616.
- Fakan, S., G. Leser, and T. E. Martin. 1984. Ultrastructural distribution of a nuclear ribonucleoprotein as visualized by immunocytochemistry on thin sections. *J. Cell Biol.* 98:358-363.
- Fisher, D., D. Weisenberger, and U. Scheer. 1991. Assigning functions to nucleolar structures. *Chromosoma*. 101:133-140.
- Fu, X.-D., and T. Maniatis. 1990. Factor required for mammalian spliceosome assembly is localized to discrete regions in the nucleus. *Nature (Lond.)*. 343:437-441.
- Gall, J. G. 1954. Lampbrush chromosomes from oocyte nuclei of the newt. *J. Morphol.* 94:283-352.
- Gall, J. G. 1968. Differential synthesis of the genes for ribosomal RNA during amphibian oogenesis. *Proc. Natl. Acad. Sci. USA*. 60:553-560.
- Gall, J. G., and H. G. Callan. 1989. The sphere organelle contains small nuclear ribonucleoproteins. *Proc. Natl. Acad. Sci. USA*. 86:6635-6639.
- Goessens, G. 1984. Nucleolar structure. *Int. Rev. Cytol.* 87:107-158.
- Gurdon, J. B., C. D. Lane, H. R. Woodland, and G. Maribaux. 1971. Use of frog eggs and oocytes for the study of messenger RNA and its translation in living cells. *Nature (Lond.)*. 233:177-182.
- Harlow, E., and D. Lane. 1988. *Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. 726 pp.
- Henikoff, S. 1987. Unidirectional digestion with exonuclease III in DNA sequence analysis. *Methods Enzymol.* 155:156-165.
- Henikoff, S., and J. G. Henikoff. 1991. Automated assembly of protein blocks for database searching. *Nucleic Acids Res.* 19:6565-6572.

1. Abbreviation used in this paper: RNP, ribonucleoprotein.

- Hervás, J. P., J. Villegas, D. Crespo, and M. LaFarga. 1980. Coiled bodies in supraoptic nuclei of the rat hypothalamus during the post natal period. *Am. J. Anat.* 159:447-454.
- Kezer, J., P. E. León, and S. K. Sessions. 1980. Structural differentiation of the meiotic and mitotic chromosomes of the salamander, *Ambystoma macrodactylum*. *Chromosoma (Berlin)*. 81:177-197.
- Krainer, A. 1988. Pre-mRNA splicing by complementation with purified human U1, U2, U4/U6 and U5 snRNPs. *Nucleic Acids Res.* 16:9415-9429.
- Lacroix, J. C., R. Azzouz, D. Boucher, C. Abbadie, C. K. Pyne, and J. Charlemagne. 1985. Monoclonal antibodies to lampbrush chromosome antigens of *Pleurodeles waltlii*. *Chromosoma (Berlin)*. 92:69-80.
- LaFarga, M., J. P. Hervás, M. C. Santa-Cruz, J. Villegas, and D. Crespo. 1983. The "accessory body" of Cajal in the neuronal nucleus. *Anat. Embryol.* 166:19-30.
- Lerner, E. A., M. R. Lerner, C. A. Janeway, and J. A. Steitz. 1981. Monoclonal antibodies to nucleic acid-containing cellular constituents: probes for molecular biology and autoimmune disease. *Proc. Natl. Acad. Sci. USA*. 78:2737-2741.
- Lischwe, M. A., R. L. Ochs, R. Reddy, R. G. Cook, L. C. Yeoman, E. M. Tan, M. Reichlin, and H. Busch. 1985. Purification and partial characterization of a nucleolar scleroderma antigen ($M_r = 34,000$; pI, 8.5) rich in N_G,N_G -dimethylarginine. *J. Biol. Chem.* 260:14304-14310.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York. 545 pp.
- McStay, B., and R. H. Reeder. 1990. An RNA polymerase I termination site can stimulate the adjacent ribosomal gene promoter by two distinct mechanisms. *Genes Dev.* 4:1240-1252.
- Munro, S., and H. R. B. Pelham. 1984. Use of peptide tagging to detect proteins expressed from cloned genes: deletion mapping functional domains of *Drosophila hsp70*. *EMBO (Eur. Mol. Biol. Organ.) J.* 3:3087-3093.
- Rebagliati, M. R., D. L. Weeks, R. P. Harvey, and D. A. Melton. 1985. Identification and cloning of localized maternal RNAs from *Xenopus* eggs. *Cell*. 42:769-777.
- Roth, M. B., and J. G. Gall. 1987. Monoclonal antibodies that recognize transcription unit proteins on newt lampbrush chromosomes. *J. Cell Biol.* 105:1047-1054.
- Roth, M. B., and J. G. Gall. 1989. Targeting of a chromosomal protein to the nucleus and to lampbrush chromosome loops. *Proc. Natl. Acad. Sci. USA*. 86:1269-1272.
- Roth, M. B., A. M. Zahler, and J. A. Stolk. 1991. A conserved family of nuclear phosphoproteins localized to sites of polymerase II transcription. *J. Cell Biol.* 115: 587-596.
- Sheer, U., and R. Benavente. 1990. Functional and dynamic aspects of the mammalian nucleolus. *Bioessays*. 12:14-21.
- Short, J. M., J. M. Fernandez, J. A. Sorge, and W. D. Huse. 1988. Lambda zap: a bacteriophage lambda expression vector with in vitro excision properties. *Nucleic Acids Res.* 16:7583-7600.
- Taggart, R. T., and I. M. Samloff. 1983. Stable antibody-producing murine hybridomas. *Science (Wash. DC)*. 219:1228-1230.
- Wu, Z., C. Murphy, H. G. Callan, and J. Gall. 1991. Small nuclear ribonucleoproteins and heterogeneous nuclear ribonucleoproteins in the amphibian germinal vesicle: loops, spheres, and snurposomes. *J. Cell Biol.* 113:465-483.
- Zahler, A. M., W. S. Lane, J. A. Stolk, and M. B. Roth. 1992. SR proteins: a conserved family of pre-mRNA splicing factors. *Genes Dev.* 6:837-847.