

Insulin Resistance in Obese Children and Adolescents: HOMA-IR Cut-Off Levels in the Prepubertal and Pubertal Periods

Selim Kurtoğlu¹, Nihal Hatipoğlu¹, Mümtaz Mazıcıoğlu², Mustafa Kendirci¹,
Mehmet Keskin³, Meda Kondolot⁴

¹Erciyes University Faculty of Medicine, Department of Pediatric Endocrinology, Kayseri, Turkey

²Erciyes University Faculty of Medicine, Department of Family Medicine, Kayseri, Turkey

³Gaziantep University Faculty of Medicine, Department of Pediatric Endocrinology, Gaziantep, Turkey

⁴Erciyes University Faculty of Medicine, Department of Pediatrics, Unit of Social Pediatrics, Kayseri, Turkey

ABSTRACT

Objective: Childhood obesity is associated with an increased risk for insulin resistance. The underlying mechanism for the physiological increase in insulin levels in puberty is not clearly understood. The aim of the present study was to determine the cut-off values for homeostasis model assessment for insulin resistance (HOMA-IR) in obese children and adolescents according to gender and pubertal status.

Methods: Two hundred and eight obese children and adolescents (141 girls, 127 boys) aged between 5 and 18 years were included in the study. The children were divided into prepubertal and pubertal groups. A standard oral glucose tolerance test (OGTT) was carried out in all children. A total insulin level exceeding 300 µU/mL in the blood samples, collected during the test period, was taken as the insulin resistance criterion. Cut-off values for HOMA-IR were calculated by receiver operating characteristic (ROC) analysis.

Results: In the prepubertal period, the rate of insulin resistance was found to be 37% in boys and 27.8% in girls, while in the pubertal period, this rate was 61.7% in boys and 66.7% in girls. HOMA-IR cut-off values for insulin resistance in the prepubertal period were calculated to be 2.67 (sensitivity 88.2%, specificity 65.5%) in boys and 2.22 (sensitivity 100%, specificity 42.3%) in girls, and in the pubertal period, they were 5.22 (sensitivity 56%, specificity 93.3%) in boys and 3.82 (sensitivity 77.1%, specificity 71.4%) in girls.

Conclusions: Since gender, obesity and pubertal status are factors affecting insulin resistance, cut-off values which depend on gender and pubertal status, should be used in evaluation of insulin resistance.

Key words: Obesity, insulin resistance, pubertal status, gender, HOMA-IR cut-off values

Conflict of interest: None declared

Received: 17.03.2010

Accepted: 23.06.2010

Introduction

Insulin resistance and glucose intolerance are frequent in obese children and adolescents and lead to a significant risk for hypertension and cardiovascular diseases, as well as for type 2 diabetes (1,2,3,4). The development of type 2 diabetes mellitus is induced by the decreased insulin sensitivity, which leads to increased insulin production. This imbalance causes a predisposition to several metabolic disorders such as early atherosclerosis, progressive obesity, acanthosis nigricans, increase in skin tags, hypertension, dyslipidemia, fatty liver and polycystic ovarian syndrome (5). Obesity and insulin resistance, previously considered as a problem of older ages, are becoming serious issue also in the pediatric age group. Early detection of insulin resistance is important for prevention of these complications.

Various methods to define insulin sensitivity have been developed. Among other models, euglycemic clamp and modified minimal model are considered to be gold standards. However, they are complex and invasive tests, which can be used for research purposes only (6,7). Oral glucose tolerance test (OGTT) has been shown to be as reliable as a frequently sampled intravenous glucose tolerance test (FSIVGTT) and a clamp test in determining insulin sensitivity (8,9). However,

Address for Correspondence

Nihal Hatipoğlu, Department of Pediatric Endocrinology, Erciyes University Faculty of Medicine, Kayseri, Turkey

Phone: +90 536 323 03 02 Fax: +90 352 437 52 85 E-mail: nihalhatipoglu@yahoo.com

© Journal of Clinical Research in Pediatric Endocrinology, Published by Galenos Publishing.

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

the use of OGTT in large populations is limited. Therefore, methods such as fasting insulin level, fasting glucose/insulin ratio (FGIR), homeostasis model assessment for insulin resistance (HOMA-IR) and quantitative insulin-sensitivity check index (QUICKI) are frequently used in population screening (7,10,11,12). HOMA-IR was found to be much more reliable than FGIR and QUICKI in determining insulin resistance in obese children (13). HOMA-IR is a frequently used parameter in clinical research (12,14).

Studies have shown that a physiological transient insulin resistance develops in the pubertal period although its cause is not fully understood. Despite the fact that HOMA-IR cut-off limit for insulin resistance is accepted as 2.5 in adults, corresponding values in prepubertal and pubertal children and adolescents have not been reported (10). It is known that the frequency of insulin resistance varies between genders and also among races (15,16,17,18,19,20). However, to our knowledge, there are no studies defining cut-off levels for HOMA-IR in children of prepubertal and pubertal ages (14,21,22,23).

The aim of the present study was to determine HOMA-IR cut-off values in obese children and adolescents according to gender and pubertal status.

Methods

Two hundred and sixty eight obese children and adolescents (141 girls, 127 boys) aged between 5 and 18 years were included in the study. All patients had presented with obesity to the Pediatric Endocrinology Clinic at Erciyes University, Faculty of Medicine. Those with an underlying endocrinologic disease or/and those under medication were excluded from the study. The children included in the study had normal thyroid function tests and morning fasting cortisol levels. Ethical approval was obtained from the Institutional Review Board of Erciyes University, Faculty of Medicine. Parental consent was taken for additional blood sampling to determine insulin levels. Body mass index (BMI) of the patients was calculated using the equation: $BMI = \text{Body weight (Kg)}/\text{Height (m)}^2$. Subjects, whose BMI values were above the 95th percentile of the BMI reference curve adjusted to Turkish children according to age and gender, were accepted as obese (24). Detailed history was taken from the parents, and the subjects underwent a detailed physical examination and pubertal evaluation performed by the same pediatric endocrinologist. A testicular volume equal to or greater than 4 ml in boys and onset of breast development in girls were accepted as the criteria for onset of puberty.

OGTT was carried out in order to determine insulin resistance. Following a 3-day high carbohydrate diet (300 g/day) and overnight fasting, an oral dose of 1.75 g/kg (maximum 75 g) glucose was given, and blood samples were taken at 0, 30, 60, 90 and 120 minutes from a venous catheter for glucose and insulin assessments (25,26).

A total (the sum of insulin levels at 0, 30, 60, 90, and 120 minutes during the OGTT) insulin level exceeding 300 $\mu\text{U}/\text{mL}$ was taken as hyperinsulinemia. This method was first introduced by Maruhama et al (25). The main reason for selecting this criterion for hyperinsulinemia was to determine a cut-off for HOMA-IR as an index of insulin resistance. At each point in OGTT, both glucose and insulin levels were measured, and then, total insulin levels exceeding 300 $\mu\text{U}/\text{mL}$ were recorded as hyperinsulinemia. Glucose levels at 120th minute were taken as a criterion for impaired glucose tolerance or diabetes mellitus. HOMA-IR was calculated using the equation: $\text{HOMA-IR} = \text{Fasting insulin } (\mu\text{U}/\text{mL}) \times \text{Fasting glucose (mg/dL)} / 405$ (10).

Biochemical Analysis

Plasma glucose levels were measured by the glucose oxidase method and the modified Trinder color reaction catalyzed by the peroxidase enzyme. Insulin levels were measured with a immunoradiometric assay kit (INS-Irma Biosource, Nivelles, Belgium). The assay detection limit was 1 $\mu\text{U}/\text{mL}$, and intra- and interassay coefficients of variation were 2.2% and 6.5%, respectively.

Statistical Analysis

All statistical analyses were carried out using the Statistical Package for Social Sciences (SPSS version 15.0 for Windows, Chicago, IL) and MedCalc (version 7.4.1.1, written by Frank Schoonjans, Mariakerke, Belgium). For the diagnosis of insulin resistance, HOMA-IR values were obtained with the receiver operating characteristic (ROC) analysis. Total insulin values exceeding 300 $\mu\text{U}/\text{mL}$ during the OGTT were accepted as insulin resistance, and ROC analysis was undertaken to determine HOMA-IR cut-off values accordingly. ROC analysis is used to assess the actual correctness of research results. Among the terms used in this analysis, sensitivity is the ratio of patients with positive results (TP) to the number of patients, while specificity is the ratio of healthy individuals with negative results (TN) to the number of healthy individuals (27). ROC analysis is a standard approach used for determining the sensitivity and specificity of the diagnostic procedure. ROC curves to find out the relationship between the sensitivity and specificity of the diagnosis are used for this purpose. The curves are between the limits 0 and 1, and while proximity to the y coordinate and the upper limit shows a successful test, curves with slopes like 45° show a failed test. Thus, the success of the test can be evaluated by examining the ROC curves. In a successful test, the area under the curve is expected to be great.

For comparisons between those with and without insulin resistance, the Student t-test was used in parametric groups and Mann-Whitney U test in non-parametric groups. Level of significance was accepted as $p < 0.05$.

Results

Forty-six boys (46.2%) were evaluated as prepubertal and 81 (63.8%) as pubertal. Of the girls, 36 (25.5%) were

evaluated as prepubertal and 105 (74.5%) as pubertal. In the prepubertal groups, the mean age was 8.9±1.8 years in boys and 8.3±1.4 years in girls, and the mean BMI was 28.2±5.4 kg/m² in boys and 26.2±5.8 kg/m² in girls. In the pubertal groups, the mean age was 13.6±1.6 years in boys and 13.2±2.0 years in girls, and the mean BMI was 30.9±4.9 kg/m² in boys and 30.4±5.0 kg/m² in girls. Chronological ages, BMI values, fasting blood sugar and insulin values, blood sugar and insulin values at 120 minutes, total insulin values measured during OGTT, FGIR and HOMA-IR values calculated according to gender and pubertal status are given in Tables 1 and 2.

Following OGTT, the rate of insulin resistance in the prepubertal period was 37% (n=17) in boys and 27.8% (n=10) in girls. In the pubertal children, these rates were 61.7% (n=50) in boys and 66.7% (n=70) in girls.

When the groups were separated as those with and without insulin resistance, there were no differences regarding age and BMI values, except for the girls in the pubertal period (Tables 1,2).

We also compared the hyperinsulinemic and nonhyperinsulinemic groups for blood glucose levels at 0 and 120th minutes of OGTT. We could not find any difference in pre- and post-prandial blood glucose level in boys neither in the prepubertal nor in the pubertal groups. In prepubertal girls, we could not find any difference in fasting blood glucose level, but detected a difference in the 120-minute glucose level. In pubertal girls, there was a significant difference between 0- and 120-minute blood glucose levels between insulin resistant and nonresistant groups (Table 2).

HOMA-IR, fasting and 120-minute insulin values, FGIR and total insulin values were significantly different between the subjects of both sexes with and without insulin resistance both in the prepubertal and pubertal groups (Tables 1, 2).

HOMA-IR cut-off values for insulin resistance were calculated to be 2.67 (sensitivity 88.2%, specificity 65.5%) in boys and 2.22 (sensitivity 100%, specificity 42.3%) in girls in the prepubertal period, and 5.22 (sensitivity 56%, specificity 93.3%) in boys and 3.82 (sensitivity 77.1%, specificity 71.4%) in girls in the pubertal period.

Fasting insulin levels above 15 µU/mL in the prepubertal period, 30 µU/mL in the pubertal period and 20 µU/mL in the postpubertal period, FGIR above 6, 120-minute insulin >75 µU/mL during OGTT and peak insulin above 150 µU/mL are recommended as cut-off levels for hyperinsulinism and consequently as parameters showing insulin resistance (5,28,29). High blood sugar values at minute 120 of OGTT is an indicator of impaired glucose tolerance and therefore of insulin resistance (5). Each of these parameters (fasting insulin, insulin and blood sugar at 120th minute, FGIR and HOMA-IR values) were compared using the ROC analysis regarding their importance in determination of insulin resistance according to pubertal status and gender (Figures 1, 2). With the exception of prepubertal girls, HOMA-IR index was found to be the best determinant of insulin resistance in sub groups. On the other hand, insulin level at 120th minute was the best indicator of insulin resistance in the prepubertal girls. Areas under the curves in the ROC analysis of each indicator of insulin resistance according to pubertal status and gender, and 95% confidence intervals are given in Table 3.

Table 1. Findings in prepubertal boys and girls with and without hyperinsulinism

PREPUBERTAL	BOYS				P	GIRLS				P
	Hyperinsulinism (n=17)		No hyperinsulinism (n=29)			Hyperinsulinism (n=10)		No hyperinsulinism (n=26)		
	Mean±SD	Range	Mean±SD	Range		Mean±SD	Range	Mean±SD	Range	
Age (years)	9.0±1.3	7-11	8.8±2.0	5.00-11.1	0.936	8.4±1.6	5-10	8.1±1.4	5.7-10	0.355
BMI (kg/m ²)	29.5±7.2	20.9-51.7	27.4±3.9	20.1-37.1	0.419	27.3±5.6	23.0-41.2	25.7±5.9	19.7-40.8	0.109
HOMA-IR	8.8±7.6	0.7-24.8	2.9±2.0	0.4-10.7	<0.001	5.0±2.1	2.3-37.5	3.3±2.1	0.7-8.2	0.034
Fasting insulin (µU/mL)	42.0±36.6	4.2-118.7	14.7±9.6	2.2-51.0	0.001	25.3±9.0	12.0-37.5	17.1±11.0	3.4-44.0	0.026
120-min insulin (µU/mL)	135.9±174.7	4.6-581	35.9±24.1	4.0-90.9	0.015	113.8±36.9	38.7-180.3	34.5±18.1	1.5-74.1	<0.001
Fasting blood sugar (mg/dL)	84.6±8.9	67.0-100.0	80.1±8.7	60.0-96.0	0.116	79.4±7.4	68-89	78.1±9.2	66.0-97.0	0.497
120-min blood sugar (mg/dL)	111.1±15.8	86.0-148.0	100.6±19.4	64-138	0.053	125.4±16.8	105.0-162.1	99.8±11.5	79.0-120.0	<0.001
FGIR	4.1± 3.8	0.7-16.1	8.8±8.6	1.7-37.1	0.004	4.0±1.7	2.2-6.5	7.3±5.9	1.5-23.9	0.028
Total insulin (µU/mL)	530.4±311.6	312.4-1333.2	159.4±71.1	18.4-298.6	<0.001	578.1±130.5	320.3-769.7	170.3±70.3	10.9-249.9	<0.001

BMI: Body mass index, HOMA-IR:Homeostasis model assessment for insulin resistance, FGIR: Fasting glucose/ insulin ratio, Range:Minimum-maximum values.

Discussion

Obesity is a critical risk factor for the development of insulin resistance (30, 31). Total body fat is important in the development of insulin resistance in children (30,32). Previous studies have examined the relationship between BMI and

insulin sensitivity and have shown a strong negative correlation between the values found with the euglycemic clamp test and FSIVGTT and BMI (18, 33). In our study, no difference was found in BMI values between groups with and without insulin resistance, except for pubertal girls. This finding indicates that BMI values are not always correlated with body fat ratio.

Table 2. Findings in pubertal boys and girls with and without hyperinsulinism

PUBERTAL	Boys				P	Girls				P
	Hyperinsulinism (n=50)		No hyperinsulinism (n=31)			Hyperinsulinism (n=70)		No hyperinsulinism (n=35)		
	Mean±SD	Range	Mean±SD	Range		Mean±SD	Range	Mean±SD	Range	
Age (years)	13.7±1.6	11.3-17.5	13.6±1.7	11.3-18.0	0.889	13.2±2.0	10.2-17.5	13.1±2.0	10.1-18.0	0.804
BMI (kg/m ²)	30.8±4.6	21.9-41.9	31.1±5.3	24.6-46.5	0.817	31.1±5.3	22.5-49.3	28.9±4.4	22.6-43.0	0.043
HOMA-IR	7.2±4.6	1.3-19.1	3.4±1.7	0.8-7.9	<0.001	7.4±5.5	1.6-28.8	3.2±1.7	0.1-8.4	<0.001
Fasting insulin (µU/mL)	35.3±21.6	7.9-91.5	18.1±9.3	4.2-40.2	<0.001	35.3±25.5	8.6-131.0	16.8±8.5	0.6-39.3	<0.001
120-min insulin (µU/mL)	113.5±114.3	6.3-516.9	34.9±31.6	2.3-127.0	<0.001	126.5±132.5	7.6-858.2	39.2±28.5	1.5-129.4	<0.001
Fasting blood sugar(mg/dL)	82.0±10.1	64.0-106.0	77.2±12.1	42.0-97.0	0.054	83.9±11.2	61.0-122.0	76.9±9.0	54-97	0.002
120-min blood sugar(mg/dL)	112.9±24.0	61.0-165	109.3±22.3	60.0-146.0	0.509	111.1±24.6	60.0-195.0	100.6±18.6	63.3-147	0.037
FGIR	3.2±1.9	0.8-8.9	5.6±3.5	1.3-17.5	<0.001	3.3±1.7	0.6-9.0	8.5±16.1	2.2-98.2	0.064
Total insulin (µU/mL)	600.0±38.1	309.7±2136	197.1±71.6	29.3-293.3	<0.001	655.0±490.0	302.0-3628.0	202.4±74.6	39.5-287.5	<0.001

BMI: Body mass index, HOMA-IR:Homeostasis model assessment for insulin resistance, FGIR: Fasting glucose/ insulin ratio, Range: Minimum-maximum values

Table 3. Areas under the curves in the ROC analysis of each indicator of insulin resistance according to pubertal status and gender, and 95% confidence intervals.

	Boys		Girls		
	AUC	95% CI	AUC	95% CI	
Prepubertal	HOMA-IR	0.800	0.653-0.904	0.729	0.550-0.866
	Insulin 0 th minute (µU/mL)	0.778	0.629-0.888	0.729	0.550-0.866
	Insulin 120 th minute (µU/mL)	0.698	0.543-0.826	0.954	0.821-0.994
	Glucose 120 th minute (mg/dL)	0.676	0.520-0.807	0.921	0.775-0.984
	FGIR	0.739	0.587-0.858	0.725	0.545-0.863
Pubertal	HOMA-IR	0.784	0.673-0.865	0.836	0.748-0.903
	Insulin 0 th minute (µU/mL)	0.766	0.658-0.853	0.803	0.711-0.876
	Insulin 120 th minute (µU/mL)	0.784	0.678-0.868	0.798	0.705-0.872
	Glucose 120 th minute (mg/dL)	0.531	0.416-0.644	0.619	0.516-0.715
	FGIR	0.745	0.635-0.836	0.535	0.432-0.636

AUC: Areas under the curves in the ROC analysis
FGIR: Fasting glucose/ insulin ratio
HOMA-IR: homeostasis model assessment for insulin resistance, CI: Confidence Interval

A transient insulin resistance develops in children during puberty. This insulin resistance emerging during pubertal maturation is accepted as a physiological condition rather than pathologic (32,34). Decreased insulin sensitivity in the pubertal period causes a compensatory increase in insulin secretion (20). Cross-sectional studies have shown that insulin resistance increases with the onset of puberty, makes a peak at Tanner stage 3 and recedes to prepubertal levels at the end of puberty (15,16,18). Longitudinal studies have found a 30% decrease in insulin sensitivity between

Tanner stages 1 and 5 (29). However, this decrease was found to return to normal at the end of puberty (35). The reasons for these changes in insulin activation and secretion are not fully understood, but are thought to be related to a mechanism enhancing the anabolic effect of insulin and growth hormone during rapid somatic growth (15,36).

Although some studies have suggested that this change in insulin sensitivity is due to the alterations in the pubertal distribution of fat, subsequent studies have shown similar decreases in insulin sensitivity both in thin and obese children




Figure 1. Comparison of the parameters used in determination of insulin resistance with ROC analysis in prepubertal subjects




Figure 2. Comparison of the parameters used in determination of insulin resistance with ROC analysis in pubertal subjects

(20, 37). Furthermore, it was shown that while body fat continuously increased before and during puberty, transient insulin elevation was seen in mid puberty with a return to normal levels at the end of puberty (16). Therefore, it is not possible to explain insulin insensitivity with the change in fat distribution only.

While type 2 diabetes was previously thought to occur in elderly adults, it is now known that it also occurs in children with an increased incidence of obesity. It is claimed that adolescence is a risky period for the development of type 2 diabetes, because the transient physiological status in insulin resistance induces an extra stress on the beta cells in the pancreas (32). Therefore, adolescence is even more important for obese children regarding the risk of developing type 2 diabetes. In children of pubertal age, evaluation of insulin resistance as pathological or physiological and early intervention will help preserve the beta cell function of the pancreas. Since both obesity and puberty have a role in the development of insulin resistance, detection of increased insulin resistance of a pathological degree is of great importance, especially in obese adolescents. HOMA-IR cut-off values, which are among the most important indicators in determining insulin resistance in obese individuals, are expected to be different in prepubertal and pubertal children. In our study, when the results were compared according to the pubertal status, HOMA-IR threshold values were found to be significantly higher in the pubertal period than before puberty, in both genders. Gender has also been shown to have an effect on insulin sensitivity. Moran et al (16) reported greater insulin resistance in girls compared to boys in a large group of 357 children with the euglycemic clamp test even after adjusting for triceps and subcapsular skin thickness, BMI, waist circumference and hip circumference. Also, in a population-based study by Lee et al (38), insulin resistance was found to be greater in girls than in boys after adjusting for weight, race and race/ethnicity.

In our study, while no difference for gender was detected in HOMA-IR cut-off values in the prepubertal period, cut-off values in the pubertal period were higher in boys than in girls. Because insulin resistance is greater in girls, HOMA-IR cut-off values are expected to be lower in boys. We do not have an explanation for this gender differences. However, it is important that, in the evaluation of insulin resistance in the pubertal period, different threshold values should be used for boys and girls.

Moran et al (16) reported that gender differences for insulin resistance disappeared among children with BMI values greater than 27 kg/m² and therefore, gender difference can become insignificant in morbid obesity. In our study, no difference between genders was observed despite the mean BMI values around 30 kg/m² in all children.

Due to the fact that our sample size was small, we are not able to propose precise cut-off limits based on the results of this study. However, based on this experience, we believe that determination of fasting glucose and insulin levels in epidemiological studies will provide much more accurate results with which we can calculate HOMA-IR.

Another weak point of this study is that we have not performed a clamp study, which is the gold standard. Thirdly, we have taken the same criteria for insulin resistance in pre-and pubertal children on OGTT. However, there are no established separate values for these periods.

In conclusion, obesity and puberty are important factors for the development of insulin resistance. Both gender and pubertal status should be considered when evaluating insulin resistance in obese children, and risk groups should be identified accordingly. Gender differences should also be considered in the evaluation of insulin resistance. We recommend that HOMA-IR cut-off values, which are among the most widely used indicators of insulin resistance, be used in this evaluation, also taking into account both pubertal status and gender differences. Thus, particularly in a period critical for the development of type 2 diabetes such as the pubertal period, risk groups can be determined in a more reliable manner, and measures can be taken.

References

1. Invitti C, Guzzaloni G, Gilardini L, Morabito F, Viberti G. Prevalence and concomitants of glucose intolerance in European obese children and adolescents. *Diabetes Care* 2003;26:118-124. [Abstract] / [Full Text] / [PDF]
2. Ferrannini E, Buzzigoli G, Bonadonna R, Giorico MA, Oleggini M, Graziadei L, Pedrinelli R, Brandi L, Bevilacqua S. Insulin resistance in essential hypertension. *N Engl J Med* 1987;317:350-357. [Abstract]
3. Martin BC, Warram JH, Krolewski AS, Bergman RN, Soeldner JS, Kahn CR. Role of glucose and insulin resistance in development of type 2 diabetes mellitus: results of a 25-year follow-up study. *Lancet* 1992;340:925-929. [Abstract] / [PDF]
4. Kahn SE, Prigeon RL, McCulloch DK, Boyko EJ, Bergman RN, Schwartz MW, Neifing JL, Ward WK, Beard JC, Palmer JP. Quantification of the relationship between insulin sensitivity and beta-cell function in human subjects. Evidence for a hyperbolic function. *Diabetes* 1993;42:1663-1672. [Abstract]
5. Ten S, Maclaren N. Insulin resistance syndrome in children. *J Clin Endocrinol Metab* 2004;89:2526-2539. [Abstract] / [Full Text] / [PDF]
6. DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 1979;237:214-223. [Abstract] / [PDF]
7. Bergman RN. Towards physiological understanding of glucose tolerance-minimal model approach. *Diabetes* 1989;38:1512-1527. [Abstract]
8. Avignon A, Boegner C, Mariano-Goulart D, Colette C, Monnier L. Assessment of insulin sensitivity from plasma insulin and glucose in the fasting or post oral glucose-load state. *Int J Obes Relat Metab Disord* 1999;23:512-517. [Abstract] / [PDF]

9. Matsuda M, DeFronzo RA. Insulin sensitivity indices obtained from oral glucose tolerance testing: comparison with the euglycemic insulin clamp. *Diabetes Care* 1999;22:1462-1470. [[Abstract](#)] / [[PDF](#)]
10. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412-419. [[Abstract](#)]
11. Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, Quon MJ. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 2000;85:2402-2410. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
12. Cutfield WS, Jefferies CA, Jackson WE, Robinson EM, Hofman PL. Evaluation of HOMA and QUICKI as measures of insulin sensitivity in prepubertal children. *Pediatr Diabetes* 2003;4: 119-125. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
13. Keskin M, Kurtoglu S, Kendirci M, Atabek ME, Yazici C. Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics* 2005;115:500-503. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
14. Wallace TM, Matthews DR. The assessment of insulin resistance in man. *Diabet Med* 2002; 19: 527-534. [[Abstract](#)]
15. Caprio S, Cline G, Boulware S, Permanente C, Shulman GI, Sherwin RS, Taborlane WV. Effects of puberty and diabetes on metabolism of insulin-sensitive fuels. *Am J Physiol* 1994;266:885-891. [[Abstract](#)] / [[PDF](#)]
16. Moran A, Jacobs DR, Steinberger J, Hong CP, Prineas R, Luepker R, Sinaiko AR. Insulin resistance during puberty: results from clamp studies in 357 children. *Diabetes* 1999;48:2039-2044. [[Abstract](#)] / [[PDF](#)]
17. Bloch CA, Clemons P, Sperling MA. Puberty decreases insulin sensitivity. *J Pediatr* 1987;110:481-487. [[Abstract](#)]
18. Amiel SA, Sherwin RS, Simonson DC, Lauritano AA, Tamborlane WV. Impaired insulin action in puberty. A contributing factor to poor glycemic control in adolescents with diabetes. *N Engl J Med* 1986;315:215-219. [[Abstract](#)]
19. Travers SH, Jeffers BW, Bloch CA, Hill JO, Eckel RH. Gender and Tanner stage differences in body composition and insulin sensitivity in early pubertal children. *J Clin Endocrinol Metab* 1995;80:172-178. [[Abstract](#)] / [[PDF](#)]
20. Caprio S, Plewe G, Diamond MP, Simonson DC, Boulware SD, Sherwin RS, Tamborlane WV. Increased insulin secretion in puberty: a compensatory response to reductions in insulin sensitivity. *J Pediatr* 1989;114:963-967. [[Abstract](#)]
21. Tresaco B, Bueno G, Pineda I, Moreno LA, Garagorri JM, Bueno M. Homeostatic model assessment (HOMA) index cut-off values to identify the metabolic syndrome in children. *J Physiol Biochem* 2005;61:381-388. [[Abstract](#)]
22. Atabek ME, Pirgon O. Assessment of insulin sensitivity from measurements in fasting state and during an oral glucose tolerance test in obese children. *J Pediatr Endocrinol Metab* 2007;20:187-195. [[Abstract](#)]
23. Radikova Z, Koska J, Huckova M, Ksinantova L, Imrich R, Vigas M, Trnovec T, Langer P, Sebkova E, Klimes I. Insulin sensitivity indices: a proposal of cut-off points for simple identification of insulin-resistant subjects. *Exp Clin Endocrinol Diabetes* 2006;114:249-256. [[Abstract](#)]
24. Ozturk A, Mazicioglu MM, Hatipoglu N, Budak N, Keskin G, Yazlak Z, Balci N, Yildiz H, Yildiz K, Ustunbas HB, Kurtoglu S. Reference body mass index curves for Turkish children 6 to 18 years of age. *J Pediatr Endocrinol Metab* 2008;21:827-836. [[Abstract](#)]
25. Maruhama Y, Abe R. A familial form of obesity without hyperinsulinism at the outset. *Diabetes* 1981;30:14-18. [[Abstract](#)]
26. Zannoli R, Rebggiani A, Chiarelli F, Morgese G. Hyperinsulinism as a marker in obese children. *Am J Dis Child* 1993;147:837-841. [[Abstract](#)]
27. Metz CE. "Basic principles of ROC analysis", *Sem Nuc Med* 1978;8:283-298. [[Abstract](#)]
28. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539-553. [[Abstract](#)]
29. Goran MI, Gower BA. Longitudinal study on pubertal insulin resistance. *Diabetes* 2001;50:2444-2450. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
30. Weiss R. Insulin sensitivity and secretion: swaying the pendulum. *J Pediatr* 2006;148:3-4. [[Full Text](#)] / [[PDF](#)]
31. Haymond MW. Measuring insulin resistance: a task worth doing. But how? *Pediatr Diabetes* 2003;4:115-118. [[Abstract](#)] / [[PDF](#)]
32. Goran MI, Ball GD, Cruz ML. Obesity and risk of type 2 diabetes and cardiovascular disease in children and adolescents. *J Clin Endocrinol Metab* 2003;88:1417-1427. [[Abstract](#)] / [[PDF](#)]
33. Cook JS, Hoffman RP, Stene MA, Hansen JR. Effects of maturational stage on insulin sensitivity during puberty. *J Clin Endocrinol Metab* 1993;77:725-730. [[Abstract](#)] / [[PDF](#)]
34. Arslanian SA, Kalhan SC. Protein turnover during puberty in normal children. *Am J Physiol* 1996;270:79-84. [[Abstract](#)] / [[PDF](#)]
35. Ball GD, Huang TT, Gower BA, Cruz ML, Shaibi GQ, Weigensberg MJ, Goran MI. Longitudinal changes in insulin sensitivity, insulin secretion, and beta-cell function during puberty. *J Pediatr* 2006;148:16-22. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
36. Caprio S, Jones J, Tamborlane W. Developmental changes in insulin action and secretion in childhood health and disease. *Adv Endocrinol Metab* 1994;5:171-201.
37. Hoffman RP, Vicini P, Sivitz WI, Cobelli C. Pubertal adolescent male-female differences in insulin sensitivity and glucose effectiveness determined by the one compartment minimal model. *Pediatr Res* 2000;48:384-388. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]
38. Lee JM, Okumura MJ, Davis MM, Herman WH, Gurney JG. Prevalence and determinants of insulin resistance among U.S. adolescents: a population based study. *Diabetes Care* 2006;29:2427-2432. [[Abstract](#)] / [[Full Text](#)] / [[PDF](#)]