

## Oncology

## Angiosarcoma of the Adrenal Gland Treated Using a Multimodal Approach



Jay G. Fuletra<sup>a</sup>, Benjamin T. Ristau<sup>a</sup>, Barton Milestone<sup>b</sup>, Harry S. Cooper<sup>c</sup>, Alysia Browne<sup>c</sup>, Sujana Movva<sup>d</sup>, Thomas J. Galloway<sup>e</sup>, Randall Lee<sup>a</sup>, Nikhil Waingankar<sup>a</sup>, Alexander Kutikov<sup>a,\*</sup>

<sup>a</sup> Division of Urologic Oncology, Fox Chase Cancer Center-Temple University Health System, Philadelphia, PA, USA

<sup>b</sup> Department of Radiology, Fox Chase Cancer Center-Temple University Health System, Philadelphia, PA, USA

<sup>c</sup> Department of Pathology, Fox Chase Cancer Center-Temple University Health System, Philadelphia, PA, USA

<sup>d</sup> Department of Medical Oncology, Fox Chase Cancer Center-Temple University Health System, Philadelphia, PA, USA

<sup>e</sup> Department of Radiation Oncology, Fox Chase Cancer Center-Temple University Health System, Philadelphia, PA, USA

## ARTICLE INFO

## Article history:

Received 15 October 2016

Accepted 1 November 2016

## Keywords:

Angiosarcoma

Adrenal cancer

Adjuvant radiation

Adjuvant chemotherapy

## ABSTRACT

Angiosarcoma of the adrenal gland is an extremely rare malignancy. We report a case of a 59-year-old female who presented with abdominal pain and profound anemia. A 7 cm enhancing, lipid-poor adrenal mass with calcifications that extended posterior to the vena cava was identified on imaging. Patient underwent right adrenalectomy with retroperitoneal lymph node dissection. Pathology demonstrated angiosarcoma of the adrenal gland. Consolidative treatment included adjuvant radiation and chemotherapy. Patient remains disease free 1.5 years following treatment. Prior reported literature on the diagnosis and management of adrenal angiosarcoma is reviewed.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction


Angiosarcoma is a malignant endothelial-derived, soft-tissue sarcoma that arises from vascular or lymphatic origin.<sup>1</sup> The majority is thought to arise spontaneously; however, malignant transformation from within benign vascular lesions has also been reported.<sup>1</sup> The most common locations for this tumor are the head and neck (27.0%), breast (19.7%), and the extremities (15.3%).<sup>1</sup>

Angiosarcoma of the adrenal gland is an exceedingly rare malignancy with less than 50 reported cases in the literature to date.<sup>2–27</sup> The patient described herein presented initially with abdominal pain and severe anemia, was found to have a right adrenal mass upon a diagnostic workup, and then received multimodal therapy in the form of surgery, adjuvant chemotherapy and radiation.

## Case presentation

A 59-year-old female with no history of prior malignant disease presented with a chief complaint of abdominal pain and anemia.


Her hemoglobin (HGB) 10 weeks prior to presentation was 7.8 gm/dL, and leukocytosis with a white blood cell count of  $\sim 18$  k/mm<sup>3</sup>



**Figure 1.** Enhancing, lipid-poor mass with coarse areas of calcification measuring 7.0 × 6.7 cm in the right adrenal gland. Note relationship of the mass to the vena cava.

\* Corresponding author. Fax: +1 888 751 6615.

E-mail address: [alexander.kutikov@fccc.edu](mailto:alexander.kutikov@fccc.edu) (A. Kutikov).



**Figure 2.** (A) Angiosarcoma of the adrenal gland. High power magnification demonstrates a sheet of large pleomorphic, round epithelioid cells with eosinophilic cytoplasm and large hyperchromatic nuclei with prominent nucleoli. Scattered intracytoplasmic lumina contain erythrocytes. Numerous mitotic figures are present (H & E, 40× magnification). (B) CD31 immunohistochemical stain. Diffuse intense membranous and cytoplasmic tumor cell staining with this vascular marker is apparent. (40× magnification).

was also present. An abdominal ultrasound revealed a 7 cm (cm) retroperitoneal mass arising from the right adrenal gland. To better characterize the mass, a computerized tomography (CT) scan was performed, which demonstrated a 7 cm enhancing, lipid-poor adrenal mass with coarse areas of calcification concerning for a primary adrenal cortical carcinoma (Fig. 1). Serum aldosterone, renin, testosterone, DHEA-S, estradiol, androstenedione, 17-OH progesterone, metanephrine, and normetanephrine levels were within normal limits, and serum cortisol was slightly elevated at 7.6 µg/dL upon low dose dexamethasone suppression testing. The patient underwent an open right adrenalectomy with lymph node dissection. Pathology demonstrated an 11.5 × 8.5 × 6.0 cm adrenal mass with invasion into the peri-adrenal fat. The tumor was within <0.01 cm margin of resection where it was densely adherent to the vena cava. Paracaval lymph nodes were negative for malignancy. Immunohistochemistry was positive for CD31, chromogranin, vimentin, and epithelial membrane antigen, and negative for calretinin, synaptophysin, TTF-1, GATA3, A103, AE1/AE3, CD5, and inhibin (Fig. 2). As such, these findings were consistent with angiosarcoma.

After a multidisciplinary discussion, patient underwent 5 of 6 planned cycles of adjuvant chemotherapy with doxorubicin, ifosfamide, and mesna (AIM) followed by external beam radiotherapy (XRT) in 25 fractions for a total of 50 Gray (Gy). Her anemia improved upon resection of the mass, with a HGB increase to 11.5 gm/dL 7 weeks after surgery and resolution of leukocytosis (WBC count of 6.5 k/mm<sup>3</sup> on postoperative day 20). Morning cortisol on postoperative day 4 was normal at 14.6 µg/dL. At 18 months following completion of treatment, the patient remains without radiographic evidence of disease.

## Discussion

We report adrenalectomy followed by adjuvant XRT and chemotherapy for localized adrenal angiosarcoma. At eighteen months, the patient remains disease-free.

Angiosarcoma of the adrenal gland was first described in 1988 by Kareti and colleagues.<sup>2</sup> Since then, an additional 40 cases have been reported in the literature with a median age at presentation of 60 years (range 34–85 years).<sup>3–27</sup> Patients present in a variety of ways including flank or abdominal pain, palpable mass, fatigue, weight loss, fever, or clinically asymptomatic with a mass detected on incidental imaging. Table 1 summarizes the age, sex, type of presentation, treatment, and follow-up for previously reported cases.<sup>2–27</sup> Patients who present with systemic symptoms tend to have worse outcomes than those with localized symptomatology.

Given the rarity of the disease, there is no consensus guideline on the best treatment algorithm for adrenal angiosarcomas. Adrenalectomy serves both diagnostic and therapeutic purposes. For patients with localized disease, the median survival is 7 months,<sup>1</sup> while one study reported a 2-year overall survival of 21%.<sup>28</sup> Adjuvant radiation therapy has been previously reported and was elected for our patient due to proximity of the surgical margin at the posterior vena cava. Previously reported multimodal treatment approaches have included doxorubicin-based chemotherapeutic regimens and adjuvant radiation therapy (XRT).<sup>28</sup> These data suggested risk reduction of systemic progression and improved 2-year survival in patients with visceral angiosarcoma who undergo multimodal treatment consisting of extirpation, adjuvant chemotherapy, and in some cases radiation.<sup>28,29</sup> Furthermore, a meta-analysis assessing the use of adjuvant doxorubicin-based chemotherapy for localized resectable soft-tissue sarcoma demonstrated a decreased risk of local, distant, and overall recurrence, and improved survival.<sup>30</sup> For our patient, restaging scans after completion of systemic therapy demonstrated no signs of local recurrence or distant metastatic disease. Adjuvant XRT has been shown to potentially improve disease-free survival in patients with angiosarcoma.<sup>31</sup> Thus, given the focally close margin at the posterior vena cava and no evidence of progression, 50 Gy of radiation was delivered to the operative bed using a modulated technique respecting normal tissue thresholds of the adjacent bowel and kidneys. Treatment was tolerated without event.

**Table 1**  
Reported cases of angiosarcoma of the adrenal gland.

Author	Age	Sex	Presentation	Treatment	Follow-up
Present case	59	F	Abdominal pain, anemia	Adrenalectomy + AIM + XRT	NER after 18 months
Ref. 2	54	M	Left upper quadrant pain	Adrenalectomy	Recurrence 7 months after surgery → en bloc resection of lateral gastric wall, tail of pancreas, left kidney, spleen. NER after 1 year NER after 6 months
Ref. 3	54	M	Fatigue, malaise, headache, cough, fever	Adrenalectomy	
Ref. 4	Unknown	Unknown	Unknown	Surgery	Presented with metastatic disease; died from disease 20 months after surgery
Ref. 5	59	M	Right upper quadrant pain, vomiting, fatigue	Adrenalectomy + nephrectomy	Died 8 days postoperatively due to bleeding
Ref. 6	41	M	Abdominal pain, loss of appetite, palpable abdominal mass	Adrenalectomy + XRT	Unknown
Ref. 7	45	F	Right-sided abdominal pain, cushingoid appearance	Adrenalectomy	NER after 11 years
	54 <sup>a</sup>	M	Left upper quadrant pain	Adrenalectomy	Recurrence 6 months after surgery → en bloc resection of lateral gastric wall, tail of pancreas, left kidney, spleen. Adjuvant therapy with cytoxan, doxorubicin, methotrexate, vincristine. DNER after 4 years
	56	F	Left-sided flank and shoulder pain	Adrenalectomy + doxorubicin	NER after 6 years
	56	M	Fever and weakness	Adrenalectomy	Died with pulmonary metastasis
	60	F	Right upper quadrant pain	Adrenalectomy	NER after 13 years
	60	F	Left flank pain	Adrenalectomy	Died from postoperative complications
	64	M	40–60 pound weight loss, hepatosplenomegaly	Adrenalectomy + splenectomy	Died with pulmonary metastasis 1 year later
	82	M	Incidental finding	Adrenalectomy	Died with pulmonary metastasis 2 years later
	85	M	Incidental finding at autopsy		
Ref. 8	Unknown	Unknown	Unknown	Unknown	Unknown
Ref. 9	63	M	Incidental finding	Adrenalectomy	Unknown
Ref. 10	67	M	Left flank pain	Adrenalectomy	Died 2 months postoperatively from metastatic disease
	60	F	Right upper quadrant pain	Adrenalectomy	Unknown
Ref. 11	34	M	Cushingoid appearance, recent hypertension and diabetes mellitus	Adrenalectomy	NER after 2 years
Ref. 12	70	M	Right upper quadrant pain	Adrenalectomy	Died with intestinal infarction and acute renal failure after 3 weeks
Ref. 13	70	F	Right-sided flank pain	Adrenalectomy	NER after 18 months
Ref. 14	50	M	Left-sided flank pain	Adrenalectomy	NER after 12 years
Ref. 15	49	F	Blunt abdominal trauma	Adrenalectomy	NER after 1 year
Ref. 16	61	M	Fatigue, weight loss, night sweats, elevated inflammatory markers	Adrenalectomy	Unknown
Ref. 17	35	M	Hypokalemic hypertension	Adrenalectomy + doxorubicin/ifosfamide	NER after 2 years
Ref. 18	69	F	Bilateral lower extremity swelling	Adrenalectomy and nephrectomy	Unknown
Ref. 19	55	M	Left-sided back pain	Adrenalectomy	Unknown
Ref. 20	62	F	Low back pain, anemia	Adrenalectomy	Unknown
Ref. 21	42	M	Right flank pain, lower back pain, weight loss	Adrenalectomy + XRT	Recurrence 9 months after completion adjuvant XRT → hepatic segmentectomy, excision of chest and right upper quadrant metastasis. 3 months later, 2.1 cm mid abdominal wall mass → refused further treatment
Ref. 22	64	F	Abdominal pain, weight loss	Adrenalectomy	Died with pulmonary and hepatic metastasis 2 years later
Ref. 23	68	M	Left thoracic pain	Ablation + anthracyclines + XRT	Died with vertebral and costal metastasis after 4 months
Ref. 24	63	M	Abdominal pain	Adrenalectomy	Unknown
Ref. 25	60	F	Incidental finding	Adrenalectomy	NER after 9 months
Ref. 26	67	M	Right upper quadrant pain	Neoadjuvant sunitinib + adrenalectomy + nephrectomy	Two 4 week courses neoadjuvant sunitinib followed by adrenalectomy. Nephrectomy performed for renal cell carcinoma. NER after 1 year
Ref. 27	53	F	Unknown	Unknown	Unknown
	74	M	Unknown	Unknown	Unknown
	75	F	Unknown	Unknown	Unknown
	75	F	Unknown	Unknown	Unknown

DNER: died, no evidence of recurrence; F: female; M: male; NER: no evidence of recurrence.

<sup>a</sup> Previously discussed in Kareti et al.<sup>2</sup>

## Conclusion

In summary, we present a case of a 59-year-old female who was found to have a large localized angiosarcoma of the adrenal gland upon presentation with abdominal pain, severe anemia, and leukocytosis. She underwent adrenalectomy with adjuvant chemotherapy and XRT. Selecting ideal treatment regimens in these rare patients remains challenging, and pooled reports such as this one seek to help guide appropriate clinical decision-making in patients with this rare malignancy.

## Conflicts of interest

None.

## References

- Young RJ, Brown NJ, Reed MW, et al. Angiosarcoma. *Lancet Oncol*. 2010;11(10):983–991.
- Kareti LR, Kattlein S, Siew S, et al. Angiosarcoma of the adrenal gland. *Arch Pathol Lab Med*. 1988;112(11):1163–1165.
- Bosco PJ, Silverman ML, Zinman LM. Primary angiosarcoma of adrenal gland presenting as paraneoplastic syndrome: case report. *J Urol*. 1991;146(4):1101–1103.
- Caplan RH, Kiskan WA, Huiras CM. Incidentally discovered adrenal masses. *Minn Med*. 1991;74(8):23–26.
- Livaditou A, Alexiou G, Floros D, et al. Epithelioid angiosarcoma of the adrenal gland associated with chronic arsenical intoxication? *Pathol Res Pract*. 1991;187(2–3):284–289.
- Ben-Izhak O, Auslander L, Rabinson S, et al. Epithelioid angiosarcoma of the adrenal gland with cytokeratin expression. Report of a case with accompanying mesenteric fibromatosis. *Cancer*. 1992;69(7):1808–1812.
- Wenig BM, Abbondanzo SL, Heffess CS. Epithelioid angiosarcoma of the adrenal glands. A clinicopathologic study of nine cases with a discussion of the implications of finding “epithelial-specific” markers. *Am J Surg Pathol*. 1994;18(1):62–73.
- Naka N, Ohsawa M, Tomita Y, et al. Angiosarcoma in Japan. A review of 99 cases. *Cancer*. 1995;75(4):989–996.
- Croitoru AG, Klausner AP, McWilliams G, et al. Primary epithelioid angiosarcoma of the adrenal gland. *Ann Diagn Pathol*. 2001;5(5):300–303.
- Ferrozzi F, Tognini G, Bova D, et al. Hemangiosarcoma of the adrenal glands: CT findings in two cases. *Abdom Imaging*. 2001;26(3):336–339.
- Invitti C, Pecori Giraldi F, Cavagnini F, et al. Unusual association of adrenal angiosarcoma and Cushing’s disease. *Horm Res*. 2001;56(3–4):124–129.
- Kruger S, Kujath P, Johannisson R, et al. Primary epithelioid angiosarcoma of the adrenal gland case report and review of the literature. *Tumori*. 2001;87(4):262–265.
- Pasqual E, Bertolissi F, Grimaldi F, et al. Adrenal angiosarcoma: report of a case. *Surg Today*. 2002;32(6):563–565.
- Al-Meshan MK, Katchy KC. An unusual angiosarcoma. A case report. *Med Princ Pract*. 2004;13(5):295–297.
- Gambino G, Mannone T, Rizzo A, et al. Adrenal epithelioid angiosarcoma: a case report. *Chir Ital*. 2008;60(3):463–467.
- Derlin T, Clauditz TS, Habermann CR. Adrenal epithelioid angiosarcoma metastatic to the epicardium: diagnosis by 18F-FDG PET/CT. *Clin Nucl Med*. 2012;37(9):914–915.
- Lepoutre-Lussey C, Rousseau A, Al Ghuzlan A A, et al. Primary adrenal angiosarcoma and functioning adrenocortical adenoma: an exceptional combined tumor. *Eur J Endocrinol*. 2012;166(1):131–135.
- Schreiner AM, Hoda RS. Primary adrenal epithelioid angiosarcoma showing rhabdoid morphology on air-dried smears. *Diagn Cytopathol*. 2012;40(suppl 2):E162–E164.
- Kedzierski L, Hawrot-Kawecka A, Holecki M, et al. Angiosarcoma of the adrenal gland. *Pol Arch Med Wewn*. 2013;123(9):502–503.
- Sebastiano C, Zhao X, Deng FM, et al. Cystic lesions of the adrenal gland: our experience over the last 20 years. *Hum Pathol*. 2013;44(9):1797–1803.
- Sung JY, Ahn S, Kim SJ, et al. Angiosarcoma arising within a long-standing cystic lesion of the adrenal gland: a case report. *J Clin Oncol*. 2013;31(9):e132–e136.
- Babinska A, Peksa R, Swiatkowska-Stodulska R, et al. The collection of five interesting cases of adrenal tumors from one medical center. *World J Surg Oncol*. 2014;12:377.
- Criscuolo M, Valerio J, Gianicolo ME, et al. A vinyl chloride-exposed worker with an adrenal gland angiosarcoma: a case report. *Ind Health*. 2014;52(1):66–70.
- Hayashi T, Gucer H, Mete O. A mimic of sarcomatoid adrenal cortical carcinoma: epithelioid angiosarcoma occurring in adrenal cortical adenoma. *Endocr Pathol*. 2014;25(4):404–409.
- Hendry S, Forrest C. Epithelioid angiosarcoma arising in an adrenal cortical adenoma: a case report and review of the literature. *Int J Surg Pathol*. 2014;22(8):744–748.
- Gusenbauer K, Ruzhynsky V, Kak I, et al. Angiosarcoma of the adrenal gland with concurrent contralateral advanced renal cell carcinoma: a diagnostic and management dilemma. *Can Urol Assoc*. 2015;9(5–6):E302–E305.
- Cornejo KM, Hutchinson L, Cyr MS, et al. MYC Analysis by fluorescent *in situ* hybridization and immunohistochemistry in primary adrenal angiosarcoma (PAA): a series of four cases. *Endocr Pathol*. 2015;26(4):334–341.
- Naka N, Ohsawa M, Tomita Y, et al. Prognostic factors in angiosarcoma: a multivariate analysis of 55 cases. *J Surg Oncol*. 1996;61(3):170–176.
- Movva S, von Mehren M, Ross EA, et al. Patterns of chemotherapy administration in high-risk soft tissue sarcoma and impact on overall survival. *J Natl Compr Cancer Netw*. 2015;13(11):1366–1374.
- Pervaiz N, Colterjohn N, Farrokhlyar F, et al. A systematic meta-analysis of randomized controlled trials of adjuvant chemotherapy for localized resectable soft-tissue sarcoma. *Cancer*. 2008;113(3):573–581.
- Mark RJ, Poen JC, Tran LM, et al. Angiosarcoma. A report of 67 patients and a review of the literature. *Cancer*. 1996;77(11):2400–2406.