



# The antiviral treatment should be initiated before the development of cirrhosis

Qiao Tang, Peng Hu

Department of Infectious Diseases, Institute for Viral Hepatitis, The Key Laboratory of Molecular Biology for Infectious Diseases, Chinese Ministry of Education, The Second Affiliated Hospital of Chongqing Medical University, Chongqing, China

*Correspondence to:* Peng Hu, PhD. Department of Infectious Diseases, Institute for Viral Hepatitis, The Key Laboratory of Molecular Biology for Infectious Diseases, Chinese Ministry of Education, The Second Affiliated Hospital of Chongqing Medical University, 74# Linjiang Road, Yuzhong District, Chongqing 400010, China. Email: hp\_cq@163.com.

*Comment on:* Huang DQ, Tamaki N, Lee HW, *et al.* Outcome of untreated low-level viremia versus antiviral therapy-induced or spontaneous undetectable HBV-DNA in compensated cirrhosis. *Hepatology* 2023;77:1746-56.

**Keywords:** Low-level viraemia; maintained virological response (MVR); untreated; hepatocellular carcinoma (HCC); decompensated cirrhosis

Submitted Nov 05, 2023. Accepted for publication Dec 06, 2023. Published online Jan 12, 2024.

doi: 10.21037/hbsn-23-580

**View this article at:** <https://dx.doi.org/10.21037/hbsn-23-580>

Hepatitis B virus (HBV) infection increases the risk of liver fibrosis, cirrhosis and hepatocellular carcinoma (HCC) (1). International guidelines recommend antiviral treatment for eligible patients to prevent the development of HCC and the progression of liver disease. However, there is discrepancy in these guidelines regarding treatment recommendations for patients with compensated cirrhosis and low-level viremia (LLV), defined as serum HBV-DNA level of 20–2,000 IU/mL (2–4). And data on the natural history of untreated compensated cirrhotic patients with LLV is limited.

Recently, Huang *et al.* performed a multi-ethnic, multi-center and retrospective study, enrolling 2,316 patients, explored the nature history of untreated compensated cirrhotic patients with LLV compared to spontaneous-maintained virological response (MVR) and antiviral therapy (AVT)-MVR (5). The annual HCC incidence was 2.7/100 person-years (PYs), 2.6/100 PYs, and 3.3/100 PYs for LLV (n=742), Spontaneous-MVR (n=333), and AVT-MVR (n=1,241) groups, respectively. No significant difference was found in the three groups, which indicated that LLV in HBV-infected compensated cirrhosis did not increase the risk of disease progression compared to spontaneous—MVR and AVT-MVR (5). Current guidelines by the American Association for the Study of the Liver Diseases (AASLD) and the European Association for the

Study of the Liver (EASL) recommend AVT for cirrhotic patients with LLV, but the Asia-Pacific Association for the Study of the Liver (APASL) guidelines recommend AVT when HBV-DNA level is >2,000 IU/mL. Present study indicated a similar risk of HCC in cirrhotic patients with LLV compared to patients with spontaneous-MVR and AVT-MVR, which is different from our previous knowledge that LLV increases the risk of developing HCC (6). On the one hand, this may be due to the different study populations included in these studies. The LLV population in present study was the untreated cirrhotic population, whereas previous studies referred to patients with chronic hepatitis B (CHB) on treatment (including cirrhotic and non-cirrhotic patients). On the other hand, previous studies have shown no significant difference in the risk of HCC between non-cirrhotic patients with LLV and MVR, but higher risk of HCC in cirrhotic patients with LLV compared to cirrhotic patients with MVR (7). Moreover, REVEAL study indicated low risk of HCC in CHB patients with viral load between 300 and 9,999 copies/mL (including a small number of cirrhotic patients), and the annual HCC incidence was 0.11/100 PYs, which is lower than the HCC incidence in present study (1). The different prognosis of cirrhotic patients with LLV may suggest the potential virologic and immunologic differences, which need to be further explored.

First, LLV mentioned in present study occurred in CHB patients without treatment is not the same as the identified LLV definition [the definition of LLV in CHB was borrowed from the concept of LLV in human immunodeficiency virus (HIV)], which occurred in CHB patients received AVT more than 48 weeks, and might be associated with stability of covalently closed circular DNA (cccDNA) pool, drug resistance, proliferation of hepatocytes, host's immune function and genetic background (2,8). And LLV occurred after AVT is generally considered to increase the risk of cirrhosis and HCC. However, the LLV occurred in untreated patients suggests effective suppression of viral replication achieved by host's immune system and low risk of disease progression. Therefore, the untreated LLV is a new definition and we need to distinguish between untreated LLV and LLV occurred after AVT. The potential virologic and immune differences between untreated LLV and LLV occurred after AVT needed to be further explored.

Second, in present study, more than 40% of the LLV patients in present study received AVT and were terminated from follow-up, but the specific reasons for AVT were not mentioned. According to antiviral indications and clinical practice, patients receiving AVT are generally at high risk of disease progression. The authors artificially screened out the population at low risk of disease progression for follow-up, which may result in low risk of HCC in cirrhotic patients with LLV. On the other hand, HBV integration in these patients was ignored by authors, which occurred in all the phases of chronic infection and closely related to the occurrence of HCC (9). And viral integration has occurred in these patients prior to the occurrence of cirrhosis and may have partially initiated the oncogene pathway. Moreover, AVT did not directly target the integrated HBV DNA, so it did not reduce the pre-existing HBV integration before AVT (10) Therefore, the risk of HCC maybe comparable in three cirrhotic populations. It is worth noting that since AVT did not decrease the risk of HCC after patients developing cirrhosis, AVT should be initiated as early as possible (before the development of cirrhosis), not after the development of cirrhosis.

Third, the methods evaluating the risk of HCC may not be appropriate in present study, although multivariable Cox-regression analyses, as well as propensity score matching (PSM), stratified and sensitivity analyses were used to demonstrate the similar risk of HCC in patients with LLV, spontaneous MVR and AVT MVR (5). However, the occurrence of HCC is associated with many variables, the authors should use the risk model for screening for

populations at high risk of developing HCC and then further compared the risk of developing HCC among the three populations. Moreover, AVT is an important disease modifier, the HCC risk models are different in patients without treatment and received AVT. The risk of occurrence of HCC should be evaluated by HCC risk models CU-HCC (The Chinese University HCC score) and GAG-HCC (The Guide with Age, Gender, HBV DNA, Core promoter mutations and Cirrhosis), which are applicable to the patients without treatment, and PAGE-B (platelet age gender-B), which is applicable to the patients received treatment (11-13).

In summary, the current international guidelines are controversial about whether to initiate AVT in cirrhotic patients with LLV (2-4), but the Chinese guideline is more positive about initiating AVT in these patients (14). Because studies have shown that there are racial differences in the risk of HCC, and that Asian populations are at a higher risk of HCC and have a poorer prognosis than Western Europe and the United States populations (15). Therefore, more robust evidence is still needed to explore the prognosis of untreated cirrhotic patients with LLV. Based on the findings in this study that AVT does not decrease the risk of HCC in cirrhotic patients, and the viral integration occurred before the development of cirrhosis, the AVT should be initiated as early as possible (before the development of cirrhosis). Moreover, what we commonly think of as LLV (occurred after treatment) is different from the LLV that occurred during the course of natural history and the potential virologic and immunologic differences between them need to be further explored to better explain the different prognosis in cirrhotic patients.

## Acknowledgments

*Funding:* This work was supported in part by grants from the Chongqing Talents Project (No. cstc2021ycjhb-gzxm0150); the First batch of key Disciplines on Public Health in Chongqing, Health Commission of Chongqing, China; the Remarkable Innovation-Clinical Research Project, The Second Affiliated Hospital of Chongqing Medical University; Scientific and Technological Research Program of Chongqing Municipal Education Commission, The Second Affiliated Hospital of Chongqing Medical University (No. KJZD-K202300404).

## Footnote

*Provenance and Peer Review:* This article was commissioned

by the editorial office, *Hepatobiliary Surgery and Nutrition*. The article did not undergo external peer review.

**Conflicts of Interest:** Both authors have completed the ICMJE uniform disclosure form (available at <https://hbsn.amegroups.com/article/view/10.21037/hbsn-23-580/coif>). The authors report that this work was supported in part by grants from the Chongqing Talents Project (No. cstc2021ycjh-bgzxm0150); the First Batch of Key Disciplines on Public Health in Chongqing, Health Commission of Chongqing, China; the Remarkable Innovation–Clinical Research Project, The Second Affiliated Hospital of Chongqing Medical University and Scientific; Technological Research Program of Chongqing Municipal Education Commission, The Second Affiliated Hospital of Chongqing Medical University (No. KJZD-K202300404). The authors have no other conflicts of interest to declare.

**Ethical Statement:** The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

**Open Access Statement:** This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

- Chen CJ, Yang HI, Su J, et al. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006;295:65-73.
- Terrault NA, Lok ASF, McMahon BJ, et al. Update on prevention, diagnosis, and treatment of chronic hepatitis B: AASLD 2018 hepatitis B guidance. *Hepatology* 2018;67:1560-99.
- European Association for the Study of the Liver. Electronic address: easloffice@easloffice; . EASL 2017 Clinical Practice Guidelines on the management of hepatitis B virus infection. *J Hepatol* 2017;67:370-98.
- Sarin SK, Kumar M, Lau GK, et al. Asian-Pacific clinical practice guidelines on the management of hepatitis B: a 2015 update. *Hepato Int* 2016;10:1-98.
- Huang DQ, Tamaki N, Lee HW, et al. Outcome of untreated low-level viremia versus antiviral therapy-induced or spontaneous undetectable HBV-DNA in compensated cirrhosis. *Hepatology* 2023;77:1746-56.
- Kim JH, Sinn DH, Kang W, et al. Low-level viremia and the increased risk of hepatocellular carcinoma in patients receiving entecavir treatment. *Hepatology* 2017;66:335-43.
- Zhang Q, Peng H, Liu X, et al. Chronic Hepatitis B Infection with Low Level Viremia Correlates with the Progression of the Liver Disease. *J Clin Transl Hepatol* 2021;9:850-9.
- Zhang Q, Cai DC, Hu P, et al. Low-level viremia in nucleoside analog-treated chronic hepatitis B patients. *Chin Med J (Engl)* 2021;134:2810-7.
- Pollicino T, Caminiti G. HBV-Integration Studies in the Clinic: Role in the Natural History of Infection. *Viruses* 2021;13:368.
- Chow N, Wong D, Lai CL, et al. Effect of Antiviral Treatment on Hepatitis B Virus Integration and Hepatocyte Clonal Expansion. *Clin Infect Dis* 2023;76:e801-9.
- Wong VW, Chan SL, Mo F, et al. Clinical scoring system to predict hepatocellular carcinoma in chronic hepatitis B carriers. *J Clin Oncol* 2010;28:1660-5.
- Yuen MF, Tanaka Y, Fong DY, et al. Independent risk factors and predictive score for the development of hepatocellular carcinoma in chronic hepatitis B. *J Hepatol* 2009;50:80-8.
- Papatheodoridis G, Dalekos G, Sypsa V, et al. PAGE-B predicts the risk of developing hepatocellular carcinoma in Caucasians with chronic hepatitis B on 5-year antiviral therapy. *J Hepatol* 2016;64:800-6.
- You H, Wang F, Li T, et al. Guidelines for the Prevention and Treatment of Chronic Hepatitis B (version 2022). *J Clin Transl Hepatol* 2023;11:1425-42.
- Fattovich G, Bortolotti F, Donato F. Natural history of chronic hepatitis B: special emphasis on disease progression and prognostic factors. *J Hepatol* 2008;48:335-52.

**Cite this article as:** Tang Q, Hu P. The antiviral treatment should be initiated before the development of cirrhosis. *HepatoBiliary Surg Nutr* 2024;13(1):136-138. doi: 10.21037/hbsn-23-580