



Article

Antioxidant Effect of *Chrysanthemum morifolium* (Chuju) Extract on H₂O₂-Treated L-O₂ Cells as Revealed by LC/MS-Based Metabolic Profiling

Ge Zhan ¹, Men Long ¹, Kai Shan ², Chong Xie ² and Runqiang Yang ^{2,*}

¹ School of Biological Science and Food Engineering, Chuzhou University, Chuzhou 239000, China; zhangezczxy@163.com (G.Z.); czxy_lm@163.com (M.L.)

² Whole Grain Food Engineering Research Center, College of Food Science and Technology, Nanjing Agricultural University, Nanjing 210095, China; shankai@njau.edu.cn (K.S.); xiechong@njau.edu.cn (C.X.)

* Correspondence: yangrq@njau.edu.cn; Tel./Fax: +86-025-84396293

Abstract: Chrysanthemum has a long history of being used to attenuate various oxidative stress-related discomforts and diseases; however, its mechanisms remain unclear. In this study, the antioxidant effect of chrysanthemum aqueous extract was investigated, and the potential mechanisms were explored via a metabolomics study. Chrysanthemum extract could significantly inhibit hydrogen peroxide (H₂O₂)-mediated cell death in L-O₂ hepatocytes. Propidium iodide staining and annexin V-PI dual staining revealed that the antioxidant effect of chrysanthemum extract was related to the relief of cell cycle arrest and inhibition of non-apoptotic cell damage. The activities of antioxidant enzymes including superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GSH-Px) were also upregulated by chrysanthemum extract. Through metabolomics studies, it was found that chrysanthemum extract mainly targeted the arginine synthesis pathway and purine metabolism pathway, in which antioxidation-related endogenous substrates including L-arginosuccinate, citrulline and inositol monophosphate were significantly upregulated by chrysanthemum extract. These results indicated that chrysanthemum extract can antagonize oxidative stress through multiple pathways and have potential therapeutic applications.

Keywords: chrysanthemum; antioxidant; L-O₂ cells; metabolites



Citation: Zhan, G.; Long, M.; Shan, K.; Xie, C.; Yang, R. Antioxidant Effect of *Chrysanthemum morifolium* (Chuju) Extract on H₂O₂-Treated L-O₂ Cells as Revealed by LC/MS-Based Metabolic Profiling. *Antioxidants* **2022**, *11*, 1068. <https://doi.org/10.3390/antiox11061068>

Academic Editor: David Arráez-Román

Received: 19 April 2022

Accepted: 23 May 2022

Published: 27 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Natural products have a long history of use as drugs, drug precursors, and/or complementary health adjuvants due to their abundant health benefits. Among them, plants are the most important source of natural products due to their wide availability. *Chrysanthemum morifolium* (Chuju) is one of the four top cut-flowers in the world. In addition to being an ornamental plant, the flowers of chrysanthemum also show potential medicinal effects [1]. For a long time, chrysanthemum and its extracts have been used to relieve multiple discomfort symptoms in Asia, such as improving liver function and alleviating red and itchy eyes [2]. Recent studies indicated that *C. morifolium* ethanol extract attenuated lipopolysaccharide-induced acute lung injury in mice by increasing the activity of total-antioxidant capacity and decreasing the contents of malondialdehyde [3]. In addition, *Chrysanthemi* was also found to alleviate acetaminophen-induced rat liver injury via inhibiting oxidative stress and apoptosis [4]. Although numerous studies have demonstrated the antioxidant effects of chrysanthemum in vitro and in vivo and some potential antioxidant substances including caffeic acid, apigenin, luteolin as well as glucosides have been identified, the molecular mechanism is still poorly studied [5].

Isolation and identification of the bioactive molecules in a mixture is a classical strategy for studying natural products. A lot of natural compounds including alkaloids, phenolics,

and terpenes in chrysanthemum have been identified as antioxidants [6]. For them, the predominant effects may involve the activation of Nrf2-related antioxidant pathways and scavenging of free radicals [7–9]. However, researchers found that the biological activities of natural extracts are much higher than those of single bioactive molecules due to the interaction between molecules at pharmacodynamic and pharmacokinetic levels, suggesting that the synergistic effect between multiple compounds cannot be overlooked [10]. Examples exist in which *Hypericum perforatum* (*H. perforatum*) extract-derived procyanidin B2 and hyperocidin can significantly improve the bioavailability and benefits of hypericin from *H. perforatum* [11]. This indicates the significance of systematic studies on the investigation of antioxidant mechanisms of natural products.

The redox system in humans is a homeostatic network coupling with material and energy metabolism. Multiple metabolic participants such as respiratory chain, NADPH oxidase and lipoxygenase generate oxides and peroxides, while GSH and NADPH play a central role in the body's antioxidant response as substrates of multiple enzymes [12]. This suggests that changes in overall metabolic pattern can be meaningful to identify the redox state during cellular oxidative stress. It has been found that oxidative stress may contribute to the reduced influx of L-arginine in platelets, while L-arginine could lower lipid peroxidation and maintain mitochondrial functions and neuronal survival [13,14]. Moreover, antioxidants contribute to decreased levels of creatine kinase activity in serum; creatine and its metabolite creatine phosphate can also act as antioxidants and prevent behavioral and oxidative stress alterations. Their metabolism is disrupted by redox disorders [15,16]. It follows that the profile of metabolites can accurately reflect the body's redox levels and help reveal the involved molecular mechanisms. Simultaneously, metabolomics can provide a good scenario for the study of antioxidant mechanisms by tracking the metabolic flux of antioxidants, oxides and endogenous metabolites.

In this study, we found that chrysanthemum extract can significantly inhibit oxidative stress in L-O2 hepatocytes in a dose-dependent manner. Through metabolomic analysis, we further found that chrysanthemum extract exerted antioxidant effects by regulating the activity of antioxidant enzyme and targeting the arginine synthesis pathway and purine metabolism pathway. This study provides a new perspective for understanding the regulation of redox by plant extracts at the metabolic level, which may help to better clarify the medicinal mechanism of chrysanthemum.

2. Materials and Methods

2.1. Materials

The chrysanthemum (chuju) extract was obtained by aqueous extraction. In brief, after drying, chrysanthemum was extracted at 80 °C for 5 min by adding deionized water at 1:20 (*w/w*). Extractions were performed three times in total. The raw extract was pooled and filtered through a 0.22 µm filter. The filtrate was then freeze-dried. L-O2 human hepatocyte was commercially acquired from Zhong Qiao Xin Zhou Biotechnology Co., Ltd. (Shanghai, China). Hydrogen peroxide (H₂O₂) was purchased from Hushi (Shanghai, China). Both the 3-(4,5)-dimethylthiazoliazol(-z-y1)-3,5-di-phenyltetrazolium bromide (MTT) cell proliferation and cytotoxicity assay kit (M1020) and 2',7'-dichlorofluorescein diacetate (4091-99-0) were purchased from Solarbio Science & Technology Co., Ltd. (Beijing, China). The malondialdehyde (MDA) assay kit (TBA method, A003-1-2), superoxide dismutase (SOD) assay kit (WST-1 method, A001-3-2), glutathione peroxidase (GSH-Px) assay kit (A005-1-2), catalase (CAT) assay kit (A007-1-1) and annexin V cell apoptosis assay kit (G003-1-2) were purchased from JianCheng Bioengineering Institute (Nanjing, China). The cell cycle assay kit (propidium iodide staining method) was purchased from Zhi En biology (Hefei, China). The mitochondrial membrane potential assay kit with JC-1 (C2006) was purchased from Beyotime (Shanghai, China).

2.2. Cell Culture

L-O2 human hepatocyte was maintained in RPMI-1640 supplemented with 10% fetal bovine serum, 100 U/mL penicillin and 100 µg/mL streptomycin. Cells were cultured at 37 °C in a humidified incubator with 5% CO₂. To induce H₂O₂-mediated oxidative stress, 2 × 10⁴/well cells were plated in a 96-well plate and cultured for 24 h. After removing the medium, H₂O₂ at final concentrations of 0, 1.0, 1.2, 1.4, 1.6, 1.8, 2.0, 3.0, 4.0, 5.0, 6.0 and 7.0 mM in medium was added into different wells. After that, the incubation continued for 4 h. To explore the effect of chrysanthemum (chuju) extract, 2 × 10⁴/well cells were plated in a 96-well plate and cultured for 24 h. After removing the medium, chrysanthemum extract at final concentrations of 0, 50, 100, 200, 400, 800, 1600 and 2400 µg/mL was added into different wells and vitamin C (50.0 µg/mL) was also added as a positive control. After 12 h of treatment, H₂O₂ was added at a final concentration of 7.0 mM and cells were cultured for another 4 h.

3. Test Method

3.1. Cell Viability Assay

For each well of 96 well plates, 25 µL of 5 mg/mL MTT solution was added. After 4 h of incubation, MTT solution was removed and 150 µL per well dimethyl sulfoxide was added to dissolve formazan. The absorbance values were measured using a microplate reader at a wavelength of 490 nm. Each group contained four replicates. Cell viability = (A_{intervention group} - A_{blank group})/(A_{control group} - A_{blank group}) × 100%.

3.2. Measurement of MDA Content and SOD, GSH Px, CAT Activities

The medium from cultured cells was removed and cells were digested using 0.25% trypsin without EDTA. Cell pellets were collected through centrifuge at 2000 × g for 5 min. After one wash using PBS, cells were lysed with lysis buffer (150 mM NaCl, 50 mM Tris HCl, 1 mM EDTA, 1% TritonX-100, pH 7.4). After a centrifuge at 12,000 × g for 20 min, supernatant was collected and protein concentration was determined using the BCA method. According to the instructions of the kits, the content of MDA and the activities of SOD, GSH Px and CAT were measured. The content of MDA is expressed in nmol/mg protein, and the activities of SOD, GSH Px and CAT are expressed in U/mg protein.

3.3. Cell Cycle Analysis

The single-cell suspension was washed using PBS, and cells were fixed by icy 70% ethanol for 2 h. After two washes, cells were incubated with 100 µL RNase (20 µg/mL) at 37 °C, then stained with 400 µL PI solution (10 µg/mL) at 4 °C for 30 min. Then, cells were analyzed through a BD Accuri™ C6 flow cytometer. The data were processed by Flowjo software.

3.4. Mitochondrial Membrane Potential Analysis

After digestion, cell pellets were collected through centrifuge at 2000 × g for 5 min. The 100 to 600 thousand cells were resuspended in 0.5 mL medium containing serum and stained according to the instruction. Then, the cells were analyzed through a BD Accuri™ C6 flow cytometer. The data were processed by Flowjo software.

3.5. Apoptosis Assay

The single-cell suspension was washed using PBS, and cells were resuspended in 0.5 mL binding buffer. Then, cells were stained with Annexin-V and PI for 15 min and analyzed through a BD Accuri™ C6 flow cytometer. The data were processed by Flowjo software.

3.6. Reactive Oxygen (ROS) Detection

Cells were stained with 15 µM 2',7'-dichlorofluorescein diacetate (DCF) for 30 min. After washing with PBS, cells were treated with H₂O₂ and chrysanthemum extract. Then, the cells were digested with trypsin. After centrifugation, cell pellets were resuspended

in 0.5 mL PBS and analyzed through a BD Accuri™ C6 flow cytometer. The data were processed by Flowjo software.

3.7. UHPLC-QE-MS Untargeted Metabolomics

Metabolites Extraction: 200 µL of water was added to the samples. After 30 s vortex, the samples were frozen and thawed with liquid nitrogen 3 times. The samples were sonicated for 10 min in an ice-water bath. Fifty microliters of homogenate was used to measure protein concentration. Then, 600 µL acetonitrile:methanol = 1:1 was added to the remaining part and transferred to a 2 mL EP tube. After 30 s vortex, the samples were sonicated for 10 min in an ice-water bath and were incubated at −40 °C for 1 h and centrifuged at 12,000 rpm for 15 min at 4 °C. The 700 µL of supernatant was transferred to an EP tube and dried in a vacuum concentrator. Then, acetonitrile:methanol:water = 2:2:1, with isotopically labelled internal standard mixture, was added in proportion. After 30 s vortex, the samples were sonicated for 10 min in an ice-water bath. Then, the samples were centrifuged at 12,000 rpm for 15 min at 4 °C. The resulting supernatant was transferred to a fresh glass vial for analysis. The quality control (QC) sample was prepared by mixing an equal aliquot of the supernatants from all of the samples.

LC-MS/MS Analysis: LC-MS/MS analyses were performed using a UHPLC system (Vanquish, Thermo Fisher Scientific, Waltham, MA, USA) with a UPLC BEH Amide column (2.1 mm × 100 mm, 1.7 µm) coupled to a Q Exactive HFX mass spectrometer (Orbitrap MS, Thermo). The mobile phase consisted of 25 mmol/L ammonium acetate and 25 ammonia hydroxide in water (pH = 9.75) (A) and acetonitrile (B). The autosampler temperature was 4 °C, and the injection volume was 3 µL. The QE HFX mass spectrometer was used for its ability to acquire MS/MS spectra on information-dependent acquisition (IDA) mode in the control of the acquisition software (Xcalibur, Thermo). In this mode, the acquisition software continuously evaluated the full scan MS spectrum. The ESI source conditions were set as follows: sheath gas flow rate as 30 Arb, Aux gas flow rate as 25 Arb, capillary temperature 350 °C, full MS resolution as 60,000, MS/MS resolution as 7500, collision energy as 10/30/60 in NCE mode, spray voltage as 3.6 kV (positive) or −3.2 kV (negative), respectively.

Data preprocessing and annotation: The raw data were converted to the mzXML format using ProteoWizard and processed with an in-house program, which was developed using R and based on XCMS, for peak detection, extraction, alignment, and integration. Then, an in-house MS2 database (BiotreeDB) was applied for metabolite annotation. The cutoff for annotation was set at 0.3.

3.8. Statistics

For all bar graphs, data are presented as mean ± SD. Two-tailed Student's *t*-test was used for the statistical comparison of two groups and one-way analysis of variance (ANOVA) was used for multiple comparisons. In cases where all groups shared an identical sample size, the Tukey test was adopted. *p* < 0.05 was considered significant. Statistical differences between the groups were marked with different letters.

4. Results and Analysis

4.1. The Effect of H₂O₂ and Chrysanthemum Extract on Cell Viability

Here, we employed a H₂O₂–induced cell death model to research the antioxidant effects of chrysanthemum extracts. Through the MTT assay, it was found that the cytotoxicity of H₂O₂ was significantly correlated with its concentration and 7.0 mM H₂O₂ can induce the strongest cell death (Figure 1A). Then, we tried to rescue the cell death with chrysanthemum extract and vitamin C (as a positive control). As expected, chrysanthemum extract attenuated H₂O₂-induced cell death, and its effects exhibited a concentration-dependent manner (Figure 1B). Notably, chrysanthemum extract of 800 µg/mL and above exhibited better protective abilities than vitamin C, implying the powerful antioxidant properties of chrysanthemum.




Figure 1. H₂O₂ and chrysanthemum extract on cell viability. **(A)** LO2 cells were treated using different concentrations of H₂O₂ for 4 h and the cell viability was determined by MTT assay. **(B)** LO2 cell was pretreated with different concentrations of chrysanthemum extract for 12 h and H₂O₂ for another 4 h. Cell viability was determined by MTT assay. Ext.: chrysanthemum extract, Vc: vitamin C. Three independent experiments were performed. The lower case letters indicate significant difference at $p < 0.05$ for different treatments.

4.2. Protective Effect of Chrysanthemum Extract against Oxidative Damage of Cells

4.2.1. Cell Cycle and Apoptosis

H₂O₂ induces damage of DNA and proteins, then initiates subsequent cellular responses. Through propidium iodide (PI) staining, we found that H₂O₂ significantly induced the cell cycle arrest as indicated by an increased cell number in G₀/G₁ phase and a decreased cell number in S phase as well as G₂/M phase (Figure 2A). In contrast, chrysanthemum extracts can effectively antagonize the effect of H₂O₂ (Figure 2A). On the other hand, H₂O₂ also caused severe non-apoptotic cell death, which was shown by a significant annexinV and PI double positive staining and a low annexin–V single positive staining (Figure 2B). Chrysanthemum extract can also inhibit such cell death in a concentration–dependent manner.




Figure 2. Chrysanthemum extract inhibited H₂O₂-induced cell cycle arrest and non-apoptotic cell death. L-O2 cell was pre-treated with different concentrations of chrysanthemum extract and treated with H₂O₂. (A) Cells were stained with PI and RNase, then analyzed with a flow cytometer through the BL-2 channel. (B) Cells were stained with PI and annexin-V-FITC, then analyzed with a flow cytometer through the BL-1 and BL-2 channels. Ext.: chrysanthemum extract. Blue represents 2N (G₀, G₁ phase), green represents S phase, and red represents 4N (G₂ phase).

4.2.2. MDA, ROS Content and Mitochondrial Membrane Potential

Since the cytotoxicity of H₂O₂ depends on oxidative stress induction, it was speculated that the protective effect of chrysanthemum extract may come from its antioxidant effect. TBARS showed a significant lipid peroxidation induced by H₂O₂, while chrysanthemum extract can effectively suppress it (Figure 3A). Consistent with the changes in lipid peroxidation levels, H₂O₂ can also induce the boost in intracellular ROS, which was shown by a high DCF peak (Figure 3B). However, chrysanthemum extract strongly inhibited the upregulation of intracellular ROS (Figure 3B). ROS can induce a decrease in mitochondrial membrane potential and JC-1 staining can characterize such change through the decrease in J-aggregates (BL-2) (Figure 3C). Chrysanthemum extract significantly rescued the mitochon-

drial membrane potential dysfunction, indicating the functional effect of chrysanthemum extract (Figure 3C).




Figure 3. MDA, ROS content and mitochondrial membrane potential. (A) TBARS was performed to evaluate the lipid peroxides in L-O2 cells treated with chrysanthemum extract and H₂O₂. (B) ROS content was measured by DCF staining followed by flow cytometry. A small peak to the right of the main peak indicates a cell population containing high levels of ROS. (C) Mitochondrial membrane potential was measured by JC-1 staining followed by flow cytometry. FITC (BL1) indicates monomeric JC1, while red fluorescence (BL2) indicates J-aggregates. Ext.: chrysanthemum extract. The lower case letters in figure (A) indicate significant difference at *p* < 0.05 for different treatments.

4.3. Antioxidant Enzyme Activities

In addition to targeting oxides and peroxides directly, modulation of antioxidant enzyme activities is also a potential mechanism of action of natural products. We found that H_2O_2 significantly disrupted the activities of three important antioxidant enzymes—SOD, CAT and GSH-Px—while chrysanthemum extract effectively restored them (Figure 4).




Figure 4. Antioxidant enzyme activities. L-O2 cells received different treatments and were lysed to evaluate the activities of SOD (A), CAT (B) and GSH-Px (C). Ext.: chrysanthemum extract. The lower case letters indicate significant difference at $p < 0.05$ for different treatments.

4.4. Untargeted Metabolomics Analysis

4.4.1. Principal Component Analysis

To explore the antioxidant mechanism of chrysanthemum extracts, we performed an untargeted metabolomics analysis. Whether in positive or negative ion mode, we can observe that H_2O_2 induced significant metabolic changes in L-O2 cells (Figure 5). Through a principal component analysis, we found that the antioxidant effect of chrysanthemum extract was accompanied by the change in cell metabolic profile (Figure 5A,B). Significantly, the metabolic profiles shaped by different concentrations of chrysanthemum extracts were similar, implying low side effects of chrysanthemum extract. Moreover, compared with the difference between the H_2O_2 group and the control group, chrysanthemum extract introduced more significant metabolic changes (Figure 5C-F).




Figure 5. H₂O₂ and chrysanthemum extract induced significant metabolic changes. L–O2 cell treated with H₂O₂ and chrysanthemum extract were lysed for a metabolomic analysis. According to the profile of metabolites, principal component analysis was performed (A,B). The volcano plot was generated according to the differences in metabolite abundance (C–F). POS: data achieved from positive ion mode of mass spectrum; NEG: data achieved from negative ion mode of mass spectrum. Con.: control; H₂O₂: 7 mM H₂O₂; 2400 + H₂O₂: 2400 μg/mL chrysanthemum extract and 7 mM H₂O₂.

4.4.2. Pathway Analysis

To evaluate the potential functions of these metabolites, we observed the quantity of metabolite that changed. The Venn plot showed that most of the metabolites influenced by H₂O₂ were also regulated by chrysanthemum extracts (Figure 6A). This suggested that the effects of chrysanthemum extract and H₂O₂ would share a similar profile of target. The main co-targets include the metabolites involved in metabolism of nucleotide bases and amino acid, especially the pathway of purine metabolism, arginine and proline metabolism (Figure 6B). Furthermore, the bar chart showed the co-targets of chrysanthemum extract with H₂O₂ (Figure 6C), implying the high specificity of chrysanthemum extract in antioxidation.




Figure 6. Metabolic pathway analysis. **(A)** Venn diagram showing differential metabolites of H₂O₂/Con. and chrysanthemum extract/H₂O₂. **(B)** Bubble plot showing the differential metabolic pathway enriched by metabolites. **(C)** Bar chart showing the differential metabolic pathway enriched by metabolites. POS: data achieved from positive ion mode of mass spectrum; NEG: data achieved from negative ion mode of mass spectrum. Con.: control; H₂O₂: 7 mM H₂O₂; 2400 + H₂O₂: 2400 µg/mL chrysanthemum extract and 7 mM H₂O₂. In figure (A), “a” represents the positive ion mode and “b” represents the negative ion mode; In figure (B), “a,b” represent the positive ion mode, and “c,d” represent the negative ion mode.

Further analyses found three types of metabolites: (1) chrysanthemum extract showed no effect on H₂O₂–treated cells including creatine and creatinine; (2) chrysanthemum extract enhanced the effect of H₂O₂ including L-Ornithine, phosphocreatine and D-ribose 5-phosphate; (3) chrysanthemum extract reversed the effect of H₂O₂, including L-citrulline, N-(L-arginino) succinate, N-acetyloronithine, N-acetyl-L-glutamate and inosine monophosphate (IMP) (Table 1).

Table 1. Metabolites regulated by H₂O₂ and chrysanthemum extract.

Metabolites	No.	RT (s)	Related Pathway	H ₂ O ₂ /Con.		2400 + H ₂ O ₂ /H ₂ O ₂	
				<i>p</i> Value ^a	FC ^b	<i>p</i> Value	FC
L-Ornithine	C00077	515.79	Arginine biosynthesis	4.59 × 10 ^{−5}	0.07	5.16 × 10 ^{−5}	0.38
L-Citrulline	C00327	402.96	Arginine biosynthesis	5.16 × 10 ^{−5}	0.38	2.11 × 10 ^{−2}	1.40
N-(L-Arginino)succinate	C03406	477.96	Arginine biosynthesis	2.22 × 10 ^{−3}	0.86	1.55 × 10 ^{−4}	1.69
N-Acetyloronithine	C00437	403.96	Arginine biosynthesis	5.71 × 10 ^{−4}	0.57	6.40 × 10 ^{−3}	1.40
N-Acetyl-L-glutamate	C00624	318.93	Arginine biosynthesis	3.07 × 10 ^{−4}	0.36	1.40 × 10 ^{−2}	1.72
Creatine	C00300	360.33	Arginine and proline metabolism	4.37 × 10 ^{−5}	1.28	1.40 × 10 ^{−2}	1.06
Creatinine	C00791	174.29	Arginine and proline metabolism	6.30 × 10 ^{−6}	1.43	1.36 × 10 ^{−3}	1.11
Phosphocreatine	C02305	450.62	Arginine and proline metabolism	3.41 × 10 ^{−6}	0.29	8.53 × 10 ^{−5}	0.39
D-Ribose 5-phosphate	C00117	410.74	Purine metabolism	3.72 × 10 ^{−4}	3.03	8.85 × 10 ^{−3}	1.49
Inosine monophosphate (IMP)	C00130	22.17	Purine metabolism	2.25 × 10 ^{−2}	0.60	4.80 × 10 ^{−5}	1.58

RT: retention time; FC: fold change; Con.: control; 2400 extract: 2400 mg/mL. ^a The *p* value was calculated by one-way ANOVA and then by Tukey's post hoc test for comparisons of multiple groups (after correction for multiple hypothesis testing and false discovery rate, 0.05). ^b The fold change (FC) of relative amounts of the drug treatment groups (GLA, gemcitabine and hydroxycamptothecin groups) compared with the normal control.

Mapping metabolites into pathways helps to understand the role of H₂O₂ and chrysanthemum extract on cell metabolism. In the arginine biosynthesis pathway, H₂O₂ induced a reduced reaction chain composed of N-Acetyl-L-glutamate, N-acetyloronithine, L-Ornithine, L-citrulline and N-(L-arginino) succinate, while chrysanthemum extract reversed it except L-Ornithine. Moreover, chrysanthemum extract increased IMP which is decreased by H₂O₂, whereas in arginine and proline metabolism, chrysanthemum extract did not influence the effect of H₂O₂ which is indicated by the changes in creatine and creatinine and the reduction in phosphocreatine was more severe (Figure 7). Therefore, the antioxidant effects of chrysanthemum extract may be associated with regulating the arginine biosynthesis pathway and the generation of IMP.




Figure 7. H_2O_2 and chrysanthemum extract regulated metabolic pathway. FC: fold change; Con.: control; H_2O_2 : 7 mM H_2O_2 ; 2400 + H_2O_2 : 2400 μ g/mL chrysanthemum extract and 7 mM H_2O_2 . Green represents the presence of enzymes in the human body.

5. Discussion

Chrysanthemum extracts exhibited significant hepatoprotective effects, but the mechanism was still unclear. In this study, through in vitro experiments, it was found that chrysanthemum aqueous extract exhibited significant antioxidant functions by effectively downregulating ROS content and lipid peroxidation. Then, chrysanthemum extract further inhibited H_2O_2 -induced decreases in mitochondrial membrane potential and hepatocyte death. H_2O_2 is a well-established inducer of oxidative stress damage which can induce an increase in ROS [17]. ROS can damage DNA and interfere with DNA replication, thus causing cell cycle arrest in the G1 phase [18]. Meanwhile, cell cycle arrest and damage to membrane phospholipids by ROS can further cause cell necrosis or necroptosis [19,20]. This fully indicated that the cytoprotective effects of chrysanthemum extract mainly involved its antioxidant effects targeting ROS. Meanwhile, this also explained the protective effect of chrysanthemum extract on mitochondria, as ROS is also a disruptor of mitochondrial function [21].

Physiologically, the intracellular oxides and peroxides are constantly produced, while a series of antioxidant enzymes can scavenge most of these cytotoxic substances and maintain the redox balance. SOD, CAT and GSH-Px constitute the basic antioxidant system in cells. SOD converts the highly reactive superoxide anion into relatively stable H_2O_2 ,

while CAT further reduces H_2O_2 to water. The family of GSH-Px, i.e., GPX, contains seven reviewed members catalyzing the reduction of a wide range of substrates including H_2O_2 and lipid peroxides. Here, chrysanthemum extract increased the total activity of SOD, CAT and GSH-Px underlying its antioxidant mechanism. Previous works have provided some possible mechanisms to explain how antioxidant enzymes are regulated by small molecule compounds. α -Tocopherol can increase the expression of superoxide dismutase [22], pioglitazone can increase catalase activity by changing the BLC structure [23] and selenium can enable the activity of GSH-Px as a cofactor [24]. Given that, it was supposed that there might also be similar molecules in chrysanthemum extract able to regulate the expression of antioxidant enzymes or directly enhance the activity of enzymes.

Previous studies often focused on the discovery and purification of bioactive substances from natural products. Flavonoids and triterpenes are common natural compounds found in chrysanthemum in which apigenin, apigenidin and faradiol, etc., have been identified as antioxidant and anti-inflammatory factors [25–28]. This is consistent with our findings on the antioxidant effect of chrysanthemum. Despite all of this, more than the study of the action of a single substance, the synergy of bioactive substances and the joint action on multiple targets cannot be ignored. Thus, we believe that, for natural extracts with complex components, it is more meaningful to study the collection of functions by means of omics.

Through metabolomics, we found that arginine biosynthesis, arginine and proline metabolism and purine metabolism were the main targets of chrysanthemum extract. As precursors of arginine, L-arginosuccinate and citrulline were inhibited by H_2O_2 but upregulated by chrysanthemum extract. As L-arginine induces antioxidant response via stimulation of glutathione synthesis and activation of the Nrf2 pathway [29], we speculate that the effect of chrysanthemum extract may be related to promoting arginine synthesis and downstream metabolism. Although citrulline was upregulated, ornithine, which is the precursor of citrulline, was downregulated by chrysanthemum extract. Previous work showed that aminoacylase-1 (3.5.1.14) was increased during oxidative stress, suggesting that chrysanthemum extract might inhibit the aminoacylase-1 and therefore cause the accumulation of N-acetyloronithine and further decreases in ornithine [30]. Significantly, H_2O_2 also decreased ornithine. However, this may be caused by a deficiency of the precursor N-acetyloronithine.

Inosine monophosphate (IMP) is an essential compound for de novo nucleotide biosynthesis and metabolism of energy, proteins, and antioxidants [31]. Mitochondrial DNA is susceptible to oxidation because of the frequent exposure to ROS. Meanwhile, ATP may promote benefits to oxidative phosphorylation and help repair ROS-induced damage to mitochondrial DNA. IMP can be converted to ADP or GDP and, subsequently, to ATP or GTP, thus playing an important role in DNA repair [32]. Here, H_2O_2 caused the exhaustion of IMP, while chrysanthemum extract could significantly restore it, suggesting an additional antioxidation mechanism. The increased IMP guarantees the energy of cells during oxidative stress and helps the scavenging of oxides. In addition, it is also able to ensure the repair of ROS-damaged DNA and this may be the potential mechanism by which chrysanthemum extract can unblock ROS-mediated cell cycle arrest.

6. Conclusions

In L-O2 hepatocytes, chrysanthemum extract can effectively inhibit oxidative stress and cell death caused by H_2O_2 . During this process, the levels of ROS and lipid peroxide decreased, the mitochondrial membrane potential recovered and the activity of antioxidant enzymes increased. Mechanistically, chrysanthemum-extract-mediated upregulation of arginine biosynthesis and IMP synthesis may contribute to the enhancement of antioxidant capacity.

Author Contributions: G.Z.: Data acquisition, writing—original draft, conceptualization. M.L. and K.S.: writing and revision. C.X.: writing—review and editing. R.Y.: software, Data analysis, visualization and formal analysis. All authors have read and agreed to the published version of the manuscript.

Funding: Financial support was provided by the Natural Science Research Project of Anhui Education Department (KJ2019A0638) and A Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Yuan, H.; Jiang, S.; Liu, Y.; Daniyal, M.; Jian, Y.; Peng, C.; Shen, J.; Liu, S.; Wang, W. The flower head of *Chrysanthemum morifolium* Ramat. (Juhua): A paradigm of flowers serving as Chinese dietary herbal medicine. *J. Ethnopharmacol.* **2020**, *261*, 113043. [[CrossRef](#)] [[PubMed](#)]
2. Jeong, S.C.; Kim, S.M.; Jeong, Y.T.; Song, C.H. Hepatoprotective effect of water extract from *Chrysanthemum indicum* L. flower. *Chin. Med.* **2013**, *8*, 7. [[CrossRef](#)] [[PubMed](#)]
3. Liu, G.; Zheng, Q.; Pan, K.; Xu, X. Protective effect of *Chrysanthemum morifolium* Ramat. ethanol extract on lipopolysaccharide induced acute lung injury in mice. *BMC Complement. Med. Ther.* **2020**, *20*, 235. [[CrossRef](#)] [[PubMed](#)]
4. Zhou, Y.; Wang, C.; Kou, J.; Wang, M.; Rong, X.; Pu, X.; Xie, X.; Han, G.; Pang, X. *Chrysanthemi Flos* extract alleviated acetaminophen-induced rat liver injury via inhibiting oxidative stress and apoptosis based on network pharmacology analysis. *Pharm. Biol.* **2021**, *59*, 1378–1387. [[CrossRef](#)] [[PubMed](#)]
5. Chen, Z.; Kong, S.; Song, F.; Li, L.; Jiang, H. Pharmacokinetic study of luteolin, apigenin, chrysoeriol and diosmetin after oral administration of Flos *Chrysanthemi* extract in rats. *Fitoterapia* **2012**, *83*, 1616–1622. [[CrossRef](#)] [[PubMed](#)]
6. Youssef, F.S.; Eid, S.Y.; Alshammari, E.; Ashour, M.L.; Wink, M.; El-Readi, M.Z. *Chrysanthemum indicum* and *Chrysanthemum morifolium*: Chemical composition of their essential oils and their potential use as natural preservatives with antimicrobial and antioxidant activities. *Foods* **2020**, *9*, 1460. [[CrossRef](#)] [[PubMed](#)]
7. Maione, F.; Cicala, C.; Musciacco, G.; De Feo, V.; Amat, A.G.; Ialenti, A.; Mascolo, N. Phenols, alkaloids and terpenes from medicinal plants with antihypertensive and vasorelaxant activities. A review of natural products as leads to potential therapeutic agents. *Nat. Prod. Commun.* **2013**, *8*, 539–544. [[CrossRef](#)]
8. Wang, Z.; Wang, L.; Luo, J.; Zhang, J. Protection against acute renal injury by naturally occurring medicines which act through Nrf2 signaling pathway. *J. Food Biochem.* **2021**, *45*, e13556. [[CrossRef](#)]
9. Jarco, S.; Pilawa, B.; Ramos, P. Free radical scavenging activity of infusions of different medicinal plants for use in obstetrics. *Plants* **2021**, *10*, 2016. [[CrossRef](#)]
10. Caesar, L.K.; Cech, N.B. Synergy and antagonism in natural product extracts: When 1 + 1 does not equal. *Nat. Prod. Rep.* **2019**, *36*, 869–888. [[CrossRef](#)]
11. Butterweck, V.; Lieflander-Wulf, U.; Winterhoff, H.; Nahrstedt, A. Plasma levels of hypericin in presence of procyanidin B2 and hyperoside: A pharmacokinetic study in rats. *Planta Med.* **2003**, *69*, 189–192. [[CrossRef](#)] [[PubMed](#)]
12. Snezhkina, A.V.; Kudryavtseva, A.V.; Kardymon, O.L.; Savvateeva, M.V.; Melnikova, N.V.; Krasnov, G.S.; Dmitriev, A.A. ROS generation and antioxidant defense systems in normal and malignant cells. *Oxidative Med. Cell. Longev.* **2019**, *2019*, 6175804. [[CrossRef](#)] [[PubMed](#)]
13. Pimentel, A.M.; Pereira, N.R.; Costa, C.A.; Mann, G.E.; Cordeiro, V.S.; de Moura, R.S.; Brunini, T.M.; Mendes-Ribeiro, A.C.; Resende, A.C. L-arginine-nitric oxide pathway and oxidative stress in plasma and platelets of patients with pre-eclampsia. *Hypertens. Res.* **2013**, *36*, 783–788. [[CrossRef](#)] [[PubMed](#)]
14. Pervin, M.; Unno, K.; Konishi, T.; Nakamura, Y. L-arginine exerts excellent anti-stress effects on stress-induced shortened lifespan, cognitive decline and depression. *Int. J. Mol. Sci.* **2021**, *22*, 508. [[CrossRef](#)] [[PubMed](#)]
15. Andrade, V.S.; Rojas, D.B.; Oliveira, L.; Nunes, M.L.; de Castro, F.L.; Garcia, C.; Gemelli, T.; de Andrade, R.B.; Wannmacher, C.M. Creatine and pyruvate prevent behavioral and oxidative stress alterations caused by hypertryptophanemia in rats. *Mol. Cell. Biochem.* **2012**, *362*, 225–232. [[CrossRef](#)] [[PubMed](#)]
16. Watanabe, L.M.; Fernandes de Lima, L.; Ferraz-Bannitz, R.; Takaara, D.; Coimbra Romano, B.; Braga Costa, T.M.; Foss de Freitas, M.C.; Bueno, A.C.; Barbosa Junior, F.; Marliere Navarro, A. Association between creatine kinase activity, oxidative stress and selenoproteins mRNA expression changes after Brazil nut consumption of patients using statins. *Clin. Nutr.* **2020**, *39*, 3175–3181. [[CrossRef](#)]
17. Evangelista-Vargas, S.; Santiani, A. Detection of intracellular reactive oxygen species (superoxide anion and hydrogen peroxide) and lipid peroxidation during cryopreservation of alpaca spermatozoa. *Reprod. Domest. Anim.* **2017**, *52*, 819–824. [[CrossRef](#)]

18. Rancourt, R.C.; Hayes, D.D.; Chess, P.R.; Keng, P.C.; O'Reilly, M.A. Growth arrest in G1 protects against oxygen-induced DNA damage and cell death. *J. Cell. Physiol.* **2002**, *193*, 26–36. [[CrossRef](#)]
19. Zhang, Y.; Su, S.S.; Zhao, S.; Yang, Z.; Zhong, C.Q.; Chen, X.; Cai, Q.; Yang, Z.H.; Huang, D.; Wu, R.; et al. RIP1 autophosphorylation is promoted by mitochondrial ROS and is essential for RIP3 recruitment into necrosome. *Nat. Commun.* **2017**, *8*, 14329. [[CrossRef](#)]
20. Csordas, A.; Wick, G.; Bernhard, D. Hydrogen peroxide-mediated necrosis induction in HUVECs is associated with an atypical pattern of caspase-3 cleavage. *Exp. Cell Res.* **2006**, *312*, 1753–1764. [[CrossRef](#)]
21. Fan, W.; Shen, T.; Ding, Q.; Lv, Y.; Li, L.; Huang, K.; Yan, L.; Song, S. Zearalenone induces ROS-mediated mitochondrial damage in porcine IPEC-J2 cells. *J. Biochem. Mol. Toxicol.* **2017**, *31*, e21944. [[CrossRef](#)]
22. Feng, Z.; Liu, Z.; Li, X.; Jia, H.; Sun, L.; Tian, C.; Jia, L.; Liu, J. alpha-Tocopherol is an effective Phase II enzyme inducer: Protective effects on acrolein-induced oxidative stress and mitochondrial dysfunction in human retinal pigment epithelial cells. *J. Nutr. Biochem.* **2010**, *21*, 1222–1231. [[CrossRef](#)] [[PubMed](#)]
23. Yekta, R.; Dehghan, G.; Rashtbari, S.; Sheibani, N.; Moosavi-Movahedi, A.A. Activation of catalase by pioglitazone: Multiple spectroscopic methods combined with molecular docking studies. *J. Mol. Recognit.* **2017**, *30*, e2648. [[CrossRef](#)] [[PubMed](#)]
24. Hariharan, S.; Dharmaraj, S. Selenium and selenoproteins: It's role in regulation of inflammation. *Inflammopharmacology* **2020**, *28*, 667–695. [[CrossRef](#)] [[PubMed](#)]
25. Lin, L.Z.; Harnly, J.M. Identification of the phenolic components of chrysanthemum flower (*Chrysanthemum morifolium* Ramat). *Food Chem.* **2010**, *120*, 319–326. [[CrossRef](#)]
26. Wang, D.; Yang, Y.; Zou, X.; Zhang, J.; Zheng, Z.; Wang, Z. Antioxidant apigenin relieves age-related muscle atrophy by inhibiting oxidative stress and hyperactive mitophagy and *Apoptosis* in skeletal muscle of mice. *J. Gerontol. Ser. A Biol. Sci. Med. Sci.* **2020**, *75*, 2081–2088. [[CrossRef](#)] [[PubMed](#)]
27. Amarowicz, R.; Pegg, R.B. The potential protective effects of phenolic compounds against low-density lipoprotein oxidation. *Curr. Pharm. Des.* **2017**, *23*, 2754–2766. [[CrossRef](#)] [[PubMed](#)]
28. Neukirch, H.; D'Ambrosio, M.; Sosa, S.; Altinier, G.; Della Loggia, R.; Guerriero, A. Improved anti-inflammatory activity of three new terpenoids derived, by systematic chemical modifications, from the abundant triterpenes of the flowery plant *Calendula officinalis*. *Chem. Biodivers.* **2005**, *2*, 657–671. [[CrossRef](#)]
29. Liang, M.; Wang, Z.; Li, H.; Cai, L.; Pan, J.; He, H.; Wu, Q.; Tang, Y.; Ma, J.; Yang, L. l-Arginine induces antioxidant response to prevent oxidative stress via stimulation of glutathione synthesis and activation of Nrf2 pathway. *Food Chem. Toxicol.* **2018**, *115*, 315–328. [[CrossRef](#)]
30. Cavalloni, G.; Peraldo-Neia, C.; Massa, A.; Bergamini, C.; Trentini, A.; De Rosa, G.; Daniele, L.; Ciccocanti, F.; Cervellati, C.; Leone, F.; et al. Proteomic analysis identifies deregulated metabolic and oxidative-associated proteins in Italian intrahepatic cholangiocarcinoma patients. *BMC Cancer* **2021**, *21*, 865. [[CrossRef](#)]
31. Popova, N.R.; Gudkov, S.V.; Bruskov, V.I. Natural purine compounds as radioprotective agents. *Radiatsionnaya Biol. Radioekol.* **2014**, *54*, 38–49.
32. Srinivasan, S.; Torres, A.G.; Ribas de Pouplana, L. Inosine in biology and disease. *Genes* **2021**, *12*, 600. [[CrossRef](#)] [[PubMed](#)]