



# Can manipulation under anesthesia alone provide clinical outcomes similar to arthroscopic circumferential capsular release in primary frozen shoulder (FS)?: the necessity of arthroscopic capsular release in primary FS

Seung-Jin Lee, Jun-Hyuk Jang, Yoon-Suk Hyun

Department of Orthopedic Surgery, Kangdong Sacred Heart Hospital, Seoul, Korea

**Background:** We evaluated the need for arthroscopic capsular release (ACR) in refractory primary frozen shoulder (FS) by comparing clinical outcomes of patients treated with ACR and manipulation under anesthesia (MUA).

**Methods:** We assessed patients with refractory primary FS, 57 patients (group A) who were treated with MUA and 22 patients (group B) who were treated with ACR. In group A, manipulation including a backside arm-curl maneuver was performed under interscalene brachial block. In group B, manipulation was performed only to release the inferior capsule before arthroscopic circumferential capsular release, which was carried out for the unreleased capsule after manipulation. Pain, range of shoulder motion, and American Shoulder and Elbow Surgeons score were recorded at 1 week, 3 months, 6 months, and 1 year after surgery. We compared outcome variables between treatment groups and between diabetics and non-diabetics and also evaluated the numbers of patients receiving additional intra-articular steroid injection.

**Results:** Outcome variables at 3 months after surgery and improvements in outcome variables did not differ between groups. Group A showed significantly better results than group B in the evaluation of pain and range of motion at 1 week. Diabetics showed comparable outcomes to non-diabetics for most variables. Eleven patients required additional steroid injections between 8 to 16 weeks after surgery: 12.2% in group A, 18.2% in group B. Additional injections were given three times more often in diabetics compared to non-diabetics.

**Conclusions:** MUA alone can yield similar clinical outcomes to ACR in refractory FS.

**Keywords:** Frozen shoulder; Manipulation; Capsular release; Diabetes

## INTRODUCTION

Even if the natural course of primary frozen shoulder (FS) is mostly self-limiting, some patients experience prolonged disability with considerable pain and disability that affect activities of

daily life [1,2]. Moreover, some patients fail to achieve desired outcomes with non-operative management. Given the natural history of adhesive capsulitis and the high proportion of patients who do well with nonsurgical management, a trial of at least 6 months of nonsurgical management is normally recommended

Received: September 30, 2020 Revised: November 1, 2020 Accepted: November 4, 2020

Correspondence to: Yoon-Suk Hyun

Department of Orthopedic Surgery, Kangdong Sacred Heart Hospital, 150 Seongan-ro, Gangdong-gu, Seoul 05355, Korea

Tel: +82-2-2224-2230, Fax: +82-2-489-4391, E-mail: [yshyun72@gmail.com](mailto:yshyun72@gmail.com), ORCID: <https://orcid.org/0000-0003-3826-2527>

IRB approval: Kangdong Sacred Heart Hospital (No. 2019-09-016).

Financial support: None.

Conflict of interest: None.

Copyright© 2020 Korean Shoulder and Elbow Society. All Rights Reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

before considering surgical management options: manipulation under anesthesia (MUA), arthroscopic capsular release (ACR), or the combination of both [3].

Although ACR is gaining in popularity with recent advances in arthroscopic technique and has shown promising results comparable to those of other treatment modalities [4-6], MUA is a traditionally well-established treatment for FS that is nevertheless controversial due to potential complications (e.g., proximal humerus fractures, shoulder dislocation, brachial plexus stretching injury, rotator cuff injury, and recurrent stiffness) [7-10]. There are no good quality randomized controlled trials in favor of ACR in comparison to MUA [11]; in two previous studies, the superior treatment was not identified [11,12] and manipulation was performed under general anesthesia. Manipulation for FS is usually performed under general anesthesia, but is also performed under interscalene brachial plexus block (ISB) anesthesia and obtains favorable outcomes [13-16].

No previous studies have compared the clinical outcomes of manipulation under ISB anesthesia and ACR. We evaluated differences in clinical outcomes between manipulation under ISB and ACR in refractory primary FS to determine whether ACR is necessary if manipulation under interscalene brachial plexus anesthesia is performed following our novel method. We hypothesized that MUA alone would provide similar clinical outcomes as ACR in primary FS.

## METHODS

### Study Design and Participants

We performed a retrospective analysis of a prospectively collected, single surgeon (YSH), single institution, consecutive series of patients with FS. The protocol of this study was reviewed and approved by Institutional Review Board of Kangdong Sacred Heart Hospital (IRB No. 2019-09-016). Written informed consents were obtained. From March 2015 to Mar 2018, 79 patients who were diagnosed with primary FS in our hospital were treated with MUA or ACR. The definition of FS in this study followed the American Shoulder and Elbow Surgeons (ASES) consensus study by Zuckerman and Rokito [17] and was characterized by functional restriction of both active and passive shoulder motion for which radiographs of the glenohumeral joint were essentially unremarkable except for the possible presence of osteopenia. In all patients, shoulder magnetic resonance imaging was used to screen for the coexistence of any other shoulder lesions before indicated management was performed. Patients with rotator cuff tear, shoulder osteoarthritis, calcified tendinitis, hemiplegia after stroke, bone metastasis in the shoulder region, history of shoul-

der fractures, and history of shoulder surgeries were excluded. Diabetic FS patients were not excluded because while diabetes is a possible predisposing factor based on statistical data, it is not known to be a cause of FS [18]. Refractory FS was defined as follows: refractory to conservative treatment (intra-articular steroid injections and physical therapy) for at least 6 months and documented restriction of both passive and active glenohumeral and scapulothoracic motion of equal to or less than 100° of elevation, and less than 50% of external rotation, as compared to the contralateral side [11].

Both treatment modalities were indicated by refractory FS. There were no differences in indications between the two modalities. ACR after MUA was used from January 2015 to May 2017, and the two modalities were used randomly without special indications for either during the 5 months from January 2017 onward. After this period of overlap, we recognized that the two modalities showed similar clinical outcomes. MUA alone was used after May 2017 to facilitate comparisons of the modalities. All 79 patients treated during this period who were followed closely over 6 months after treatments were reviewed. A total of 57 patients (group A) were treated with MUA and 22 patients (group B) were treated with ACR after manipulation. We evaluated range of motion (ROM; passive forward elevation, external rotation arm at side, and thumb reach along the vertebral spine, in which the thumb points up), visual analog pain score, and ASES score preoperatively and at 1 week, 3 months, 6 months, and 1 year after surgery. All outcome variables in this study were measured by trained residents and physician assistants under the supervision of the senior author (YSH). We consider the number of vertebral spines in which the patient's thumb can reach up, which we refer to as thumb-to-spine, to be better than degree of internal rotation for the evaluation of surgical outcomes. When the patient's thumb reaches up over the thoracolumbar junction, we defined it as a pass. Regarding the evaluation of pain severity, we asked the patient for an average value, taking into account pain while sleeping at night, pain while resting, and pain during everyday activities. We compared results for the two groups and analyzed them statistically. We compared outcome measurements, pretreatment period, follow-up period, and number of additional intra-articular steroid injections between groups. We also compared all variables between diabetic and non-diabetic patients.

### Study Procedures

In group A, all MUA procedures were performed by a single surgeon (YSH) following a standardized, identical protocol. MUA was performed in outpatient settings. The patient was made to lie

down in a supine position after interscalene plexus block was performed by the anesthesiologist. Before MUA, intra-articular steroid injections were performed to control post-MUA inflammation and pain (triamcinolone 40 mg, 1 mL+lidocaine 0.1%, 5 mL+sterile normal saline, 4 mL). Manipulations were first performed at forward flexion with scapular stabilization by an assistant, and then at external rotation with the arm at the side, and finally at 90° abduction (Fig. 1) [16]. Next, internal rotation was performed with the arm at 90° abduction, while the assistant pushed the shoulder girdle downward to the floor (Fig. 1A) [16]. During 90° abduction with full internal rotation, further abduction can lead to further capsular release indicated by an audible sound (Fig. 1B). Next, horizontal adduction with scapular stabilization was performed. After the patient was made to sit, we placed the patient's hand behind their back and then pushed the patient's arm backwards, while pushing the patient's hand up with

forced adduction (Fig. 2). This last step is the “backside arm-curl maneuver.” Most of the time, a recognizable tearing sound was heard during each step. The sequence was repeated until the maximum ROM was acquired. A short lever arm with the elbow flexed at 90° is used to prevent fractures and brachial plexus traction injuries. After MUA completion, the recovered range of shoulder motion was confirmed with the patient (Fig. 3). The patient was instructed to start passive exercise programs immediately after MUA to maintain the restored ROM, in which forward elevation and the backside arm-curl maneuver were emphasized (Fig. 4). The patient was also instructed to perform a self-assisted stretch for 5 minutes every hour until the next visit. Patients were allowed to go home after recovery from ISB anesthesia.

In group B, manipulation preceded arthroscopic procedures and was performed at forward elevation only for the safe release of the inferior capsular area after interscalene plexus block and




**Fig. 1.** (A) Internal rotation with the arm at 90° abduction, with the assistant pushing the shoulder girdle downward toward the floor was performed to complete the posterior and inferior capsule tear. (B) With 90° abduction with full internal rotation, further abduction can result in further capsular release with an audible sound.



**Fig. 2.** Backside arm-curl maneuver. (A) It is often hard to place the patient's hand on the midline of the back in the sitting position even after full restoration of internal rotation in the supine position. (B, C) After the patient was made to sit, we placed the patient's hand behind their back and then pushed the patient's arm backwards, while pushing the patient's hand up with forced arm adduction.



general anesthesia. Therefore, the backside arm-curl maneuver in the sitting position was omitted in group B. During ACR, the unreleased inferior capsule, anterior capsule, posterior capsule, and anterior part of the superior capsule (superior glenohumeral ligament and coracohumeral ligament) were released. After arthroscopic 360° circumferential capsular release, an epidural catheter was inserted into the glenohumeral joint space for intra-articular steroid injection. The day after surgery, the passive exercise program was started after intra-articular steroid injection. The exercise program in group B also emphasized forward elevation and the backside arm curl maneuver. The patients remained in hospital until they showed more than 135° of forward elevation and could reach higher than the fifth lumbar vertebra during the backside arm curl maneuver at least once. The length of hospital stay after surgery averaged 1 or 2 days. We informed the patient that the sooner they began exercise the better regarding the result of ROM restoration. The patient was also instructed to perform a self-assisted stretch for 5 minutes every hour until the next visit. All patients received instructions to perform rehabilitation on their own without help.



**Fig. 3.** (A, B) After manipulation under anesthesia completion, range of motion was confirmed with the patient.



**Fig. 4.** (A, B) The patient was instructed to initiate passive exercise programs right after manipulation under anesthesia to maintain the restored range of motion, in which forward elevation and the backside arm-curl maneuver were emphasized.

Within 1 week after discharge, all patients in both groups visited the outpatient clinic to check whether they had maintained the restored ROM in group A and how much ROM had been restored in group B. All patients were evaluated according to our follow-up schedule. In group B, additional visits during the month after surgery were performed to encourage patients to perform rehabilitation exercises because most group B patients did not show significant improvements in pain and ROM at the first visit. During the follow up period, we performed intra-articular steroid injections (triamcinolone 40 mg, 1 mL +lidocaine 0.1%, 5 mL+sterile normal saline, 4 mL) if the patient complained of aggravated pain impeding ROM exercises and night sleep.

For statistical analysis, independent t-tests (age), Fisher's exact tests (sex) and Mann-Whitney U-tests (ASES scores, ROM, pre-treatment periods, follow-up periods, pain VAS) were used with significance set at the 5% level (IBM SPSS 22.0; IBM Corp., Armonk, NY, USA).

## RESULTS

There were no significant differences in demographic data between groups (Table 1). Twenty-four patients had diabetes mellitus (30.4%, 24 of 79 patients). Three patients had thyroid disease in group A, while none did in group B. Preoperative outcome

**Table 1.** Demographic data for both study groups

Variable	Group A	Group B	p-value
Number of patients	57	22	
Age (yr)	55.3 ± 8.5	53.9 ± 6.4	0.474
Male:female	24:33	8:14	
Diabetes	17 (29)	7 (32)	0.443
Thyroid disease case	3	0	
Pretreatment period (mo)	6.4 ± 3.7	6.6 ± 4.1	0.391
Follow-up period (mo)	7.68 ± 1.7	7.22 ± 1.6	0.285

Values are presented as mean ± standard deviation or number (%).

variables, outcome variables at 3 months after surgery, and amount of improvement in all variables (ASES score, pain VAS, and ROM) did not differ between groups (Tables 1 and 2). However, at the last visit, the degree of forward elevation was better in group A than group B (p = 0.029) although other outcome variables did not differ between groups.

No patients were able to raise their thumbs higher than the thoracolumbar junction. At the first visit after surgery, all patients in group A were graded as passing. In later evaluations, all patients in group A continued to be graded as passing. However, in the ACR group, only five patients passed at their first visit. The numbers of patients who passed were 12 at 1 month, 21 at 3 months, and 22 (all patients) at the last visit. Group A showed significantly better results than group B for pain and ROM (Table 1).

No patients in either group (manipulation under ISB only and ACR with MUA) showed decreased ROM during the follow-up period. Seven patients among 57 treated by MUA (12.2%) and four by ACR (18.2%) complained of aggravated pain between 8 and 16 weeks (median value, 10 weeks) after surgery. The pain in these 11 patients subsided after a single intra-articular steroid injection and did not recur during the follow-up period. These 11 patients did not show any significant differences in ROM, ASES score, or severity of pain at final evaluation. No serious complications after manipulation, such as shoulder dislocation, rotator cuff tear, or brachial plexus palsy were reported in our patients.

We included 24 diabetic patients (30.4%, 24/79) in this study. When outcome variables were evaluated without grouping, diabetic patients showed slightly worse outcomes at 3 months after surgery, but no significant little differences in last values or improvement of outcome variables (Table 3). The results were different when each group was analyzed separately. There were no differences in clinical outcomes between diabetics and non-diabetics in group A (Table 3). In group B, diabetics had worse outcomes at 3 months after surgery and at the last visit, but there were no differences in improvement of ASES score or ROM except for pain severity (Table 3). Among patients who needed additional intra-articular steroid injections, 41.2% of diabetic patients (7/17) and 7.5% of non-diabetic patients (3/40) in group A needed additional injections, and 57.1% of diabetic patients (4/7) and 11.1% of non-diabetic patients (2/18) in group B needed additional injections.

## DISCUSSION

We did not detect differences in overall outcomes between patients treated with MUA and ACR. Interestingly, pain severity and ROM scores were better in the MUA group than the ACR

**Table 2.** Outcome variables in groups A and B

Variable	Preoperative		p-value	First visit within 1 week		p-value	After 3 months		p-value	Last visit		Improvement*		p-value
	Group A	Group B		Group A	Group B		Group A	Group B		Group A	Group B	Group A	Group B	
Pain VAS	56.3 ± 12.3	58.63 ± 13.5	0.466	28.9 ± 12.5	55.9 ± 15.6	<0.001	19.9 ± 20.0	19.8 ± 10.5	0.944	16.3 ± 9.7	16.1 ± 11.2	40.0 ± 16.0	42.0 ± 15.6	0.533
ASES score	48.7 ± 8.2	49.7 ± 7.7	0.596	No data	No data		79.5 ± 7.2	79.0 ± 8.4	0.757	84.5 ± 6.9	84.9 ± 6.8	35.9 ± 11.4	35.1 ± 7.3	0.760
Range of motion														
FE	87.5 ± 9.7	84.3 ± 9.7	0.263	148.7 ± 10.6	126.8 ± 13.0	<0.001	155.4 ± 5.5	153.4 ± 7.5	0.190	160 ± 5.8	156.1 ± 9.2	72.4 ± 13.3	71.8 ± 14.2	0.852
ERside	6.3 ± 7.4	6.1 ± 8.0	0.925	48.3 ± 6.4	27.3 ± 9.7	<0.001	50.1 ± 7.7	46.8 ± 8.8	0.100	52.7 ± 8.2	51.6 ± 8.5	46.4 ± 11.1	45.5 ± 11.9	0.739
Thumb to spine <sup>†</sup>	0:0			57:5			57:21			57:22				

Values are presented as mean ± standard deviation.

VAS: visual analog scale, ASES: American Shoulder and Elbow Surgeons, FE: forward elevation, ERside: external rotation arm at side.

\*The amount of improvement of outcome variables between preoperative values and values at the last visit; improvement = value at last visit - preoperative value. †The ratio of the number of patients whose thumb could reach up over the thoracolumbar junction, group A:group B.

**Table 3.** Outcome variables in diabetic patients and non-diabetic patients among all 79 patients

Variable	Group A+B						Group A						Group B																										
	Preoperative		After 3 months		Last visit		Improvement*		Preoperative		After 3 months		Last visit		Improvement*		Preoperative		After 3 months		Last visit		Improvement*																
	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value	Non-DM	DM	P-value															
Pain VAS	56.4 ± 1.2	57.2 ± 1.1	0.818	24.6 ± 1.1	17.9 ± 1.0	0.033	17.5 ± 1.0	15.7 ± 1.0	0.474	38.9 ± 1.1	41.5 ± 1.1	0.523	55.6 ± 1.1	56.6 ± 1.1	0.774	23.8 ± 1.1	18.3 ± 1.1	0.267	14.1 ± 1.1	17.3 ± 1.1	0.267	41.5 ± 1.1	39.4 ± 1.1	0.655	58.6 ± 1.1	58.7 ± 1.1	0.988	26.4 ± 1.1	16.7 ± 1.1	0.039	25.7 ± 1.1	11.7 ± 1.1	0.003	32.8 ± 1.1	47 ± 1.1	0.045			
ASES score	49.1 ± 1.2	48.9 ± 1.1	0.899	76 ± 1.1	80.9 ± 1.1	0.007	82 ± 1.1	85.8 ± 1.1	0.023	32.9 ± 1.1	36.9 ± 1.1	0.114	49.6 ± 1.1	48.2 ± 1.1	0.555	13 ± 1.1	18.3 ± 1.1	0.163	8.7 ± 1.1	8.6 ± 1.1	0.163	82.6 ± 1.1	85.4 ± 1.1	0.163	32.9 ± 1.1	37.2 ± 1.1	0.205	47.9 ± 1.1	50.6 ± 1.1	0.449	73.3 ± 1.1	81.8 ± 1.1	0.027	80.6 ± 1.1	86.8 ± 1.1	0.043	32.7 ± 1.1	36.2 ± 1.1	0.309
Range of motion	7.4 ± 1.1	8.4 ± 1.1	0.493	7 ± 1.1	7 ± 1.1	0.205	6.4 ± 1.1	6.8 ± 1.1	0.129	10.5 ± 1.1	10.2 ± 1.1	0.792	8.4 ± 1.1	8.2 ± 1.1	0.342	6.4 ± 1.1	7.4 ± 1.1	0.678	6.1 ± 1.1	7.1 ± 1.1	0.678	160 ± 1.1	160 ± 1.1	0.411	11.7 ± 1.1	11.2 ± 1.1	0.411	4.1 ± 1.1	8.9 ± 1.1	0.655	10.6 ± 1.1	6 ± 1.1	0.076	7.3 ± 1.1	5.8 ± 1.1	0.089	64.3 ± 1.1	75.3 ± 1.1	0.089
ERside	8.1 ± 1.1	5.5 ± 1.1	0.191	46.3 ± 1.1	50.5 ± 1.1	0.020	50.4 ± 1.1	53.3 ± 1.1	0.160	42.3 ± 1.1	47.8 ± 1.1	0.044	7.9 ± 1.1	5.7 ± 1.1	0.284	48.2 ± 1.1	51 ± 1.1	0.573	51.8 ± 1.1	53.1 ± 1.1	0.573	5.6 ± 1.1	6 ± 1.1	0.258	11.9 ± 1.1	13.9 ± 1.1	0.99	8.6 ± 1.1	5 ± 1.1	0.342	41.4 ± 1.1	49.3 ± 1.1	0.047	47.1 ± 1.1	53.7 ± 1.1	0.094	38.6 ± 1.1	48.7 ± 1.1	0.061
	8.7 ± 1.1	6.9 ± 1.1		8 ± 1.1	7.9 ± 1.1		8.3 ± 1.1	7.9 ± 1.1		7.3 ± 1.1	7.9 ± 1.1		8.1 ± 1.1	8.4 ± 1.1		8.1 ± 1.1	8.2 ± 1.1		8.1 ± 1.1	8.4 ± 1.1		11.9 ± 1.1	10.8 ± 1.1		10.3 ± 1.1	6.8 ± 1.1		8 ± 1.1	8.2 ± 1.1		8.1 ± 1.1	8.1 ± 1.1		15.7 ± 1.1	8.3 ± 1.1				

Values are presented as mean±standard deviation.

DM: diabetes mellitus, VAS: visual analog scale, ASES: American Shoulder and Elbow Surgeons, FE: forward elevation, ERside: external rotation arm at side.  
\*The amount of improvement of outcome variables between preoperative values and values at the last visit; improvement=value at last visit-preoperative value.

group at the first visit, but there were no differences in final outcomes or improvement between groups.

Eleven patients complained of aggravated pain between 8 and 16 weeks (median, 10 weeks) after each procedure: 7 of 57 patients (12.2%) in the MUA group and 4 of 22 patients (18.2%) in the ACR group. Regarding the recurrence of pain, the number of diabetic patients requiring additional injections was three times higher than non-diabetic patients.

Given the natural history of adhesive capsulitis and the high proportion of patients who do well with nonsurgical management, a trial of at least 6 months of nonsurgical management is generally recommended before considering surgical management options [3]. MUA and ACR are the two most common surgical management strategies. There are no good quality randomized controlled trials with findings showing the superiority of ACR in comparison to MUA [11], but ACR is favored over MUA because it is believed to allow a more controlled and complete release of the contracted capsule, to reduce the risk of iatrogenic injury compared to MUA, and also resulted in favorable short-term and long-term outcomes [11,19-24]. In our study, patients treated with MUA showed similar outcomes to ACR and did not report any serious iatrogenic complications.

There is controversy regarding the optimal method of ACR. Some authors have recommended 360° capsular release to restore normal ROM, whereas others have determined that partial capsular release (anterior-inferior capsule) is sufficient [4,22,24,25]. In group B, we performed 360° circumferential capsular release, in which the coracohumeral ligament and superior glenohumeral ligament as well as the anterior-inferior-posterior capsule were released, because we believe that release of the coracohumeral ligament and superior glenohumeral ligament is important [4,26-28]. In three patients who were not included in this study, we performed arthroscopic examinations to determine the extent of capsular release after MUA that included the backside arm-curl maneuver. We found that the anterior middle glenohumeral ligament, inferior glenohumeral ligament (anterior and posterior band), and posterior capsule below the equator were successfully released. Previous studies supporting the efficacy of arthroscopic anteroinferior release and our clinical findings that MUA with backside arm-curl could restore nearly normal ROM, constitute evidence that ACR is unnecessary.

Recent developments in ultrasound technology have enabled its use with brachial plexus anesthesia or cervical root block [13-16,29]. Previous reports described good results like those of the present study in primary FS patients with or without diabetes. However, no previous studies have compared outcomes of manipulation under ISB and ACR in patients with refractory prima-

ry FS. Our study may be the first to compare the outcomes of patients treated by both modalities in a single institution. Our novel manipulation techniques resulted in comparable clinical outcomes to ACR.

Only two previous studies compared outcomes of MUA and ACR in patients with refractory FS [11,12]. Neither determined which of the two treatment modalities is superior. Grant et al. [11] conducted a systematic review and concluded that given the low level of evidence and lack of direct comparisons, there are no clear differences in shoulder ROM or patient-reported outcomes when comparing MUA to ACR for the treatment of refractory primary FS. Kim et al. also reported similar results, that MUA (general anesthesia) offered equivalent clinical outcomes compared with ACR (360° circumferential release) in the early period after the procedure [12]. However, they recommended that clinicians should consider MUA as an option before choosing ACR in patients with refractory FS because MUA is simple, safe, and cost-effective.

In this study, we introduced and described the novel backside arm-curl maneuver in the sitting position. In previous studies of MUA in primary FS, the descriptions of manipulation procedures are brief or variable, and procedures are generally not conducted in the sitting position [12,15,18,20,25]. For the successful restoration of ROM in MUA, the restoration of the height of thumb reach up the spine as well as forward elevation is very important. We found that sufficient shoulder extension, backward elevation, and adduction in the sitting position are critical for the restoration of the height of the thumb reaching up the spine to normal range and tearing sounds are audible during this procedure. Performance of the backside arm-curl maneuver in the sitting position is extremely important.

Woods and Loganathan [30] found that patients with successful outcomes have significant improvement in pain within three to four days after MUA, and improvement in ROM within about three weeks. Kraal et al. [29] also found that an initial period of one to 2 weeks of intensive physiotherapy after MUA is essential to prevent recurrence of restrictions and advocated more aggressive rehabilitation with intensive stretching and ROM exercises in the first weeks after MUA to preserve the obtained ROM. The importance of early rehabilitation is the same in ACR and MUA in terms of surgical treatment for joint stiffness [20]. We also emphasized the early restoration of ROM in both groups. Patients in the MUA-only group showed earlier improvements in pain and ROM at first visit than the ACR group, although there was no significant difference at 3 months between groups. Final outcomes are important, but it is also important to treat pain quickly and restore ROM. The faster recovery of ROM and pain, the bet-

ter the patient's treatment compliance.

When using our method of performing MUA, we were able to immediately confirm ROM recovery during manipulation under ISB. ISB and intra-articular steroid injections minimized pain during the rehabilitation exercise, which was carried out immediately after MUA, so ROM restored with the procedure was well maintained. Immediate initiation of exercise after manipulation for 6 to 8 hours becomes possible thanks to the use of ISB, and intra-articular steroid injections also reduce pain on exercise after ISB wears off. Less invasive procedures than ACR are also helpful to reduce pain on rehabilitation. Patients in the ACR group reported more pain during joint exercise compared to MUA at their first outpatient visits.

There are clear differences difference between presence and absence of restored ROM. In manipulation under ISB, there is no ROM to restore, and full restoration is accomplished as soon as manipulation is complete. In MUA or ACR under general anesthesia, the immediate confirmation of restoration of ROM with the patient is difficult or impossible, and some patients doubt the success of manipulation if rehabilitation is onerous or they do not restore normal ROM.

Regarding the recurrence of FS after MUA, two previous studies of MUA reported rates of recurrence of between 10% and 40% [30,31]. Woods and Loganathan [30] performed further MUA if there were no improvements in ROM or pain 3 months after MUA. They also reported a period of recurrence in some patients (17.8%, 141 of 792 shoulders) between 6 and 8 weeks after treatment, which could be accounted for by the loss of the effect of the intra-articular steroid injection. In our study, no patients experienced decreased ROM during the follow-up period, while 11 patients complained of aggravated pain between 8 and 16 weeks after the procedure: seven of 57 patients (12.2%) in the MUA group and four of 22 patients (18.2%) in the ACR group.

FS is strongly associated with diabetes and is two to four times more common in diabetic patients than in the general population [32-36]. FS in patients with diabetes tends to have a more severe and protracted course, and to be difficult to treat [32,37,38]. Some researchers have suggested that patients with diabetes should consider ACR, but most of our patients experienced good outcomes after manipulation alone [17]. However, in our study, diabetic patients showed slightly worse outcomes compared with non-diabetic patients although outcome variables in the ACR group were not significantly different. We observed no differences in any outcome variables in the MUA group, a finding that is consistent with those of other reports and suggests that diabetes is not a contraindication for MUA [16,18,31,34]. Regarding recurrence after surgery, Jenkins et al. reported that re-



peat MUA was required in 36% of diabetic patients compared with 15% of nondiabetic patients, and Woods and Loganathan [30] reported that patients with type-1 diabetes mellitus were at 38% increased risk of requiring further MUA, compared with 18% increased risk among a group that included both diabetics and non-diabetic patients [31]. Regarding recurrence pain, diabetic patients were more than three times as likely to experience recurrence than non-diabetics (41.2% vs. 7.5% in group A and 57.1% vs. 11.1% in group B) in our study.

This study has several limitations. It is a retrospective, non-randomized study with a relatively small sample size. Sample sizes are also different between groups. There may be subtle differences in the indications of the two procedures because ACR requires general anesthesia, so it can be more difficult for patients to accept than MUA, which is performed under regional anesthesia. We developed and introduced the backside arm-curl maneuver with good success, but further research is needed to validate this procedure. Future studies should include evaluations of ROM improvement after manipulation with and without the backside arm-curl maneuver. Comparisons between manipulation under ISB and general anesthesia should also be performed in the future. Despite these limitations, our study is the first to compare the clinical results of manipulation under ISB with ACR.

In primary FS, manipulation under ISB alone can result in similar and favorable clinical outcomes with ACR in refractory cases. Regardless of type of surgery, clinical outcomes in patients with diabetes were similar to those without diabetes. However, diabetic patients often require additional intra-articular steroid injections.

## ORCID

Seung-Jin Lee <https://orcid.org/0000-0001-8951-0039>  
 Jun-Hyuk Jang <https://orcid.org/0000-0002-9640-1700>  
 Yoon-Suk Hyun <https://orcid.org/0000-0003-3826-2527>

## REFERENCES

1. Reeves B. The natural history of the frozen shoulder syndrome. *Scand J Rheumatol* 1975;4:193-6.
2. Shaffer B, Tibone JE, Kerlan RK. Frozen shoulder: a long-term follow-up. *J Bone Joint Surg Am* 1992;74:738-46.
3. Hsu JE, Anakwenze OA, Warrender WJ, Abboud JA. Current review of adhesive capsulitis. *J Shoulder Elbow Surg* 2011; 20:502-14.
4. Cvetanovich GL, Leroux TS, Bernardoni ED, et al. Clinical outcomes of arthroscopic 360° capsular release for idiopathic adhesive capsulitis in the lateral decubitus position. *Arthroscopy* 2018;34:764-70.
5. Gallacher S, Beazley JC, Evans J, et al. A randomized controlled trial of arthroscopic capsular release versus hydrodilatation in the treatment of primary frozen shoulder. *J Shoulder Elbow Surg* 2018;27:1401-6.
6. De Carli A, Vadalà A, Perugia D, et al. Shoulder adhesive capsulitis: manipulation and arthroscopic arthrolysis or intra-articular steroid injections. *Int Orthop* 2012;36:101-6.
7. Loew M, Heichel TO, Lehner B. Intraarticular lesions in primary frozen shoulder after manipulation under general anesthesia. *J Shoulder Elbow Surg* 2005;14:16-21.
8. Amir-Us-Saqblain H, Zubairi A, Taufiq I. Functional outcome of frozen shoulder after manipulation under anaesthesia. *J Pak Med Assoc* 2007;57:181-5.
9. Milch H. Brachial palsy after manipulation of frozen shoulder. *N Engl J Med* 1954;250:429-30.
10. Uppal HS, Evans JP, Smith C. Frozen shoulder: a systematic review of therapeutic options. *World J Orthop* 2015;6:263-8.
11. Grant JA, Schroeder N, Miller BS, Carpenter JE. Comparison of manipulation and arthroscopic capsular release for adhesive capsulitis: a systematic review. *J Shoulder Elbow Surg* 2013; 22:1135-45.
12. Kim DH, Song KS, Min BW, Bae KC, Lim YJ, Cho CH. Early clinical outcomes of manipulation under anesthesia for refractory adhesive capsulitis: comparison with arthroscopic capsular release. *Clin Orthop Surg* 2020;12:217-23.
13. Ando A, Hamada J, Hagiwara Y, Sekiguchi T, Koide M, Itoi E. Short-term clinical results of manipulation under ultrasound-guided brachial plexus block in patients with idiopathic frozen shoulder and diabetic secondary frozen shoulder. *Open Orthop J* 2018;12:99-104.
14. Saito T, Sasanuma H, Iijima Y, et al. Short-term clinical results of frozen shoulder treated with shoulder manipulation under ultrasound-guided cervical nerve root block at outpatient setting: a case series. *J Orthop Sci* 2017;22:275-80.
15. Kraal T, Van der Meer O, Van den Borne M, Koenraadt K, Eygendaal D, Boer R. Manipulation under anesthesia for frozen shoulders: a retrospective cohort study. *Acta Orthop Belg* 2019; 85:400-5.
16. Takahashi R, Kajita Y, Harada Y, Iwahori Y, Deie M. Clinical results of shoulder manipulation under ultrasound-guided cervical nerve root block for frozen shoulder in patients with diabetes. *J Orthop* 2020;21:297-301.
17. Zuckerman JD, Rokito A. Frozen shoulder: a consensus definition. *J Shoulder Elbow Surg* 2011;20:322-5.



18. Wang JP, Huang TF, Ma HL, Hung SC, Chen TH, Liu CL. Manipulation under anaesthesia for frozen shoulder in patients with and without non-insulin dependent diabetes mellitus. *Int Orthop* 2010;34:1227-32.
19. Neviaser AS, Hannafin JA. Adhesive capsulitis: a review of current treatment. *Am J Sports Med* 2010;38:2346-56.
20. Neviaser AS, Neviaser RJ. Adhesive capsulitis of the shoulder. *J Am Acad Orthop Surg* 2011;19:536-42.
21. Farrell CM, Sperling JW, Cofield RH. Manipulation for frozen shoulder: long-term results. *J Shoulder Elbow Surg* 2005;14:480-4.
22. Le Lievre HM, Murrell GA. Long-term outcomes after arthroscopic capsular release for idiopathic adhesive capsulitis. *J Bone Joint Surg Am* 2012;94:1208-16.
23. Barnes CP, Lam PH, Murrell GA. Short-term outcomes after arthroscopic capsular release for adhesive capsulitis. *J Shoulder Elbow Surg* 2016;25:e256-64.
24. Ranalletta M, Rossi LA, Zaidenberg EE, et al. Midterm outcomes after arthroscopic anteroinferior capsular release for the treatment of idiopathic adhesive capsulitis. *Arthroscopy* 2017; 33:503-8.
25. Kim YS, Lee HJ. Essential surgical technique for arthroscopic capsular release in the treatment of shoulder stiffness. *JBJS Essent Surg Tech* 2015;5:e14.
26. Kim YS, Chung SW, Kim JY, Ok JH, Park I, Oh JH. Is early passive motion exercise necessary after arthroscopic rotator cuff repair. *Am J Sports Med* 2012;40:815-21.
27. Ozaki J, Nakagawa Y, Sakurai G, Tamai S. Recalcitrant chronic adhesive capsulitis of the shoulder. Role of contracture of the coracohumeral ligament and rotator interval in pathogenesis and treatment. *J Bone Joint Surg Am* 1989;71:1511-5.
28. Pouliart N, Somers K, Eid S, Gagey O. Variations in the superior capsuloligamentous complex and description of a new ligament. *J Shoulder Elbow Surg* 2007;16:821-36.
29. Kraal T, The B, Boer R, et al. Manipulation under anesthesia versus physiotherapy treatment in stage two of a frozen shoulder: a study protocol for a randomized controlled trial. *BMC Musculoskelet Disord* 2017;18:412.
30. Woods DA, Loganathan K. Recurrence of frozen shoulder after manipulation under anaesthetic (MUA): the results of repeating the MUA. *Bone Joint J* 2017;99-B:812-7.
31. Jenkins EF, Thomas WJ, Corcoran JP, et al. The outcome of manipulation under general anesthesia for the management of frozen shoulder in patients with diabetes mellitus. *J Shoulder Elbow Surg* 2012;21:1492-8.
32. Wang K, Ho V, Hunter-Smith DJ, Beh PS, Smith KM, Weber AB. Risk factors in idiopathic adhesive capsulitis: a case control study. *J Shoulder Elbow Surg* 2013;22:e24-9.
33. Thomas SJ, McDougall C, Brown ID, et al. Prevalence of symptoms and signs of shoulder problems in people with diabetes mellitus. *J Shoulder Elbow Surg* 2007;16:748-51.
34. Theodorides AA, Owen JM, Sayers AE, Woods DA. Factors affecting short- and long-term outcomes of manipulation under anaesthesia in patients with adhesive capsulitis of the shoulder. *Shoulder Elbow* 2014;6:245-56.
35. Bridgman JF. Periarthritis of the shoulder and diabetes mellitus. *Ann Rheum Dis* 1972;31:69-71.
36. Arkkila PE, Kantola IM, Viikari JS, Rönnemaa T. Shoulder capsulitis in type I and II diabetic patients: association with diabetic complications and related diseases. *Ann Rheum Dis* 1996;55: 907-14.
37. Janda DH, Hawkins RJ. Shoulder manipulation in patients with adhesive capsulitis and diabetes mellitus: a clinical note. *J Shoulder Elbow Surg* 1993;2:36-8.
38. Griggs SM, Ahn A, Green A. Idiopathic adhesive capsulitis. A prospective functional outcome study of nonoperative treatment. *J Bone Joint Surg Am* 2000;82:1398-407.