



RESEARCH ARTICLE

REVISED Australian state influenza notifications and school holiday closures in 2019 [version 3; peer review: 2 approved, 1 approved with reservations]

Previously titled: Australian state influenza notifications and school closures in 2019

Anna Mae Scott ¹, Mina Bakhit ¹, Justin Clark ¹, Melanie Vermeulen¹, Mark Jones¹, David Looke^{2,3}, Chris Del Mar¹, Paul Glasziou¹

¹Institute for Evidence-Based Healthcare, Bond University, Robina, Queensland, 4226, Australia

²Department of Medicine, University of Queensland, St Lucia, Queensland, 4072, Australia

³Infectious Disease and Clinical Microbiology, Princess Alexandra Hospital, Brisbane, Queensland, Australia

V3 First published: 11 Nov 2019, 8:1899
<https://doi.org/10.12688/f1000research.21145.1>

Second version: 08 Oct 2020, 8:1899
<https://doi.org/10.12688/f1000research.21145.2>

Latest published: 26 May 2021, 8:1899
<https://doi.org/10.12688/f1000research.21145.3>

Abstract

Background: The impact of school holidays on influenza rates has been sparsely documented in Australia. In 2019, the early winter influenza season coincided with mid-year school breaks, enabling us the unusual opportunity to examine how influenza incidence changed during school holiday closure dates.

Methods: The weekly influenza data from five Australian state and one territory health departments for the period of week 19 (mid-May) to week 39 (early October) 2019 were compared to each state's public-school holiday closure dates. We used segmented regression to model the weekly counts and a negative binomial distribution to account for overdispersion due to autocorrelation. The models' goodness-of-fit was assessed by plots of observed versus expected counts, plots of residuals versus predicted values, and Pearson's Chi-square test. The main exposure was the July two-week school holiday period, using a lag of one week. The effect is estimated as a percent change in incidence level, and in slope.

Results: School holidays were associated with significant declines in influenza incidence in three states and one territory by between 41% and 65%. Two states did not show evidence of declines although one of those states had already passed its peak by the time of the school holidays. The models showed acceptable goodness-of-fit. The first decline during school holidays is seen in the school aged (5-19 years) population, with the declines in the adult and infant populations being smaller and following a week later.

Conclusions: Given the significant and rapid reductions in incidence,

Open Peer Review

Reviewer Status ? ✓ ✓

	Invited Reviewers		
	1	2	3
version 3 (revision) 26 May 2021			
version 2 (revision) 08 Oct 2020		report ✓	report ✓
version 1 11 Nov 2019	report ?	report ?	


- Casey Zipfel** , Georgetown University, Washington, USA
- Shweta Bansal** , Georgetown University, Washington, USA
- Núria Torner** , University of Barcelona, Barcelona, Spain
- August Wrotek** , Centre of Postgraduate Medical Education, Warsaw, Poland

these results have important public health implications. Closure or extension of holiday periods could be an emergency option for state governments.

Keywords

Influenza, influenza epidemic, child, adolescent, school closure, school holidays, absenteeism

Any reports and responses or comments on the article can be found at the end of the article.



This article is included in the [Disease Outbreaks](#) gateway.

Corresponding author: Anna Mae Scott (ascott@bond.edu.au)

Author roles: **Scott AM:** Data Curation, Investigation, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing; **Bakhit M:** Data Curation, Investigation, Writing – Review & Editing; **Clark J:** Data Curation, Investigation, Writing – Review & Editing; **Vermeulen M:** Data Curation, Investigation, Visualization, Writing – Review & Editing; **Jones M:** Formal Analysis, Methodology, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Looke D:** Conceptualization, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Del Mar C:** Conceptualization, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Glasziou P:** Conceptualization, Formal Analysis, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: The author(s) declared that no grants were involved in supporting this work.

Copyright: © 2021 Scott AM *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Scott AM, Bakhit M, Clark J *et al.* **Australian state influenza notifications and school holiday closures in 2019 [version 3; peer review: 2 approved, 1 approved with reservations]** F1000Research 2021, 8:1899 <https://doi.org/10.12688/f1000research.21145.3>

First published: 11 Nov 2019, 8:1899 <https://doi.org/10.12688/f1000research.21145.1>

REVISED Amendments from Version 2

At the reviewer's suggestion, we have rephrased a sentence in the discussion from: "no other intervention has comparable effects" to "no other intervention may have comparable effects."

Any further responses from the reviewers can be found at the end of the article

Introduction

In 2009, the United Kingdom experienced a summer influenza epidemic, with the National Health Service resorting to pharmacy dispensed oseltamivir to slow the growth. However, when schools closed in August for the six-week summer break, the epidemic dropped to almost zero within a few weeks¹. This was an extreme example of an association that has been documented in other countries, though generally for shorter closures²⁻⁴. A 2013 review of both deliberate and non-deliberate school closures concluded that even without co-interventions, closure of schools could reduce influenza transmission during an outbreak^{3,5}.

The association of school closure and influenza rates has been only sparsely documented in Australia, partly because school holidays generally fall outside the peak influenza period. However, 2019 saw an early epidemic of influenza in Australia with rates around five times normal for the May-July period, with consequent hospitalisation and deaths also increased. Because of the early high rates, winter influenza in 2019 also coincided with the mid-year school breaks, which appeared to be associated with a dip in influenza incidence. Since each country's school system differs in features such as age range, classroom structure and sizes, and class mixing, we thought it important to document the size of any impact in Australia. Furthermore, the variation by states, and the ability to examine delay to changes in the non-school population would extend the range of sub-questions we could address.

To explore any relationship between the school holiday closure and influenza cases, we examined the relationship between changes in influenza incidence with the school holiday closure dates in 2019 using the different holiday dates that apply in the Australian states and territories.

Methods

Setting

Influenza is a notifiable disease in all Australian states and territories⁶. We collected influenza data notified weekly to state and territory health departments, for the period of week 19 (mid-May) to week 39 (early October) 2019, which corresponds to the influenza season in Australia.

We included data from five states (New South Wales, Queensland, Victoria, South Australia, Western Australia) and one territory (Australian Capital Territory); we excluded data from one state (Tasmania) and one territory (Northern Territory) due to paucity of available data and small population sizes.

Data sources and extraction

States and territories differ in how they collect and report data on influenza cases. New South Wales⁷ and Queensland⁸ report the number of samples that test positive for Influenza A and B (lab-confirmed); Australian Capital Territory⁹ reports the number of influenza (lab) notifications to the state of influenza A, influenza B, and co-infections of influenza A and B; whilst Western Australia¹⁰, Victoria¹¹ and South Australia¹² report the number of lab-confirmed influenza cases (the specific strains are unspecified).

We collected influenza data as it was reported by each jurisdiction. We collected data on public school holiday closures between week 19 and week 39 from each state or territory's education department website.

Weekly, numerical influenza data were available from reports produced by the Health Departments in New South Wales, Queensland, Western Australia, Victoria, and Australian Capital Territory. For South Australia, the number of lab-confirmed influenza cases for weeks 19–24 were extracted from a figure (using *webplotdigitiser*¹³) since weekly numerical data were available only for weeks 25–35. Data on school holidays were extracted from each state or territory's website. Raw data are provided (see *Data availability* section, Appendix 1)¹⁴.

Statistical methods

Weekly counts of influenza cases were collected from five states and one territory across Australia for the first 10 months of 2019. We restricted analysis to the period from week 19 (beginning of school term 2) to week 39 of 2019 (end of school term 3). Due to differences in the population numbers of each state and territory, differences in the methods of data collection (e.g. some states use only sentinel sites), and insufficient number of states to reliably fit a random effects model, the analysis was conducted separately by state.

We used segmented regression¹⁵ to model the weekly counts and specified a negative binomial distribution to account for overdispersion. Explanatory factors included the linear effect of week (slope), initial effect of holiday (change in level), change in linear effect of week after start of holiday (change in slope), initial effect of returning to school, and change in linear effect of week after school return (Appendix 2)¹⁶. Goodness-of-fit of the models was assessed by plots of observed versus expected counts, plots of residuals versus fitted values, and Pearson's Chi-square test. The main exposure for our analysis was the July school holiday period of two weeks. Due to the delay from exposure to the virus to confirmation of infection status, we included a lag of one week¹⁷. The effects are reported as a % change in level (with 95% confidence interval) as well as a percentage change in slope (with 95% confidence interval). We hypothesized that the holiday period (lagged by one week) would lead to reductions in weekly counts of influenza cases.

Statistical analysis was conducted using SAS University Edition 9.4 for Windows.

Results

Figure 1 Influenza cases for the included states and territory. Figures for each individual state or territory are provided in Appendix 3¹⁶.

The estimates for the initial effect and the subsequent slopes (Table 1) show significant declines for all states except South Australia, which had already passed its peak by the time of the school holidays, and Victoria. The models showed acceptable goodness-of-fit (Appendix 3 and 4) with Pearson's Chi-square tests all indicating insufficient evidence of lack of fit ($P > 0.05$).

Results shown in the table can be interpreted as follows, using NSW as an example: In NSW prior to the school holiday the rate of influenza cases was increasing by 26% (+22%, +31%) per week. The school holiday was associated with a 50% decrease (-27%, -66%) in level as well as a 27% decrease (-53%, +13%) in slope. After the students returned to school, there was a 38% increase (-33%, +185% increase) in level and 6% decrease (-40%, +46%) in slope. QLD, ACT and WA showed a similar pattern whereas school holiday appeared to have little impact in SA and VIC.

The data on influenza rates by age group (Figure 2) show the first decline during school holidays is seen in the school aged (5–19 years) population, with the decline in the adult (20–64 years) population being smaller and following a further week later, and with even smaller and delayed drops in the infant (1–4) and elderly (65+ years) age groups. We did not have access to the state specific age bands, and hence have indicated the school holiday closure dates with the largest population groups.

Discussion

The 2019 Australia influenza notification data show a significant association between school holidays and declines in influenza incidence in most states in Australia. We also found the earliest and largest declines in influenza incidence were in the school aged group (5–19 years), with still later and smaller declines in the adult group (20–64 years), and least impact on the preschool (0–4 years) and over 65's.

Our findings are consistent with previous reports³ of school closure for usual holiday periods and emergency closure for epidemics as well as studies of how much worse epidemics would have been if schools had *not* been closed¹⁸.

The size of the declines is also consistent with those predicted by models of transmission for school closures¹⁹. Some




Figure 1. Rates of influenza for the included Australian states and territory (boxes show school holidays periods). NSW, New South Wales; QLD, Queensland; WA, Western Australia; ACT, Australian Capital Territory; VIC, Victoria; SA, Southern Australia.

Table 1. Estimates (95% CI) from segmented regression models of weekly influenza counts.

State	Pre-holiday linear effect of week	Change in level due to school holiday*	Change in slope due to holiday*	Change in level after school return^	Change in slope after school return^
NSW	+26% (+22%, +31%); P<.0001	-50% (-66%, -27%); P=.0004	-27% (-53%, +13%); P=.15	+38% (-33%, +185%); P=.38	-6% (-40%, +46%); P=.78
QLD	+24% (+19%, +29%); P<.0001	-41% (-59%, -16%); P=.004	-10% (-40%, +36%); P=.63	+79% (-8%, +150%); P=.087	-19% (-46%, +22%); P=.31
VIC	+13% (+7%, +18%); P<.0001	-2% (-39%, +57%); P=.93	-21% (-54%, +36%); P=.40	+1% (-58%, +143%); P=.98	+7% (-37%, +84%); P=.80
WA	+10% (+1%, +19%); P=.03	-65% (-84%, -21%); P=.012	-44% (-88%, +41%); P=.22	+7% (-77%, +390%); P=.93	+32% (-48%, +231%); P=.56
SA	-17% (-20%, -14%); P<.0001	+28% (-14%, +90%); P=.22	-6% (-41%, +51%); P=.81	+21% (-44%, +163%); P=.63	+33% (-17%, +114%); P=.23
ACT	+20% (+14%, +26%); P<.0001	-42% (-64%, -6%); P=.025	-35% (-62%, +13%); P=.13	+125% (-9%, +455%); P=.079	+8% (-38%, +87%); P=.79

*Compared to pre-holiday period

^Compared to holiday period

NSW, New South Wales; QLD, Queensland; VIC, Victoria; WA, Western Australia; SA, South Australia; ACT, Australian Capital Territory.

Influenza cases by age group




Figure 2. Influenza cases by age group (box shows the school holidays period).

states experienced a rebound within weeks of school restarting whereas others saw a continued decline. We do not have a simple explanation for this difference.

The association of school holiday closure with declines in influenza has several potential explanations besides a causal effect. First, cases of influenza might be underreported because of delayed presentation or non-presentation during the school holiday period, for example, because with parents able to care for them they do not attend for medical certificates. However, if this were true, we would expect to see an immediate restart after school returns, which is not the case. Second, it could be because of other societal changes, such as parents also being on holiday and hence less transmission at work. However, if this were true, we would expect to see a simultaneous reduction in both child and adult cases. Hence, the most likely remaining explanation seems to be causal reduction in transmission from school holidays. However, a final issue is that, even if reduction in incidence is real, it not clear whether there is a net annual decrease or merely a delay in total annual cases.

There are some limitations to both our data and the analysis. Good quality, numerical weekly influenza data were unavailable for all states – e.g., South Australia reports its data as a mix of figures and numbers. This may have introduced some errors into the accuracy of the numbers for those states. We contacted the Australian National Notifiable Diseases Surveillance System, which coordinates the national surveillance of influenza in Australia, to obtain raw data for each state and territory. However, the raw data underlying the notifications for 2019 will not be available for release until July 2020. There is also likely to be differences in the accuracy of different states' data, due to the different methods of collection. For the analysis, we used a lag of one week to allow for the delay from exposure to the virus to confirmation of infection status¹⁷. In a sensitivity analysis we refitted the model without a lag and produced consistent results (Appendix 5, *Extended data*)¹⁶. The school holiday only lasted 2 weeks hence our estimates of change in slope are imprecise. We used a lag of 1 week to allow for the delay from exposure to the virus to confirmation of infection status. In a sensitivity analyses we refitted the model without a lag. Finally, we did not include a control group (i.e. previous years' data), although existing data on notifications of laboratory-confirmed influenza in Australia between 2013 and 2019 shows the notifications in 2019 to be higher than the 5-year average, and higher than all of the years other than 2017²⁰.

Given the size of the effect, these results have important public health implications as no other intervention may have comparable effects. Hence closure or extension of holiday periods could be an emergency option for state governments. Vaccinating children against influenza has also been shown to substantially reduce influenza attack rates in *both* children and adults²¹. Thus, in addition to encouraging influenza vaccination, the Centres for Disease Control has a number of suggestions in their guidance, such as encouraging students and staff to stay home when sick, liberalising sickness policies during epidemics, encouraging respiratory etiquette, encouraging hand hygiene, regular cleaning of shared surfaces such as door handles and faucets, and providing a “sick room” to quarantine students with flu-like illnesses²². An additional strategy is to consider face masks, which, with hand hygiene, appear to substantially reduce transmission. All these strategies would need to be triggered by health departments to schools at an appropriate point in an epidemic or pandemic.

Data availability

Underlying data

This project contains the following underlying data: Australian state influenza notifications and holiday school closures in 2019: Appendix 1 – Underlying data DOI: [10.26139/5c47ae4cd8e16](https://doi.org/10.26139/5c47ae4cd8e16)¹⁴

Data can be accessed at: <https://research.bond.edu.au/en/datasets/australian-state-influenza-notifications-and-school-closures-in-2>

Extended data

This project contains the following extended data: Australian state influenza notifications and school holiday closures in 2019: Appendix 2-5 – Extended data DOI: [10.4225/57/555d781f8f2a3](https://doi.org/10.4225/57/555d781f8f2a3)¹⁶

Data can be accessed at: <https://research.bond.edu.au/en/datasets/australian-state-influenza-notifications-and-school-closures-in-2-2>

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/) (CC-BY 4.0).

Acknowledgements

We would like thank Nancy Glasziou for pointing out the potential of school holiday closure, and for comments on the manuscript. We would also like to thank August Wrotek, Casey Zipfel, Shweta Bansal, and Núria Torner for their helpful feedback.

References

1. Health Protection Agency: **Weekly pandemic flu media update: 17 December 2009**. 2009; [Accessed: 18 October 2019]. [Reference Source](#)
2. Ewing A, Lee EC, Viboud C, et al.: **Contact, Travel, and Transmission: The Impact of Winter Holidays on Influenza Dynamics in the United States**. *J Infect Dis*. 2017; **215**(5): 732–739. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Jackson C, Vynnycky E, Hawker J, et al.: **School closures and influenza:**

- systematic review of epidemiological studies. *BMJ Open*. 2013; 3(2): e002149.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
4. Luca GD, Kerckhove KV, Coletti P, *et al.*: **The impact of regular school closure on seasonal influenza epidemics: a data-driven spatial transmission model for Belgium.** *BMC Infect Dis*. 2018; 18(1): 29.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 5. Bin Nafisah S, Alamery AH, Al Nafesa A, *et al.*: **School closure during novel influenza: A systematic review.** *J Infect Public Health*. 2018; 11(5): 657–661.
[PubMed Abstract](#) | [Publisher Full Text](#)
 6. Australian Government: **Australian national notifiable diseases and case definitions.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 7. New South Wales Dept of Health: **Influenza Surveillance Report.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 8. Queensland Health: **Influenza surveillance reporting.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 9. ACT Government Department of Health: **Flu in the ACT.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 10. Government of Western Australia Dept of Health: **Virus WAtch.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 11. Victoria Department of Health: **Seasonal Influenza Reports for 2019.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 12. South Australia Health: **Surveillance of notifiable conditions.** 2019; [Accessed: 20 September 2019].
[Reference Source](#)
 13. Rohatgi A: **WebPlotDigitizer (Version 4.2).** 2019; [Accessed: 25 September 2019].
[Reference Source](#)
 14. Scott A, Bakhit M, Clark J, *et al.*: **Australian state influenza notifications and school closures in 2019: Appendix 1 – Underlying data.** Bond University. Scott_et_al_Australian_state_influenza_notifications_and_school_closures_in_2019_Appendix_1_Underlying_data.xlsx, 2019.
<http://www.doi.org/10.26139/5c47ae4cd8e16>
 15. Gebiski V, Ellingson K, Edwards J, *et al.*: **Modelling interrupted time series to evaluate prevention and control of infection in healthcare.** *Epidemiol Infect*. 2012; 140(12): 2131–41.
[PubMed Abstract](#) | [Publisher Full Text](#)
 16. Scott A, Bakhit M, Clark J, *et al.*: **Australian state influenza notifications and school closures in 2019: Appendix 2-5 – Extended data.** Bond University. Appendix_2_5_Extended_data.pdf, 2019.
<http://www.doi.org/10.4225/57/555d781f8f2a3>
 17. Centers for Disease Control and Prevention: **Seasonal Influenza (Flu): How Flu Spreads.** 2019a; [Accessed: 25 September 2019].
[Reference Source](#)
 18. Eames KTD: **The influence of school holiday timing on epidemic impact.** *Epidemiol Infect*. 2014; 142(9): 1963–71.
[PubMed Abstract](#) | [Publisher Full Text](#)
 19. Jackson C, Mangtani P, Hawker J, *et al.*: **The effects of school closures on influenza outbreaks and pandemics: systematic review of simulation studies.** *PLoS One*. 2014; 9(5): e97297.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Australian Government: **Australian Influenza Surveillance Report: No. 12, 2019.** 23 September to 6 October 2019. 2019
[Reference Source](#)
 21. Basta NE, Chao DL, Halloran ME, *et al.*: **Strategies for pandemic and seasonal influenza vaccination of schoolchildren in the United States.** *Am J Epidemiol*. 2009; 170(6): 679–86.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Centers for Disease Control and Prevention: **Guidance for School Administrators to Help Reduce the Spread of Seasonal Influenza in K-12 Schools.** 2019b; [Accessed: 25 September 2019].
[Reference Source](#)

Open Peer Review

Current Peer Review Status: ? ✓ ✓

Version 2

Reviewer Report 04 May 2021

<https://doi.org/10.5256/f1000research.29863.r83446>

© 2021 Wrotek A. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

✓ **August Wrotek** 

Department of Pediatrics, Centre of Postgraduate Medical Education, Warsaw, Poland

Influenza is one of the most common paediatric global health concerns, and various strategies have been suggested to decrease the spread of the disease. The authors of the study aimed to address the question of the impact of temporary school closing on the epidemics. In 2019, Australia faced an unexpected situation when the beginning of the winter season coincided with holiday school closing. The methodology of the study based upon segmented regression modelling in order to assess the effect of the school closure. The authors showed a strong significant decrease (ranging from 41% to 65%) in 3 out of 5 analysed states and in 1 territory (ranging from 41% to 65%). Of note, a decrease in the incidence was first noticed in the school-aged children, followed by the decrease in adults and infants (yet, to a smaller degree, and observed a week later). That indirectly proves the role of school-aged children as vectors who increase the disease spread. The results of the study may in fact have important public health implications. After the comments of the first two reviewers, the doubts have been cleared, and appropriate corrections have been made. Although, for example, a sentence "as no other intervention has comparable effects" may be somehow controversial (thus might be considered to be deleted), yet the general results and conclusions of the study may resume the "old-fashioned" approach based upon temporary closing of the school in the periods of highest influenza virus activity, especially since "lockdown"-based strategies have been applied globally in the COVID-19 era. Certainly, the strategies to decrease influenza cases based on school closing need to be well documented, but this study indeed expands our knowledge on this topic.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Paediatrics, infectious diseases, pulmonology, particularly influenza, RSV, epidemiology, air pollution, inflammatory markers.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 11 May 2021

Anna Scott, Bond University, Robina, Australia

Dear Doctor Wrotek,

Thank you very much for your kind comments.

Following on your suggestion, we have rephrased the sentence "no other intervention has comparable effects" to "no other intervention may have comparable effects."

With warmest wishes,

Anna Scott, on behalf of the authors

Competing Interests: No competing interests.

Reviewer Report 12 October 2020

<https://doi.org/10.5256/f1000research.29863.r72716>

© 2020 Torner N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Núria Torner 

CIBER Epidemiology and Public Health CIBERESP, University of Barcelona, Barcelona, Spain

The author has satisfactorily replied to my queries.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Surveillance, prevention and control of infectious disease outbreaks, with specific experience in seasonal influenza.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 27 July 2020

<https://doi.org/10.5256/f1000research.23276.r66760>

© 2020 Torner N. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

? **Núria Torner** 

CIBER Epidemiology and Public Health CIBERESP, University of Barcelona, Barcelona, Spain

Seasonal Influenza Surveillance deploying epidemiological and virological data on patients with influenza virus infection is a global task performed in coordination with WHO requirements. These include sentinel surveillance and hospital surveillance for severe cases of laboratory confirmed influenza. Even though influenza has been studied in depth, there are still some issues that need to be addressed, at least to encourage increase in vaccine uptake and recommendations.

There are several issues that are not clear and being this a point observation of just one season, inference is not sound enough to ascertain as evidence. Comparison with previous seasons would be desirable.

Comments:

1. One highly contrasted issue is the amplification that children trigger as to transmission of infection to their siblings and household members from the school setting. And this might be prevented by vaccination. (Basta *et al.* 2009)¹. So this is not new but it might be relevant for Australia to implement preventive measures, if so this should be stated.
2. During epidemic periods transmission is high and the number of syndromic cases make laboratory confirmation out of range of health systems. Is this so in Australia? As stated in data sources only laboratory confirmed cases are taken into account for reporting: *States and territories differ in how they collect and report data on flu cases. New South Wales⁷ and Queensland⁸ report the number of samples that test positive for Influenza A and B (lab-confirmed); Australian Capital Territory⁹ reports the number of influenza (lab) notifications to the state; and Western Australia¹⁰, Victoria¹¹ and South Australia¹² report the number of lab-confirmed influenza cases (strains unspecified).*

What is the difference between the three (besides type identification)?

Positive for Influenza A and B (lab-confirmed); number of influenza (lab) notifications; and number of lab-confirmed influenza cases (strains unspecified). Do the authors mean type and subtype by strain? Or does it make reference to the strain type?

Type A

Subtype A H1N1

Strain: A/Brisbane/59/2007 (H1N1)

Use the correct term.

3. There is also the climate factor to be taken into account. Do all regions included in the study have the same temperature, humidity range? Some differences in decrease of transmission could also be due to a lesser degree of transmission such as in Victoria and Southern Australia for example. Did this particular season have a different climatology that made it coincide with the school holiday period?
4. The results of the pre-school closure period refer to the baseline epidemic burden of disease, if so then there is a decline before school closure? By the way, the term school closure can be misleading as to a containment preventive measure that can be implemented to stop transmission and what is referred to in the study should be specific to school holiday closure. What has happened in other epidemic seasons should be included to be able to compare and strengthen the results obtained.
5. In the introduction first line: A pandemic is global, so it wasn't just the UK that was affected. As a matter of fact school closure emerged as one of the non-pharmaceutical strategies during that pandemic and has been implemented in the present COVID-19 pandemic. Sentence should be rephrased.

The authors findings could be in the same line as Eames (2013)² who stated that allowing some flexibility in holiday timing would provide maximum health benefit from the holiday period, yet this is difficult to carry out because of the different social and labor implications it has for schools, parents, and care givers. Was this the aim of the study?

6. In the conclusions the authors state that: no other intervention has comparable effects. There is no comparison in the study to other non-pharmaceutical mitigation measures. Can you include a reference to this statement?

Minor comments:

- Include label on Figure 1, are they number of cases in total for all states or incidence rates? Please complete.
- Why does Fig 1 have dates in the x-axis and Figure 2 the week number? Why different, they should be the same.

References

1. Basta NE, Chao DL, Halloran ME, Matrajt L, et al.: Strategies for pandemic and seasonal influenza vaccination of schoolchildren in the United States. *Am J Epidemiol.* 2009; **170** (6): 679-86 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Eames KT: The influence of school holiday timing on epidemic impact. *Epidemiol Infect.* 2014; **142** (9): 1963-71 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

No

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Surveillance, prevention and control of infectious disease outbreaks, with specific experience in seasonal influenza.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 24 Sep 2020

Anna Scott, Bond University, Robina, Australia

Thank you very much for your comments. We provide our responses below:

1. We agree that a multi-year comparison would be preferable. However, for 2020, the number of flu infections is atypical (because of COVID), and for years prior to 2019, raw data is very difficult to obtain as it requires multiple data access applications to the individual states and to the Commonwealth. We have included the lack of control group as a limitation of the study, and provided a reference to a Government document comparing the trend from 2019 to years going back to 2013.
2. Thank you for providing this reference. We added the following sentence to the Discussion: "Vaccinating children against influenza has also been shown to substantially reduce influenza attack rates in *both* children and adults (Basta et al 2009)."
3. The relatively small size of the Australian population (approximately 25 million) makes laboratory confirmation a bit easier than in many other, more populous countries (e.g. in Europe, North America, South America). As we note in the methods section, we report lab-confirmed cases.
4. States/territories report influenzas a bit differently from each other. States of NSW and QLD report influenza A and influenza B only. ACT reports influenza A, influenza B, and coinfections with influenza A and B. The remaining states report lab-confirmed influenzas but do not specify strains. We have made this clearer in the text.
5. The temperatures were 1.6 deg above the long-term average, in line with global warming but with no unusual anomaly in July 2019 (<http://www.bom.gov.au/climate/mwr/aus/mwr-aus-201907.pdf>). We do not have humidity data, but rainfall was lower than usual, suggesting lower humidity.
6. There was no decline before school closure – as Figure 1 shows, the decline only occurs within the grey boxes (the holiday periods). Any apparent decline – e.g. NSW – is because we have joined the dots of the weekly reported numbers.
7. We have modified the title, and edited the body of the manuscript to refer to school holiday closures, to make this clearer.
8. We have replaced the term 'pandemic' with the term 'epidemic' in the first line of the introduction.
9. Eames' study used modelling to estimate a counterfactual: what would have happened if the UK had NOT closed the schools during the 2009 epidemic - and found the epidemic would have been much worse than it was. We estimated what happened with the actual school holiday closure in Australia - and found the influenza incidence decreased. Our study and Eames study are therefore consistent. We have added this point to the discussion section.
10. We added references to the newly updated Cochrane review by Tom Jefferson, on physical interventions to reduce the spread of respiratory viruses.
11. They are cases - we edited the figure and its title to clarify this.
12. We edited the figures so they both refer to dates for consistency.

Competing Interests: n/a

Reviewer Report 16 January 2020

<https://doi.org/10.5256/f1000research.23276.r57787>

© 2020 Bansal S et al. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Casey Zipfel

Department of Biology, Georgetown University, Washington, DC, USA

Shweta Bansal

Department of Biology, Georgetown University, Washington, DC, USA

Summary:

Australia's influenza season does not typically coincide with school holidays. However, in 2019, the influenza season occurred earlier than usual, allowing the opportunity to evaluate the impacts of school closure on influenza dynamics in Australia. In this article, the authors present state and territory-level influenza data by week, and perform segmented regression, comparing influenza incidence before and after the school holiday. They also count weekly increases and decreases before, during, and after the school holiday, and show age-based dynamics. This article asks an important question of public health relevance to non-pharmaceutical interventions in a setting where it has not been previously investigated in a concise manner. However, this article could be strengthened through additional framing, increased statistical rigor, and more nuanced interpretation.

Comments:

- The introduction leaves several unanswered questions that would help to more clearly identify the importance of the study. The relationship between school closure and influenza dynamics has been explored in a number of previous papers, with both epidemiological data and mechanistic models. Thus, it is unclear to me why is it important to consider this relationship in Australia, specifically. Do the authors have hypotheses regarding trends or mechanisms that might be different from other countries? Is it important for public health planning? This question seems especially salient to me since school closures and the influenza season in Australia do not typically coincide, so these epidemiological trends may not be typical for Australia.
- The article concludes that there is a significant reduction in influenza in most states in Australia based on the results of segmented regression comparing influenza incidence level and slope before and after the holiday period. I am not totally convinced by this conclusion, as Table 1 shows that the change in level is not significant for Victoria, Australian Capital Territory, and Southern Australia, and the change in slope is not significant for Queensland, Victoria, Western Australia, and Southern Australia. This is contrary to the section of the results that states that "the estimates for the initial effect and the subsequent slope show significant declines for all states except South Australia". These results could be more carefully interpreted. This would give rise to an interesting question: what are the differences between the states/territories that result in different dynamics?

- I think this article could be strengthened with a more robust time series modeling approach, like ARIMA modeling. This would be particularly useful because the predicted vs. residual plot in Appendix 4 appears to indicate heteroskedasticity in the data. Including model terms in a time series model could handle this issue, as well as temporal autocorrelation.
- The results also provide the percent of weeks in the pre-vacation, during-vacation, and post-vacation period in which there is a weekly decrease in influenza incidence, indicating that there is a higher proportion of decreases in the during and post-vacation period. However, the pre-vacation period seems to include all of the early part of the season. If there were numerous decreases in this time period, then the onset of the influenza season would not have occurred. Thus, this finding may be a bit circular.
- It would be useful to compare this influenza season, which co-occurs with the holiday period, with another influenza season that does not co-occur with the holiday period, as a control. This could help to isolate the impact of school closure on influenza dynamics.
- The authors specify mid-May to early September as their study period. However, it's unclear what total population and age-specific dynamics looked like post week 35. Past studies have shown that school holidays simply delay dynamics so that the epidemic wave recovers to pre-holiday rates after the holiday. I would be curious if this is the case here.
- In the model, a one-week lag was assumed. Is there evidence about care-seeking and influenza surveillance/reporting to support this? The discussion mentions a sensitivity analysis performed with no lag, but I wonder if the lag could be longer due to a combination of delay to seeking healthcare and delay in reporting.
- The fourth paragraph of the discussion comments on the possible causes of the association between school closure and influenza. This paragraph rules out several possible explanations but does not then provide a plausible explanation.

Minor comments:

- A y-axis label on Figure 1 and Figure 2 would be helpful. The Figure 1 caption says it is the influenza rate, so I assume it is influenza cases per week, but this could be clearer.
- Is the x-axis in Figure 2 the week number? It would be easier to interpret if it was put into the same units as Figure 1, with dates instead of week counts.
- Does Figure 2 include cases for all of the states and territories combined?
- In Table 1, it is unclear whether the changes described (in both slope and level) are comparing the during-vacation period or the post-vacation period to the pre-vacation period. The observed and predicted plots in Appendix 3 appear to show 3 different segments fit for each of these periods, but it is unclear what change is reported in Table 1.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Partly

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Modeling social and spatio-temporal dynamics of infectious disease, with specific experience in seasonal influenza.

We confirm that we have read this submission and believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Author Response 24 Sep 2020

Anna Scott, Bond University, Robina, Australia

Thank you very much for your comments. We provide our responses to each comment below:

1. We have added to the introduction the following: "Since each country's school system differs in features such as age range, classroom structure and sizes, and class mixing, we thought it important to document the size of any impact in Australia. Furthermore, the variation by states, and the ability to examine delay to changes in the non-school population would extend the range of sub-questions we could address."
2. We agree, and have revised our interpretation of the results.
3. We are unfamiliar with ARIMA modelling, but generally prefer simpler approaches. Our main purpose for the statistical analysis is to test hypotheses for each state on whether the school closures have resulted in changes in levels and/or changes in slopes with regards to influenza cases. The goodness-of-fit measures suggest the regression models provide adequate fit hence we believe our results are valid.
4. We agree, and have removed this analysis from the manuscript.
5. Unfortunately we were unable to formally compare as the previous years' raw numbers are not readily available. We have included the lack of control group as a limitation of the study, and provided a reference to a Government document comparing the trend from 2019 to years going back to 2013.

6. We have included 1 additional month of data – up until the start of the October school holiday period.
7. A lag of 2 weeks would only be plausible if a person being exposed to influenza on the last week of school prior to the vacation would generally be reported at least 2 weeks and 3 days later. Given “symptoms can begin about 2 days (but can range from 1 to 4 days) after the virus enters the body” this delay seems unrealistic.
8. We agree, and have added the following: "Hence, the most likely remaining explanation seems to be causal reduction in transmission from school holidays."
9. Comments: 9-12. We have revised the Figures and Tables to address these comments.

Competing Interests: n/a

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research