



Research paper

Gabapentin increases expression of δ subunit-containing GABA_A receptors



Jieying Yu^{a,1}, Dian-Shi Wang^{a,1}, Robert P. Bonin^{b,1}, Antonello Penna^{a,c}, Ali Alavian-Ghavanini^a, Agnieszka A. Zurek^a, Gail Rauw^d, Glen B. Baker^d, Beverley A. Orser^{a,e,f,*}

^a Department of Physiology, University of Toronto, Toronto, ON M5S 1A8, Canada

^b Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON M5S 3M2, Canada

^c Department of Anesthesia and Centro de Investigación Clínica Avanzada, Universidad de Chile, Santiago, 838 0456, Chile

^d Neurochemical Research Unit, Department of Psychiatry, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, AB T6G 2G3, Canada

^e Department of Anesthesia, University of Toronto, Toronto, ON M5G 1E2, Canada

^f Department of Anesthesia, Sunnybrook Health Sciences Centre, Toronto, ON M4N 3M5, Canada

ARTICLE INFO

Article history:

Received 10 December 2018

Received in revised form 22 February 2019

Accepted 4 March 2019

Available online 14 March 2019

Keywords:

Gabapentin
 GABA_A receptor
 Tonic inhibition
 Cerebellum
 Hippocampus

ABSTRACT

Background: Gabapentin is a structural analog of the inhibitory neurotransmitter γ -aminobutyric acid (GABA). Its anticonvulsant, analgesic and anxiolytic properties suggest that it increases GABAergic inhibition; however, the molecular basis for these effects is unknown as gabapentin does not directly modify GABA type A (GABA_A) receptor function, nor does it modify synaptic inhibition. Here, we postulated that gabapentin increases expression of δ subunit-containing GABA_A (δ GABA_A) receptors that generate a tonic inhibitory conductance in multiple brain regions including the cerebellum and hippocampus.

Methods: Cell-surface biotinylation, Western blotting, electrophysiologic recordings, behavioral assays, high-performance liquid chromatography and gas chromatography-mass spectrometry studies were performed using mouse models.

Findings: Gabapentin enhanced expression of δ GABA_A receptors and increased a tonic inhibitory conductance in neurons. This increased expression likely contributes to GABAergic effects as gabapentin caused ataxia and anxiolysis in wild-type mice but not δ subunit null-mutant mice. In contrast, the antinociceptive properties of gabapentin were observed in both genotypes. Levels of GABA_A receptor agonists and neurosteroids in the brain were not altered by gabapentin.

Interpretation: These results provide compelling evidence to account for the GABAergic properties of gabapentin. Since reduced expression of δ GABA_A receptor occurs in several disorders, gabapentin may have much broader therapeutic applications than is currently recognized.

Fund: Supported by a Foundation Grant (FDN-154312) from the Canadian Institutes of Health Research (to B.A.O.); a NSERC Discovery Grant (RGPIN-2016-05538), a Canada Research Chair in Sensory Plasticity and Reconsolidation, and funding from the University of Toronto Centre for the Study of Pain (to R.P.B.).

© 2019 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Gabapentin (1-(aminomethyl)cyclohexaneacetic acid) was approved by the US Food and Drug Administration, initially for the treatment of seizures in 1993 and subsequently for the treatment of postherpetic neuralgia in 2002 [1]. Over the past two decades, the indications for gabapentin have expanded widely and it is now one of the

most commonly prescribed medications in the United States [1]. Gabapentin is primarily used “off-label” for the treatment of acute and neuropathic pain [2]. Also, due to its anxiolytic and sedative effects, gabapentin is widely prescribed for sleep disturbances, bipolar disorder and restless leg syndrome [3]. Despite its widespread use, gabapentin causes serious adverse effects including drowsiness, ataxia, mental confusion and respiratory depression [3,4]. Given its increasing use and multiple adverse effects, it is important to understand gabapentin’s molecular mechanisms of action [1,4,5].

The behavioral properties of gabapentin have been attributed to inhibition of the auxiliary $\alpha_2\delta$ subunit of voltage-dependent calcium channels, which reduces calcium channel function and the release of

* Corresponding author at: Department of Physiology, University of Toronto, Room 3318, Medical Sciences Building, 1 King’s College Circle, Toronto, Ontario M5S 1A8, Canada.

E-mail address: beverley.orser@utoronto.ca (B.A. Orser).

¹ J.Y., D.S.W., and R.P.B. contributed equally to this work.

Research in context

Evidence before this study

While the behavioral properties of gabapentin suggest that it increases GABAergic inhibition, gabapentin is not a positive allosteric modulator of GABA_A receptors, nor does it increase inhibitory synaptic transmission. Thus, the basis of gabapentin's GABAergic effects is unknown. Here, we tested the hypothesis that gabapentin increases GABAergic inhibition by stimulating the expression of δ subunit-containing GABA_A (δ GABA_A) receptors in neurons.

Added value of this study

Gabapentin robustly increases cell-surface expression of δ GABA_A receptors and increases a tonic inhibitory conductance in neurons. This enhanced δ GABA_A receptor function contributes to the ataxic and anxiolytic but not antinociceptive properties of gabapentin. Gabapentin does not increase levels of GABA_A receptor agonists or several neurosteroids in the brain.

Implications of all the available evidence

These findings first identify δ GABA_A receptors as targets of gabapentin and challenge the widely-held notion that inhibition of $\alpha_2\delta$ subunits of voltage-dependent calcium channels is gabapentin's primary mechanism of action. Also, since dysregulation of δ GABA_A receptors contributes to several cognitive and psychiatric disorders, the therapeutic applications of gabapentin may be broader than currently recognized.

excitatory neurotransmitters from presynaptic terminals [6,7]. Interestingly, many of the behavioral properties of gabapentin suggest that it increases the function of inhibitory γ -aminobutyric acid (GABA) type A (GABA_A) receptors. Similar to many drugs that are positive allosteric modulators of GABA_A receptors including benzodiazepines, ethanol and many general anesthetic drugs, gabapentin causes sedation, somnolence, hypnosis, memory deficits, ataxia and respiratory depression [3,4]. Gabapentin is a structural analog of the inhibitory neurotransmitter GABA, yet it has no direct effects on GABA_A receptor function, nor does it increase inhibitory synaptic transmission [1,8]. Thus, the molecular basis of gabapentin's GABAergic properties has remained enigmatic.

GABA_A receptors are pentameric transmitter-gated ion channels formed from 19 different subunits [9,10]. The combinations of subunits produce receptor subtypes with different pharmacological and physiological properties. We hypothesized that gabapentin selectively modifies a subtype of GABA_A receptors through mechanisms that have not been identified.

GABA_A receptors are classified into two major groups: synaptic and extrasynaptic where synaptic receptors generate transient inhibitory postsynaptic current and extrasynaptic receptors generate a tonic inhibitory current that regulates neuronal excitability and network plasticity [9,10]. Extrasynaptic GABA_A receptors usually contain either a δ subunit or $\alpha 5$ subunits in combination with $\beta 2$ or $\beta 3$ subunits [9–11].

Results from our previous study of hippocampal neurons suggested that gabapentin targets extrasynaptic GABA_A receptors [8]. In that earlier study, treating cultured hippocampal neurons with gabapentin for 36 to 48 h increased a tonic inhibitory current but did not change the amplitude, frequency or time course of miniature inhibitory

postsynaptic currents. These results were unexpected as the tonic inhibitory current in hippocampal neurons is primarily generated by $\alpha 5$ subunit-containing GABA_A receptors and gabapentin did not up-regulate the function or expression of these receptors. In that report, it was assumed that the increase in GABA_A receptor activity was caused by increasing levels of endogenous agonists, as suggested by others [7,12,13].

In the current study, we consider an alternative novel hypothesis. We postulate that gabapentin increases the cell-surface expression of extrasynaptic δ subunit-containing GABA_A (δ GABA_A) receptors in neurons. We focus on δ GABA_A receptors because these receptors are the most widely expressed subtype of extrasynaptic GABA_A receptors in the mammalian brain [9,10]. These receptors are normally expressed at low levels in cultured hippocampal neurons. However, the localization of GABA_A receptors on the cell surface can be dynamically regulated by complex trafficking mechanisms that can be pharmacologically modified [14]. The net movement of GABA_A receptors to and from the plasma membrane determines the number of receptors that accumulate on the cell surface, and these processes could be altered by gabapentin [15].

Our results first demonstrate that gabapentin increases expression of δ GABA_A receptors, an effect that would account for at least some of the drug's GABAergic properties. Given that reduced expression of δ GABA_A receptors contributes to a variety of cognitive and psychiatric disorders [16], these results have broad implications.

2. Methods

2.1. Experimental animals

Experiments were approved by the Animal Care Committee of the University of Toronto. All studies were performed in accordance with guidelines from the Canadian Council on Animal Care. GABA_A receptor δ subunit null-mutant (*Gabrd* $-/-$) mice and wild-type (WT) mice (C57BL/6J \times 129Sv/SvJ) aged 2.5 to 4.5 months old were used. These mice were bred from a stock that was generously provided by Dr. Gregg E. Homanics (University of Pittsburgh, Pittsburgh, Pennsylvania, USA). The generation, genotyping, and characterization of *Gabrd* $-/-$ mice have been previously described [17]. In addition, adult Swiss Webster mice (3–4 months old, Charles River, Montreal, Quebec, Canada) were used for measuring the levels of agonists and neurosteroids that modify GABA_A receptors. Only male mice were selected for the studies because the estrous cycle alters the expression of extrasynaptic δ GABA_A receptors [18,19]. Mice were housed in a pathogen-free facility at the University of Toronto (25 \pm 1 °C) on a 14-h light/10-h dark cycle (lights on 6:00 AM to 8:00 PM). Experimenters were blinded to genotypes and treatment conditions.

2.2. Preparation of brain slices

After live decapitation, the brain was removed from 10- to 14-week-old WT mice and placed in ice-cold, oxygenated (95% O₂, 5% CO₂) artificial cerebrospinal fluid (ACSF) that contained (in mM): 124 NaCl, 3 KCl, 1.3 MgCl₂, 2.6 CaCl₂, 1.25 NaH₂PO₄, 26 NaHCO₃ and 10 D-glucose with solution osmolarity adjusted to 300–310 mOsm. Coronal brain slices (350 μ m) were prepared with a VT1200S vibratome (Leica, Deerfield, Illinois, USA). Cerebellar and hippocampal slices were then dissected out in ice-cold oxygenated ACSF. For *ex vivo* experiments, the slices were collected 2 h post-treatment and transferred to small chambers that were filled with ice-cold oxygenated ACSF. Slices were equilibrated for 1 h before undergoing cell-surface biotinylation and Western blotting procedures. For *in vitro* experiments, individual cerebellar or hippocampal slices were equilibrated at room temperature (23–25 °C) for 1 h. Slices were then treated with gabapentin (300 μ M) that was dissolved in oxygenated ACSF or vehicle control at 35–37 °C. This concentration of gabapentin increases the tonic current after a prolonged treatment of cultured hippocampal neurons [8].

2.3. Cell-surface biotinylation and western blotting

Slices were placed on ice and incubated twice for 30 min with 0.75 mg/ml NHS-SS-biotin (Thermo Scientific, Rockford, Illinois, USA) that was dissolved in DPBS (Gibco, Burlington, Ontario, Canada). Excess biotin was quenched and removed by washing slices 6 times with ice-cold modified TBS solution containing (in mM): 25 Tris-Cl, 137 NaCl, 1 KCl, 2.3 CaCl₂, pH 7.4. Slices were then placed in lysis buffer (pH 7.4) containing complete protease inhibitor cocktail (Roche, Laval, Quebec, Canada) for homogenization. Insoluble material was removed by centrifugation. Bicinchoninic acid assay (Bio-Rad, Hercules, California, USA) was performed to determine the protein concentration. Supernatant lysates were incubated with Hi-Capacity NeutrAvidin beads (Thermo Scientific, Rockford, Illinois, USA) for 16–18 h at 4 °C. The beads were washed with PBS containing 0.05% SDS. Bound material was eluted with elution buffer containing (in mM): 50 Tris-Cl, 2% SDS, 2 DTT; protein concentration was determined using DC™ Protein Assay (Bio-Rad, Hercules, California, USA) and subjected to SDS-PAGE analysis. Western blot analyses with antibodies for δ (Millipore, Billerica, Massachusetts, USA), $\alpha 1$ (Abcam, Cambridge, Massachusetts or Millipore, Billerica, Massachusetts, USA), and $\alpha 5$ (PhosphoSolutions, Aurora, Colorado, USA) GABA_A receptor subunits were performed. Anti- β -actin antibody (Millipore, Billerica, Massachusetts, USA) and anti-Na⁺/K⁺ ATPase antibody (Developmental Studies Hybridoma Bank, Iowa City, Iowa, USA) were also used. Blots were imaged using the Chemidoc XRS+ system (Bio-Rad, Hercules, California, USA) and quantified using Image Lab software (Bio-Rad, Hercules, California, USA). All receptor bands were normalized to the loading control, Na⁺/K⁺ ATPase. Blots containing surface protein were probed for β -actin to determine the purity of the isolated biotinylated surface protein. Samples for surface proteins were analyzed using two replicate blots to control for transfer and loading errors. The two normalized values were averaged to obtain a single value for each sample. Data for each sample were presented as a percentage of the mean of the control sample.

2.4. Whole-cell voltage clamp recordings from cerebellar slices

After live decapitation, the cerebellum was quickly removed from 10- to 14-week-old mice and submerged in oxygenated (95% O₂, 5% CO₂) ice-cold cutting solution, which contained (in mM): 235 sucrose, 2.5 KCl, 1.25 NaH₂PO₄, 28 NaHCO₃, 0.5 CaCl₂, 7 MgSO₄, 28 D-glucose (pH 7.4; 305–315 mOsm). The tissue was maintained in ice-cold solution while sagittal slices of cerebellar vermis (300 μ m) were cut using a VT1200S vibratome. The slices were transferred to oxygenated ACSF and allowed to recover for at least 1 h at room temperature before being transferred to a submersion recording chamber, where they were perfused with ACSF at 3–4 ml/min.

Experiments were performed on an Olympus BX51 upright microscope equipped with differential interference contrast/infrared optics. Conventional whole-cell voltage clamp recordings were carried out at room temperature from granule neurons located in the granule cell layer, which were visually identified. 6-Cyano-7-nitroquinoxaline-2,3-dione (CNQX, 10 μ M) and (2R)-amino-5-phosphonovaleric acid (APV, 20 μ M) were added to the extracellular solution to block ionotropic glutamate receptors and tetrodotoxin (0.3 μ M) was used to block voltage-dependent sodium channels. Patch pipettes (4–6 M Ω) were fabricated from borosilicate glass capillary tubing and filled with an intracellular solution containing (in mM): 140 CsCl, 10 HEPES, 11 EGTA, 4 MgATP, 2 MgCl₂, 1 CaCl₂ and 10 TEA (pH 7.3 with CsOH, 285–295 mOsm). Whole-cell currents were recorded using the MultiClamp 700B amplifier (Molecular Devices, Sunnyvale, California, USA) that was controlled with pClamp 9.0 software via a Digidata 1322 interface (Molecular Devices). Membrane capacitance was measured using the membrane test protocol in pClamp 9.0. Currents were sampled at 10 kHz and filtered at 2 kHz with an eight-pole low-pass Bessel filter. All cells were recorded at a holding potential of –60 mV. To measure tonic current, the

GABA_A receptor competitive antagonist bicuculline (10 μ M) was applied. Tonic current was quantified by measuring the change in the holding current calculated from a Gaussian fitting of 30-s segments.

2.5. Rotarod assay

The rotarod assay is commonly used to assess motor coordination in rodents. The rotarod apparatus (Econometex, Columbus Instruments, Columbus, Ohio, USA) consists of an elevated rotating rod. Mice placed on the rod must actively move in a coordinated manner to prevent themselves from falling. Mice were placed in the experimental room 1 h prior to training or testing in order to acclimatize to the environment. Prior to the testing day, the mice were trained on the rotarod for 4 times a day with 15-min intervals between trials, until they could stay on the rod consistently for a minimum of 300 s. The rotation speed of the rod was set to 12 rotations per minute. Training took approximately 2 weeks. For testing, each mouse was placed on the rotarod and the time spent on the rotating rod was recorded. The maximally allowed time on the rod was 300 s, and each mouse was given 4 trials with 15-min intervals. The time spent on the rotarod (latency to fall) from the 4 trials was averaged.

2.6. Elevated plus maze assay

The elevated plus maze assay is used to test anxiety-like behavior in rodents [20]. The maze contains four arms (14 in. \times 2 in.) that were made of transparent plexiglass. Two arms were open and two were enclosed (6-in.-high walls). The arms were joined by a middle square (2 in. \times 2 in.). The entire maze was positioned 50 cm above the floor. Mice generally consider the dark or enclosed space as “safe” and tend to stay in these areas when anxious. Exploration of the open areas was deemed to reflect low levels of anxiety. Each mouse was placed in the center square facing an open arm, in the opposite position to where the experimenter was located and was allowed to explore the maze for 5 min. The time spent and number of entries in open and closed arms were manually scored in real-time using custom written scripts in Microsoft Excel.

2.7. Tail flick assay

The tail flick assay is used to measure acute nociception in rodents. Acute thermal nociception was assessed using a radiant heat tail flick apparatus (Columbus Instruments, Columbus, Ohio, USA). Each mouse was gently restrained and the midpoint of the tail was placed over the apparatus. The light intensity was fixed to produce an average tail-flick latency of 3 to 4 s in control mice. The latency to flick the tail away from the activated light source (latency to tail flick) was automatically measured to the nearest 1/100th of a second using an integrated light beam detector. A maximal cutoff time to respond was set at 10 s to prevent tissue damage.

2.8. Formalin nociception assay

The formalin assay was used to assess nociception-related behaviors in freely moving mice [21]. Briefly, mice were acclimatized to the testing chamber for 20 min before receiving an injection of formalin. The chamber consisted of a round, acrylic cylinder (15 cm diameter and 30 cm tall). The cylinder was suspended above an angled mirror to facilitate observation from all vantage points. Formalin (5%, 20 μ l) was injected subcutaneously into the plantar surface of the right hind paw, and the time spent licking the injected hind paw was recorded. The first two minutes immediately after the formalin injection were not scored. Subsequently, the mice were observed for 2-min intervals every 4 min for the next 44 min. Two mice were tested concurrently and staggered 2 min apart. Nociceptive hind paw licking was classified as occurring in 1 of

2 phases: Phase 1 from 0 to 6 min and Phase 2 from 6 to 46 min after injection of formalin.

2.9. Measurement of GABA_A receptor agonist levels with high-performance liquid chromatography

Adult Swiss Webster mice (3–4 months old, Charles River, Montreal, Quebec, Canada) were treated with gabapentin (30 mg/kg, I.P.) or normal saline (0.9% w/v of NaCl, I.P.) every 12 h for 4 consecutive days. Two hours after the final treatment, the whole brain and spinal cord extracts were collected for measurements of GABA, taurine and alanine concentrations using high-performance liquid chromatography (HPLC), as previously described [22,23]. The mice were anesthetized with isoflurane using a bell jar apparatus and decapitated. The whole brain was removed and flash frozen in isopentane that was cooled with a slurry of dry ice and ethanol. The spinal column was cut at the end of the thoracic and mid-sacral segments, then was extracted by hydraulic extrusion using ice-cold saline and snap frozen in an Eppendorf tube using isopentane that was cooled with a slurry of dry ice and ethanol. The frozen tissues were homogenized in 80% (v/v) methanol at 0 °C. An aliquot of the homogenate was diluted a further 5× in ice-cold methanol and left on ice for 10 min prior to centrifugation at 12,000 g for 4 min at 4 °C. The supernatant was diluted ten-fold with ice-cold deionized water and transferred to a HPLC vial.

Derivatizing reagent (5 µl) containing 2-mercaptoethanol and o-phthalaldehyde was mixed with the 5 µl of sample or standard, held in the instrument sample loop for 1.5 min, and injected into an Alliance 2690XE HPLC system (Waters, Mississauga, Ontario, Canada). The system employed a Symmetry C18 column (4.5 × 100 mm, 5 µm) that was coupled to a guard column and a Waters 474 programmable fluorescence detector (Waters, Mississauga, Ontario, Canada). All the data were acquired and processed using Empower Pro software (Waters, Mississauga, Ontario, Canada).

A mobile phase gradient consisting of varying amounts of sodium phosphate buffer and methanol was used to separate the amino acids. The samples were maintained at 4 °C and the column compartments were maintained at 30 °C. The excitation and emission wavelengths were 260 nm and 455 nm, respectively.

2.10. Measurement of neurosteroid levels with gas chromatography-mass spectrometry

Alphaxalone was added to the residual 80% methanol homogenate as an internal standard. The mixture was vortexed and left on ice for 10 min prior to centrifugation at 12,000 g for 5 min at 4 °C. Solid-phase extractions on the supernatants were performed using an Oasis HLB 30 mg extraction plate (Waters, Mississauga, Ontario, Canada). The wells were washed with 5% (v/v) methanol and the neurosteroids were eluted using a 9:1 (v/v) methylene dichloride:methanol solution. The eluents were dried and the analytes were derivatized for 1 h at 45 °C with 10 µl of n-heptafluorobutyrylimidazole in 50 µl of ethyl acetate. The samples were dried and reconstituted in 50 µl of toluene, washed with 50 µl of deionized water, and centrifuged at 12,000 g for 2 min. The toluene layer was analyzed with gas chromatography using an Agilent 6890 gas chromatograph (Mississauga, ON, Canada) with a HP-MS5 5% phenylmethylsiloxane column (30 cm × 250 µm, 0.25 µm film thickness) coupled to an Agilent 5973 mass spectrometer (Mississauga, Ontario, Canada) in the negative ion-chemical ionization mode. The inlet temperature was 250 °C. The oven temperature was initialized at 100 °C, increased to 295 °C at a rate of 8 °C/min, and maintained at 295 °C for 5 min.

The mass spectrometer was operated in the single ion monitoring mode and standard curves were run in parallel with each assay. The capture and analysis of data were performed using Agilent MSD Chemstation software (Mississauga, Ontario, Canada). Tissue concentrations of

neurosteroids that were below the limit of detection in a sample were assigned a value of 1 pg steroid/g tissue.

2.11. Drugs and chemicals

Gabapentin, CNQX, APV and bicuculline were purchased from Abcam (Cambridge, United Kingdom). Tetrodotoxin was purchased from Alomone Labs (Jerusalem, Israel), and formalin from Sigma Aldrich (St. Louis, Missouri, USA).

2.12. Statistical analysis

Data are presented as mean ± SEM. An unpaired Student *t*-test was used to compare two groups. The two-way analysis of variance (ANOVA, treatment × genotype) followed by the Tukey's multiple comparisons post hoc test was used to compare more than two groups. All analyses were performed using GraphPad Prism 6.01 (Graphpad Software Inc., La Jolla, California, USA). A two-tailed hypothesis test was used, and statistical significance was set at $P < 0.05$. No outliers were excluded from the statistical analyses.

2.13. Data availability

Data supporting the findings of this manuscript are available from the corresponding author upon request.

3. Results

3.1. Gabapentin increases δ GABA_A receptor expression in the cerebellum

To determine whether gabapentin modifies the expression of δ GABA_A receptors, mice were treated with gabapentin (100 mg/kg, I.P.) or vehicle (0.9% w/v of NaCl, I.P.) and 2 h later, *ex vivo* brain slices were harvested and processed for biotinylation and Western blotting (Fig. 1a). Studies were first performed with tissue from the cerebellum as this brain region has the highest expression levels of δ GABA_A receptors and cerebellar dysfunction causes ataxia [16].

Gabapentin increased cell-surface expression of δ subunits to 136.3% ± 5.4% of control ($n = 6$ per condition, $P = 0.0005$; Fig. 1b) whereas total expression levels were unchanged (109.1% ± 4.4% of control, $n = 6$ per condition, $P = 0.16$). We also studied the expression levels of $\alpha 1$ subunits as these subunits primarily contribute to synaptic GABA_A receptors. Gabapentin did not modify cell-surface expression or total expression levels of $\alpha 1$ subunits, suggesting that it selectively modifies expression of δ GABA_A receptors (Fig. 1c).

Gabapentin could increase cell-surface expression of δ GABA_A receptors by exerting either a direct effect on the brain or by stimulating systemic changes, such as an increase in levels of circulating neurosteroids or other paracrine factors that enter the brain [18,24]. To determine whether gabapentin up-regulates the expression of δ GABA_A receptors, through mechanisms that are independent of systemic actions, slices of cerebellum harvested from drug-naïve mice were directly incubated with gabapentin (300 µM) for 6 h (Fig. 1d) [25]. Gabapentin increased both cell-surface expression (167.7% ± 3.6% of control, $n = 3$ each, $P = 0.0003$) and total expression levels (179.2% ± 5.9% of control, $n = 3$ each, $P = 0.0002$) of δ subunits (Fig. 1e). Thus, treatment of the brain slices recapitulated the effect of administering gabapentin systemically. The expression of $\alpha 1$ subunits was unchanged in cerebellar slices treated under similar experimental conditions (Supplementary Fig. 1).

3.2. Gabapentin increases a tonic inhibitory current in cerebellar granule neurons

We next asked whether increased cell-surface expression of δ subunits was sufficient to increase the amplitude of the tonic current in cerebellar neurons. Whole-cell currents were recorded from cerebellar




Fig. 1. The expression of δ subunits in the cerebellum is increased by gabapentin. (a) A schematic diagram shows the experimental timeline. (b, c) Representative Western blots and summarized data of cell-surface and total expression of δ subunits (b) and $\alpha 1$ subunits (c) in the cerebellum. b: $n = 6$ for both surface and total, c: $n = 6, 5$ for surface and total, respectively. (d) A schematic diagram shows the experimental timeline. (e) Representative Western blots and summarized data of the cell-surface and total expression of δ subunits after the cerebellar slices were treated with gabapentin. $n = 3$ for each condition. *** $P < 0.001$, unpaired Student t -test. Data are Mean \pm SEM. Con = control, GBP = gabapentin, NKA = Na/K-ATPase.

granule neurons with patch clamp technique, as previously described [26]. Cerebellar slices were harvested from mice that lacked the δ GABA_A receptors (*Gabrd*^{-/-} mice) and wild-type (WT) mice. The mice were treated with gabapentin (100 mg/kg, I.P.) then sacrificed 2 h later (Fig. 2a). To measure the amplitude of the tonic current, the competitive GABA_A receptor antagonist bicuculline (10 μ M) was applied to the *ex vivo* brain slices. Current amplitude was quantified by measuring the change in the holding current before and after the application of bicuculline [27,28].

Gabapentin increased the tonic current in WT granule neurons to 153.9% of control (Gabapentin: 8.0 ± 0.8 pA/pF, $n = 11$ vs. Control: 5.2 ± 0.6 pA/pF, $n = 12$, $P = 0.02$; Fig. 2b & c) but had no effect on the tonic current in *Gabrd*^{-/-} neurons (Gabapentin: 2.3 ± 0.6 pA/pF,

$n = 8$ vs. Control: 2.2 ± 0.5 pA/pF, $n = 7$, $P = 1.0$; Fig. 2b & c). We observed that the baseline tonic current was lower in *Gabrd*^{-/-} neurons than in WT neurons (*Gabrd*^{-/-}: 2.2 ± 0.5 pA/pF, $n = 8$ vs. WT: 5.2 ± 0.6 pA/pF, $n = 12$, $P = 0.02$), as previously reported by others [29]. These results provide evidence that gabapentin increases a tonic inhibitory current in cerebellar granule neurons that is primarily generated by δ GABA_A receptors [16].

3.3. Gabapentin increases δ GABA_A receptor cell-surface expression in the hippocampus

Clinical and preclinical studies have shown that gabapentin causes memory loss, sedation and anxiolysis [3,4]. These diverse effects suggest




Fig. 2. Tonic inhibitory current in cerebellar granule neurons is enhanced by gabapentin. (a) A schematic diagram shows the experimental timeline. (b, c) Representative traces (b) and summarized data (c) for the tonic current recorded in cerebellar granule cells from WT and *Gabrd*^{-/-} slices. WT: n = 12, 11 for control and gabapentin, respectively. *Gabrd*^{-/-}: n = 8, 7 for control and gabapentin, respectively. Two-way ANOVA, effect of gabapentin treatment: $F_{(1,34)} = 4.3$, $P = 0.046$; effect of genotype: $F_{(1,34)} = 39.5$, $P < 0.0001$; effect of interaction: $F_{(1,34)} = 3.8$, $P = 0.06$; * $P < 0.05$, Tukey's multiple comparisons test. Data are Mean \pm SEM. BIC = bicuculline, Con = control, GBP = gabapentin.

that gabapentin modifies δ GABA_A receptors in regions of the central nervous system other than cerebellum. Thus, we next studied whether gabapentin-induced increased expression of δ GABA_A receptors generalizes to another brain region.

δ GABA_A receptors are highly expressed in the dentate gyrus [9,16]. Indeed, autoradiographic studies have demonstrated high levels of gabapentin binding in this subregion of the hippocampus [9,30]. To determine whether *in vivo* treatment with gabapentin increased expression levels of δ GABA_A receptors in the hippocampus, mice were injected with gabapentin (100 mg/kg, I.P.) and 2 h later, *ex vivo* hippocampal slices were harvested and processed for biotinylation analyses (Fig. 3a). Cell-surface expression of δ subunit was increased ($122.4\% \pm 3.8\%$ of control, n = 6 each, $P = 0.0013$; Fig. 3b) whereas the total expression levels were unchanged ($101.5\% \pm 0.6\%$ of control, n = 6 each, $P = 0.7$). Gabapentin did not alter expression levels of $\alpha 1$ subunits (Fig. 3c).

Since extrasynaptic $\alpha 5$ GABA_A receptors are highly expressed in the hippocampus [11], we next studied whether gabapentin increased the expression levels of this receptor subtype. Gabapentin did not modify the expression levels of $\alpha 5$ subunits (Fig. 3d),

consistent with our previous findings from studies of cultured hippocampal neurons [8].

Finally, we harvested hippocampal slices from drug-naïve WT mice and treated them directly with gabapentin (300 μ M) for 6 h (Supplementary Fig. 2a). Interestingly, cell-surface expression ($132.6\% \pm 3.3\%$ of control, n = 4 each, $P = 0.0002$) but not the total expression level of δ subunits ($106.9\% \pm 6.7\%$ of control, n = 4 each, $P = 0.4$) was increased (Supplementary Fig. 2b). Despite similar experimental conditions, the latter finding was inconsistent with those in cerebellar slices where total expression levels were increased. In other studies of hippocampal slices, direct application of gabapentin did not change the expression levels of $\alpha 1$ and $\alpha 5$ subunits (Supplementary Fig. 2c&d). Thus, the selective increase in expression of δ GABA_A receptors is not restricted to the cerebellum.

3.4. Gabapentin causes ataxia and reduces anxiety in WT mice but not *Gabrd*^{-/-} mice

We next asked whether increased expression of δ GABA_A receptors contributes to some of the behavioral properties of gabapentin. Behaviors including ataxia, anxiety and nociception, which are mediated by different regions of the central nervous system, were studied in WT and *Gabrd*^{-/-} mice (Fig. 4a). Ataxia, a lack of voluntary coordination of muscle movement that can result from cerebellum dysfunction [31], is a common adverse effect of gabapentin. Ataxia is frequently studied using the rotarod assay [31]. Mice were treated with gabapentin or vehicle and 2 h later, placed on a rotating rod. The latency to fall off the rotarod indicated voluntary motor coordination. Under baseline conditions, performance on the rotarod was similar in WT and *Gabrd*^{-/-} mice (Fig. 4b). Gabapentin (30–100 mg/kg) impaired the performance of WT mice, as evidenced by a decrease in latency time (Fig. 4b). In contrast, the same doses had no significant effects on motor performance in *Gabrd*^{-/-} mice. While there was a slight reduction at the highest dose of gabapentin in *Gabrd*^{-/-} mice, the change was not significant.

Gabapentin is often prescribed for its anxiolytic and sedative properties. Indeed, in a clinical study we showed that gabapentin reduces anxiety scores in highly anxious patients prior to surgery [32]. To probe the mechanisms of gabapentin-induced anxiolysis, WT and *Gabrd*^{-/-} mice were studied with the elevated plus maze assay. The time spent in the open arms of the maze provided a surrogate measure of anxiety caused by open spaces. Mice were treated with gabapentin (30 mg/kg, I.P.), and 2 h later were placed on the elevated plus maze. Baseline performance was similar between genotypes ($P = 0.8$, Fig. 4c). Gabapentin-treated WT mice exhibited an 11-fold increase in the time spent in the open arms (Gabapentin: $11.3 \pm 3.2\%$, n = 10 vs. Control: $1.0 \pm 0.5\%$, n = 9, $P = 0.007$; Fig. 4c). In contrast, gabapentin did not increase the open arm time of *Gabrd*^{-/-} mice (Gabapentin: $3.2 \pm 1.1\%$, n = 11 vs. Control: $3.7 \pm 2.0\%$, n = 11, $P = 1.0$). Similarly, the gabapentin-treated WT mice, but not the gabapentin-treated *Gabrd*^{-/-} mice, showed a significant increase in the number of entries into the open arms (Supplementary Fig. 3a). In both genotypes, gabapentin treatment did not alter the time spent in the closed arms, or the number of entries into the closed arms (Supplementary Fig. 3b). Thus, while baseline anxiety was not influenced by the expression of δ GABA_A receptors, δ GABA_A receptors were necessary for the anxiolytic effects of gabapentin.

3.5. Gabapentin modulation of nociception in WT mice and *Gabrd*^{-/-} mice

We and others have shown that δ GABA_A receptors regulate nociception [33,34]. We next asked whether gabapentin modulates nociception in WT and *Gabrd*^{-/-} mice. Tail flick and formalin assays were used to study acute and persistent nociception, respectively. The tail flick assay assesses nociceptive threshold by measuring the latency

Ex vivo hippocampal slices




Fig. 3. The cell-surface expression of δ subunits in hippocampus is increased by gabapentin. (a) A schematic diagram shows the experimental timeline. (b, c, d) Representative Western blots and summarized data of the cell-surface and total expression of the δ (b), $\alpha 1$ (c), and $\alpha 5$ (d) subunits in the hippocampus. b: $n = 6$ for both surface and total; c: $n = 3, 5$ for surface and total, respectively; d: $n = 6, 4$ for surface and total, respectively. ** $P < 0.01$, unpaired Student t -test. Data are Mean \pm SEM. Con = control, GBP = gabapentin, NKA = Na/K-ATPase.

time to tail flick in response to a thermal stimulus. The formalin assay involves both acute and tonic phases. The acute phase (Phase 1) occurs within the first 6 min after formalin injection and responses are thought to reflect the direct activation of nociceptors by the noxious chemical. The tonic phase (Phase 2) is measured 6 min to 46 min after the formalin injection. The licking behavior exhibited by the mice is thought to result from central sensitization and inflammation at the level of the spinal dorsal horn [33].

Under baseline conditions, the latency to tail flick was similar in the two genotypes (Fig. 4d). Gabapentin (60 mg/kg, I.P.), administered at a dose that was anxiolytic in WT mice, caused no change in the tail flick response in either group (Fig. 4d). For Phase 1 of the formalin test, under baseline conditions, no differences were observed between genotypes for the licking time. Gabapentin (60 mg/kg, I.P.) did not alter the licking time in either genotype (Fig. 4e). For Phase 2, the baseline response of *Gabrd*^{-/-} mice was greater than WT mice (WT control: 99.0 ± 9.4 s, $n = 10$ vs. *Gabrd*^{-/-} control: 135.1 ± 5.4 s, $n = 8$, $P = 0.006$; Fig. 4e). Gabapentin reduced the Phase 2 response of both WT and *Gabrd*^{-/-} mice and abolished the differences between the two genotypes (Fig. 4e). These results suggest that gabapentin attenuation of chronic or inflammatory nociception is independent of the expression of δ GABA_A receptors.

3.6. Longer-term administration of gabapentin increases expression of δ GABA_A receptors in the cerebellum

In the studies described above, the mice were treated with a single dose of gabapentin. However, patients are typically treated with gabapentin for days to months [4]. Such long-term exposure to other GABAergic drugs including benzodiazepines and barbiturates can cause tachyphylaxis or drug resistance due to internalization of GABA_A receptors from the cell surface [35]. Thus, we next studied whether a longer exposure to gabapentin would trigger a sustained increase in the cell-surface expression of δ GABA_A receptors.

Mice were treated with gabapentin (100 mg/kg/day, I.P.) or vehicle for 6 consecutive days. Two hours after the final injection, *ex vivo* cerebellar slices were harvested and processed for biotinylation and Western blotting (Supplementary Fig. 4a). Gabapentin increased cell-surface expression ($144.4 \pm 17.4\%$ of control, $n = 4$ each, $P = 0.046$; Supplementary Fig. 4b). The total expression was also increased but the change did not reach statistical significance ($149.3 \pm 23.9\%$ of control, $n = 4$ each, $P = 0.1$; Supplementary Fig. 4b). Thus, similar to acute gabapentin treatment, longer-term treatment increased expression of δ GABA_A receptors.




Fig. 4. δ GABA_A receptors contribute to the ataxic and anxiolytic, but not the analgesic, properties of gabapentin. (a) A schematic diagram shows the experimental timeline. (b) Gabapentin dose-dependently reduces the latency to fall in WT but not in *Gabrd*^{-/-} mice. $n = 6$ for each group in WT mice and $n = 5$ for each group in *Gabrd*^{-/-} mice. Two-way ANOVA, effect of gabapentin treatment: $F_{(3,36)} = 22.7$, $P < 0.0001$; effect of genotype: $F_{(1,36)} = 29.0$, $P < 0.0001$; effect of interaction: $F_{(3,36)} = 5.9$, $P = 0.002$. $***P < 0.001$ compared with control, Tukey's multiple comparisons test. (c) Gabapentin increases the time that WT but not *Gabrd*^{-/-} mice spent in the open arms. $n = 9$, 10 for control and gabapentin, respectively in WT mice and $n = 11$ for each group in *Gabrd*^{-/-} mice. Two-way ANOVA, effect of gabapentin treatment: $F_{(1,37)} = 5.9$, $P = 0.02$; effect of genotype: $F_{(1,37)} = 1.7$, $P = 0.2$; effect of interaction: $F_{(1,37)} = 7.2$, $P = 0.01$. $**P < 0.01$, N.S.: no significance, Tukey's multiple comparisons test. (d) gabapentin does not change the latency to tail flick in either genotype. $n = 13$, 14 for control and gabapentin, respectively in WT mice and $n = 13$, 12 for control and gabapentin, respectively in *Gabrd*^{-/-} mice. Two-way ANOVA, effect of gabapentin treatment: $F_{(1,48)} = 1.3$, $P = 0.3$; effect of genotype: $F_{(1,48)} = 0.005$, $P = 0.9$; effect of interaction: $F_{(1,48)} = 0.16$, $P = 0.7$. (e) Gabapentin does not alter licking time in either genotype for Phase 1 but reduced licking time for both genotypes for Phase 2. $n = 10$ for each treatment group in WT mice and $n = 8$ for each treatment group in *Gabrd*^{-/-} mice. Phase 1: Two-way ANOVA, effect of gabapentin treatment: $F_{(1,32)} = 4.1$, $P < 0.05$; effect of genotype: $F_{(1,32)} = 4.8$, $P = 0.04$; effect of interaction: $F_{(1,32)} = 0.12$, $P = 0.73$. Phase 2: Two-way ANOVA, effect of gabapentin treatment: $F_{(1,32)} = 114.4$, $P < 0.0001$; effect of genotype: $F_{(1,32)} = 12.9$, $P = 0.001$; effect of interaction: $F_{(1,32)} = 2.0$, $p = 0.16$. $**P < 0.01$, $***P < 0.001$, Tukey's multiple comparisons test. Data are Mean \pm SEM. Con = control, GBP = gabapentin.

3.7. Gabapentin does not increase the levels of GABA_A receptor agonists in the brain or spinal cord

Several endogenous agonists including the primary neurotransmitter GABA, as well as taurine and alanine, activate GABA_A receptors. δ GABA_A receptors have a high affinity for GABA compared to other subtypes of native GABA_A receptors [36]. Furthermore, taurine is a potent agonist of extrasynaptic GABA_A receptors [37]. Alanine, which is also present at relatively high concentrations in the brain [38–40], evokes bicuculline-sensitive Cl⁻ currents in hippocampal neurons (unpublished results). Thus, gabapentin could stimulate small increases in levels of GABA, taurine and alanine in the brain or spinal cord and thereby increase the function of high-affinity δ GABA_A receptors. Such small changes in agonist levels could

contribute to the gabapentin's behavioral properties, as suggested by previous studies [7,12,13].

We next measured the levels of GABA, taurine and alanine in the brain or spinal cord of gabapentin- and vehicle-treated mice. A 4-day treatment paradigm was used for these studies to reduce the risk of failing to detect any transient or modest changes in neurotransmitter levels. Mice were treated with gabapentin (30 mg/kg, I.P.) or vehicle every 12 h for 4 consecutive days (Fig. 5a). Two hours after the final treatment, the whole brain, and the lumbar and upper sacral regions of the spinal cord were extracted. Levels of GABA, taurine and alanine were measured using high-performance liquid chromatography. The results showed that gabapentin failed to modify the levels of GABA, taurine and alanine (Fig. 5b), suggesting that gabapentin does not increase agonist concentrations after chronic treatment.




Fig. 5. Levels of GABA_A receptor agonists are not increased by gabapentin. (a) A schematic diagram shows the experimental timeline. (b) Gabapentin does not change levels of GABA_A receptor agonists in the brain and spinal cord. $n = 8$ and 6 for control and gabapentin, respectively; $P > 0.1$ for each agonist, unpaired Student t -test. Data are mean \pm SEM. Con = control, GBP = gabapentin.

An increase in the levels of neurosteroids could also account for gabapentin-induced increase in the function of δ GABA_A receptors, as these receptors are highly sensitive to certain neurosteroids [41,42]. We next measured the levels of the neurosteroids 5-dehydroepiandrosterone (DHEA), $3\alpha,5\alpha$ -THPROG, $3\alpha,5\beta$ -THPROG, pregnenolone, $3\beta,5\alpha$ -THPROG, and THDOC in the brain after 4 days of same treatment with gabapentin. Gabapentin did not alter the levels of these neurosteroids in the brain (Supplementary Fig. 5).

4. Discussion

The results show that gabapentin increases expression of δ GABA_A receptors in the cerebellum and hippocampus. This enhanced expression of δ GABA_A receptors is associated with an increase in the tonic current in cerebellar neurons. Behavioral studies show that δ GABA_A receptors are necessary for the ataxia and anxiolysis but not anti-nociceptive properties of gabapentin. Collectively, these results provide the first evidence of a clinically-approved drug that rapidly increases cell-surface expression levels of δ GABA_A receptors within a period of hours. The findings challenge the widely accepted notion that modulation of $\alpha_2\delta$ subunits of voltage-dependent calcium channels alone accounts for many of the behavioral effects of gabapentin.

Increased expression of δ GABA_A receptors on the cell surface is due to either enhanced trafficking of δ GABA_A receptors and/or stabilization of receptors at the plasma membrane. The increase in total levels of δ GABA_A receptors suggests *de novo* synthesis of new receptors [43].

Thus, more than one mechanism could account for the increase in receptor expression. Interestingly, whether gabapentin increases cell-surface as well as total expression of δ GABA_A receptors depends on the brain region and experimental paradigm. As an example, treatment *in vivo* with gabapentin increases cell-surface expression but not total expression levels in the cerebellum, whereas direct application of gabapentin to cerebellar slices *in vitro* increases both cell-surface and total expression. The factors that account for such differences in cell-surface and total expression of δ GABA_A receptors are uncertain. Little is known about regulation of the expression and trafficking of δ GABA_A receptors. The signaling pathways, transcription factors, kinases and phosphatases that regulate the gene encoding the δ subunits (*Gabrd*) could differ between brain regions. Interestingly, the trafficking and expression of synaptic GABA_A receptors are differently modified by protein kinase A, protein kinase C, p38 mitogen-activated protein kinases and phosphatases [15,43]. Additional studies are needed to determine how gabapentin modifies cell-surface and total expression of δ GABA_A receptors.

Several lines of evidence support the notion that increased activity of δ GABA_A receptors contributes to gabapentin-induced ataxia and anxiolysis. Modulators that increase the function of δ GABA_A receptors, including increased endogenous hormones in the estrus cycle, exogenous neurosteroids and ethanol all cause ataxia [44] and anxiolysis [18,24]. Similarly, the δ GABA_A receptor-selective super-agonist 4,5,6,7-tetrahydroisoxazolo(5,4-c) pyridin-3-ol (gaboxadol or THIP) causes ataxia and anxiolysis. Gaboxadol is considered to be a “super-agonist” as it generates a current that is larger than current evoked by a saturating concentration of GABA [36]. In both humans and laboratory animals, gaboxadol, similar to gabapentin, produces anxiolysis and impairs motor coordination [45–49].

The results show that δ GABA_A receptors do not contribute to gabapentin's antinociceptive properties. Gabapentin does not modify responses in the tail flick assay or Phase 1 of the formalin assay. In a previous study, we showed that gaboxadol attenuates responses in Phase 1 of the formalin test in WT but not *Gabrd*^{−/−} mice [33]. These differences in behavioral properties may be due to differences in the efficacy of gaboxadol versus gabapentin. Gaboxadol, which acts as an allosteric modulator and super-agonist, may cause a greater increase in δ GABA_A receptor function than gabapentin, which is not a positive allosteric modulator but rather increases cell-surface expression. In addition, under baseline conditions, expression levels of δ GABA_A receptors are low in dorsal root ganglion neurons and the spinal cord [16]. Thus, the gabapentin-induced increase in cell-surface expression may not be sufficient to modify antinociception. We also showed that gabapentin reduces responses in Phase 2 of the formalin test in both WT and *Gabrd*^{−/−} mice. This effect does not depend on δ GABA_A receptors but may involve other receptor targets, including $\alpha_2\delta$ subunit of voltage-dependent calcium channels, $\alpha_2\delta$ -NMDA receptor complexes, hyperpolarization-activated cation channels or sodium channels [6,50–52].

Gabapentin caused ataxia in WT but not *Gabrd*^{−/−} mice raising the concern that ataxia potentially confounded the effects of gabapentin on anxiety and nociception. However, the low dose of gabapentin (30 mg/kg, I.P.) that was selected for elevated plus maze studies did not impair performance on the rotarod. Also, gabapentin increased the number of entries into the open arms. Ataxia would likely have produced the opposite effect and reduced the number of entries. Finally, while gabapentin produced ataxia only in WT mice, it had similar antinociceptive effects in WT and *Gabrd*^{−/−} mice. These results are consistent with previous findings showing that deficits in motor coordination do not impair primary sensory function [53]. Overall, it is unlikely that gabapentin-induced ataxia confounded the responses in the elevated plus maze and formalin assays.

Gabapentin does not increase levels of GABA, alanine, taurine or neurosteroids in the brain or spinal cord. Similar results of GABA levels have been reported by others [54–56]. However, in other studies

[7,12,13], gabapentin increases GABA levels in both rodents and humans, either by increasing GABA synthesis via the activation of GABA-synthesizing enzyme glutamic acid decarboxylase or by decreasing GABA metabolism via the inhibition of GABA transaminase. Even small increases in agonist levels could enhance activity of high-affinity δ GABA_A receptors. One limitation of our study is that agonist and neurosteroid levels were measured in the whole brain and spinal cord. This approach may not be sufficiently sensitive to detect small changes in discrete, yet functionally important pools of agonists or neurosteroids. Indeed, gabapentin increases GABA turnover in some, but not all, brain regions [57].

Our results raise several questions that are worthy of future study. It would be of interest to investigate whether δ subunit binding partners including the $\alpha 4$ and $\alpha 6$ subunits are also upregulated by gabapentin. In cerebellum, the predominant form of δ GABA_A receptors is $\alpha 6\beta 2/3\delta$, with low expression levels of $\alpha 4\beta 2/3\delta$. In the hippocampus and thalamus, $\alpha 4\beta 2/3\delta$ GABA_A receptors are predominantly expressed [9,10,16]. $\beta 2$ or $\beta 3$ subunits can also partner with subunits other than the δ subunit, to form either synaptic ($\alpha 1\beta 2/3\gamma$) or extrasynaptic ($\alpha 5\beta 3\gamma$) GABA_A receptors [9–11]. Interestingly, the effects of gabapentin are selective for the δ subunit as expression of both $\alpha 1$ subunit and $\alpha 5$ subunit was unchanged, through mechanisms that are unknown.

Finally, at a broader translational level, one of the most serious and potentially life-threatening adverse effects of gabapentin is respiratory depression [58–60]. The association of the use of gabapentin and a 60% increase in narcotic-related death is of great concern [59]. In a collaborative preclinical study, we showed that increased δ GABA_A receptor function reduces ventilation and upper airway muscle tone [61]. Thus, it will be of interest to determine whether respiratory depression due to coadministration of gabapentin and a narcotic is mediated by increased expression of δ GABA_A receptors.

Overall, the results identify increased expression of δ GABA_A receptors as a novel mechanism to account for gabapentin's GABAergic properties. Rapid upregulation of δ GABA_A receptors may be helpful in treating disorders associated with reduced expression of δ GABA_A receptors including alcoholism, fragile X syndrome and neuropsychiatric disorders including depression and schizophrenia [16].

Conflict of interest

The authors declare no conflict of interest.

Author contributions

J.Y., R.P.B., G.B.B., and B.A.O. designed research; J.Y., R.P.B., A.P., A.A.G., A.A.Z., and G.R. performed research; J.Y., D.S.W., R.P.B., A.P., and G.R. analyzed data; D.S.W. and B.A.O. wrote the manuscript.

Acknowledgments

This work was supported by a Foundation Grant (FDN-154312) from the Canadian Institutes of Health Research (to B.A.O.); a NSERC Discovery Grant (RGPIN-2016-05538), a Canada Research Chair in Sensory Plasticity and Reconsolidation and funding from the University of Toronto Centre for the Study of Pain (to R.P.B.). The authors thank Mr. Shahin Khodaei and Mr. Marc A. Manzo for their help during the writing of the manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ebiom.2019.03.008>.

References

- Wallach JD, Ross JS. Gabapentin approvals, off-label use, and lessons for postmarketing evaluation efforts. *JAMA* 2018;319(8):776–8.
- Wiffen PJ, Derry S, Bell RF, Rice AS, Tolle TR, Phillips T, et al. Gabapentin for chronic neuropathic pain in adults. *Cochrane Database Syst Rev* 2017(6):CD007938.
- Quintero GC. Review about gabapentin misuse, interactions, contraindications and side effects. *J Exp Pharmacol* 2017;9:13–21.
- Smith RV, Havens JR, Walsh SL. Gabapentin misuse, abuse and diversion: a systematic review. *Addiction* 2016;111(7):1160–74.
- Goodman CW, Brett AS. Gabapentin and pregabalin for pain - is increased prescribing a cause for concern? *N Engl J Med* 2017;377(5):411–4.
- Dolphin AC. Voltage-gated calcium channels and their auxiliary subunits: physiology and pathophysiology and pharmacology. *J Physiol* 2016;594(19):5369–90.
- Taylor CP, Gee NS, Su TZ, Kocsis JD, Welty DF, Brown JP, et al. A summary of mechanistic hypotheses of gabapentin pharmacology. *Epilepsy Res* 1998;29(3):233–49.
- Cheng VY, Bonin RP, Chiu MW, Newell JG, MacDonald JF, Orser BA. Gabapentin increases a tonic inhibitory conductance in hippocampal pyramidal neurons. *Anesthesiology* 2006;105(2):325–33.
- Brickley SG, Mody I. Extrasynaptic GABA_A receptors: their function in the CNS and implications for disease. *Neuron* 2012;73(1):23–34.
- Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of GABA_A receptors. *Nat Rev Neurosci* 2005;6(3):215–29.
- Caraiscos VB, Elliott EM, You-Ten KE, Cheng VY, Belelli D, Newell JG, et al. Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by $\alpha 5$ subunit-containing γ -aminobutyric acid type A receptors. *Proc Natl Acad Sci U S A* 2004;101(10):3662–7.
- Errante LD, Williamson A, Spencer DD, Petroff OA. Gabapentin and vigabatrin increase GABA in the human neocortical slice. *Epilepsy Res* 2002;49(3):203–10.
- Petroff OA, Hyder F, Rothman DL, Mattson RH. Effects of gabapentin on brain GABA, homocarnosine, and pyrrolidinone in epilepsy patients. *Epilepsia* 2000;41(6):675–80.
- Luscher B, Fuchs T, Kilpatrick CL. GABA_A receptor trafficking-mediated plasticity of inhibitory synapses. *Neuron* 2011;70(3):385–409.
- Uusi-Oukari M, Korpi ER. Regulation of GABA_A receptor subunit expression by pharmacological agents. *Pharmacol Rev* 2010;62(1):97–135.
- Whissell PD, Lecker I, Wang DS, Yu J, Orser BA. Altered expression of δ GABA_A receptors in health and disease. *Neuropharmacology* 2015;88:24–35.
- Mihalek RM, Banerjee PK, Korpi ER, Quinlan JJ, Firestone LL, Mi ZP, et al. Attenuated sensitivity to neuroactive steroids in γ -aminobutyrate type A receptor δ subunit knockout mice. *Proc Natl Acad Sci U S A* 1999;96(22):12905–10.
- Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA_A receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nat Neurosci* 2005;8(6):797–804.
- Maguire J, Ferando I, Simonsen C, Mody I. Excitability changes related to GABA_A receptor plasticity during pregnancy. *J Neurosci* 2009;29(30):9592–601.
- Hogg S. A review of the validity and variability of the elevated plus-maze as an animal model of anxiety. *Pharmacol Biochem Behav* 1996;54(1):21–30.
- Tjolsen A, Berge OG, Hunskaar S, Rosland JH, Hole K. The formalin test: an evaluation of the method. *Pain* 1992;51(1):5–17.
- Parent M, Bush D, Rauw G, Master S, Vaccarino F, Baker G. Analysis of amino acids and catecholamines, 5-hydroxytryptamine and their metabolites in brain areas in the rat using *in vivo* microdialysis. *Methods* 2001;23(1):11–20.
- Sloley BD, Kah O, Trudeau VL, Dulka JG, Peter RE. Amino acid neurotransmitters and dopamine in brain and pituitary of the goldfish: involvement in the regulation of gonadotropin secretion. *J Neurochem* 1992;58(6):2254–62.
- Shen H, Gong QH, Yuan M, Smith SS. Short-term steroid treatment increases δ GABA_A receptor subunit expression in rat CA1 hippocampus: pharmacological and behavioral effects. *Neuropharmacology* 2005;49(5):573–86.
- Cho S, Wood A, Bowlby MR. Brain slices as models for neurodegenerative disease and screening platforms to identify novel therapeutics. *Curr Neuropharmacol* 2007;5(1):19–33.
- Accardi MV, Brown PM, Miraucourt LS, Orser BA, Bowie D. $\alpha 6$ -Containing GABA_A receptors are the principal mediators of inhibitory synapse strengthening by insulin in cerebellar granule cells. *J Neurosci* 2015;35(26):9676–88.
- Wang DS, Zurek AA, Lecker I, Yu J, Abramian AM, Avramescu S, et al. Memory deficits induced by inflammation are regulated by $\alpha 5$ -subunit-containing GABA_A receptors. *Cell Rep* 2012;2(3):488–96.
- Zurek AA, Yu J, Wang DS, Haffey SC, Bridgwater EM, Penna A, et al. Sustained increase in $\alpha 5$ GABA_A receptor function impairs memory after anesthesia. *J Clin Invest* 2014;124(12):5437–41.
- Glykys J, Mann EO, Mody I. Which GABA_A receptor subunits are necessary for tonic inhibition in the hippocampus? *J Neurosci* 2008;28(6):1421–6.
- Hill DR, Suman-Chauhan N, Woodruff GN. Localization of [³H]gabapentin to a novel site in rat brain: autoradiographic studies. *Eur J Pharmacol* 1993;244(3):303–9.
- Schmahmann JD. Disorders of the cerebellum: ataxia, dysmetria of thought, and the cerebellar cognitive affective syndrome. *J Neuropsychiatry Clin Neurosci* 2004;16(3):367–78.
- Clarke H, Kirkham KR, Orser BA, Katznelson R, Mitsakakis N, Ko R, et al. Gabapentin reduces preoperative anxiety and pain catastrophizing in highly anxious patients prior to major surgery: a blinded randomized placebo-controlled trial. *Can J Anaesth* 2013;60(5):432–43.
- Bonin RP, Labrakakis C, Eng DG, Whissell PD, De Koninck Y, Orser BA. Pharmacological enhancement of δ -subunit-containing GABA_A receptors that generate a tonic inhibitory conductance in spinal neurons attenuates acute nociception in mice. *Pain* 2011;152(6):1317–26.

- [34] Iura A, Takahashi A, Hakata S, Mashimo T, Fujino Y. Reductions in tonic GABAergic current in substantia gelatinosa neurons and GABA_A receptor δ subunit expression after chronic constriction injury of the sciatic nerve in mice. *Eur J Pain* 2016;20(10):1678–88.
- [35] Barnes Jr EM. Use-dependent regulation of GABA_A receptors. *Int Rev Neurobiol* 1996;39:53–76.
- [36] Brown N, Kerby J, Bonnert TP, Whiting PJ, Wafford KA. Pharmacological characterization of a novel cell line expressing human $\alpha_4\beta_3\delta$ GABA_A receptors. *Br J Pharmacol* 2002;136(7):965–74.
- [37] Jia F, Yue M, Chandra D, Keramidis A, Goldstein PA, Homanics GE, et al. Taurine is a potent activator of extrasynaptic GABA_A receptors in the thalamus. *J Neurosci* 2008;28(1):106–15.
- [38] Golden GT, Ferraro TN, Fariello RG, Hare TA. Amino acid profiles in long-Evans rat superior colliculus, visual cortex, and inferior colliculus. *Neurochem Res* 1989;14(5):465–72.
- [39] Honda T. Amino acid metabolism in the brain with convulsive disorders. Part I: free amino acid patterns in the brain of E1 mouse with convulsive seizure. *Brain Dev* 1984;6(1):17–21.
- [40] Perry TL, Hansen S, Berry K, Mok C, Lesk D. Free amino acids and related compounds in biopsies of human brain. *J Neurochem* 1971;18(3):521–8.
- [41] Abramian AM, Comenencia-Ortiz E, Modgil A, Vien TN, Nakamura Y, Moore YE, et al. Neurosteroids promote phosphorylation and membrane insertion of extrasynaptic GABA_A receptors. *Proc Natl Acad Sci U S A* 2014;111(19):7132–7.
- [42] Belelli D, Lambert JJ. Neurosteroids: endogenous regulators of the GABA_A receptor. *Nat Rev Neurosci* 2005;6(7):565–75.
- [43] Jacob TC, Moss SJ, Jurd R. GABA_A receptor trafficking and its role in the dynamic modulation of neuronal inhibition. *Nat Rev Neurosci* 2008;9(5):331–43.
- [44] Hancher HJ, Dodson PD, Olsen RW, Otis TS, Wallner M. Alcohol-induced motor impairment caused by increased extrasynaptic GABA_A receptor activity. *Nat Neurosci* 2005;8(3):339–45.
- [45] Agmo A, Soria P. GABAergic drugs and sexual motivation, receptivity and exploratory behaviors in the female rat. *Psychopharmacology (Berl)* 1997;129(4):372–81.
- [46] Hoehn-Saric R. Effects of THIP on chronic anxiety. *Psychopharmacology (Berl)* 1983;80(4):338–41.
- [47] Leufkens TR, Lund JS, Vermeeren A. Highway driving performance and cognitive functioning the morning after bedtime and middle-of-the-night use of gaboxadol, zopiclone and zolpidem. *J Sleep Res* 2009;18(4):387–96.
- [48] Saarelainen KS, Ranna M, Rabe H, Sinkkonen ST, Moykkynen T, Uusi-Oukari M, et al. Enhanced behavioral sensitivity to the competitive GABA agonist, gaboxadol, in transgenic mice over-expressing hippocampal extrasynaptic $\alpha_6\beta$ GABA_A receptors. *J Neurochem* 2008;105(2):338–50.
- [49] Voss J, Sanchez C, Michelsen S, Ebert B. Rotarod studies in the rat of the GABA_A receptor agonist gaboxadol: lack of ethanol potentiation and benzodiazepine cross-tolerance. *Eur J Pharmacol* 2003;482(1–3):215–22.
- [50] Chen J, Li L, Chen SR, Chen H, Xie JD, Sirrieh RE, et al. The $\alpha_2\delta$ -1-NMDA receptor complex is critically involved in neuropathic pain development and gabapentin therapeutic actions. *Cell Rep* 2018;22(9):2307–21.
- [51] Tae HS, Smith KM, Phillips AM, Boyle KA, Li M, Forster IC, et al. Gabapentin modulates HCN4 channel voltage-dependence. *Front Pharmacol* 2017;8:554.
- [52] Yang RH, Wang WT, Chen JY, Xie RG, Hu SJ. Gabapentin selectively reduces persistent sodium current in injured type-A dorsal root ganglion neurons. *Pain* 2009;143(1–2):48–55.
- [53] Bastian AJ. Moving, sensing and learning with cerebellar damage. *Curr Opin Neurobiol* 2011;21(4):596–601.
- [54] Errante LD, Petroff OA. Acute effects of gabapentin and pregabalin on rat forebrain cellular GABA, glutamate, and glutamine concentrations. *Seizure* 2003;12(5):300–6.
- [55] Leach JP, Sills GJ, Butler E, Forrest G, Thompson GG, Brodie MJ. Neurochemical actions of gabapentin in mouse brain. *Epilepsy Res* 1997;27(3):175–80.
- [56] Meyerhoff DJ, Murray DE, Durazzo TC, Pennington DL. Brain GABA and glutamate concentrations following chronic gabapentin administration: a convenience sample studied during early abstinence from alcohol. *Front Psych* 2018;9:78.
- [57] Loscher W, Honack D, Taylor CP. Gabapentin increases aminoxyacetic acid-induced GABA accumulation in several regions of rat brain. *Neurosci Lett* 1991;128(2):150–4.
- [58] Cavalcante AN, Sprung J, Schroeder DR, Weingarten TN. Multimodal analgesic therapy with gabapentin and its association with postoperative respiratory depression. *Anesth Analg* 2017;125(1):141–6.
- [59] Gomes T, Juurlink DN, Antoniou T, Mamdani MM, Paterson JM, van den Brink W. Gabapentin, opioids, and the risk of opioid-related death: A population-based nested case-control study. *PLoS Med* 2017;14(10):e1002396.
- [60] Keskinbora K, Pekel AF, Aydinli I. Gabapentin and an opioid combination versus opioid alone for the management of neuropathic cancer pain: a randomized open trial. *J Pain Symptom Manage* 2007;34(2):183–9.
- [61] Montandon G, Wu H, Liu H, Vu MT, Orser BA, Horner RL. δ -Subunit containing GABA_A receptors modulate respiratory networks. *Sci Rep* 2017;7(1):18105.