

MEETING ABSTRACT

Open Access

Recovery from general anaesthesia in the elderly patient: sevoflurane vs propofol

B Bastianini, L Vescicchio, F Grassia, B Lettieri*

From XXIII Annual Meeting of the Italian Society of Geriatric Surgery
Lecce, Italy. 2-4 December 2010

Background

Pharmacokinetics of anaesthetic drugs is greatly influenced by physiological modifications due to advanced age:

- Slower drug metabolism
- Decrease in TBW
- Increase in LBM/FBM
- Decrease in kidney function

This leads to the need to use short half-life drugs, to prevent drug stacking phenomena. Both sevoflurane and propofol meet these requirements.

Materials and methods

This study compares recovery times from general anaesthesia between the selected drugs in patients over 65yo who underwent surgery.

In 3 months 41 patients were selected, similar in age (65-78yo), sex and ASA class, undergoing elective surgery. Patients with lung, liver, kidney, brain and coronary dysfunctions with ASA III-IV were excluded from the study.

In this single-blind, prospective study patients were randomized in 2 groups: patients in P-group were treated with propofol (5-8 mg/kg/h), while patients in S-group were treated with sevoflurane (1-2%).

Dosage was managed so that CF and AP wouldn't shift more than 20% their basal values. All patients were pre-treated with midazolam (0.03mg/kg), fentanyl (1 μ g/kg) and atropine (up to 0.01ng/kg). Anaesthesia was induced using propofol (2mg/kg) and cisatracurium (0.2mg/kg). Patients were intubated and ventilated using O₂ (33%) and N₂O (66%). Analgesia during surgery was achieved using fentanyl (1 μ g/kg) 60min after induction and then using boli every 45min. Parameters monitored

Table 1 Prospective, randomized study, patients selectedPatients total 41

S	P
21 pz. (9M/12F)	20 pz. (10M/10F)

for each patient were: AP, CF, ECG, PO₂, ET CO₂, BIS and diuresis. Times for extubation, recovery room monitoring, eye opening and oriented motory and verbal response were recorded. MMSE was also taken.

Results

41 Patients were selected (table 1), age 65-78, ASA II; groups were homogeneous in type and duration of surgery (120±67min). Induction and intra-surgery dosages of drugs were similar between patients, as well as 2-hours post-operation pain-score. Extubation was quicker in S-group (6.8±4min) than in P-group (9.8±5min). BIS system score has shown quicker recover from sleep in S-group (table 2). Simple commands execution (eye opening, hand movement) and time-space orientation recovery (saying one's own name and birthday) were quicker in the S-group as well (table 3). Discharge times from recovery room were similar between groups, as were results in MMSE.

Table 2 recovery after sevoflurane or propofol suspension

	S	P
Extubation (min)	6.8±4	9.8±5
Eye Opening	5.9±2	9.3±2
Punch (orientated motory response)	10.2±2	14±4
Orientated verbal response	12±8	18±5
MMSE	24±4	24±4
Observation time (min)	39±9	40±21
Pain score (2 hours after surgery)	3±2	3±3

* Correspondence: biagio.lettieri@unina2.it

Department of Anaesthesia, Surgical and Emergency Science, Second University of Naples, Italy

Table 3 monitored parameters

AP	S	P
Pre-operative (mmHg)	94±4	94±2
Induction (mmHg)	116±5	100±8
Incision (mmHg)	80±6	109±9.5
Maintenance (mmHg)	83±2	74±7
Recovery (mmHg)	105±5	107±11

CF	S	P
Pre-operative (b/min)	72±7	72±7
Intubation (b/min)	100±10	87±6
Incision (b/min)	81±9	88±11
Maintenance (b/min)	78±5	67±13
Recovery (b/min)	88±7	90±8

Conclusions

Sevoflurane shows the lowest periféric blood solubility and the lowest partition ratio amongst inhaled anaesthetics. This grants both quick induction and recovery. These properties associated with better hemodynamic stability during maintenance of general anaesthesia make this drug favorable for the elderly patient, who often has deficitary cardiac homeostatic mechanisms, both due to advanced age and to other associated diseases.

Published: 24 August 2011

References

1. Mathur S, Farhangkhgoee P, Karmazyn M: Cardioprotective effects of propofol and sevoflurane in ischemic and reperfused rat hearts: Role of K(ATP) channels and interaction with the sodium-hydrogen exchange inhibitor HOE 642 (cariporide). *Anesthesiology* 1999, 91:1349-60.
2. Ghatge S, Lee J, Smith I: Sevoflurane: an ideal agent for adult day-case anaesthesia? *Acta Anaesthesiol Scand* 2003, 47(8):917-31.
3. Yildirim H, Adanir T, Atay A, Katircioğlu K, Savaci S: The effects of sevoflurane, isoflurane and desflurane. *Eur J Anaesthesiol* 2004, 21:566-70.
4. Mourisse J, Lerou J, Struys M, Zwarts M, Booij L: Multi-level approach of anaesthetic effects caused by sevoflurane or propofol in humans: 1. BIS and blink reflex. *Br J Anaesth* 2007, 98:737-45.
5. Ghatge S, Lee J, Smith I: Propofol injection pain: is it still an issue? The effect of premedication. *Minerva Anestesiol* 2010, 76(9):720-4.

doi:10.1186/1471-2318-11-S1-A3

Cite this article as: Bastianini et al.: Recovery from general anaesthesia in the elderly patient: sevoflurane vs propofol. *BMC Geriatrics* 2011 11 (Suppl 1):A3.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

