

POSTER PRESENTATION

Open Access

Expression of auxin carrier genes during adventitious rooting in *Eucalyptus globulus*

Arthur Fett-Neto*, Marcia De Almeida, Carolina Ruedell

From IUFRO Tree Biotechnology Conference 2011: From Genomes to Integration and Delivery
Arraial d'Ajuda, Bahia, Brazil. 26 June - 2 July 2011

Background

Eucalyptus globulus and its hybrids are important for the cellulose and paper industry, mainly due to their relatively low lignin content. However, rooting of cuttings of this species is often recalcitrant and exogenous auxin application is necessary for adventitious root development. Auxin plays a central role in rooting capacity, which is particularly affected by its endogenous content and transport rate. The shoot apex is a major source of endogenous auxin, which is mainly transported by both influx (*AUX1*) and efflux (*PIN*) carriers in a specific basipetal active transport through the vascular parenchyma in stems. As part of a larger study to investigate the causes of low rooting in *E. globulus* microcuttings without exogenous auxin, we evaluated the expression profiles of *AUX1* and *PIN1* during the process of adventitious rooting using qPCR.

Material and methods

E. globulus *in vitro* tip microcuttings obtained from 14 week-old seedlings were submitted to a culture system consisting of a two-step protocol: an initial step of induction, which lasted 96 h (induction medium composition: 0.3x MS salt concentration, 0.4 mg l⁻¹ thiamine HCl, 100 mg l⁻¹ inositol, zero (control) or 10 mg l⁻¹ indolyl-butyric acid (IBA – root promoting auxin), equivalent to 49.3 µM, 30 g l⁻¹ sucrose and 6 g l⁻¹ agar, followed by a formation step (same composition of induction medium, but devoid of auxin and supplemented with 1 g l⁻¹ activated charcoal). The expression analysis of the selected genes was monitored along the rooting process and the harvest of microcuttings for RNA extraction was done at 6, 12, 24, 48 and 96 h of exposure to induction medium and 24 and 48 h after transfer to formation medium (formation

step), for both treatments (with and without auxin in the first step). For the formation step harvest, the microcuttings remained for 96 h in the induction medium before transfer to formation medium. Total RNA was extracted, and the first strand cDNA synthesis was performed for all of the samples starting from about 500 ng total RNA, using oligo-dT primers and reverse transcriptase M-MLV (Invitrogen) in a final volume of 20 µl. The final cDNA products were diluted 50-fold in RNase-free distilled water prior to use in qPCR. The analysis was carried out using specific primers for *Arabidopsis thaliana* orthologue genes in eucalypt and both *Histone H2B* (*H2B*) and *Alpha-Tubulin* (*TUA*) genes were used as references [1]. The data were analysed with the comparative Ct method [2].

Results and conclusion

The gene encoding the auxin influx carrier (*AUX1*) did not show differences in expression profile between treatments (with and without exogenous auxin), suggesting that *AUX1* is not critical to the process of adventitious rooting promoted by exogenous auxins in microcuttings. This would be in line with the fact that the rate of endogenous auxin transport is probably not limiting under exogenous auxin supply. The auxin efflux carrier gene encoding *PIN1* showed an expression increase during the first 24 hours of the induction step in microcuttings exposed to exogenous auxin when compared with the control treatment (without exogenous auxin supply). This result seems to indicate a requirement of *PIN1* to redistribute and perhaps concentrate auxin, possibly IBA-derived IAA (indolyl-3-acetic acid), in specific areas of the base of microcuttings, in order to allow root development. Although IBA is a natural auxin recognized as an IAA precursor, its use as exogenous auxin instead of IAA, a common practice in clonal propagation by cuttings, may have also involved the

* Correspondence: fettneto@cbiot.ufrgs.br
Gabriela Pereira Center for Biotechnology, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil

expression of other sets of recently described IBA-specific transporters [3]. In conclusion, despite the unusual exogenous auxin entrance pathway into the microcutting compared to the endogenous auxin fluxes, *PIN1* likely takes part in the process of auxin concentration required to program founder cells involved in the establishment of new root meristems.

Acknowledgements

Funding for this investigation was provided by the Brazilian agencies National Council for Scientific and Technological Development (CNPq) and Rio Grande do Sul State Foundation for Research Support (Fapergs).

Published: 13 September 2011

References

1. De Almeida MR, Ruedell CM, Ricachenevsky FK, Sperotto RA, Pasquali G, Fett-Neto AG: Reference gene selection for quantitative reverse transcription-polymerase chain reaction normalization during *in vitro* adventitious rooting in *Eucalyptus globulus* Labill. *BMC Molecular Biology* 2010, 11:73.
2. Livak KJ, Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta CT$ method. *Methods* 2001, 25:402-408.
3. Strader LC, Bartel B: Transport and metabolism of the endogenous auxin precursor Indole-3-Butyric Acid. *Molecular Plant* 2011, 1-10.

doi:10.1186/1753-6561-5-S7-P64

Cite this article as: Fett-Neto et al.: Expression of auxin carrier genes during adventitious rooting in *Eucalyptus globulus*. *BMC Proceedings* 2011 5(Suppl 7):P64.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

