

SCIENTIFIC REPORTS

**OPEN**

Author Correction: Effect of BI-1 on insulin resistance through regulation of CYP2E1

Geum-Hwa Lee¹, Kyoung-Jin Oh^{2,3}, Hyung-Ryong Kim⁴, Hye-Sook Han², Hwa-Young Lee¹, Keun-Gyu Park⁵, Ki-Hoan Nam⁶, Seung-Hoi Koo² & Han-Jung Chae¹

Correction to: *Scientific Reports* <https://doi.org/10.1038/srep32229>, published online 31 August 2016

This Article contains an error in the labelling of the middle and lower panels of Supplementary Figure 11, which were incorrectly labelled as 'IP: IRE-1 α ' and 'IP: CPR' respectively. Both panels should be labelled as 'IP:HA'. In addition, the 'IRE-1 α ' and 'HA' labels in the middle panel and the 'CPR' and 'HA' labels in the lower panel were inverted. The correct Supplementary Figure 11 appears below as Figure 1.

¹Department of Pharmacology and New Drug Development Institute, Medical School, Chonbuk National University, Jeonju, 561-181, Republic of Korea. ²Division of Life Sciences, Korea University, 145 Anam-Ro, Seongbuk-Gu, Seoul, 136-713, Republic of Korea. ³Metabolic Regulation Research Center, Korea Research Institute of Bioscience and Biotechnology (KRIBB), Daejeon, 305-806, Republic of Korea. ⁴Department of Dental Pharmacology and Wonkwang Dental Research Institute, School of Dentistry, Wonkwang University, Iksan, 570-749, Republic of Korea. ⁵Department of Internal Medicine, Kyungpook National University School of Medicine, Daegu, 700-721, Republic of Korea. ⁶Laboratory Animal Resource Center, KRIBB, Ochang-eup, 363-883, Republic of Korea. Geum-Hwa Lee, Kyoung-Jin Oh and Hyung-Ryong Kim contributed equally. Correspondence and requests for materials should be addressed to S.-H.K. (email: koohei@korea.ac.kr) or H.-J.C. (email: hjchae@jbnu.ac.kr)





Figure 1. The interaction of BI-1 with CPR and IRE-1 α . Immunoprecipitation assay demonstrating the effects of BI-1 on palmitate-induced interactions with CPR or IRE-1 α . Neo and BI-1 cells were treated with 250 μ M palmitate for the indicated periods. Immunoprecipitation was performed with anti-HA, IRE-1 α , or CPR antibody. Western blotting was performed with anti-CPR, IRE-1 α or HA antibody. CPR, NADPH-dependent CYP reductase.

 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018