

Article

Phenoxyacetohydrazide Schiff Bases: β -Glucuronidase Inhibitors

Waqas Jamil ^{1,2}, Shagufta Perveen ¹, Syed Adnan Ali Shah ^{3,4}, Muhammad Taha ^{4,5},
Nor Hadiani Ismail ^{4,5}, Shahnaz Perveen ⁶, Nida Ambreen ¹, Khalid M. Khan ^{1,*} and
Muhammad I. Choudhary ¹

¹ H. E. J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Karachi 75270, Pakistan; E-Mails: waqas143kh@yahoo.com (W.J.); sarosh.iqbal@yahoo.com (S.P.); nida_mishi2002@yahoo.com (N.A.); hassaan2@super.net.pk (M.I.C.)

² Institute of Advance Research Studies in Chemical Sciences, University of Sindh Jamshoro, Hyderabad 76080, Pakistan

³ Faculty of Pharmacy, Universiti Teknologi MARA (UiTM), Puncak Alam Campus, Bandar Puncak Alam, Selangor Darul Ehsan 42300, Malaysia; E-Mail: benzene301@yahoo.com or syedadnan@salam.uitm.edu.my

⁴ Atta-ur-Rahman Institute for Natural Products Discovery (AuRIns), Level 9, FF3, Universiti Teknologi MARA (UiTM), Puncak Alam Campus, Bandar Puncak Alam, Selangor Darul Ehsan 42300, Malaysia; E-Mails: taha_hej@yahoo.com (M.T.); norhadiani@puncakalam.uitm.edu.my (N.H.I.)

⁵ Faculty of Applied Science UiTM, Shah Alam, Selangor 40450, Malaysia

⁶ PCSIR Laboratories Complex, Shahrah-e-Dr. Salimuzzaman, Karachi 75280, Pakistan; E-Mail: shamim_302@yahoo.com

* Author to whom correspondence should be addressed; E-Mail: khalid.khan@iccs.edu; Tel.: +92-213-482-4910; Fax: +92-213-481-9018.

Received: 7 May 2014; in revised form: 6 June 2014 / Accepted: 11 June 2014 /

Published: 25 June 2014

Abstract: Phenoxyacetohydrazide Schiff base analogs **1–28** have been synthesized and their *in vitro* β -glucouronidase inhibition potential studied. Compounds **1** ($IC_{50} = 9.20 \pm 0.32 \mu M$), **5** ($IC_{50} = 9.47 \pm 0.16 \mu M$), **7** ($IC_{50} = 14.7 \pm 0.19 \mu M$), **8** ($IC_{50} = 15.4 \pm 1.56 \mu M$), **11** ($IC_{50} = 19.6 \pm 0.62 \mu M$), **12** ($IC_{50} = 30.7 \pm 1.49 \mu M$), **15** ($IC_{50} = 12.0 \pm 0.16 \mu M$), **21** ($IC_{50} = 13.7 \pm 0.40 \mu M$) and **22** ($IC_{50} = 22.0 \pm 0.14 \mu M$) showed promising β -glucuronidase inhibition activity, better than the standard (D-saccharic acid-1,4-lactone, $IC_{50} = 48.4 \pm 1.25 \mu M$).

Keywords: phenoxyacetohydrazide; Schiff bases; β -glucouoronidase; glucuronosyl-*O*-bonds, D-saccharic acid-1,4-lactone

1. Introduction

A range of bioactivities are reported for hydrazide-hydrazone compounds, such as antibacterial, anticonvulsant, antimalarial, analgesic, antiinflammatory, antiplatelets, antifungal, antituberculosis, and anticancer activities [1–6]. A variety of semicarbazones, thiosemicarbazones and guanyl hydrazones are found to be key compounds for drug design [7], for metal complexes [8], organocatalysis [9], and are used for the preparation of heterocyclic rings [10]. A few pyrazole carbohydrazide hydrazone derivatives [11] and novel 3-aryl-1-arylmethyl-1*H*-pyrazole-5-carbohydrazide hydrazones were found to be proliferation inhibitors of A549 cells [12,13]. Some evidence proposes a pharmacophoric character for the hydrazone moiety present in phenylhydrazone derivatives in the inhibition of cyclooxygenase [14]. Antioxidant [15–18], antiglycation [19–22] and antileishmanial [23] activity have recently been reported, as well as applications in mass spectrometry [24].

The present work aimed to investigate the potential activity of a series of aryl hydrazide-hydrazones as *in vitro* β -glucouoronidase inhibitors. In our designed analogues substituted phenoxy-acetohydrazides were treated with different aromatic aldehydes to scrutinize their potential activity. The earlier reported literature [25] showed that β -glucouoronidase is a lysosomal enzyme, present in many organs like the spleen, kidney, lung, bile, serum and urine, *etc.*, where its specific task is to catalyze the cleavage of glucuronosyl-*O*-bonds [26–28]. It degrades glucuronic acid-containing glycosaminoglycans, like heparan sulfate, chondroitin sulfate and dermatan sulfate [29]. An elevated level of β -glucouoronidase was observed in various types of malignancies, such as breast, lung and gastrointestinal tract carcinomas, and melanomas. Its high expression also observed in bronchial tumors [30]. On the other hand, mucopolysaccharidosis type VII (MPS VII; Sly Syndrome) is caused by the deficiency of human β -glucuronidase [31]. The circulating level of β -glucuronidase is also useful as a lysosomal enzyme in children affected by leprosy. In borderline tuberculoid patients and lepromatous patients higher activity of this enzyme was also observed.

2. Results and Discussion

2.1. Chemistry

Lead identification is a well defined tool in drug design and discovery. Our research group has been involved for a decade in lead discovery programs in search of novel therapeutic agents. We have earlier reported Schiff bases of different classes of organic compounds in the search for lead molecules with different biological activities [32–34]. Earlier, our group reported the leishmanicidal and β -glucuronidase inhibition potential of hydrazides derived from the corresponding esters [35–38]. In view of the formerly reported work we synthesized hydrazide Schiff bases and screened their potential biological activities [39–42]. Acylhydrazide Schiff base derivatives **1–28** were synthesized from an acylhydrazide by condensing it with different aromatic aldehydes and acetophenones under reflux

conditions in ethanol for 2–3 h (Scheme 1). The crude products (Table 1) were further recrystallized from methanol and needle-like crystals were obtained in most of the cases. The starting acylhydrazide was synthesized from ester of ethyl 2-(4-chloro-2-methylphenoxy) acetate by refluxing with hydrazine hydrate.

Scheme 1. Synthetic scheme for benzohydrazide followed by synthesis of Schiff bases 1–28.
























Table 1. Synthesis of acylhydrazide Schiff base derivatives 1–28.

Compound No.	R ₁	R ₂	Yield (%)	Compound No.	R ₁	R ₂	Yield (%)
1		H	81	15		H	88
2		H	85	16		H	92
3		H	93	17		CH ₃	94
4		H	87	18		H	89

Table 1. Cont.

Compound No.	R ₁	R ₂	Yield (%)	Compound No.	R ₁	R ₂	Yield (%)
5		H	83	19		CH ₃	87
6		H	86	20		H	84
7		H	89	21		H	94
8		H	91	22		H	93
9		H	94	23		H	91
10		H	82	24		CH ₃	95
11		H	88	25		H	91
12		H	91	26		H	88
13		H	86	27		H	81
14		H	93	28		H	93

2.2. β -Glucuronidase

Synthetic acyl hydrazides Schiff bases **1–28** were screened for their *in vitro* potential as β -glucuronidase inhibitors. The *in vitro* β -glucuronidase inhibitory potential was evaluated by using the literature protocol [43]. Compounds **1–28** showed diversified β -glucuronidase inhibitory activities, with IC_{50} values ranging between 9.20–30.7 μ M. Compounds **1, 5, 7, 8, 11, 12, 15, 21, and 22** showed excellent β -glucuronidase inhibitory activities, with IC_{50} values of 9.20 ± 0.32 , 9.47 ± 0.16 , 14.7 ± 0.19 , 15.4 ± 1.56 , 19.6 ± 0.62 , 30.7 ± 1.49 , 12.0 ± 0.16 , 13.7 ± 0.40 , and 22.0 ± 0.14 μ M, respectively, and the remaining compounds exhibited no activity (Table 2).


Table 2. *In vitro* β -glucuronidase activity of compounds **1–28**.

Compounds	IC_{50} (μ M \pm SEM ^a)	Compounds	IC_{50} (μ M \pm SEM ^a)
1	9.20 ± 0.32	15	12.0 ± 0.16
2	NA ^b	16	NA ^b
3	NA ^b	17	NA ^b
4	NA ^b	18	NA ^b
5	9.47 ± 0.16	19	NA ^b
6	NA ^b	20	NA ^b
7	14.7 ± 0.19	21	13.7 ± 0.40
8	15.4 ± 1.56	22	22.0 ± 0.14
9	NA ^b	23	NA ^b
10	NA ^b	24	NA ^b
11	19.6 ± 0.62	25	NA ^b
12	30.7 ± 1.49	26	NA ^b
13	NA ^b	27	NA ^b
14	NA ^b	28	NA ^b
D-saccharic acid-1,4-lactone ^c	48.4 ± 1.25	-	-

SEM ^a is the standard error of the mean; NA ^b Not active; ^c standard inhibitor for β -glucuronidase.

It was observed that both the substituents' nature and their position at the benzilidene part have great importance in the β -glucuronidase inhibition activity of a compound, and apparently the acylium part does not take part in the activity (Figure 1).

Figure 1. The two parts of molecule on which activity is based.



The best activity was shown by compound **1** ($IC_{50} = 9.20 \pm 0.32$ μ M, fivefold better than the standard D-saccharic acid-1,4-lactone, IC_{50} value 48.4 ± 1.25 μ M) which has a methoxy group at the *ortho* position. Surprisingly, a marked decline in activity (to the point of being inactive) was observed

in analog **2** which has a methoxy residue at the *para* position instead of the *ortho* position as in compound **1**. This huge difference in the activities of compounds **1** and **2** clearly indicates that a specific group at a specific position of the benzylidene phenyl ring part plays a vital role in making a potent β -glucuronidase inhibitor in this type of compounds. The excellent activity of compound **5** ($IC_{50} = 9.47 \pm 0.16 \mu\text{M}$) having an *ortho* nitro group on the phenyl ring as compared to the inactivity of its closely related derivative **6** having a *meta* nitro group on phenyl ring proves our hypothesis that a suitable group at a suitable position of the phenyl ring of benzylidene part of molecules is a prerequisite for β -glucuronidase inhibitory potential in these *N*-acylhydrazone Schiff bases. Comparison of activity of chloro-containing compounds **7** ($IC_{50} = 14.7 \pm 0.19 \mu\text{M}$), **8** ($IC_{50} = 15.4 \pm 1.56 \mu\text{M}$), and **9** (inactive) demonstrated that the nature and location of a substitution is important for β -glucuronidase inhibitory potential. Dichloro-substituted compounds **9** and **10** were found to be completely inactive which further proves our hypothesis. Compound **11** ($IC_{50} = 19.6 \pm 0.62 \mu\text{M}$) having an *ortho* fluoro group showed excellent activity, but a little less than analogous chloro compounds **7** and **8**. We also evaluated the effect of heterocyclic ring-containing derivatives, and it was observed that the five membered heterocyclic thiophene ring-containing derivative **12** ($IC_{50} 30.7 \pm 1.49 \mu\text{M}$) produced remarkable activity, while on the other hand five membered heterocyclic rings like furan and its methyl derivatives **13** and **14** were found to be completely inactive. Almost all mono-, di- and trihydroxy substituted compounds **16**, **17**, **18**, **19**, and **20** found to be completely inactive, but unexpectedly compound **15** ($IC_{50} = 12.0 \pm 0.16 \mu\text{M}$) which bears 2,3-dihydroxy substitution, was found to be very efficient and displayed remarkable activity, better than the standard, but, compound **16**, also a 3,4-dihydroxy derivative did not show any activity. *N*-acylhydrazones Schiff base **21** ($IC_{50} = 13.7 \pm 0.40 \mu\text{M}$) synthesized from 1-naphthaldehyde was found to be more active than **22** ($IC_{50} = 22.0 \pm 0.14 \mu\text{M}$) which was synthesized from 2-naphthaldehyde, both without any substitution. Remaining compounds **23–28** were found to be completely inactive. This pattern of activity reveals that the substituent and its position on the phenyl ring of benzylidene part is a driving force for β -glucuronidase inhibition activity.

In conclusion, a number of potential lead molecules has been identified as β -glucuronidase inhibitors. Compounds **1**, **5**, **7**, **8**, **11**, **12**, **15**, **21**, and **22** demonstrated excellent activity and it is anticipated that by slight synthetic modification in these molecules, some new most active β -glucuronidase inhibitors can be developed.

3. Experimental

3.1. General Information

$^1\text{H-NMR}$ experiments were performed on an Avance-Bruker AM 300 MHz instrument (Wissembourg Cedex, France). A Carlo Erba Strumentazione-Mod-1106 (Milan, Italy) used to measure CHN analysis. EI MS was performed on a Finnigan MAT-311A (Bremen, Germany). Thin layer chromatography (TLC) was carried out on pre-coated silica gel glass plates (Kieselgel 60, 254, E. Merck, Darmstadt, Germany). The chromatograms were visualized by UV at 254 and 365 nm or iodine vapours.

3.2. Biological Assays

β -Glucuronidase (E.C. 3.2.1.31 from bovine liver, G-0251) and *p*-nitrophenyl- β -D-glucuronide (N-1627) were purchased From Sigma Chemical Co. (St. Louis, MO, USA). Anhydrous Na₂CO₃ and all other reagents of standard grade were obtained from E. Merck. The anhydrous EtOH and CHCl₃ used in the experiments were dried employing the standard methods. All other solvents and reagents like the benzoyl chloride were of standard grade.

3.3. Assay for β -D-Glucuronidase

β -D-Glucuronidase inhibition was determined by measuring the absorbance of the *p*-nitrophenol which is produce from the substrate at 405 nm. The total reaction volume was 250 μ L. The reaction mixture contains 5 μ L of test compound solution, 185 μ L of 0.1 M acetate buffer, and 10 μ L of enzyme, and it was incubated at 37 °C for 30 min. The plates were read on a multiplate reader at 405 nm after the addition of 50 μ L of 0.4 mM *p*-nitrophenyl- β -D-glucuronide. All assays were performed in triplicate.

3.4. Typical Method for the Synthesis of Compounds 1–28

To a mixture of 2-(4-chloro-2-methylphenoxyacetic acid) hydrazide (1 mmol) in methanol (25 mL) was added a substituted aldehyde (1 mmol) and 3 drops of glacial acetic acid and the mixture was refluxed for 3 h. After completion of the reaction (TLC analysis), it was cooled and evaporated on a rotary evaporator. The resultant crude product was crystallized from methanol to afford 80%–90% yields of pure product. The structures of synthetic compounds 1–28 were determined by different spectroscopic techniques, including ¹H-NMR, and EI MS spectroscopy.

2-(4-Chloro-2-methylphenoxy)-N'-[(3-(2-methoxyphenyl)-2-propenylidene]acetohydrazide (1). Yield: 68%; ¹H-NMR (DMSO-*d*₆) δ : 8.04 (d, 1H, $J = 9.3$ Hz, N=CH-CH), 7.63 (dd, 1H, $J_{5',6'} = 7.8$ Hz, $J_{5',3'} = 1.2$ Hz, H-6'), 7.34 (dd, 1H, $J_{\text{CH-HC=CH}} = 8.1$ Hz, CH-HC=CH), 7.23 (d, 1H, $J_{\text{HC=CH}} = 6$ Hz, HC=CH), 7.12–7.20 (m, 2H, H-3/5) 6.85–7.06 (m, 3H, H- 3'/4'/5'), 6.80 (d, 1H, $J_{6,5} = 9$ Hz, H-6), 5.17 (s, 2H, -OCH₂), 3.84 (s, 3H, -OCH₃), 2.23 (s, 3H, CH₃); EI MS: m/z (%) 358 (M⁺, 30), 327 (100), 189 (40), 175 (55), 159 (100), 125 (75.0), Anal. Calcd for C₁₉H₁₉ClN₂O₃, C = 63.60, H = 5.34, N = 7.81. Found: C = 63.55, H = 5.31, N = 7.80.

2-(4-Chloro-2-methylphenoxy)-N'-[(3-(4-methoxyphenyl)-2-propenylidene]acetohydrazide (2). Yield: 70%; ¹H-NMR (DMSO-*d*₆) δ : 8.02 (d, 1H, $J = 9.3$ Hz, N=CH-CH), 7.55 (d, 2H, $J_{2,3'} = J_{6',5'} = 8.7$ Hz, H-2'/6'), 7.12–7.23 (m, 2H, H-3/5), 7.01 (d, 2H, $J_{3',2'} = J_{5',6'} = 7.0$ Hz, H-3'/5'), 6.76–6.87 (m, 2H, HC = CH), 6.92 (d, 1H, $J_{5,6} = 8.4$ Hz, H-6), 5.03 (s, 2H, -OCH₂), 3.76 (s, 3H, OCH₃), 2.21 (s, 3H, CH₃); m/z (%) 358 (M⁺, 95), 189 (40), 175 (85), 159 (100), 125 (60). Anal. Calcd for C₁₉H₁₉ClN₂O₃, C = 63.60, H = 5.34, N = 7.81. Found: C = 63.57, H = 5.32, N = 7.79.

2-(4-Chloro-2-methylphenoxy)-N'-[(4-ethoxyphenyl)methylidene]acetohydrazide (3). Yield: 80%; ¹H-NMR (DMSO-*d*₆) δ : 8.20 (s, 1H, -N=CH), 7.62 (d, 2H, $J_{2,3'} = J_{6',5'} = 8.7$ Hz, H-2'/6'), 7.23–7.12 (m, 2H, H-3/5), 6.84 (d, 2H, $J_{3',2'} = J_{5',6'} = 8.7$ Hz, H-3'/5'), 6.84 (d, 1H, $J_{6,5} = 8.7$ Hz, H-6), 5.13 (s, 2H, OCH₂), 4.04 (q,

2H, $J = 6.9$ Hz, $-\text{CH}_2$), 2.22 (s, 3H, $-\text{CH}_3$), 1.32 (t, 3H, $J = 6.9$ Hz, $-\text{CH}_3$); EI MS: m/z (%) 347 (M^+ , 56), 205 (95), 155 (89), 147 (92), 119 (73). Anal. Calcd for $\text{C}_{18}\text{H}_{19}\text{ClN}_2\text{O}_3$, C = 62.34, H = 5.52, N = 8.08. Found: C = 62.33, H = 5.50, N = 8.03.

2-(4-Chloro-2-methylphenoxy)-N'-[(3,4,5-trimethoxyphenyl)methylidene]acetohydrazide (4). Yield: 72%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.43 (s, 1H, $-\text{N}=\text{CH}$), 7.56 (d, 1H, $J_{2',6'} = 2.7$ Hz, H-2'), 7.23 (m, 1H, H-5), 7.13 (d, 1H, $J_{3,5} = 2.7$ Hz, H-3), 6.92 (s, 1H, H-6'), 6.83 (d, 1H, $J_{5,6} = 8.7$ Hz, H-6), 5.13 (s, 2H, $-\text{OCH}_2$), 3.82 (s, 9H, $-\text{OCH}_3$), 2.22 (s, 3H, $-\text{CH}_3$); EI MS: m/z (%) 392 (M^+ , 62.1), 251 (15.5), 193 (100), 179 (88.9), 155 (26.8). Anal. Calcd for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_5$, C = 58.09, H = 5.39, N = 7.13. Found: C = 58.05, H = 5.37, N = 7.11.

2-(4-Chloro-2-methylphenoxy)-N'-[(2-nitrophenyl)methylidene]acetohydrazide (5). Yield: 65%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.09 (s, 1H, $\text{N}=\text{CH}$), 8.29 (m, 2H, H-4'/5'), 7.98 (d, 2H, $J_{3',4'} = J_{6',5'} = 9$, H-3'/6'), 7.24 (m, 1H, H-5), 7.13 (d, 1H, $J_{3,5} = 2.7$ Hz, H-3), 6.89 (d, 1H, $J_{6,5} = 8.7$, H-6), 5.22 (s, 2H, $-\text{OCH}_2$), 2.22 (s, 3H, $-\text{CH}_3$); EI MS: m/z (%) 347 (M^+ , 72), 206 (100), 155 (83.3), 125 (75.8). Anal. Calcd for $\text{C}_{16}\text{H}_{14}\text{ClN}_3\text{O}_4$, C = 55.26, H = 4.06, N = 12.08. Found: C = 55.23, H = 4.02, N = 12.04.

2-(4-Chloro-2-methylphenoxy)-N'-[(3-nitrophenyl)methylidene]acetohydrazide (6). Yield: 70%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.40 (s, 1H, $-\text{N}=\text{CH}$), 8.51 (d, 1H, $J_{5',6'} = 7.2$ Hz, H-6'), 8.12–8.25 (m, 2H, H-2'/5'), 7.76 (m, 1H, H-4'), 7.12–7.24 (m, 2H, H-3/5), 6.90 (dd, 1H, $J_{6,5} = 6.2$ Hz, H-6), 5.22 (s, 2H, OCH_2), 2.23 (s, 3H, CH_3); EI MS: m/z (%) 347 (M^+ , 92), 206 (100), 178 (28.4), 141 (17.5), 125 (39.9). Anal. Calcd for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_5$, C = 58.09, H = 5.39, N = 7.13. Found: C = 58.06, H = 5.37, N = 7.12.

2-(4-Chloro-2-methylphenoxy)-N'-[(4-chlorophenyl)methylidene]acetohydrazide (7). Yield: 63%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.27 (s, 1H, $-\text{N}=\text{CH}$), 7.73 (d, 2H, $J_{2',3'} = J_{6',5'} = 8.7$ Hz, H-2'/6'), 7.51 (d, 2H, $J_{3,2'} = J_{5,6'} = 8.1$ Hz, H-3'/5'), 7.13–7.23 (m, 2H, H-3/5), 6.89 (d, $J_{6,5} = 7.8$ Hz, 1H, H-6), 5.15 (s, 2H, $-\text{OCH}_2$), 2.19 (s, 3H, CH_3); EI MS: m/z (%) 336 (M^+ , 20), 195 (35), 155 (55), 125 (80), 89 (100). Anal. Calcd for $\text{C}_{16}\text{H}_{14}\text{Cl}_2\text{N}_2\text{O}_2$, C = 56.99, H = 4.18, N = 8.31. Found: C = 56.95, H = 4.15, N = 8.28.

2-(4-Chloro-2-methylphenoxy)-N'-[(2-chlorophenyl)methylidene]acetohydrazide (8). Yield: 66%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.26 (s, 1H, $-\text{N}=\text{CH}$), 7.77 (m, 2H, H-3'/6'), 7.48 (br. t, 2H, $J_{4',5'} = J_{5',4'} = 8.5$ Hz, H-4'/5'), 7.13–7.24 (m, 1H, H-3/5), 6.89 (d, 1H, $J_{6,5} = 8.7$ Hz, H-6), 5.15 (s, 2H, $-\text{OCH}_2$), 2.23 (s, 3H, $-\text{CH}_3$), 2.19 (s, 3H, CH_3); EI MS: m/z (%) 336 (M^+ , 10), 195 (30), 155 (40), 125 (75), 89 (100). Anal. Calcd for $\text{C}_{16}\text{H}_{14}\text{Cl}_2\text{N}_2\text{O}_2$, C = 56.99, H = 4.18, N = 8.31. Found: C = 56.96, H = 4.15, N = 8.27.

2-(4-Chloro-2-methylphenoxy)-N'-[(3,4-dichlorophenyl)methylidene]acetohydrazide (9). Yield: 70%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.25 (s, 1H, $-\text{N}=\text{CH}$), 7.96 (s, 1H, H-2'), 7.68 (br. s, 2H, 5'/6'), 7.12–7.23 (m, 2H, H-3/5), 6.88 (d, 1H, $J_{6,5} = 7.0$ Hz, H-6), 5.08 (s, 2H, OCH_2), 2.22 (s, 3H, CH_3); EI MS: m/z (%) 371 (M^+ , 92.2), 228 (100), 155 (87.1), 125 (89.8). Anal. Calcd for $\text{C}_{16}\text{H}_{13}\text{Cl}_3\text{N}_2\text{O}_2$, C = 51.71, H = 3.53, N = 7.54. Found: C = 51.68, H = 3.50, N = 7.52.

2-(4-Chloro-2-methylphenoxy)-N'-[(2,6-dichlorophenyl)methylidene]acetohydrazide (10). Yield: 60%; $^1\text{H-NMR}$ ($\text{DMSO}-d_6$) δ : 8.48 (s, 1H, $-\text{N}=\text{CH}$), 7.57 (d, 2H, $J_{3',4'} = J_{5',4'} = 8.7$ Hz, H-3'/5'), 7.44 (dd, 1H, $J_{4,5} = 7.2$ Hz, H-4'), 7.24–7.11 (m, 2H, H-3/5), 6.78 (d, 1H, $J_{6,5} = 8.7$ Hz, H-6), 5.09 (s, 2H, $-\text{OCH}_2$), 2.18

(s, 3H, CH₃); EI MS: *m/z* (%) 370 (M⁺, 17.3), 229 (39.5), 155 (60.1), 125 (100), 89 (62). Anal. Calcd for C₁₆H₁₃Cl₃N₂O₂, C = 51.71, H = 3.53, N = 7.54. Found: C = 51.66, H = 3.51, N = 7.52.

2-(4-Chloro-2-methylphenoxy)-N'-[-(2-fluorophenyl)methylidene]acetohydrazide (11). Yield: 74%; ¹H-NMR (DMSO-*d*₆) δ: 8.52 (s, 1H, -N=CH), 7.95 (dd, 1H, *J*_{4',5'} = 8.1 Hz, H-5'), 7.48 (dd, 1H, *J*_{6',5'} = 6.9 Hz, H-6'), 7.25–7.31 (m, 2H, H-3'/4'), 7.12–7.21 (m, 2H, H-3/5), 6.90 (d, 1H, *J*_{6,5} = 8.7 Hz, H-6), 5.17 (s, 2H, -OCH₂), 2.23 (s, 3H, CH₃); EI MS: *m/z* (%) 320 (M⁺, 12), 179 (80), 125 (100), 89 (75.8). Anal. Calcd for C₁₆H₁₄ClFN₂O₂, C = 59.91, H = 4.40, N = 8.73. Found: C = 59.89, H = 4.39, N = 8.72.

2-(4-Chloro-2-methylphenoxy)-N'-[3-thienylmethylidene]acetohydrazide (12). Yield: 68%; ¹H-NMR (DMSO-*d*₆) δ: 8.07 (s, 1H, -N=CH), 7.12–7.24 (m, 3H, H-3, 5, 6), 6.73–9.92 (m, 3H, H-2'/4'/5'), 5.10 (s, 2H, -OCH₂), 2.22 (s, 3H, CH₃); EI MS: *m/z* (%) 308 (M⁺, 15), 199 (10), 155 (45), 125 (100). Anal. Calcd for C₁₄H₁₃ClN₂O₂S, C = 54.46, H = 4.24, N = 9.07. Found: C = 54.45, H = 4.22, N = 9.04.

2-(4-Chloro-2-methylphenoxy)-N'-[-2-furylmethylidene]acetohydrazide (13). Yield: 70%; ¹H-NMR (DMSO-*d*₆) δ: 8.17 (s, 1H, -N=CH), 7.11–7.23 (m, 2H, H-3,5), 6.90 (d, 2H, *J*_{3',4'} = *J*_{5',4'} = 3.3 Hz, H-3'/5'), 6.81 (d, 1H, *J*_{6,5} = 8.7 Hz, 6), 6.62 (dd, 1H, *J*_{4,3} = *J*_{4,5} = 3.3 Hz, 4'), 5.07 (s, 2H, -OCH₂), 2.22 (s, 3H, -CH₃); EI MS: *m/z* (%) 292 (M⁺, 35), 155 (50), 151 (80.), 125 (100). Anal. Calcd for C₁₄H₁₃ClN₂O₃, C = 57.44, H = 4.48, N = 9.57. Found: C = 57.44, H = 4.46, N = 9.56.

2-(4-Chloro-2-methylphenoxy)-N'-[-(5-methyl-2-furyl)methylidene]acetohydrazide (14). Yield: 72%; ¹H-NMR (DMSO-*d*₆) δ: 8.07 (s, 1H, -N=CH), 7.11–7.23 (m, 2H, H-3,5), 6.88 (d, 1H, *J*_{5,6} = 8.7 Hz, H-6), 6.80 (d, 1H, *J*_{3',4'} = 5.4 Hz, 3'), 6.24 (d, 1H, *J*_{4',3'} = 2.7 Hz, 4'), 5.07 (s, 2H, -OCH₂), 2.21 (s, 3H, -CH₃) 2.19 (s, 3H, -CH₃); EI MS: *m/z* (%) 306 (M⁺, 20), 65 (30), 137 (100), 125 (40). Anal. Calcd for C₁₅H₁₅ClN₂O₃, C = 58.73, H = 4.93, N = 9.13. Found: C = 58.70, H = 4.96, N = 9.10.

2-(4-Chloro-2-methylphenoxy)-N'-[-(2,3-dihydroxyphenyl)methylidene]acetohydrazide (15). Yield: 76%; ¹H-NMR (DMSO-*d*₆) δ: 8.41 (s, 1H, -N=CH), 7.24 (m, 1H, H-5), 7.13 (d, 1H, *J*_{3,5} = 2.7 Hz, H-3), 6.89 (d, 1H, *J*_{5',6'} = 8.7 Hz, H-6'), 6.83 (d, 1H, *J*_{6,5} = 8.7 Hz, H-6), 6.71 (m, 2H, H-3'/4'), 5.15 (s, 2H, -OCH₂), 2.19 (s, 3H, -CH₃); EI MS: *m/z* (%) 334 (M⁺, 100), 300 (9), 175 (89.7), 193 (35), 179 (50), 137 (65), 125 (60). Anal. Calcd for C₁₆H₁₅ClN₂O₄, C = 57.41, H = 4.5, N = 8.37. Found: C = 57.40, H = 3.50, N = 8.36.

2-(4-Chloro-2-methylphenoxy)-N'-[-(3,4-dihydroxyphenyl)methylidene]acetohydrazide (16). Yield: 68%; ¹H-NMR (DMSO-*d*₆) δ: 8.06 (s, 1H, -N=CH), 7.12–7.19 (m, 2H, H-3/5), 7.24 (d, 1H, *J*_{2',6'} = 2.7 Hz, H-2'), 6.85–6.92 (m, 2H, H-5'/6'), 6.76 (d, *J*_{6,5} = 8.1 Hz, 1H, H-6), 5.10 (s, 2H, -OCH₂), 2.22 (s, 3H, CH₃); EI MS: *m/z* (%) 334 (M, 36.8), 193 (23.9), 155 (71.6), 125 (88.8), 77 (100). Anal. Calcd for C₁₆H₁₅ClN₂O₄, C = 57.41, H = 4.5, N = 8.37. Found: C = 57.39, H = 3.50, N = 8.37.

2-(4-Chloro-2-methylphenoxy)-N'-[1-(4-hydroxyphenyl)ethylidene]acetohydrazide (17). Yield: 70%; ¹H-NMR (DMSO-*d*₆) δ: 7.65 (d, 2H, *J*_{2',3'} = *J*_{6',5'} = 8.7 Hz, H-2'/6'), 7.23 (m, 1H, Hz H-6), 7.12 (d, 1H, *J* = 2.4 Hz, H-3), 6.90 (d, *J*_{3',2'} = *J*_{5',6'} = 8.7 Hz, 2H, H-3'/5'), 6.77 (d, 1H, *J*_{6,5} = 8.7 Hz, H-6), 5.16 (s, 2H, -OCH₂), 2.19 (s, 6H, CH₃); EI MS: *m/z* (%) 332 (M⁺, 56.7), 191 (100), 177 (24.5), 149 (34.5), 134

(37.9). Anal. Calcd for $C_{17}H_{16}ClN_2O_3$, C = 61.36, H = 5.15, N = 8.42. Found: C = 61.34, H = 5.14, N = 8.40.

2-(4-Chloro-2-methylphenoxy)-N'-[(3-hydroxyphenyl)methylidene]acetohydrazide (**18**). Yield: 61%; 1H -NMR (DMSO- d_6) δ : 8.18 (s, 1H, -N=CH), 7.04–7.24 (m, 5H, Ar-H), 6.79–6.89 (m, 2H, H-5-6), 5.14 (s, 2H, -OCH₂), 2.22 (s, 3H, -CH₃); EI MS: m/z (%) 318 (M^+ , 97.6), 177 (100), 155 (71), 125 (65.3). Anal. Calcd for $C_{16}H_{15}ClN_2O_3$, C = 60.29, H = 4.74, N = 8.79. Found: C = 69.27, H = 3.49, N = 8.34.

2-(4-Chloro-2-methylphenoxy)-N'-[1-(2,4,6-trihydroxyphenyl)ethylidene]acetohydrazide (**19**). Yield: 62%; 1H -NMR (DMSO- d_6) δ : 8.16 (s, 1H, -N=CH), 7.21 (br.s, 2H, H-3'/5'), 7.16–7.23 (m, 2H, H-3/5), 6.86 (d, $J_{6,5} = 8.7$ Hz, 1H, H-6), 4.98 (s, 2H, -OCH₂), 2.18 (s, 3H, CH₃), 1.93 (s, 3H, CH₃); EI MS: m/z (%) 254 (M^+ , 100), 155 (91.2), 142 (75.5), 125 (100), 113 (100), 99 (86.6).

2-(4-Chloro-2-methylphenoxy)-N'-[(4-hydroxyphenyl)methylidene]acetohydrazide (**20**). Yield: 72%; 1H -NMR (DMSO- d_6) δ : 7.88 (s, 1H, -N=CH), 7.52 (d, 2H, d, 2H, $J_{2,3'} = J_{6,5'} = 8.7$ Hz, H-2'/6'), 7.24 (d, 1H, $J_{5,6} = 8.1$ Hz, H-5), 7.13 (d, 1H, $J = 2.4$ Hz, H-3), 6.88 (d, 2H, $J_{3,2'} = J_{5',6'} = 8.7$ Hz, H-3'/5'), 6.79 (d, 1H, $J_{6,5} = 8.7$ Hz, H-6), 5.11 (s, 2H, -OCH₂), 2.22 (s, 3H, CH₃); EI MS: m/z (%) 318 (M^+ , 9.30), 199 (10.78), 177 (27.43), 155 (44), 125 (67.50), 77 (100). Anal. Calcd for $C_{16}H_{15}ClN_2O_3$, C = 60.29, H = 4.74, N = 8.79. Found: C = 69.26, H = 3.50, N = 8.34.

2-(4-Chloro-2-methylphenoxy)-N'-[1-naphthylmethylidene]acetohydrazide (**21**). Yield: 68%; 1H -NMR (DMSO- d_6) δ : 8.66 (s, 1H, N=CH), 8.83 (d, $J = 8.1$ Hz 1H, H-8'), 8.61 (d, 1H, $J_{4,3'} = 8.4$ Hz, H-4'), 8.02 (br, d, 2H, $J = 7.8$ Hz, H-2'/5'), 7.93 (d, 1H, $J_{3',4'} = 7.8$ Hz, H-3') 7.65 (m, 2H, H-6'/7'), 7.14–7.25 (m, 2H, H-3/5), 6.95 (d, $J_{6,5} = 8.4$ Hz, 1H, H-6), 5.24 (s, 2H, -OCH₂), 2.26 (s, 3H, -CH₃); EI MS: m/z (%) 352 (M^+ , 100), 211(65), 199. Anal. Calcd for $C_{20}H_{17}ClN_2O_2$, C = 68.09, H = 4.86, N = 7.94. Found: C = 68.06, H = 4.84, N = 7.92.

2-(4-Chloro-2-methylphenoxy)-N'-[2-naphthylmethylidene]acetohydrazide (**22**). Yield: 70%; 1H -NMR (DMSO- d_6) δ : 8.43 (s, 1H, N=CH), 8.16 (t, 2H, $J = 8.4$ Hz, H-3',6'), 7.94 (br, s, 2H, H-5/4') 7.56 (t, 2H, $J_{5,6'} = J_{8,7'} = 9.6$ Hz, H-5'/8'), 7.22 (s, 1H, H-1'), 7.17 (d, 1H, $J_{7,8'} = 8.7$ Hz, H-7'), 6.92 (m, 2H, H-5/6), 5.23 (s, 2H, OCH₂), 2.23 (s, 3H, CH₃); EI MS: m/z (%) 352 (M^+ , 72), 211 (60) 199 (35), 169 (45), 153 (100), 127 (60). Anal. Calcd for $C_{20}H_{17}ClN_2O_2$, C = 68.09, H = 4.86, N = 7.94. Found: C = 68.04, H = 4.85, N = 7.91.

2-(4-Chloro-2-methylphenoxy)-N'-[(5-Chloro-2-hydroxyphenyl)methylidene]acetohydrazide (**23**). Yield: 66%; 1H -NMR (DMSO- d_6) δ : 8.23 (s,1H, -N=CH), 7.70 (d, 1H, $J_{6,4'} = 2.7$ Hz H-6'),7.12–7.23 (m, 2H, H-3/5), 7.31 (m, 1H, H-4'), 6.93 (d, 1H, $J_{3',4'} = 7.8$ Hz, H-3'),6.86 (d, 1H, $J_{6,5} = 8.7$ Hz, H-6), 5.17 (s, 2H, -OCH₂), 2.22 (s, 3H, -CH₃); EI MS: m/z (%) 352 (M^+ , 83.8), 210 (7.4), 197 (62.6), 155 (100), 141 (46.8). Anal. Calcd for $C_{16}H_{14}Cl_2N_2O_3$, C = 54.41, H = 4.00, N = 7.93. Found: C = 54.39, H = 4.00, N = 7.92.

2-(4-Chloro-2-methylphenoxy)-N'-[1(2,4-dihydroxy-5-nitrophenyl)ethylidene]acetohydrazide (**24**). Yield: 70%; 1H -NMR (DMSO- d_6) δ : 8.23 (s, 1H, H-6'), 7.17–7.23 (m, 2H, H-3/5), 6.89 (d, 1H, $J_{6,5} = 8.7$ Hz,

H-6), 6.51 (s, 1H, H-3'), 4.82 (s, 2H, -OCH₂), 2.26 (s, 3H, CH₃), 2.21 (s, 3H, CH₃); EI MS: *m/z* (%) 392 (M⁺, 64.4), 251 (72.1), 237 (100), 125 (89.9), 77 (68.5). Anal. Calcd for C₁₆H₁₄ClN₃O₆, C = 50.60, H = 3.72, N = 11.07. Found: C = 50.57, H = 3.70, N = 11.05.

2-(4-Chloro-2-methylphenoxy)-N'-[(4-hydroxy-3-methoxyphenyl)methylidene]acetohydrazide (25). Yield: 59%; ¹H-NMR (DMSO-*d*₆) δ : 8.11 (s, 1H, -N=CH), 7.12–7.23 (m, 2H, H-3/5), 7.03 (s, 1H, H-2'), 6.86–6.96 (m, 2H, H-5'/6') 6.83 (d, 1H, *J*_{6,5} = 8.7 Hz, H-6), 5.12 (s, 2H, -OCH₂), 3.78 (s, 3H, OCH₃), 2.23 (s, 3H, CH₃); EI MS: *m/z* (%) 348 (M⁺, 88.5), 207 (51), 193 (100), 155 (43), 125 (46.9). Anal. Calcd for C₁₇H₁₇ClN₂O₄, C = 58.54, H = 4.91, N = 8.03. Found: C = 58.53, H = 4.90, N = 8.02.

2-(4-Chloro-2-methylphenoxy)-N'-{[4-(methylsulfonyl)phenyl]methylidene}acetohydrazide (26). Yield: 63%; ¹H-NMR (DMSO-*d*₆) δ : 8.22 (s, 1H, -N=CH), 7.63 (d, 2H, *J*_{2,3} = *J*_{6,5} = 8.4 Hz, H-2'/6'), 7.31 (d, 2H, *J*_{3,2} = *J*_{5,6} = 8.4 Hz, H-3'/5'), 7.23 (m, 1H, H-5), 7.13 (d, 1H, *J* = 2.7 Hz, H-3), 6.89 (d, *J*_{6,5} = 8.4 Hz, 1H, H-6), 5.15 (s, 2H, -OCH₂), 2.19 (s, 3H, CH₃); EI MS: *m/z* (%) 348 (M⁺, 60), 207 (25), 149 (100), 125 (35), 118 (35). Anal. Calcd for C₁₇H₁₇ClN₂O₂S, C = 58.53, H = 4.91, N = 8.03. Found: C = 58.51, H = 4.90, N = 8.02.

2-(4-Chloro-2-methylphenoxy)-N'-[(2-methylphenyl)methylidene]acetohydrazide (27). Yield: 67%; ¹H-NMR (DMSO-*d*₆) δ : 8.25 (s, 1H, -N=CH), 7.78 (d, *J* = 7.5 Hz, 1H, H-6'), 7.30–7.21 (m, 3H, H-3'/4'/5'), 7.13–7.17 (m, 2H, H-3/5), 6.84 (d, 1H, *J*_{6,5} = 8.7 Hz, H-6), 5.15 (s, 2H, -OCH₂), 2.23 (s, 3H, -CH₃), 2.19 (s, 3H, CH₃); EI MS: *m/z* (%) 316 (M⁺, 79.4), 199 (78.9), 175 (89.7), 155 (100), 141 (93.2), 118 (94.2). Anal. Calcd for C₁₇H₁₇ClN₂O₂, C = 64.46, H = 4.41, N = 8.84. Found: C = 64.45, H = 4.40, N = 8.83.

2-(4-Chloro-2-methylphenoxy)-N'-[(4-methylphenyl)methylidene]acetohydrazide (28). Yield: 63%; ¹H-NMR (DMSO-*d*₆) δ : 8.23 (s, 1H, -N=CH), 7.59 (d, 2H, *J*_{2,3} = *J*_{6,5} = 7.8 Hz, H-2'/6'), 7.24 (d, 2H, *J*_{3,2} = *J*_{5,6} = 8.1 Hz, H-3'/5'), 7.13 (d, *J* = 2.7 Hz, 1H, H-3), 7.17 (m, 1H, H-5), 6.85 (d, *J*_{6,5} = 8.7 Hz, 1H, H-6), 5.15 (s, 2H, -OCH₂), 2.32 (s, 3H, CH₃), 2.19 (s, 3H, CH₃); EI MS: *m/z* (%) 316 (M⁺, 9.61), 199 (10.80), 175 (36.26), 155 (47.70), 125 (71.62). Anal. Calcd for C₁₇H₁₇ClN₂O₂, C = 64.46, H = 4.41, N = 8.84. Found: C = 64.44, H = 4.40, N = 8.82.

4. Conclusions

A number of potential lead molecules **1**, **5**, **7**, **8**, **11**, **12**, **15**, **21**, and **22** have been identified as β -glucuronidase inhibitors and it is anticipated that by slight synthetic modification in these molecules, some new most active β -glucuronidase inhibitors can be developed.

Supplementary Materials

Supplementary materials can be accessed at: <http://www.mdpi.com/1420-3049/19/7/8788/s1>.

Acknowledgments

Syed Adnan Ali Shah would like to acknowledge Universiti Teknologi MARA (UiTM) for the financial support under the Principal Investigator Support Initiative (PSI) Grant Scheme with reference

number UiTM 600-RMI/DANA 5/3/PSI (251/2013) and Cumulative Impact Factor Initiative (CIFI) Grant Scheme with reference number UiTM 600-RMI/DANA 5/3/CIFI (117/2013) and The authors are thankful to Organization for Prohibition of Chemical Weapons (OPCW), Netherlands (Project No. L/ICA/ICB/173681/12) and Higher Education Commission (HEC) Pakistan under “National Research Support Program for Universities” (Project No. 1910) for financial support.

Author Contributions

All authors equally contributed.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Küçükgül, S.G.; Mazi, A.; Sahin, F.; Öztürk, S.; Stables, J. Synthesis and biological activities of diflunisal hydrazide hydrazones. *Eur. J. Med. Chem.* **2003**, *38*, 1005–1013.
2. Melnyk, P.; Leroux, V.; Sergheraert, C.; Grellier, P. Design, synthesis and *in vitro* antimalarial activity of an acylhydrazone library. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 31–35.
3. Lima, P.C.; Lima, L.M.; da Silva, K.C.M.; Léda, P.H.O.; Miranda, A.L.P.; Fraga, C.A.M.; Barreiro, E.J. Synthesis and analgesic activity of novel *N*-acyl aryl hydrazones and isosters, derived from natural saffrole. *Eur. J. Med. Chem.* **2000**, *35*, 187–203.
4. Cunha, A.C.; Figueiredo, J.M.; Tributino, J.L.M.; Miranda, A.L.P.; Castro, H.C.; Zingali, R.B.; Fraga, C.A.M.; de Souza, M.C.B.V.; Ferreira, V.F.; Barreiro, E. Antiplatelet properties of novel *N*-Substituted-phenyl-1,2,3-triazole-4-acylhydrazone Derivatives. *J. Bioorg. Med. Chem.* **2003**, *11*, 2051–2059.
5. Bedia, K.K.; Elçin, O.; Seda, U.; Fatma, K.; Nathaly, S.; Sevim, R.; Dimoglo, A. Synthesis and characterization of novel hydrazide-hydrazones and the study of their structure antituberculosis activity. *Eur. J. Med. Chem.* **2006**, *41*, 1253–1261.
6. Terzioglu, N.; Gürsoy, A. Synthesis and anticancer evaluation of some new hydrazone derivatives of 2,6-dimethylimidazo[2,1-*b*][1,3,4]thiadiazole-5-carbohydrazide. *Eur. J. Med. Chem.* **2003**, *38*, 781–786.
7. Beraldo, H.; Gambino, D. The wide pharmacological versatility of semicarbazones, thiosemicarbazones and their metal complexes. *Mini-Rev. Med. Chem.* **2004**, *4*, 31–39.
8. Costa, R.F.F.; Rebolledo, A.P.; Matencio, T.; Calado, H.D.R.; Ardisson, J.D.; Cortes, M.E.; Rodrigues, B.L.; Beraldo, H. Metal complexes of 2-benzoylpyridine-derived thiosemicarbazones: Structural, electrochemical and biological studies. *J. Coord. Chem.* **2005**, *58*, 1307–1319.
9. Hussain, Z.; Khan, K.M.; Perveen, S.; Nawaz, Y.; Bukhari, I.H. Antifungal activity of the pyrolyzate of glucose, sucrose and starch in comparison to paper pyrolyzate, *J. Chem. Soc. Pak.* **2011**, *33*, 694–697.
10. Bondock, S.; Khalifa, W.; Fadda, A.A. Synthesis and antimicrobial evaluation of some new thiazole, thiazolidinone and thiazoline derivatives starting from 1-chloro-3,4-dihydronaphthalene-2-carboxaldehyde. *Eur. J. Med. Chem.* **2007**, *42*, 948–954.

11. Bernardino, A.M.R.; Gomes, A.O.; Charret, K.S.; Freitas, A.C.C.; Machado, G.M.C.; Canto-Cavalheiro, M.M.; Leon, L.L.; Amaral, V.F. Synthesis and leishmanicidal activities of 1-(4-X-phenyl)-N'-[(4-Y-phenyl)methylene]-1H-pyrazole-4-carbohydrazides *Eur. J. Med. Chem.* **2006**, *41*, 80–87.
12. Xia, Y.; Fan, C.-D.; Zhao, B.-X.; Zhao, J.; Shin, D.-S.; Miao, J.-Y. Synthesis and structure activity relationships of novel 1-arylmethyl-3-aryl-1H-pyrazole-5-carbohydrazone derivatives as potential agents against A549 lung cancer cells. *Eur. J. Med. Chem.* **2008**, *43*, 2347–2353.
13. Zheng, L.W.; Wu, L.L.; Zhao, B.-X.; Dong, W.L.; Miao, J.Y. Synthesis of novel substituted pyrazole-5-carbohydrazone derivatives and discovery of a potent apoptosis inducer in A549 lung cancer cells. *Bioorg. Med. Chem.* **2009**, *17*, 1957–1962.
14. Todeschini, A.R.; Miranda, A.L.; Silva C.M.; Parrini, S.C.; Barreiro, E.J. Synthesis and evaluation of analgesic, antiinflammatory and antiplatelet properties of new 2-pyridylarylhydrazone derivatives. *Eur. J. Med. Chem.* **1998**, *33*, 189–199.
15. Anouar, E.H.; Raweh, S.; Bayach, I.; Taha, M.; Baharudin, M.S.; Meo, F.D.; Hasan, M.H.; Adam, A.; Ismail, N.H.; Weber, J.F.; *et al.* Antioxidant properties of phenolic Schiff bases: Structure-activity relationship and mechanism of action. *J. Comput. Aided. Mol. Des.* **2013**, *27*, 951–964.
16. Taha, M.; Ismail, N.H.; Jamil, W.; Yousuf, S.; Jaafar, F.M.; Ali, M.I.; Kashif, S.M.; Hussain, E. Synthesis, evaluation of antioxidant activity and crystal structure of 2,4-dimethylbenzoylhydrazones. *Molecules* **2013**, *18*, 10912–10929.
17. Khan, K.M.; Shah, Z.; Ahmad, V.U.; Khan, M.; Taha, M.; Ali, S.; Perveen, S.; Choudhary, M.I.; Voelter, W. 2,4,6-Trichlorophenylhydrazine Schiff bases as DPPH Radical and super oxide anion scavengers. *Med. Chem.* **2012**, *8*, 452–461.
18. Khan, K.M.; Taha, M.; Naz, F.; Ali, S.; Perveen, S.; Choudhary, M.I. Synthesis of acylhydrazone Schiff bases and their anti-oxidant activity. *Med. Chem.* **2012**, *8*, 705–710.
19. Khan, K.M.; Shah, Z.; Ahmad, V.U.; Khan, M.; Taha, M.; Rahim, F.; Jahan, H.; Perveen, H.; Choudhary, M.I. Synthesis of 2,4,6-trichlorophenyl hydrazones and their inhibitory potential against glycation of protein. *Med. Chem.* **2011**, *7*, 572–580.
20. Khan, K.M.; Rahim, F.; Ambreen, N.; Taha, M.; Khan, M.; Jahan, H.; Najeebullah, U.; Shaikh, A.; Iqbal, S.; Perveen, S.; *et al.* Synthesis of benzophenonehydrazone Schiff bases and their *in vitro* antiglycating activities. *Med. Chem.* **2013**, *9*, 588–595.
21. Taha, M.; Naz, H.; Rasheed, S.; Ismail, N.H.; Rahman, A.A.; Yousuf, S.; Choudhary M.I. Synthesis of 4-Methoxybenzoylhydrazones and Evaluation of Their Antiglycation Activity. *Molecules* **2014**, *19*, 1286–1301.
22. Khan, K.M.; Taha, M.; Rahim, F.; Fakhri, M.I.; Jamil, W.; Khan, M.; Rasheed, S.; Karim, A. Perveen, S.; Choudhary, M.I. Acylhydrazone Schiff bases: Synthesis and antiglycation activity *J. Pak. Chem Soc.* **2013**, *35*, 929–937.
23. Taha, M.; Baharudin, M.S.; Ismail, N.H.; Khan, K.M.; Jaafar, F.M.; Samreen; Siddiqui, S.; Choudhary, M.I. Synthesis of 2-methoxybenzoylhydrazone and evaluation of their antileishmanial activity. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 3463–3466.

24. Musharraf, S.G.; Bibi, A.; Shahid, N.; Najam-ul-Haq, M.; Khan, M.; Taha, M.; Mughal, U.R.; Khan, K.M. Acylhydrazide and isatin Schiff bases as alternate UV-laser desorption ionization (LDI) matrices for low molecular weight (LMW) peptides analysis. *Am. J. Anal. Chem.* **2012**, *3*, 779–789.
25. De Graaf, M.; Boven, E.; Scheeren, H.W.; Haisma, H.J.; Pinedo, H.M. β -Glucuronidase-mediated drug release. *Curr. Pharm. Design.* **2002**, *8*, 1391–1403
26. Sperker, B.; Backman, J.T.; Kromer, K. The role of β -glucuronidase in drug disposition and drug targeting in humans. *Clin. Pharmacokinet.* **1997**, *33*, 18–31.
27. Wakabayashi, M. *Metabolic Conjugation and Metabolic Hydrolysis*; Fishman, W.H., Ed.; Academic Press: New York, NY, USA, 1970; p. 519.
28. Paigen, K. Mammalian β -Glucuronidase: Genetics, molecular biology, and cell biology. *Prog. Nucleic Acid Res. Mol. Biol.* **1989**, *37*, 155–205.
29. Murdter, T.E.; Sperker, B.; Kivisto, K.T.; McClellan, M.; Fritz, P.; Friedel, G.; Linder, A.; Bosslet, K.; Toomes, H.; Dierkesmann, R.; *et al.* Enhanced uptake of doxorubicin into bronchial carcinoma: α -Glucuronidase mediates release of doxorubicin from a glucuronide prodrug (hmr 1826) at the tumor site. *Cancer Res.* **1997**, *57*, 2440.
30. Sly, W.S.; Quinton, B.A.; McAlister, W.H.; Rimoin, D.L. β -Glucuronidase deficiency: Report of clinical, radiologic, and biochemical features of a new mucopolysaccharidosis. *J. Pediatr.* **1973**, *82*, 249–257.
31. Hall, C.W.; Cantz, M.; Neufeld, E.F. A β -Glucuronidase deficiency mucopolysaccharidosis: Studies in cultured fibroblasts. *Arch. Biochem. Biophys.* **1973**, *155*, 32–38.
32. Khan, K.M.; Taha, M.; Ali, M.; Perveen, S. A mild and alternative approach towards symmetrical disulfides using $H_3IO_5/NaHSO_3$ combination. *Lett. Org. Chem.* **2009**, *6*, 319–320.
33. Khan, K.M.; Khan, M.Z.; Taha, M.; Maharvi, G.M.; Saify, Z.S.; Parveen, S.; Choudhary, M.I. Leishmanicidal potential of *N*-substituted morpholine derivatives: Synthesis and structure-activity relationships. *Nat. Prod. Res.* **2009**, *23*, 479–484.
34. Taha, M.; Ismail, N.H.; Imran, S.; Khan, K.M. 4-[5-(2-Methoxyphenyl)-1,3,4-oxadiazol-2-yl] benzohydrazide. *Molbank* **2014**, *2014*, M826.
35. Khan, K.M.; Shujaat, S.; Rahat, S.; Hayat, S.; Atta-ur-Rahman; Choudhary, M.I. β -*N*-Cyanoethyl acyl hydrazide derivatives: A new class of β -glucuronidase inhibitors. *Chem. Pharm. Bull.* **2002**, *50*, 1443–1446.
36. Ahmed, W.; Rani, M.; Khan, I.A.; Iqbal, A.; Khan, K.M.; Haleem, M.A.; Azim, M.K. Characterization of hydrazides and hydrazine derivatives as novel aspartic protease inhibitors. *J. Enz. Inhib. Med. Chem.* **2010**, *25*, 673–678.
37. Khan, K.M.; Taha, M.; Naz, F.; Khan, M.; Rahim, F.; Samreen, Perveen, S.; Choudhary, M.I. Synthesis and *in vitro* leishmanicidal activity of disulfide derivatives. *Med. Chem.* **2011**, *7*, 704–710.
38. Khan, K.M.; Rasheed, M.; Zia-Ullah, Hayat, S.; Kaukab, F.; Choudhary M.I.; Atta-ur-Rahman. Synthesis and *in vitro* leishmanicidal activity of some hydrazides and their analogs. *Bioorg. Med. Chem.* **2003**, *11*, 1381–1387.
39. Khan, K.M.; Naz, F.; Taha, M.; Khan, A.; Perveen, S.; Choudhary, M.I.; Voelter, W. Synthesis and *in vitro* urease inhibitory activity of *N,N'*-disubstituted thioureas. *Eur. J. Med. Chem.* **2014**, *74*, 314–323.

40. Khan, K.M.; Rahim, F.; Wadood, A.; Taha, M.; Khan, M.; Naureen, S.; Ambreen, N.; Hussain, S.; Perveen, S.; Iqbal, M.C. Evaluation of bisindole as potent β -glucuronidase inhibitors: Synthesis and *in silico* based studies. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1825–1829.
41. Khan, K.M.; Ambreen, N.; Taha, M.; Halim, S.A.; Zaheer-ul-Haq; Naureen, S.; Rasheed, S.; Perveen, S.; Ali, S.; Choudhary, M.I. Structure-based design, synthesis and biological evaluation of β -glucuronidase Inhibitors. *J. Comput. Aided Mol. Des.* **2014**, *28*, 577–585.
42. Khan, K.M.; Rahim, F.; Wadood, A.; Kosar, N.; Taha, M.; Khan, A.; Fakhri, M.I.; Junaid, M.; Rehman, W.; Khan, M.; *et al.* Synthesis and molecular docking studies of potent α -glucosidase inhibitors based on biscoumarin skeleton. *Eur. J. Med. Chem.* **2014**, *81*, 245–252.
43. Collins, R.A.; Ng, T.B.; Fong, W.P.; Wan, C.C.; Yeung, H.W. Inhibition of glycohydrolase enzymes by aqueous extracts of chinese medicinal herbs in a microplate format. *IUBMB Life* **1997**, *42*, 1163–1169.

Sample Availability: Not available

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).