


Pharmacological aspects

Psychiatric side effects of medications prescribed in internal medicine

Rodrigo Casagrande Tango, MD



Several pharmacological treatments used in internal medicine can induce psychiatric side effects (PSEs) that mimic diagnoses seen in psychiatry. PSEs may occur upon withdrawal or intoxication, and also at usual therapeutic doses. Drugs that may lead to depressive, anxious, or psychotic syndromes include corticosteroids, isotretinoin, levodopa, mefloquine, interferon- α , and anabolic steroids, as well as some over-the-counter medications. PSEs are often difficult to diagnose and can be very harmful to patients. PSEs are discussed in this review, as well as diagnostic clues to facilitate their identification.

Dialogues Clin Neurosci. 2003;5:155-165.

Keywords: adverse effect; psychiatry; mechanism; risk factor; diagnosis; mefloquine; chloroquine; metronidazole; isotretinoin; interferon; steroid; β -blocker

Author affiliations: Unité de Psychopharmacologie Clinique, Hôpitaux Universitaires de Genève, Chêne-Bourg, Switzerland

Address for correspondence: Unité de Psychopharmacologie Clinique, 2, chemin du Petit Bel-Air 1225 Chêne-Bourg, Switzerland (e-mail: rctango@web.de)

Psychediatric side effects (PSEs) can be induced by the pharmacological treatment of physical illnesses. The clinical presentation of PSEs often resembles spontaneous psychiatric syndromes (ie, noniatrogenic, naturally occurring diseases). PSEs can occur at usual doses, in cases of intoxication, or during the days following withdrawal of a given treatment. PSEs range from short-lasting anxiety to severe confusion, and alleged cases of suicide have even been reported.

The *Diagnostic and Statistic Manual of Mental Disorders, Fourth Edition (DSM-IV)*¹ defines some dozens of categories of PSE, according to the disorder and to the incriminated substance, eg, “persisting dementia induced by sedatives, hypnotics or anxiolytics.” The *DSM-IV* categories include drugs for therapeutic purposes, medication taken abusively, and other substances. The *International Classification of Diseases*² is very similar to *DSM-IV* in its categorization, with minor differences in terms of category codes.

The challenge of PSEs in everyday practice is the difficulty in recognizing these frequent and potentially dangerous situations. The diagnosis of PSEs raises the same questions as the diagnosis of any psychiatric sign or symptom, which means that the clinician has to consider a rather long list of differential diagnoses. The following concepts refer to PSEs, as well as to other kinds of side effects:

- **Exposure:** The period of time the patient received the drug suspected of inducing a side effect.
- **Dechallenge:** The interruption of the suspected medication, regardless of the remission of adverse effects. *Positive dechallenge* means that remission was temporally associated with the interruption of medication.
- **Rechallenge:** The reintroduction of the suspected drug. *Positive rechallenge* means that symptom reappearance was temporally associated with suspected drug reintroduction.

Pharmacological aspects

A drug can be considered to have a high probability of causing side effects in cases of positive exposure, with a positive dechallenge and a positive rechallenge.

The importance of PSEs relates to the potential harm of these side effects and to their high incidence. Two examples are reserpine and corticosteroids. Reserpine, when it was prescribed, may have caused mood disorders in 10% of treated subjects. With corticosteroids, 6% of all patients develop some PSEs.³

The focus of this review is on depression, anxiety, and psychotic states, but a few other PSEs are also mentioned. A summary of the PSEs is presented and some examples are given in detail. In addition, diagnostic issues are discussed to facilitate identification of PSEs in internal medicine. PSEs secondary to psychotropic medication (such as antidepressants, anxiolytics, antipsychotics, or mood-stabilizing agents) are not described here. They probably induce more PSEs than the compounds quoted here, since they act directly on the nervous system.

Mechanisms of PSEs

As with any side effect, pharmacological mechanisms are divided according to their pharmacokinetic or pharmacodynamic nature. Patient-specific factors also lead to PSEs.

Pharmacodynamic mechanisms

Medications used in the treatment of physical disorders can modify neurotransmitter systems (as do psychotropic medications). These modes of action can imply a direct influence on neurotransmitters, as is the case for dopaminergic agents in the treatment of Parkinson's disease. Interleukin (IL) treatment is another example where there is a direct influence, since interleukins are involved in neurotransmission as well as in many other bodily functions. Other nonpsychotropic medications influence neurotransmitter systems in a more indirect manner, such as corticosteroids or sex steroids.

It is interesting to compare the mode of action of nonpsychotropic medications with what is known concerning the postulated pathophysiology of psychiatric disorders. Indeed, the mechanisms of PSEs are sometimes compatible with a given hypothesis for the corresponding spontaneous syndromes. For example, the occurrence of hallucinations and delusions on dopamine agonists (eg, levodopa) is clearly within the domain of the dopaminergic hypothesis for schizophrenia. In this case, there is a

clear relationship between the hypothesis for a disease and the postulated mechanisms of a side effect. In another example, that of depression as a side effect of interferons (IFNs), the relationship is not so easily identifiable. Stress and depression are associated with increased circulating concentrations of cytokines such as IL-1 β , IL-6, γ -IFN, and positive acute-phase proteins, and hyperactivity of the hypothalamus-pituitary-adrenal axis. Immunological activation induces "stress-like" behavioral and neurochemical changes in animals.⁴ An association of the cerebrospinal fluid (CSF) concentration of proinflammatory cytokines and major depressive disorders was reported in depressed patients with higher CSF concentrations of IL-1 β , lower IL-6, and no change in tumor necrosis factor α (TNF- α).⁵ A positive correlation was found between serum IL-1 β and the severity of depression. Other studies suggest that antidepressants can act on neuroimmunomodulation, and have been shown to shift the cytokines toward a decreased production of proinflammatory cytokines.⁶

Pharmacokinetic mechanisms

Pharmacokinetic mechanisms are relevant when the PSE is known to follow a dose-response curve. A low clearance represents the main pharmacokinetic mechanism inducing PSEs, ie, other changes in the pharmacokinetics of drugs are of little relevance. Disease states, hepatic enzyme polymorphisms, and drug interactions leading to metabolic inhibition are the main reasons for a low clearance.

Interaction by metabolic inhibition is a general principle, applicable not only to PSEs, but also to other side effects. Many drugs inhibit one or more pathways of hepatic metabolism. Cytochrome P-450 (CYP450) enzymes metabolize endogenous as well as a variety of exogenous substrates, such as toxins and drugs. Some drugs are metabolized by one metabolic pathway, others by many. When all metabolic pathways of a medication are inhibited, then the concentration of this drug will rise, favoring the occurrence of side effects.

Antifungals can inhibit some metabolic pathways, including those of mefloquine, ie, the 3A4 isoenzyme of CYP450.⁷ Mefloquine can rarely lead to serious PSEs at prophylactic doses,^{8,9} but these risks are greater at high plasma concentrations.¹⁰ The prescription of a macrolide antibiotic will probably raise concentrations of mefloquine, as most macrolides are 3A4 inhibitors. Hence, serious PSEs can occur even at usual doses of both drugs.

Risk factors

Patient-specific mechanisms of PSEs are more precisely defined as patient-related risk factors. The risk factors for developing PSEs can be medication-related or patient-related, as shown in *Table I*.

Polypharmacy is one of the most important iatrogenic risk factors for PSEs, because of the addition of pharmacological effects or due to metabolic inhibition. Addition of pharmacological effects is illustrated by the concomitant prescription of clozapine and biperiden. These drugs are both potent anticholinergics, so the risk of anticholinergic side effects is greater when they are taken together than with each medication taken alone. Polypharmacy mimics a slow metabolizer picture for many drugs, when hepatic metabolism is inhibited. There are many inhibitors of hepatic metabolism: omeprazole, cimetidine, antifungals, antivirals, HMG-CoA (3-hydroxy-3-methylglutaryl coenzyme A) reductase inhibitors (statins), antihypertensives, antiepileptics, antidepressants, grapefruit juice, and many other compounds. It is practically impossible to memorize all the CYP450 isoenzyme substrates, inhibitors, and inducers. Hence pocket tables¹¹ and software^{12,13} are useful for obtaining rapid information about such drug-drug interactions, and potentially avoiding induction of PSEs. Many PSEs are dose-dependent, so their risk increases with factors that raise the concentration of drugs. The CYP450 2D6 isoenzyme has absent or impaired activity in 7% of Caucasians^{14,15} and the 2C19 activity is absent or impaired in as many as 12% to 22% of Asians.¹⁵

Medication-related PSEs
Polypharmacy
High doses
Route of administration (eg, intravenous or intrathecal)
Faster administration (by any route)
Narrow therapeutic index
Patient-related PSEs
Present or past mental illnesses
Hepatic insufficiency, slow metabolizer, and other metabolic conditions
Augmented permeability of the blood-brain barrier (eg, meningitis or porphyria)
Very young or elderly patients
Postpartum
Other situations of stress (eg, intensive care unit)

Table I. Risk factors for psychiatric side effects (PSEs).

Diagnosis and differential diagnosis

The diagnosis of PSEs can be challenging. The clinical presentations of depressive, anxious, or psychotic PSEs meet most criteria of the *DSM-IV* for the corresponding spontaneous (noniatrogenic) syndromes. Therefore, almost any psychiatric symptom or syndrome could be considered as a potential PSE, until one has proven the contrary.

A simple case would be that of a peculiar or unusual psychiatric symptom, observed in a person who has started (or interrupted) a medical treatment recently and has no history of a previous psychiatric decompensation and no evident susceptibility to develop such a decompensation. A difficult case would be that of a person who has already suffered from many decompensations of psychiatric disorders and who develops a recurrence that presents itself clinically in a similar manner as that known for the subject. In this case, a PSE can easily be overlooked, ie, the role of a medical treatment as a relevant factor is difficult to identify. Another case is that of a physical disorder that can also induce psychiatric signs. For example, the clinician might not be able to determine whether a case of depression relates to the patient's multiple sclerosis or to the corticosteroid treatment. Another example would be a malaria patient treated with mefloquine and presenting delirium: is the delirium due to the malaria rather than to the mefloquine?

The fact of being hospitalized for a severe physical illness constitutes a strain: in an intensive care unit, the patient is exposed to pain, sleep deprivation, unusual environment, and threat of disability or even death. In this context, it is difficult to distinguish reactive or drug-induced psychological signs. Complex medical cases receive polypharmacy: some patients can receive antiarrhythmics, bronchodilators, analgesics, antibiotics, benzodiazepines, and other medications. Among these complex situations, it can become practically impossible to determine a single cause for a PSE. Yet, making such a diagnosis is necessary. For example, in systemic lupus erythematosus, the occurrence of PSEs can be due to corticosteroid treatment, but also to the lupus cerebritis; the latter is associated with high levels of antibodies to P ribosomal proteins, in both CSF and serum.¹⁶ This differential diagnosis is relevant, since the corticosteroid dose may need to be increased.

The differential diagnoses of PSEs are summarized in *Table II*. History and chronology of drug administration are first-line tools to diagnose a PSE. As already mentioned, an anamnesis with a positive exposure, positive

Pharmacological aspects

dechallenge, and positive rechallenge, indicates a high probability of a causal link between a psychiatric sign and a prescribed medication.

A PSE can differ from a spontaneous psychiatric syndrome in duration, since the duration of the PSE is more linked to the presence or withdrawal of the offending agent. Once the incriminated treatment is interrupted, behavioral symptoms usually remit within days to weeks, depending on the half-life of the substance or the presence of a withdrawal syndrome. In complex cases of polypharmacy, if the chronology of medication cannot help determine which medication caused the side effect, a trial could be done by replacing one of the suspected drugs by another with a lesser risk of PSEs.

Another issue about chronology concerns what can occur after interruption of treatment. This can be illustrated by the case of an elderly male patient, who took St John's wort for 4 months, with partial improvement of his depression. The dose was gradually increased, but without a complete remission of the depression. Travel to an endemic zone of malaria was planned and mefloquine prophylaxis was introduced. No side effect occurred during the first 10 days, until the clinician decided to replace St John's wort by citalopram, without changes in the mefloquine prophylaxis. The patient rapidly developed hallucinations after the introduction of citalopram. He had no mental status changes when he received St John's wort and mefloquine, so the clinician stopped citalopram. The hallucinations persisted. When mefloquine was discontinued, the hallucinations remitted. The message is that even the interruption of a drug can lead to an increase in the plasma concentra-

tions of another drug, causing side effects. St John's wort induces mefloquine metabolism, which means that, in this case, mefloquine concentrations were lower while St John's wort was given. Hallucinations are known side effects of mefloquine.

To improve the detection of PSEs, the physician should look for the anamnestic key factors listed below:

- Dates of occurrence of psychiatric symptoms suspected of being side effects.
- Dates of medication exposure, dechallenge, and rechallenge.
- Previous psychiatric history.
- If polypharmacy is given, dates of introduction or discontinuation of other drugs.
- Dates of factors worsening existing comorbidities.
- Plasma concentration measurements.

The most useful complementary examination for PSE investigation is generally the monitoring of plasma concentrations of suspected medications. Monitoring of drug concentration is frequently performed for some drugs with high risk of toxicity, eg, digoxin, theophylline, or lidocaine. Many other compounds can also be dosed in specialized laboratories.

If past analyses were performed for a given patient, they may also provide valuable clues. This may apply even if different medications were measured. This occurs because an abnormally high concentration of a medication may suggest a weak or absent metabolic pathway, as discussed in the mechanisms section above. Knowledge of the patient's deficiencies in metabolism allows avoidance of some PSEs by future prescriptions. Genotyping is a complementary examination to detect polymorphisms of hepatic enzymes.

Diagnoses other than medication side effects

Underlying physical illnesses with psychiatric symptoms (eg, multiple sclerosis, systemic neoplasias, electrolytic disturbances, lupus erythematosus)

Aggravation of an existing psychiatric illness

Inaugural psychiatric decompensation in individuals with no evident susceptibility

Differential diagnoses among PSEs

PSEs at usual doses

Withdrawal-related PSEs. Side effects can occur after the discontinuation of antiparkinsonian agents, benzodiazepines, antipsychotics, antidepressants, anabolic androgen steroids, etc

Intoxication-related PSEs

Table II. Differential diagnoses of psychiatric side effects (PSEs) of medications.

Description of psychiatric side effects

Table III^{3,8,9,17-197} gives a list of medications that might induce depression, mania, anxiety, or psychotic syndromes (defined by delusions and/or hallucinations). This information is qualitative, in the sense that the severity or the frequency of these side effects under each medication or class is not indicated. Specific information can be found in the bibliography. Some psychotropics, such as benzodiazepines, are listed in *Table III* because they are frequently prescribed in internal medicine. Obviously, more than one of these PSEs can occur in a given patient. For example, many depressive states are accompanied by anxiety. Some clinically relevant examples of medications presented in *Table III* are discussed below in more detail.

Mefloquine and chloroquine

Mefloquine, which is prescribed for the prophylaxis or treatment of malaria, frequently causes PSEs. These PSEs can be severe: psychosis, delusion, and even suicidal ideation. Disabling PSEs occur in less than 1% of patients under mefloquine at therapeutic doses, and in less than

1:10 000 under mefloquine prophylaxis. This indicates that the PSEs are dose-related. However, suicide attempts have even been reported at prophylactic doses.^{158,159} Mefloquine PSEs may begin some hours after the first dose.^{8,160} Their mechanism is not clearly understood; actions as a *N*-methyl-D-aspartate (NMDA) receptor antagonist or on sigma receptors have been proposed. Chloroquine, another antimalarial, also produces psychiatric and neurological side effects: agitation, aggressiveness, amnesia, confusion, depression, hallucinations, and mania. Psychiatric changes under chloroquine may develop insidiously. Memory or perception changes can be the only clues to side effects in this developing phase. The half-life of chloroquine is long at around 1 month. Therefore, remission of a chloroquine-related PSE may take days. Children appear to be at greatest risk to psychosis under chloroquine treatment.⁸¹

	Depression	Mania	Anxiety	Psychotic symptoms
Amantadine ¹⁷⁻²²	X	X	X	X
Aminoglycosides ^{23,24}				X
Amphetamines ²⁵⁻³⁰	X	X	X	X
Anabolic steroids ³¹⁻³³	X	X	X	X
Anesthetics ^{21,22,34,35}		X		X
Anticholinergics ^{21,36-39}	X		X	X
Antihistamines ^{21,22,40}	X	X		
Antitubercular agents ^{22,41-45}	X		X	X
Antivirals ⁴⁶⁻⁵³	X		X	X
Baclofen ^{22,54-58}	X	X	X	X
Barbiturates ^{22,59}	X	X	X	X
Benzodiazepines ^{21,22,60-62}		X	X	X
β-Blockers ^{21,22,63-72}	X	X	X	X
Bromocriptine ^{21,73-77}		X	X	X
Cephalosporins ^{22,78-80}		X		X
Chloroquine ⁸¹⁻⁸⁸	X	X	X	X
Clonidine ^{21,22,89-93}	X	X	X	X
Corticosteroids ^{3,21,22,94-112}	X	X	X	X
Digoxin ^{21,22,113-115}	X	X		X
Disulfiram ^{21,22,116-118}	X	X	X	X
Interferon-α ^{22,119-131}	X	X	X	X
Isotretinoin ^{22,132-137}	X			X
Levodopa ^{21,22,138-152}	X	X	X	X
Lidocaine ¹⁵³⁻¹⁵⁷	X	X	X	X
Mefloquine ^{8,9,22,158-160}	X	X	X	X
Methylidopa ^{21,22}	X		X	X
Methylphenidate ^{22,161-163}		X	X	X
Metoclopramide ^{21,22,164-169}	X	X	X	
Metronidazole ^{22,170-175}				X
Opioids ^{21,22,176-181}	X	X	X	X
Oral contraceptives ^{21,22,182,183}	X		X	
Procainamide ^{21,184-188}	X	X	X	X
Pseudoephedrine ^{22,189-192}			X	X
Quinidine ^{21,22,193,194}			X	X
Quinolones ^{22,195}	X		X	X
Thiazide diuretics ^{22,196}	X			

Table III. Psychiatric side effects potentially induced by pharmacological treatment.

Metronidazole

The common side effects of metronidazole are confusion, paresthesia, dizziness, vertigo, and syncope. Seizures and encephalopathy are rare. Psychotic symptoms have also been described.^{170,171} Brain toxicity can occur at usual doses of metronidazole. However, at larger or chronic doses, an increased prevalence (as high as 25%) has been observed. An interaction between metronidazole and alcohol, similar to the disulfiram effect, was first described in the 1960s,¹⁷² and many publications on this topic followed. One death has been attributed to this interaction.¹⁷³ This PSE was believed to result from a metronidazole blockade of hepatic aldehyde dehydrogenase, an enzyme in the metabolism of ethanol, followed by the accumulation of acetaldehyde in the blood. However, these early publications have been called into question.¹⁷⁴ A placebo-controlled study¹⁹⁷ found no effect on blood acetaldehyde concentrations when 12 subjects ingested metronidazole with ethanol; it induced no objective or subjective disulfiram-like effects.

Isotretinoin

The Food and Drug Administration (FDA) has received reports of depression and suicide attempts in patients treated with isotretinoin. The authors who commented these cases¹³² affirm that “factors suggesting a possible association between isotretinoin and depression include a temporal association between use of the drug and depression, positive dechallenges (often with psychiatric treatment), positive rechallenges, and possible biologic plausibility.

Pharmacological aspects

Compared with all drugs in the FDA's Adverse Event Reporting System database to June 2000, isotretinoin ranked within the top 10 for number of reports of depression and suicide attempt.¹⁹⁸ An American drug database¹⁹⁸ also warns that isotretinoin "may cause depression, psychosis and, rarely, suicidal ideation, suicide attempts, suicide, and aggressive and/or violent behaviors." Despite some publications contrary to this association,^{199,200} the possibility of depression and suicidal ideation should be taken into account when prescribing isotretinoin. Establishing a causality link between isotretinoin and these PSEs can be complex. Most isotretinoin-treated patients are young, and young people are also at greater risk for depression. Another aspect is that disfiguring dermatological conditions could be associated with a greater prevalence of depression and suicidal ideation,²⁰¹ while isotretinoin improves the aspect of patients suffering from severe forms of acne. The improvement in these severe forms of acne after isotretinoin treatment may reduce anxiety and depression.²⁰²

Interferons

IFN- α is prescribed in viral hepatitis and in some neoplasias; IFN- β is prescribed in multiple sclerosis and hepatitis. Frequent side effects limit their use. IFN- α induces more PSEs than IFN- β . Psychiatric manifestations of IFNs are depression, personality disorders, panic attacks, other anxiety states, manic and psychotic symptoms,¹¹⁹ impulsiveness, and aggressive behavior.²⁰³ More than 10% of patients receiving IFN- α manifest PSEs.¹²⁰⁻¹²² Depressive states related to IFNs usually occur in the first weeks of treatment. They are more prevalent and severe in people who also suffered from depression before IFN treatment. Suicidal behavior is an alleged side effect of IFN- α .¹²³⁻¹²⁷ IFN- α -related depression or suicidal behavior may continue after interruption of treatment.^{123,126} This feature has rarely been attributed to IFN- β .^{50,204} Since suicide attempts were described after withdrawal of IFN- α , even without a depressive episode during the treatment, some authors¹²³ advise psychiatric supervision "even more frequently after interferon withdrawal." A randomized controlled trial¹²¹ found a favorable effect upon the prescription of a selective serotonin reuptake inhibitor (SSRI) as prophylaxis of depression in patients who are programmed to receive IFN- α . SSRIs are also useful for treatment of depression,¹²⁸ once IFN- α is started. Paroxetine is the most studied,^{121,128} but other

SSRIs, such as sertraline,¹²⁹ citalopram,¹³⁰ fluoxetine,¹³¹ and fluvoxamine may also be effective. There is no consensus on whether SSRIs should be given as prophylaxis for all patients programmed to receive IFN- α or only to those who develop depression.

Corticosteroids

Corticosteroid treatment may lead to many PSEs. The most frequent PSEs are depression, mania, anxiety, insomnia, delusions (paranoia or other themes), hallucinations, agitation, and confusional states. Rarer ones include serious heteroaggressivity, disturbances of consciousness, and depersonalization.^{3,95,205} These PSEs can start after just 1 day of treatment. In more than half of patients, side effects usually remit after interruption of corticosteroids.²⁰⁶ However, side effects can also occur during the withdrawal period, eg, anhedonia and fatigue may last several weeks.^{207,208} The risk of PSEs is high: about 6% of patients manifest some PSEs. Prednisone is the most implicated corticosteroid, but PSEs were also described with methylprednisolone, dexamethasone, and beclomethasone. A dose-response effect is clearly seen with prednisone: doses greater than 40 mg/day are related to greater psychiatric morbidity. Moreover, hypoalbuminemia leads to an increase in plasma prednisone free fraction, potentially increasing the amount of prednisone that reaches brain. Hence, hypoalbuminemia might be associated with an increase in PSE incidence, but this has not been confirmed. Small neuroleptic doses can lead to a favorable response of corticosteroid PSEs in some days.^{94,206} Treatment with lithium may be helpful.⁹⁴ Prophylaxis of corticosteroid PSEs with lithium⁹⁶⁻⁹⁸ or valproate⁹⁹ has been described, but these approaches can be harmful to patients on corticosteroids who might not develop PSEs. Patients on corticosteroids should be monitored for psychiatric and cognitive side effects.³¹

Anabolic androgen steroids

Anabolic androgen steroids (testosterone and its synthetic derivatives) are associated with a number of PSEs: impulsiveness, irritability, and belligerence ("steroid rage"), delusions, hypersexuality, auditory and visual hallucinations, and dependence.³¹ Mania, hypomania, and major depression have been significantly associated with exposure to steroids.³² PSEs due to anabolic androgen steroids are mostly seen in abusive users. These PSEs relate to drug

concentrations in a definite pattern. Hypomania is correlated with anabolic androgen intake and major depression follows its withdrawal.³¹ In chronic users of slow-liberation forms, lassitude or depression may be seen just before administration of the next dose. The abuse of anabolic androgen steroids seems prevalent among teenagers wishing to increase muscular mass. It was found in subjects as young as 9 years old,²⁰⁹ with a possible peak at ages 15 and 16.²¹⁰ In another study, with a sample of 12 000 American high-school students, a prevalence of 4% was found in young males.²¹¹ Steroid users often seek medical care for the acne these medications induce or exacerbate. If family members complain of aggressiveness and mood changes (which are less noticed by the users themselves), the clinician might suspect of anabolic androgen abuse, especially in teenagers of male sex.

β-Adrenergic antagonists (β-blockers)

Depression, nightmares, and sexual dysfunction are commonly reported PSEs of β-blockers. Hallucinations have been attributed to propranolol.⁶³⁻⁶⁸ With oral administration, depression and agitation related to propranolol might be dose-dependent.⁶⁹ Ophthalmic preparations of β-blockers may also induce these PSEs, eg, timolol.⁷⁰ Withdrawal reactions to β-blockers can occur even with ophthalmic presentations; cases of rebound tachycardia were reported after ophthalmic timolol interruption. It has long been recognized that β-blockers cause psychiatric and sexual side effects. However, this has become controversial, according to recent studies. In a placebo-controlled trial, the authors found no difference between propranolol and placebo groups for the occurrence of depressive symptoms or sexual dysfunction.²¹² A later review stated that “β-blockers have no significant increased risk of depressive symptoms and only small increased risks of fatigue and sexual dys-

function.”²¹³ Nevertheless, it could be that the risk of suicide increases in users of β-blockers.⁷¹ The conflicting results on β-blocker depression suggest that some may improve depression (eg, pindolol), others may worsen it, and others may have little effect.²¹⁴

Conclusion

This review shows that drug-induced PSEs may occur with several medications prescribed in internal medicine and that these side effects might be overlooked. A PSE can be a stressful and traumatic life event for patients and their families. For example, a person without known psychiatric antecedents who develops a drug-induced psychosis might suffer sequelae from the fear of having lost their mind or from hospitalization in a psychiatric ward. In given cases, a PSE might even lead to a post-traumatic stress disorder, either because of the severity of the PSE itself, or because of a deficient explanation of the side effect to the patient.

Another issue is the importance of making a correct diagnosis of a PSE. This is relevant for several reasons. The main one is probably to help patients to make adequate attributions and conclusions concerning their psychological changes. Indeed, for someone who experiences a PSE, knowing that it is a side effect has a different significance than wondering if oneself is mentally insane. Detecting a PSE avoids its confusion with a sign of a psychiatric disease; since spontaneous psychiatric diseases frequently require long-term treatment, the correct diagnosis of a PSE can spare the patient the stigma, distress, and other costs of an unjustified long-term psychiatric treatment. Finally, the correct diagnosis of a PSE also enables the prescriber to communicate suspected side effects to the organization responsible for pharmacovigilance. □

REFERENCES

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. Washington, DC: American Psychiatric Association; 1994.
2. World Health Organization. *The ICD-10 Classification of Mental and Behavioral Disorders. Clinical descriptions and diagnostic guidelines*. Geneva, Switzerland: World Health Organization; 1992.
3. Kershner P, Wang-Cheng R. Psychiatric side effects of steroid therapy. *Psychosomatics*. 1989;30:135-139.
4. Connor TJ, Leonard BE. Depression, stress and immunological activation: the role of cytokines in depressive disorders. *Life Sci*. 1998;62:583-606.
5. Levine J, Barak Y, Chengappa KN, Rapoport A, Rebey M, Barak V. Cerebrospinal cytokine levels in patients with acute depression. *Neuropsychobiology*. 1999;40:171-176.
6. Connor TJ, Harkin A, Kelly JP, Leonard BE. Olfactory bulbectomy provokes a suppression of interleukin-1beta and tumour necrosis factor-alpha production in response to an in vivo challenge with lipopolysaccharide: effect of chronic desipramine treatment. *Neuroimmunomodulation*. 2000;7:27-35.
7. Ridditid W, Wongnawa M, Mahatthanatrakul W, Chaipol P, Sunbhanich M. Effect of rifampin on plasma concentrations of mefloquine in healthy volunteers. *J Pharm*. 2000;52:1265-1269.
8. Rodor F, Bianchi G, Grignon S, Samuelian JC, Jouglard J. Recurrent psychiatric manifestations during malaria prevention with mefloquine. A case report [in French]. *Thérapie*. 1990;45:433-434.

Pharmacological aspects

Efectos secundarios psiquiátricos de los medicamentos prescritos en medicina interna

Varios tratamientos farmacológicos utilizados en medicina interna pueden inducir efectos secundarios psiquiátricos (ESP) que simulan diagnósticos observados en psiquiatría. Los ESP se pueden producir durante la retirada de un medicamento, por una intoxicación y también cuando se utiliza en dosis terapéuticas habituales. Entre los fármacos que pueden inducir síndromes depresivos, ansiosos o psicóticos están los corticoides, la isotretinoína, la levodopa, la mefloquina, el interferon alfa y los esteroides anabólicos, al igual que algunos medicamentos de venta libre. Los ESP a menudo son difíciles de diagnosticar y pueden ser muy peligrosos para los pacientes. En esta revisión se discuten los ESP y también los elementos diagnósticos que facilitan su identificación.

Effets secondaires psychiatriques des médicaments prescrits en médecine interne

Plusieurs traitements pharmacologiques en médecine interne peuvent entraîner des effets secondaires psychiatriques (ESP) simulant des pathologies vues en psychiatrie. Ces ESP peuvent survenir lors du sevrage d'un médicament, lors d'intoxication ou aux doses thérapeutiques usuelles. Parmi les médicaments qui peuvent induire des syndromes dépressifs, anxieux ou psychotiques, on note les corticoïdes, l'isotrétinoïne, la lévodopa, la méfloquine, l'interféron alpha et les stéroïdes anabolisants, tout comme quelques médicaments en vente libre. Les ESP sont souvent difficiles à diagnostiquer et peuvent être très délétères pour les patients. Cet article passe en revue les ESP, ainsi que les éléments de diagnostic qui facilitent leur identification.

- Lysack JT, Lysack CL, Kvern BL. A severe adverse reaction to mefloquine and chloroquine prophylaxis. *Aust Fam Physician*. 1998;27:1119-1120.
- Dukes MNG, Aronson JK, eds. *Meyler's Side Effects of Drugs*. 14th ed. Amsterdam, The Netherlands: Elsevier; 2000:956-957.
- Flockhart DA. Cytochrome P450 drug interaction table. Homepage of the Indiana University Department of Medicine. Available at <http://medicine.iupui.edu/flockhart>, December 2002. Accessed 18 April 2003.
- Strain JJ, Chiu NM, Brodsky M, Karim A, Caliendo G. Comparison of three methods for identifying medical drug-psychotropic drug interactions. *Gen Hosp Psychiatry*. 2002;24:311-315.
- Sittig DF, Jimison HB, Hazlehurst BL, et al. Techniques for identifying the applicability of new information management technologies in the clinical setting: an example focusing on handheld computers. *Proc AMIA Symp*. 2000:804-808.
- Marsh JC, Chowdry J, Parry-Jones N, et al. Study of the association between cytochromes P450 2D6 and 2E1 genotypes and the risk of drug and chemical induced idiosyncratic aplastic anaemia. *Br J Haematol*. 1999;104:266-270.
- Bertilsson L. Geographical/interracial differences in polymorphic drug oxidation. Current state of knowledge of cytochromes P450 (CYP) 2D6 and 2C19. *Clin Pharmacokinet*. 1995;29:192-209.
- Kohen M, Asherson RA, Gharavi AE, Lahita RG. Lupus psychosis: differentiation from the steroid-induced state. *Clin Exp Rheumatol*. 1993;11:323-326.
- Hausner RS. Amantadine-associated recurrence of psychosis. *Am J Psychiatry*. 1980;137:240-242.
- Lieb K, Hufert FT, Bechter K, Bauer J, Kornhuber J. Depression, Borna disease, and amantadine. *Lancet*. 1997;349:958.
- Rego MD, Giller EL Jr. Mania secondary to amantadine treatment of neuroleptic-induced hyperprolactinemia. *J Clin Psychiatry*. 1989;50:143-144.
- Muller HF, Dastoor DP, Klingner A, Cole M, Boillat J. Amantadine in senile dementia: electroencephalographic and clinical effects. *J Am Geriatr Soc*. 1979;27:9-16.
- Fann JR. Psychiatric effects of neuropharmacological agents. *Semin Clin Neuropsychiatry*. 2002;7:206-212.
- [No authors listed]. Drugs that may cause psychiatric symptoms. *Med Lett Drugs Ther*. 2002;44:59-62.
- Tolnai A, Barcs I, Nagy E, Kali G. Complication of aminoglycoside therapy: acute psychosis and renal insufficiency. *Orv Hetil*. 1984;125:3047-3050.
- Kane FJ Jr, Byrd G. Acute toxic psychosis associated with gentamicin therapy. *South Med J*. 1975;68:1283-1285.
- Srisurapanont M, Kittiratanapaiboon P, Jarusuraisin N. Treatment for amphetamine psychosis. *Cochrane Database Syst Rev*. 2001;4:CD003026.
- Yui K, Ikemoto S, Ishiguro T, Goto K. Studies of amphetamine or methamphetamine psychosis in Japan: relation of methamphetamine psychosis to schizophrenia. *Ann N Y Acad Sci*. 2000;914:1-12.
- Flaum M, Schultz SK. When does amphetamine-induced psychosis become schizophrenia? *Am J Psychiatry*. 1996;153:812-815.
- Caplehorn JR. Amphetamine psychosis. *Br J Addict*. 1990;85:1505-1506.
- Hall RC, Popkin MK, Beresford TP, Hall AK. Amphetamine psychosis: clinical presentations and differential diagnosis. *Psychiatr Med*. 1988;6:73-79.
- Kokkinidis L, Zacharko RM. Response sensitization and depression following long-term amphetamine treatment in a self-stimulation paradigm. *Psychopharmacology (Berl)*. 1980;68:73-76.
- Malone DA Jr, Dimeff RJ, Lombardo JA, Sample RH. Psychiatric effects and psychoactive substance use in anabolic-androgenic steroid users. *Clin J Sport Med*. 1995;5:25-31.
- Pope HG Jr, Katz DL. Psychiatric and medical effects of anabolic-androgenic steroid use. A controlled study of 160 athletes. *Arch Gen Psychiatry*. 1994;51:375-382.
- Weiss EL, Bowers MB Jr, Mazure CM. Testosterone-patch-induced psychotic mania. *Am J Psychiatry*. 1999;156:969.
- Oga K, Kojima T, Matsuura M, et al. Effects of low-dose ketamine on neuropathic pain: an electroencephalogram-electrooculogram/behavioral study. *Psychiatry Clin Neurosci*. 2002;56:355-363.
- Pal HR, Berry N, Kumar R, Ray R. Ketamine dependence. *Anaesth Intensive Care*. 2002;30:382-384.
- Christensen RC. Misdiagnosis of anticholinergic delirium as schizophrenic psychosis. *Am J Emerg Med*. 1995;13:117-118.
- Hvizdos AJ, Bennett JA, Wells BG, Rappaport KB, Mendel SA. Anticholinergic psychosis in a patient receiving usual doses of haloperidol, desipramine, and benztropine. *Clin Pharm*. 1983;2:174-178.
- Hall RC, Feinsilver DL, Holt RE. Anticholinergic psychosis: differential diagnosis and management. *Psychosomatics*. 1981;22:581-583; 587.
- Perry PJ, Wilding DC, Juhl RP. Anticholinergic psychosis. *Am J Hosp Pharm*. 1978;35:725-728.

40. Lee JH, Turndorf H, Poppers PJ. Physostigmine reversal of antihistamine-induced excitement and depression. *Anesthesiology*. 1975;43:683-684.
41. Iannaccone R, Sue YJ, Avner JR. Suicidal psychosis secondary to isoniazid. *Pediatr Emerg Care*. 2002;18:25-27.
42. Djibo A, Lawan A. Behavioral disorders after treatment with isoniazid. *Bull Soc Pathol Exot*. 2001;94:112-114.
43. Chan TY. Pyridoxine ineffective in isoniazid-induced psychosis. *Ann Pharmacother*. 1999;33:1123-1124.
44. Hsu CW, Chu KA, Lu T, Lai RS, Lu JY. Ethambutol-induced psychosis: a case report. *Zhonghua Yi Xue Za Zhi (Taipei)*. 1999;62:724-727.
45. Salafia A, Candida. Rifampicin induced flu-syndrome and toxic psychosis. *Indian J Lepr*. 1992;64:537-539.
46. Poulsen HD, Lublin HK. Efavirenz-induced psychosis leading to involuntary detention. *AIDS*. 2003;17:451-453.
47. de la Garza CL, Paoletti-Duarte S, Garcia-Martin C, Gutierrez-Casares JR. Efavirenz-induced psychosis. *AIDS*. 2001;15:1911-1912.
48. Hansen BA, Greenberg KS, Richter JA. Ganciclovir-induced psychosis. *N Engl J Med*. 1996;335:1397.
49. Bonaccorso S, Marino V, Biondi M, Grimaldi F, Ippoliti F, Maes M. Depression induced by treatment with interferon-alpha in patients affected by hepatitis C virus. *J Affect Disord*. 2002;72:237-241.
50. Lana-Peixoto MA, Teixeira AL Jr, Haase VG. Interferon beta-1a-induced depression and suicidal ideation in multiple sclerosis. *Arq Neuropsiquiatr*. 2002;60:721-724.
51. Tamam L, Yerdelen D, Ozpoyraz N. Psychosis associated with interferon-alpha therapy for chronic hepatitis B. *Ann Pharmacother*. 2003;37:384-387.
52. Garcia-Pares G, Domenech C, Gil M. Psychosis induced by interferon-alpha. *Psychosomatics*. 2002;43:428-429.
53. Kornischka J, Burtscheidt W, Gaebel W. Interferon-induced paranoid psychosis. Review of the literature and case report. *Nervenarzt*. 2002;73:463-467.
54. Nakagawa Y, Ishima T, Ishibashi Y, Tsuji M, Takashima T. Involvement of GABA_B receptor systems in experimental depression: baclofen but not bicuculline exacerbates helplessness in rats. *Brain Res*. 1996;741:240-245.
55. Sommer BR, Petrides G. A case of baclofen-induced psychotic depression. *J Clin Psychiatry*. 1992;53:211-212.
56. Yassa RY, Iskandar HL. Baclofen-induced psychosis: two cases and a review. *J Clin Psychiatry*. 1988;49:318-320.
57. Kirubakaran V, Mayfield D, Rengachary S. Dyskinesia and psychosis in a patient following baclofen withdrawal. *Am J Psychiatry*. 1984;141:692-693.
58. Arnold ES, Rudd SM, Kirshner H. Manic psychosis following rapid withdrawal from baclofen. *Am J Psychiatry*. 1980;137:1466-1467.
59. Sarrechia C, Sordillo P, Conte G, Rocchi G. Barbiturate withdrawal syndrome: a case associated with the abuse of a headache medication [in Italian]. *Ann Ital Med Int*. 1998;13:237-239.
60. Modell JG. Protracted benzodiazepine withdrawal syndrome mimicking psychotic depression. *Psychosomatics*. 1997;38:160-161.
61. Terao T, Tani Y. Two cases of psychotic state following normal-dose benzodiazepine withdrawal. *J UOEH*. 1988;10:337-340.
62. Hugues FC, Jouglard J, Le Jeune C, Moulin M, Begaud B. Psychotic disorders linked to the inhibition of benzodiazepine catabolism. *Rev Med Intern*. 1987;8:433-436.
63. Love JN, Handler JA. Toxic psychosis: an unusual presentation of propranolol intoxication. *Am J Emerg Med*. 1995;13:536-537.
64. Cunnane JG, Blackwood GW. Psychosis with propranolol: still not recognized? *Postgrad Med J*. 1987;63:57-58.
65. Parker WA. Propranolol-induced depression and psychosis. *Clin Pharm*. 1985;4:214-218.
66. McGahan DJ, Wojslaw A, Prasad V, Blankenship S. Propranolol-induced psychosis. *Drug Intell Clin Pharm*. 1984;18:601-603.
67. Remick RA, O'Kane J, Sparling TG. A case report of toxic psychosis with low-dose propranolol therapy. *Am J Psychiatry*. 1981;138:850-851.
68. Viadero JJ, Wong SH, White WB. Acute psychotic behavior associated with atenolol. *Am J Psychiatry*. 1983;140:1382.
69. Petrie WM, Maffucci RJ, Woosley RL. Propranolol and depression. *Am J Psychiatry*. 1982;139:92-94.
70. Shore JH, Fraunfelder FT, Meyer SM. Psychiatric side effects from topical ocular timolol, a beta-adrenergic blocker. *J Clin Psychopharmacol*. 1987;7:264-267.
71. Sorensen HT, Mellekjaer L, Olsen JH. Risk of suicide in users of beta-adrenoceptor blockers, calcium channel blockers and angiotensin-converting enzyme inhibitors. *Br J Clin Pharmacol*. 2001;52:313-318.
72. Schweitzer I, Maguire K, Tuckwell V. Antiglaucoma medication and clinical depression. *Aust N Z J Psychiatry*. 2001;35:569-571.
73. Reeves RR, Pinkofsky HB. Postpartum psychosis induced by bromocriptine and pseudoephedrine. *J Fam Pract*. 1997;45:164-166.
74. Jimeno Valdes A, Ruiz Otazo AM, Jimeno Bulnes N. A case of toxic psychosis caused by bromocriptine [in Spanish]. *Actas Luso Esp Neurol Psiquiatr Cienc Afines*. 1989;17:386-390.
75. Iffy L, Lindenthal J, Szodi Z, Griffin W. Puerperal psychosis following ablation with bromocriptine. *Med Law*. 1989;8:171-174.
76. Canterbury RJ, Haskins B, Kahn N, Saathoff G, Yazel JJ. Postpartum psychosis induced by bromocriptine. *South Med J*. 1987;80:1463-1464.
77. Boyd A. Bromocriptine and psychosis: a literature review. *Psychiatr Q*. 1995;66:87-95.
78. Vincken W. Psychotic reaction to cefuroxime. *Lancet*. 1984;1:965.
79. Saker BM, Musk AW, Haywood EF, Hurst PE. Reversible toxic psychosis after cephalixin. *Med J Aust*. 1973;1:497-498.
80. Heinecke G, Hoffler U, Finke K. Reversible encephalopathy following cephacetril therapy in high doses in a patient on chronic intermittent hemodialysis. *Clin Nephrol*. 1976;5:45-47.
81. Garg P, Mody P, Lall KB. Toxic psychosis due to chloroquine—not uncommon in children. *Clin Pediatr*. 1990;29:448-450.
82. Nicolas X, Granier H, Adam F, Laborde JP, Talarmin F, Klotz F. Acute psychosis following treatment with chloroquine for a primo-invasion of *Plasmodium vivax* malaria [in French]. *Presse Med*. 2003;32:117.
83. Bhatia MS. Chloroquine-induced recurrent psychosis (brief report). *Indian J Med Sci*. 1996;50:302-304.
84. Masson C, Cheron F, Gautier MS, Cambier J. Acute psychosis after ingestion of chloroquine. *Presse Med*. 1990;19:334.
85. Agrawal A, Agrawal R. Chloroquine-induced psychosis. *Indian Pediatr*. 1988;25:1226-1227.
86. Akhtar S, Mukherjee S. Chloroquine-induced mania. *Int J Psychiatry Med*. 1993;23:349-356.
87. Lovestone S. Chloroquine-induced mania. *Br J Psychiatry*. 1991;159:164-165.
88. Boudreau E, Schuster B, Sanchez J, et al. Tolerability of prophylactic Larium regimens. *Trop Med Parasitol*. 1993;44:257-265.
89. Enginar N, Eroglu L. The long-term clonidine treatment induced behavioral depression in rats. *Pol J Pharmacol Pharm*. 1990;42:409-415.
90. Kostowski W, Obersztyń M. Chronic administration of desipramine and imipramine but not zimelidine attenuates clonidine-induced depression of avoidance behavior in rats. *Pol J Pharmacol Pharm*. 1988;40:341-349.
91. Terkelsen KG. Mania after surreptitious discontinuation of clonidine. *Am J Psychiatry*. 1984;141:153.
92. Adler LE, Bell J, Kirch D, Friedrich E, Freedman R. Psychosis associated with clonidine withdrawal. *Am J Psychiatry*. 1982;139:110-112.
93. Elizur A, Liberson Z. An acute psychotic episode at the beginning of clonidine therapy. 1. *Prog Neuropsychopharmacol*. 1980;4:211-213.
94. Brown ES, Khan DA, Nejtck VA. The psychiatric side effects of corticosteroids. *Ann Allergy Asthma Immunol*. 1999;83(6 Pt 1):495-503. Quiz: 503-504.
95. Lewis DA, Smith RE. Steroid-induced psychiatric syndromes—a report of 14 cases and a review of the literature. *J Affect Disord*. 1983;5:319-332.
96. Merrill W. Case 35-1998: use of lithium to prevent corticosteroid-induced mania. *N Engl J Med*. 1999;340:1123.
97. Goggans FC, Weisberg LJ, Koran LM. Lithium prophylaxis of prednisone psychosis: a case report. *J Clin Psychiatry*. 1983;44:111-112.
98. Sabet-Sharghi F, Hutzler JC. Prophylaxis of steroid-induced psychiatric syndromes. *Psychosomatics*. 1990;31:113-114.
99. Abbas A, Styra R. Valproate prophylaxis against steroid-induced psychosis. *Can J Psychiatry*. 1994;39:188-189.
100. Patten SB. Exogenous corticosteroids and major depression in the general population. *J Psychosom Res*. 2000;49:447-449.
101. Patten SB, Williams JV, Love EJ. Self-reported depressive symptoms following treatment with corticosteroids and sedative-hypnotics. *Int J Psychiatry Med*. 1996;26:15-24.

Pharmacological aspects

102. DeSilva CC, Nurse MC, Vokey K. Steroid-induced psychosis treated with risperidone. *Can J Psychiatry*. 2002;47:388-389.
103. Lopez-Medrano F, Cervera R, Trejo O, Font J, Ingelmo M. Steroid-induced psychosis in systemic lupus erythematosus: a possible role of serum albumin level. *Ann Rheum Dis*. 2002;61:562-563.
104. Mesurolle B, Ariche M, Cohen D. Premedication before IV contrast-enhanced CT resulting in steroid-induced psychosis. *AJR Am J Roentgenol*. 2002;178:766-767.
105. Lee KM, Lin YZ, Huang FY. Steroid-induced acute psychosis in a child with asthma: report of one case. *Acta Paediatr Taiwan*. 2001;42:169-171.
106. Dawson KL, Carter ER. A steroid-induced acute psychosis in a child with asthma. *Pediatr Pulmonol*. 1998;26:362-364.
107. Brockington IF, Brownell LW. Steroid-induced prepartum psychosis. *Br J Psychiatry*. 1997;170:90.
108. Bloch M, Gur E, Shalev A. Chlorpromazine prophylaxis of steroid-induced psychosis. *Gen Hosp Psychiatry*. 1994;16:42-44.
109. Mullen RS, Romans-Clarkson SE. Behavioural sensitisation and steroid-induced psychosis. *Br J Psychiatry*. 1993;162:549-551.
110. Wada K, Yamada N, Sato T, et al. Corticosteroid-induced psychotic and mood disorders: diagnosis defined by DSM-IV and clinical pictures. *Psychosomatics*. 2001;42:461-466.
111. Sutor B, Wells LA, Rummans TA. Steroid-induced depressive psychosis responsive to electroconvulsive therapy. *Convuls Ther*. 1996;12:104-107.
112. Pies R. Differential diagnosis and treatment of steroid-induced affective syndromes. *Gen Hosp Psychiatry*. 1995;17:353-361.
113. Carney MW, Rapp S, Pearce K. Digoxin toxicity presenting with psychosis in a patient with chronic phobic anxiety. *Clin Neuropharmacol*. 1985;8:193-195.
114. Singh RB, Singh VP, Somani PN. Psychosis: a rare manifestation of digoxin intoxication. *J Indian Med Assoc*. 1977;69:62-63.
115. Chyrek R, Jablęcka A, Pupek-Musialik D, Lowicki Z. Digoxin as a cause of chromatopsia and depression in a patient with heart failure and hyperthyroidism. *Pol Merkuriusz Lek*. 2000;9:563-564.
116. Murthy KK. Psychosis during disulfiram therapy for alcoholism. *J Indian Med Assoc*. 1997;95:80-81.
117. Rossiter SK. Psychosis with disulfiram prescribed under probation order. *BMJ*. 1992;305:763.
118. Goyer PF, Brown GL, Minichiello MD, Major LF. Mood-altering effects of disulfiram in alcoholics. *J Stud Alcohol*. 1984;45:209-213.
119. Debieu C, De Chouly De Lenclave MB, Foutrein P, Bailly D. Alpha-interferon and mental disorders [in French]. *Encéphale*. 2001;27:308-317.
120. Castera L, Zigante F, Bastie A, Buffet C, Dhumeaux D, Hardy P. Incidence of interferon- α -induced depression in patients with chronic hepatitis C. *Hepatology*. 2002;35:978-979.
121. Musselman DL, Lawson DH, Gumnick JF, et al. Paroxetine for the prevention of depression induced by high-dose interferon- α . *N Engl J Med*. 2001;344:961-966.
122. Renault PF, Hoofnagle JH, Park Y, et al. Psychiatric complications of long-term interferon- α therapy. *Arch Intern Med*. 1987;147:1577-1580.
123. Koseki K, Nakano M, Takaiwa M, Kamata T, Yosida J. Suicidal attempts in three postoperative patients with renal cancer after alpha interferon withdrawal. *Nippon Hinyokika Gakkai Zasshi*. 2000;91:29-32.
124. Windemuth D, Bacharach-Buhles M, Hoffmann K, Altmeyer P. Depression and suicidal intentions as a side effect of high dosage interferon- α therapy—two cases. *Hautarzt*. 1999;50:266-269.
125. Fukunishi K, Tanaka H, Maruyama J, et al. Burns in a suicide attempt related to psychiatric side effects of interferon. *Burns*. 1998;24:581-583.
126. Rifflet H, Vuillemin E, Oberti F, et al. Suicidal impulses in patients with chronic viral hepatitis C during or after therapy with interferon alpha. *Gastroenterol Clin Biol*. 1998;22:353-357.
127. Bacq Y, Tantaoui Elaraki A, Metman EH. Suicide in a patient with chronic viral hepatitis C treated with interferon- α : favoring role of alcohol? *Gastroenterol Clin Biol*. 1997;21:797-798.
128. Kraus MR, Schafer A, Faller H, Csef H, Scheurlen M. Paroxetine for the treatment of interferon- α -induced depression in chronic hepatitis C. *Aliment Pharmacol Ther*. 2002;16:1091-1099.
129. Schramm TM, Lawford BR, Macdonald GA, Cooksley WG. Sertraline treatment of interferon- α -induced depressive disorder. *Med J Aust*. 2000;173:359-361.
130. Farah A. Interferon-induced depression treated with citalopram. *J Clin Psychiatry*. 2002;63:166-167.
131. Levenson JL, Fallon HJ. Fluoxetine treatment of depression caused by interferon- α . *Am J Gastroenterol*. 1993;88:760-761.
132. Wysowski DK, Pitts M, Beitz J. An analysis of reports of depression and suicide in patients treated with isotretinoin. *J Am Acad Dermatol*. 2001;45:515-519.
133. Wysowski DK, Beitz J. Methodological limitations of the study "Isotretinoin use and risk of depression, psychotic symptoms, suicide, and attempted suicide." *Arch Dermatol*. 2001;137:1102-1103.
134. Wysowski DK, Pitts M, Beitz J. Depression and suicide in patients treated with isotretinoin. *N Engl J Med*. 2001;344:460.
135. O'Donnell J. Overview of existing research and information linking isotretinoin (Accutane), depression, psychosis, and suicide. *Am J Ther*. 2003;10:148-159.
136. Ng CH, Tam MM, Hook SJ. Acne, isotretinoin treatment and acute depression. *World J Biol Psychiatry*. 2001;2:159-161.
137. Ng CH, Schweitzer I. The association between depression and isotretinoin use in acne. *Aust N Z J Psychiatry*. 2003;37:78-84.
138. Svetel MV, Sternic NM, Filipovic SR, Vojvodic NM, Kostic VS. Clozapine in the treatment of adverse psychiatric manifestations of long-term therapy with levodopa [in Serbo-Croatian]. *Srp Arh Celok Lek*. 1997;125:203-206.
139. Bowers MB Jr. Plasma HVA and MHPG in psychiatric patients with renal failure and levodopa-treated Parkinson's disease. *J Neuropsychiatry Clin Neurosci*. 1996;8:450-452.
140. Hara K, Iijima S, Nakamura S, Kameyama M. Psychiatric symptoms and plasma levodopa dynamics in Parkinson's disease. *Rinsho Shinkeigaku*. 1987;27:846-851.
141. Nausieda PA, Glantz R, Weber S, Baum R, Klawans HL. Psychiatric complications of levodopa therapy of Parkinson's disease. *Adv Neurol*. 1984;40:271-277.
142. Nausieda PA, Tanner CM, Klawans HL. Serotonergically active agents in levodopa-induced psychiatric toxicity reactions. *Adv Neurol*. 1983;37:23-32.
143. Beasley BL, Nutt JG, Davenport RW, Chase TN. Treatment with tryptophan of levodopa-associated psychiatric disturbances. *Arch Neurol*. 1980;37:155-156.
144. Klawans HL, Moskovitz C, Nausieda PA, Weiner WJ. Levodopa-induced dopaminergic hypersensitivity in the pathogenesis of psychiatric and neurologic disorders. *Int J Neurol*. 1979;13:225-236.
145. Lloyd KG. Psychiatric disturbances occurring during levodopa therapy of Parkinson's disease. *Prim Care*. 1977;4:561-575.
146. Lin JT, Ziegler DK. Psychiatric symptoms with initiation of carbidopa-levodopa treatment. *Neurology*. 1976;26:699-700.
147. Gulmann NC, Korsgaard S. Psychiatric side-effects of levodopa treatment. *Ugeskr Laeger*. 1976;138:1105-1107.
148. Sathananthan G, Angrist BM, Gershon S. Response threshold to levodopa in psychiatric patients. *Biol Psychiatry*. 1973;7:139-146.
149. Yaryura-Tobias JA, Diamond B, Merlis S. Psychiatric manifestations of levodopa. *Can Psychiatr Assoc J*. 1972;17(suppl 2):S5123.
150. Goodwin FK. Psychiatric side effects of levodopa in man. *JAMA*. 1971;218:1915-1920.
151. Knopp W. Psychiatric changes in patients treated with levodopa. I. The clinical experiment. *Neurology*. 1970;20:23-30.
152. Celesia GG, Barr AN. Psychosis and other psychiatric manifestations of levodopa therapy. *Arch Neurol*. 1970;23:193-200.
153. Remick RA, Gimbarzevsky B. Manic reaction to lidocaine anesthesia. *J Clin Psychopharmacol*. 1990;10:442-443.
154. Basile Filho A, Passos HV, Aldrighi JR, Maciel Junior JA. Acute psychotic reactions and speech disorders induced by the rapid intravenous injection of lidocaine. *Rev Paul Med*. 1986;104:229.
155. Turner WM. Lidocaine and psychotic reactions. *Ann Intern Med*. 1982;97:149-150.
156. Humbert P, Girardin P, Aubin F. Fear of dying after intra-keloid injection of triamcinolone acetonide and lidocaine: Hoigne's syndrome. *Ann Dermatol Venereol*. 2001;128:762.
157. Saravay SM, Marke J, Steinberg MD, Rabiner CJ. "Doom anxiety" and delirium in lidocaine toxicity. *Am J Psychiatry*. 1987;144:159-163.

158. Lebain P, Juliard C, Davy JP, Dollfus S. Neuropsychiatric symptoms in preventive antimalarial treatment with mefloquine: apropos of 2 cases [in French]. *Encéphale*. 2000;26:67-70.
159. Croft AM, World MJ. Neuropsychiatric reactions with mefloquine chemoprophylaxis. *Lancet*. 1996;347:326.
160. Sowunmi A, Adio RA, Oduola AM, Ogundahunsi OA, Salako LA. Acute psychosis after mefloquine. Report of six cases. *Trop Geogr Med*. 1995;47:179-180.
161. Spensley J. Folie à deux with methylphenidate psychosis. *J Nerv Ment Dis*. 1972;155:288-290.
162. Rosenfeld AA. Depression and psychotic regression following prolonged methylphenidate use and withdrawal: case report. *Am J Psychiatry*. 1979;136:226-228.
163. Koehler-Troy C, Strober M, Malenbaum R. Methylphenidate-induced mania in a prepubertal child. *J Clin Psychiatry*. 1986;47:566-567.
164. Lu ML, Pan JJ, Teng HW, Su KP, Shen WW. Metoclopramide-induced supersensitivity psychosis. *Ann Pharmacother*. 2002;36:1387-1390.
165. Anfinson TJ. Akathisia, panic, agoraphobia, and major depression following brief exposure to metoclopramide. *Psychopharmacol Bull*. 2002;36:82-93.
166. Friend KD, Young RC. Late-onset major depression with delusions after metoclopramide treatment. *Am J Geriatr Psychiatry*. 1997;5:79-82.
167. Jibiki I, Maeda T, Yamaguchi N. Metoclopramide-induced parkinsonism and depression. *Acta Neurol (Napoli)*. 1992;14:130-133.
168. Shearer RM, Bownes IT, Curran P. Tardive akathisia and agitated depression during metoclopramide therapy. *Acta Psychiatr Scand*. 1984;70:428-431.
169. Herzog P. Depression after metoclopramide medication. *Harefuah*. 1982;103:374-375.
170. Schreiber W, Spernal J. Metronidazole-induced psychotic disorder. *Am J Psychiatry*. 1997;154:1170-1171.
171. Schentag JJ, Ziemniak JA, Greco JM, Rainstein M, Buckley RJ. Mental confusion in a patient treated with metronidazole: a concentration-related effect? *Pharmacotherapy*. 1982;2:384-387.
172. Taylor JAT. Metronidazole—a new agent for combined somatic and psychic therapy of alcoholism. *Bull Los Angeles Neurol Soc*. 1964;29:158-162.
173. Cina SJ, Russell RA, Conradi SE. Sudden death due to metronidazole/ethanol interaction. *Am J Forensic Med Pathol*. 1996;17:343-346.
174. Williams CS, Woodcock KR. Do ethanol and metronidazole interact to produce a disulfiram-like reaction? *Ann Pharmacother*. 2000;34:255-257.
175. McCahill ME, Braff DL. Atypical postpartum psychosis possibly associated with metronidazole therapy. *J Fam Pract*. 1988;27:323-325.
176. Kanof PD, Aronson MJ, Ness R. Organic mood syndrome associated with detoxification from methadone maintenance. *Am J Psychiatry*. 1993;150:423-428.
177. Camacho A, Matthews SC, Dimsdale JE. "Invisible" synthetic opiates and acute psychosis. *N Engl J Med*. 2001;345:469.
178. Shreeram SS, McDonald T, Dennison S. Psychosis after ultrarapid opiate detoxification. *Am J Psychiatry*. 2001;158:970.
179. Davis A, Yudofsky B, Quidwai S. Acute psychosis associated with butorphanol. *J Neuropsychiatry Clin Neurosci*. 1998;10:236-237.
180. Levinson I, Galynker II, Rosenthal RN. Methadone withdrawal psychosis. *J Clin Psychiatry*. 1995;56:73-76.
181. Christie JM, Meade WR, Markowsky S. Paranoid psychosis after intrathecal morphine. *Anesth Analg*. 1993;77:1298-1299.
182. Slap GB. Oral contraceptives and depression: impact, prevalence and cause. *J Adolesc Health Care*. 1981;2:53-64.
183. Moller SE. Effect of oral contraceptives on tryptophan and tyrosine availability: evidence for a possible contribution to mental depression. *Neuropsychobiology*. 1981;7:192-200.
184. Bizjak ED, Nolan PE Jr, Brody EA, Galloway JM. Procainamide-induced psychosis: a case report and review of the literature. *Ann Pharmacother*. 1999;33:948-951.
185. Harrington L. Procainamide-induced psychosis. *Crit Care Nurse*. 1993;13:70-72.
186. Schubert DS, Gabinet L, Hershey LA. Psychosis induced by sustained-release procainamide. *Can Med Assoc J*. 1984;131:1188-1190.
187. Menken M. Procainamide-induced psychosis. *JAMA*. 1979;241:1107-1108.
188. McCrum ID, Guidry JR. Procainamide-induced psychosis. *JAMA*. 1978;240:1265-1266.
189. Soutullo CA, Cottingham EM, Keck PE Jr. Psychosis associated with pseudoephedrine and dextromethorphan. *J Am Acad Child Adolesc Psychiatry*. 1999;38:1471-1472.
190. Roberge RJ, Hirani KH, Rowland PL 3rd, Berkeley R, Krenzelok EP. Dextromethorphan- and pseudoephedrine-induced agitated psychosis and ataxia: case report. *J Emerg Med*. 1999;17:285-288.
191. Reeves RR, Pinkofsky HB. Postpartum psychosis induced by bromocriptine and pseudoephedrine. *J Fam Pract*. 1997;45:164-166.
192. Taylor DM, O'Toole KS, Auble TE, Ryan CM, Sherman DR. The psychometric and cardiac effects of pseudoephedrine in the hyperbaric environment. *Pharmacotherapy*. 2000;20:1045-1050.
193. Johnson AG, Day RO, Seldon WA. A functional psychosis precipitated by quinidine. *Med J Aust*. 1990;153:47-49.
194. Deleu D, Schmedding E. Acute psychosis as idiosyncratic reaction to quinidine: report of two cases. *Br Med J (Clin Res Ed)*. 1987;294:1001-1002.
195. Defoin JF, Debonne T, Rambourg MO, et al. Acute psychiatric syndrome and quinolones. *J Toxicol Clin Exp*. 1990;10:469-472.
196. Okada F. Depression after treatment with thiazide diuretics for hypertension. *Am J Psychiatry*. 1985;142:1101-1102.
197. Visapaa JP, Tillonen JS, Kaihovaara PS, Salaspuro MP. Lack of disulfiram-like reaction with metronidazole and ethanol. *Ann Pharmacother*. 2002;36:971-974.
198. Medical Economics Staff, ed. *Physicians' Desk Reference (PDR)*. Montvale, NJ: Medical Economics Company; 2003.
199. Jacobs DG, Deutsch NL, Brewer M. Suicide, depression, and isotretinoin: is there a causal link? *J Am Acad Dermatol*. 2001;45:5168-5175.
200. Jick SS, Kremers HM, Vasilakis-Scaramozza C. Isotretinoin use and risk of depression, psychotic symptoms, suicide, and attempted suicide. *Arch Dermatol*. 2000;136:1231-1236.
201. Gupta MA, Gupta AK. Depression and suicidal ideation in dermatology patients with acne, alopecia areata, atopic dermatitis and psoriasis. *Br J Dermatol*. 1998;139:846-850.
202. Rubinow DR, Peck GL, Squillace KM, Gantt GG. Reduced anxiety and depression in cystic acne patients after successful treatment with oral isotretinoin. *J Am Acad Dermatol*. 1987;17:25-32.
203. Russo S, Kema IP, Haagsma EB, et al. Irritability rather than depression during interferon treatment is linked to increased tryptophan catabolism. Poster at the European Congress of Neuropsychopharmacology (ECNP); 2002; Barcelona, Spain.
204. O'Connor NT. Interferon beta and suicide in multiple sclerosis. *Lancet*. 1996;347:1417-1418.
205. Hall RC, Popkin MK, Stickney SK, et al. Presentation of the steroid psychoses. *J Nerv Mental Dis*. 1979;167:229-236.
206. Ahmad M, Rasul FM. Steroid-induced psychosis treated with haloperidol in a patient with active chronic obstructive pulmonary disease. *Am J Emerg Med*. 1999;17:735.
207. Kimball CP. Psychological dependency on steroids? *Ann Intern Med*. 1971;75:111-113.
208. Dixon RB, Christy NP. On the various forms of corticosteroid withdrawal syndrome. *Am J Med*. 1980;68:224-230.
209. Faigenbaum AD, Zaichkowsky LD, Gardner DE, Micheli LJ. Anabolic steroid use by male and female middle school students. *Pediatrics*. 1998;101:E6.
210. Nilsson S. Androgenic anabolic steroid use among male adolescents in Falkenberg. *Eur J Clin Pharmacol*. 1995;48:9-11.
211. DuRant RH, Escobedo LG, Heath GW. Anabolic-steroid use, strength training, and multiple drug use among adolescents in the United States. *Pediatrics*. 1995;96(1 pt 1):23-28.
212. Perez-Stable EJ, Halliday R, Gardiner PS, et al. The effects of propranolol on cognitive function and quality of life: a randomized trial among patients with diastolic hypertension. *Am J Med*. 2000;108:359-365.
213. Ko DT, Hebert PR, Coffey CS, Sedrakyan A, Curtis JP, Krumholz HM. Beta-blocker therapy and symptoms of depression, fatigue, and sexual dysfunction. *JAMA*. 2002;288:351-357.
214. Terao T. Beta-blocker therapy and depression. *JAMA*. 2002;288:1845-1846.