

# Survival significance of epidermal growth factor receptor tyrosine kinase inhibitors and current staging system for survival after recurrence in patients with completely resected lung adenocarcinoma

Hisashi Saji<sup>1,2</sup>  
Hiroki Sakai<sup>1</sup>  
Hiroyuki Kimura<sup>1</sup>  
Tomoyuki Miyazawa<sup>1</sup>  
Hideki Marushima<sup>1</sup>  
Haruhiko Nakamura<sup>1</sup>

<sup>1</sup>Department of Chest Surgery, St Marianna University School of Medicine, Miyamae-ku, Kawasaki, Kanagawa, Japan; <sup>2</sup>Department of Thoracic Surgery, Tokyo Medical University, Shinjuku-ku, Tokyo, Japan

Correspondence: Hisashi Saji  
Department of Chest Surgery,  
St Marianna University School of  
Medicine, 2-16-1 Sugao, Miyamae-ku,  
Kawasaki, Kanagawa 216-8511, Japan  
Tel +81 44 977 8111  
Fax +81 44 976 5792  
Email saji-q@ya2.so-net.ne.jp

**Objective:** We previously reported that the staging system and epidermal growth factor receptor (EGFR) mutation status are key factors for treatment strategy and predicting survival. However, the significance of these factors as predictors of overall survival (OS) and postoperative recurrence survival (PRS) has not been sufficiently elucidated. The objective here was to investigate EGFR mutation status and p-stage, which affect PRS and OS in patients with completely resected lung adenocarcinoma, using a different database.

**Patients and methods:** We retrospectively reviewed 56 consecutive lung adenocarcinoma patients with disease recurrence in St. Marianna University Hospital between January 2010 and December 2014.

**Results:** EGFR mutants (M) were detected in 16/56 patients (29%). The patients with EGFR M had a better OS than those with EGFR wild-type (WT) status (5-year survival: 50.3% vs 43.1,  $P=0.133$ ). There was no significant difference in the 3-year recurrence-free survival rate between patients with M and WT (6.3% vs 7.7%,  $P=0.656$ ), and the patients with EGFR M had a significantly better 3-year PRS than those with WT (77.4% vs 51.7%,  $P=0.033$ ). The 3-year PRS rate for patients with M/pathologic stage (p-stage) I–II (87.5%) was better than that for patients with M/p-stage III (60.0%), WT/p-stage I–II (52.7%), and WT/p-stage III (43.8%). There was a significant difference between patients with M/p-stage I and WT/p-stage I–II or WT/p-stage III ( $P=0.021$  and  $0.030$ , respectively). During the study period, of the 16 patients with mutants, 12 patients (75%) received EGFR-tyrosine kinase inhibitor (TKI) therapy and among the 40 patients with WT, no patient received EGFR-TKI therapy. Multivariate survival analysis showed that patients with EGFR-TKI therapy had a statistically significant association with favorable PRS (hazard ratio 0.271; 95% confidence interval 0.074–1.000;  $P=0.050$ ).

**Conclusion:** EGFR status and p-stage were found to be essential prognostic factors for estimating PRS using this database. The recurrent patients with EGFR M and EGFR-TKI therapy had a statistically significant association with favorable PRS.

**Keywords:** EGFR, lung adenocarcinoma, postoperative recurrence survival

## Introduction

Surgical resection with a curative intent is considered to be the standard treatment for early-stage non-small-cell lung cancer (NSCLC). However, recurrence rates after complete resection for stage I to III of NSCLC are from 30% to 75% and depend on

pathologic staging and follow-up period. Even with combined modality treatment, including chemotherapy, radiotherapy, or other therapeutic modalities, most patients with recurrence after resection have little possibility of cure.<sup>1–5</sup> With the advancement of local and systemic therapies for recurrent NSCLC, survival after disease recurrence, which we defined as postoperative recurrence survival (PRS), has extensively improved.<sup>1,6</sup> However, conclusive factors that affect PRS in NSCLC have not been discussed as widely as those that affect overall survival (OS) or recurrence-free survival (RFS) from the time of initial treatment.

The epidermal growth factor receptor (EGFR) mutation, which has been detected mostly in lung adenocarcinoma, is a predictor of response to EGFR tyrosine kinase inhibitors (TKIs), with 70%–80% of NSCLC patients receiving substantial benefits from these targeted therapies. EGFR mutation is reported to be a predictive and prognostic factor of EGFR-TKI therapy outcome.<sup>7–9</sup> Testing for these mutations in all patients with recurrent/metastatic lung adenocarcinoma is, therefore, recommended in standard practice.

We previously reported that the staging system and EGFR mutation status are key factors for treatment strategy and survival prediction.<sup>5</sup> However, the significance of these factors as predictors of OS and PRS has not been sufficiently elucidated. This study aims to further investigate EGFR mutation status and pathologic stage (p-stage), which affect PRS and OS in patients with completely resected lung adenocarcinoma, using a different database during the standard treatment with EGFR-TKIs in patients with NSCLC with EGFR mutations.

## Patients and methods

### Patient selection

We retrospectively reviewed a total of 340 consecutive patients with adenocarcinoma who underwent lobectomy, bilobectomy, or pneumonectomy with systematic lymph node dissection at St. Marianna University Hospital from January 2010 to December 2014. Of these, 75 (22%) patients had recurrence of disease after surgery during median follow-up time of 36.6 months (range, 6.0–77.0 months). We excluded six patients who had received preoperative chemotherapy, radiotherapy, or both, or patients whose tumors had EGFR mutation in exon 20 (two patients because of resistance to EGFR-TKIs) or the echinoderm microtubule-associated protein-like 4-anaplastic lymphoma kinase fusion gene (two patients). We excluded 11 patients who received good supportive care, including only palliative radiotherapy for controlling pain from bone metastasis, and no treatments such as radiotherapy, chemotherapy, or surgical treatment after

disease recurrence. We enrolled the remaining 56 patients with postoperative recurrent disease in this study.

### Data collection

Preoperative evaluation included physical examination, chest radiography, computed tomography (CT) scan of the chest and abdomen, blood examination, and positron emission tomography-CT scan. Histologic subtypes of lung cancer were determined according to the World Health Organization classification.<sup>10</sup> Determination of the tumor stages was based on the Seventh TNM Classification for Lung and Pleural Tumors.<sup>11</sup>

All patients were followed up at our hospital every 3 months in the first year, every 6 months from the second to the fifth year, and annually thereafter on an outpatient basis, and were aimed at continuing follow-up for 10 years. Follow-up procedures included physical examination, chest radiography, and blood examination (including serum tumor markers). CT scans of chest and upper abdomen were routinely done in every scheduled outpatient department visit at our hospital for follow-up. Whenever any symptoms or signs of recurrence were detected, magnetic resonance imaging of the brain and bone scintigraphy were performed. Once a metastasis was discovered by physical examination and diagnostic imaging, histologic or cytologic confirmation of the metastatic site was performed when clinically feasible. Local recurrence was defined as disease recurrence at the surgical margin, ipsilateral hemithorax, or mediastinum. Distant metastasis was defined as disease recurrence in the contralateral lung or outside the hemithorax and mediastinum. A second primary tumor was recorded when a patient presented with a new histologic type and with clinical features consistent with a new primary tumor.

The hospital charts of all patients were reviewed to collect clinicopathologic data including age, sex, smoking history, tumor location, operative procedure, histologic type, tumor size, vascular invasion, lymphatic permeation, visceral pleural invasion, EGFR mutation status, administration of EGFR-TKI therapy, recurrence pattern, OS, RFS, and PRS. The institutional review board of the St Marianna University School of Medicine approved the protocols for data collection and analyses. The requirement for written informed consent from the patients was waived because of the retrospective study design. All patient data were maintained confidentially throughout this research.

### Statistical analysis

OS, RFS, and PRS were estimated using the Kaplan–Meier method, and survival differences between patient groups

were determined using log-rank analysis. OS was defined as the time elapsed from the date of pulmonary resection to the date of death. The length of PRS was defined as the interval between the date of initial recurrence identification and the date of either death or the last follow-up. RFS was defined as the time elapsed from the date of pulmonary resection to the date of the first tumor recurrence.

Univariate analysis was conducted between the groups. Significance in a 2×2 contingency table was tested by using the  $\chi^2$  test. *P*-values and hazard ratios in the multivariate analyses were calculated using the Cox regression model. All *P*-values <0.05 were considered to indicate a statistically significant difference. All statistical calculations were performed using the SPSS statistical software package (version 21.0; SPSS, Inc., Chicago, IL, USA).

## Results

### Patient characteristics

The characteristics of the patients are listed in Table 1. Of the 56 patients, EGFR mutants (M) were detected in 16 patients (29%). The most common EGFR mutation was in-frame deletion in exon 19 (8/16; 50%). The second most common mutation was missense mutation (mostly L858R) in exon 21 in seven patients (44%). Regarding the initial recurrence patterns, 35 patients (64%) had distant or local and distant recurrence and 21 patients (36%) had only local recurrence. Of the 16 patients with mutants, 12 patients (75%) received EGFR-TKI therapy. Among the 40 patients with wild-type (WT) status, no patient received EGFR-TKI therapy.

### Survival

The median follow-up time for all 56 patients was 36.6 months (range, 6.0–77.0 months). The median PRS time for all 56 patients was 16.7 months (range, 0.7–61.0 months). The 5-year OS rates of patients with EGFR M and WT were 50.3% and 43.1%, respectively (Figure 1A). There was no significant difference between them (*P*=0.133). The 3-year RFS rates of patients with EGFR M and WT were 6.3% and 7.7%, respectively (*P*=0.656; Figure 1B). The 3-year PRS of patients with EGFR M was significantly longer than that of patients with WT (77.4% vs 51.7%, *P*=0.033; Figure 1C). Figure 1D shows the PRS curves stratified according to the EGFR status and the p-stage. The 3-year PRS rates for the patients with EGFR M with p-I–II, EGFR M with p-III, WT with p-I–II, and WT with p-III were 87.5%, 60.0%, 52.7%, and 43.8%, respectively. There were statistically significant differences in the PRS rates between the patients with EGFR M with p-I–II and those with WT with p-I–II (*P*=0.030) and between the patients with EGFR M with

**Table 1** Patient characteristics (N=56)


Variable	Category	n (%)
Sex	Male	36 (64)
	Female	20 (36)
Median age (range), years		70.3 (42–81)
Smoking habits	Ever smoker/unknown	40 (71)
	Never smoker	16 (29)
Tumor location	Right	31 (55)
	Left	25 (45)
Surgical procedure	Lobectomy/bilobectomy	54 (96)
	Pneumonectomy	2 (4)
Median tumor size (range), cm		27.4 (0.7–83)
Pathologic N status	N0	40 (71)
	N1	5 (9)
	N2	11 (20)
Pathologic stage	IA/IB	37 (66)
	IIA/IIB	8 (14)
	IIIA	11 (20)
Vascular invasion	Positive	9 (16)
Lymphatic permeation	Positive	17 (30)
Visceral pleural invasion	Positive	22 (39)
EGFR mutation status	Mutant	16 (29)
	(L858R/del 19/exon 18)	7/8/1
	Wild type	40 (71)
Treatment for initial recurrence	Chemotherapy	48 (85)
	Chemotherapy with radiotherapy	3 (5)
	Radiotherapy	5 (10)
EGFR-TKI therapy	With	12 (21)
	(Gefitinib/erlotinib/afatinib/both)	6/2/3/1
Initial recurrence pattern	Without	44 (79)
	Distant/both	35 (64)
Initial recurrence site (including duplication)	Locoregional	21 (36)
	Brain	9 (16)
	Bone	8 (14)
Pleural dissemination/malignant effusion	Pleural dissemination/malignant effusion	8 (14)
	Lung	22 (39)
	Lymph nodes	12 (21)
	Liver	3 (5)
	Adrenal gland	1 (2)

**Abbreviations:** EGFR, epidermal growth factor receptor; TKI, tyrosine kinase inhibitor.

p-I–II and those with WT with p-III (*P*=0.021). There were no significant differences between the patients with EGFR M with p-III and those with WT with p-I–II (*P*=0.468). Figure 2A and B shows that there was a significant difference in PRS between patients with and without EGFR-TKIs (*P*=0.045), but no significant difference in OS between the two groups (*P*=0.165).

### Prognostic factors

Of the 16 patients with mutants, 12 patients (75%) received EGFR-TKI therapy. Among the 40 patients with WT, no




**Figure 1** (A) Overall survival stratified according to EGFR mutants and WT. (B) RFS stratified according to EGFR mutants and WT. (C) PRS stratified according to EGFR mutants and WT. Significant differences in the PRS rates were observed between the patients with mutants and WT ( $P=0.030$ ). (D) PRS stratified according to EGFR status and p-stage. Significant differences in the PRS rates were observed between the patients with mutants/p-stage I-II and WT/p-stage I-II ( $P=0.030$ ) and between the patients with mutants/p-stage I-II and WT/p-stage III ( $P=0.021$ ).

**Abbreviations:** EGFR, epidermal growth factor receptor; PRS, postoperative recurrence survival; p-stage, pathologic stage; RFS, recurrence-free survival; WT, wild type.

patient received EGFR-TKI therapy. During the study period, EGFR-TKIs were recommended for most recurrent or metastatic patients with lung adenocarcinoma with EGFR M as the standard practice; we, therefore, selected administration of EGFR-TKI for this prognostic analysis. The potential prognostic factors for PRS were analyzed using univariate survival analysis (Table 2), which showed that sex and EGFR-TKI therapy were significant prognostic factors. Multivariate survival analysis (Table 2) showed that patients with EGFR-TKI therapy had a statistically significant association with favorable PRS (hazard ratio 0.271; 95% confidence interval 0.074–1.000;  $P=0.050$ ).

## Discussion

Personalization of lung cancer treatment requires predictive biomarkers that have been validated by correlation between tumor features and outcomes after therapy. Several mutations have been identified in *EGFR* in NSCLC. Mutation status in this gene is thought to be an important predictor of response to EGFR-TKIs, with 70%–80% of NSCLC patients receiving substantial benefits from this targeted therapy.<sup>7</sup> *EGFR* mutation is both a predictive and prognostic factor of EGFR-TKI therapy outcome.<sup>7,12</sup> Testing for these mutations in all patients with recurrent or metastatic lung adenocarcinoma is, therefore, recommended in standard practice.



**Figure 2 (A)** Overall survival stratified according to whether EGFR-TKIs were administered or not. **(B)** RFS stratified according to whether EGFR-TKIs were administered or not. Significant differences in the PRS rates were observed between the patients with and without EGFR-TKIs ( $P=0.045$ ).

**Abbreviations:** EGFR, epidermal growth factor receptor; PRS, postoperative recurrence survival; RFS, recurrence-free survival; TKI, tyrosine kinase inhibitor.

We aimed to 1) identify the prognostic factors of PRS in patients who underwent complete resection of lung adenocarcinoma, 2) clarify whether EGFR mutation status and p-stage can be considered as conclusive indicators of OS or PRS, and 3) elucidate whether the EGFR-TKI therapy and p-stage at surgery affect prediction of PRS. We found that having EGFR mutation itself was a favorable PRS factor, but neither a predicting OS factor nor a disease free survival factor. Although p-stage at surgery itself may not affect the prediction of PRS, PRS outcomes were stratified by EGFR mutation status and p-stage. Moreover, whether EGFR-TKI therapy was administered was the most independent prognostic factor in patients who underwent complete resection of lung adenocarcinoma and relapsed.

Although several studies have reported on the PRS of patients with NSCLC,<sup>1–3,6,13,14</sup> no standard treatment strategy has been established for the patients with recurrence based on prospective studies. The concept of PRS of surgically treated patients has been the focus of much attention recently because encouraging new treatments such as EGFR-TKIs,<sup>7–9</sup>

anaplastic lymphoma kinase inhibitors,<sup>15</sup> vascular endothelial growth factor antibody,<sup>16</sup> or recently, immune check point inhibitors<sup>17,18</sup> have provided long-term PRS and better quality of life to selected patients with recurrent or metastatic disease. Several retrospective studies, in addition to our own recently published study, have reported that shorter relapse-free interval, initial recurrence pattern, lymph node metastasis pattern, tumor differentiation, EGFR status, and administration of EGFR-TKIs and the p-stage are independent prognostic factors for PRS.<sup>1,5,6,13,19–23</sup> In our current series, EGFR-TKI administration was shown to be a strong independent prognostic factor for estimating PRS. Of the 16 relapse patients with EGFR mutants, 12 patients (75%) received EGFR-TKI therapy. Because our series enrolled from 2010 when analysis of EGFR status are common and EGFR-TKIs are already thought as a standard first-line therapy in practice setting for metastatic lung cancer patients with EGFR mutant.<sup>7–9,24</sup> Our results, however, demonstrated that p-stage at surgery may not affect PRS, suggesting that p-stage at surgery does not correlate with survival interval after relapse.

**Table 2** Univariate and multivariate analyses for PRS (N=56)

Variable	UVA	MVA		
	P-value	Hazard ratio	95% CI	P-value
Age <75 years (vs ≥75 years)	0.839	Not included in MVA		
Male sex (vs female sex)	0.045*	0.342	0.073–1.589	0.171
Pathologic stage III (vs I–II)	0.223	Not included in MVA		
EGFR-TKI therapy: without (vs with)	0.026*	0.271	0.074–1.000	0.050*
Recurrence-free interval <1 year (vs ≥1 year)	0.113	Not included in MVA		
Initial recurrence pattern distant or both (vs locoregional)	0.668	Not included in MVA		

**Note:** \* $P<0.05$ .

**Abbreviations:** EGFR, epidermal growth factor receptor; MVA, multivariate analysis; PRS, postoperative recurrence survival; TKI, tyrosine kinase inhibitor; UVA, univariate analysis.

Two studies have reported that EGFR mutations have an impact on prognosis after surgical resection of NSCLC<sup>25,26</sup> and that the mutation status of EGFR can serve as an independent prognostic marker associated with decreased recurrence and improved progression-free survival and OS in patients with stage I lung adenocarcinoma.<sup>25</sup> This study, however, restricted to patients who had no adjuvant or neoadjuvant systemic therapy administered and focused solely on the postoperative prognostic differences among different genotypes, showed that the significant prognostic impact of EGFR mutation was lost after adjusting for other confounding prognostic factors.<sup>26</sup> Other studies in the literature corroborate this finding.<sup>27,28</sup> Additionally, we showed in our previous study that EGFR mutation status and the use of EGFR-TKI therapy particularly affected the PRS among patients who underwent surgical resection for lung cancers,<sup>5</sup> and that the results could also affect OS in general. In this study of a small sample and a short median follow-up time, however, we found that the EGFR mutation status could affect only PRS, but not OS.

There are several limitations and biases of this study that should be taken into account for interpreting the results. As a retrospective single-institute study, patient selection bias and time trend bias regarding the treatment for recurrent diseases were inevitable. Because follow-up examination of the patients after the initial resection was comparatively uniform, systematic follow-up is suggested to increase the early detection of recurrent lesions to which we could apply intensive treatment. Recently, it has been shown there are some differences in response to EGFR-TKIs among EGFR mutation subtypes.<sup>29</sup> Because of the small size of sample in our study, we could not analyze these differences.

## Conclusion

We retrospectively reviewed 56 consecutive lung adenocarcinoma patients with disease recurrence in our hospital between January 2010 and December 2014. EGFR status and p-stage were also found to be essential prognostic factors for estimating PRS using this database. The recurrent lung adenocarcinoma patients with EGFR M and EGFR-TKI therapy had a statistically significant association with favorable PRS. Advances in treatment strategy for PRS may provide improvement in OS among patients who undergo surgery.

## Acknowledgment

This study was supported by a Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science (24592104), Ministry of Education, Culture, Sports, Science and Technology, Japan.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Yoshino I, Yohena T, Kitajima M, et al. Survival of non-small cell lung cancer patients with postoperative recurrence at distant organs. *Ann Thorac Cardiovasc Surg.* 2001;7(4):204–209.
2. Hung JJ, Hsu WH, Hsieh CC, et al. Post-recurrence survival in completely resected stage I non-small cell lung cancer with local recurrence. *Thorax.* 2009;64(3):192–196.
3. Hung JJ, Jeng WJ, Hsu WH, et al. Prognostic factors of postrecurrence survival in completely resected stage I non-small cell lung cancer with distant metastasis. *Thorax.* 2010;65(3):241–245.
4. Shimada Y, Saji H, Yoshida K, et al. Prognostic factors and the significance of treatment after recurrence in completely resected stage I non-small cell lung cancer. *Chest.* 2013;143(6):1626–1634.
5. Kudo Y, Shimada Y, Saji H, et al. Prognostic factors for survival after recurrence in patients with completely resected lung adenocarcinoma: important roles of epidermal growth factor receptor mutation status and the current staging system. *Clin Lung Cancer.* 2015;16(6):e213–e221.
6. Sugimura H, Nichols FC, Yang P, et al. Survival after recurrent nonsmall-cell lung cancer after complete pulmonary resection. *Ann Thorac Surg.* 2007;83(2):409–417; discussion 417–418.
7. Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med.* 2009;361(10):947–957.
8. Maemondo M, Inoue A, Kobayashi K, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med.* 2010;362(25):2380–2388.
9. Mitsudomi T, Morita S, Yatabe Y, et al. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol.* 2010;11(2):121–128.
10. International Agency for Research on Cancer, Travis WD, Brambilla E, Burke A, Marx A, Nicholson AG. *WHO Classification of Tumours of the Lung, Pleura, Thymus and Heart.* 4th ed. Geneva, Switzerland: World Health Organization; 2015.
11. Sobin LH, Gospodarowicz MK, Wittekind C. International Union against Cancer, ebrary Inc. *TNM Classification of Malignant Tumours.* 7th ed. Chichester, West Sussex, UK; Hoboken, NJ: Wiley-Blackwell; 2009. Available from: <http://site.ebrary.com/lib/yale/Doc?id=10342913>. Accessed June 10, 2017.
12. Eberhard DA, Johnson BE, Amler LC, et al. Mutations in the epidermal growth factor receptor and in KRAS are predictive and prognostic indicators in patients with non-small-cell lung cancer treated with chemotherapy alone and in combination with erlotinib. *J Clin Oncol.* 2005; 23(25):5900–5909.
13. Williams BA, Sugimura H, Endo C, et al. Predicting postrecurrence survival among completely resected nonsmall-cell lung cancer patients. *Ann Thorac Surg.* 2006;81(3):1021–1027.
14. Nakagawa T, Okumura N, Ohata K, Igai H, Matsuoka T, Kameyama K. Postrecurrence survival in patients with stage I non-small cell lung cancer. *Eur J Cardiothorac Surg.* 2008;34(3):499–504.
15. Kwak EL, Bang YJ, Camidge DR, et al. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med.* 2010;363(18):1693–1703.
16. Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med.* 2006; 355(24):2542–2550.
17. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med.* 2015; 373(2):123–135.
18. Reck M, Rodriguez-Abreu D, Robinson AG, et al; KEYNOTE-024 Investigators. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med.* 2016;375(19):1823–1833.

19. Okami J, Taniguchi K, Higashiyama M, et al. Prognostic factors for gefitinib-treated postoperative recurrence in non-small cell lung cancer. *Oncology*. 2007;72(3-4):234-242.
20. Saisho S, Yasuda K, Maeda A, et al. Post-recurrence survival of patients with non-small-cell lung cancer after curative resection with or without induction/adjuvant chemotherapy. *Interact Cardiovasc Thorac Surg*. 2013;16(2):166-172.
21. Jeon JH, Kang CH, Kim HS, Seong YW, Park IK, Kim YT. Prognostic and predictive role of epidermal growth factor receptor mutation in recurrent pulmonary adenocarcinoma after curative resection. *Eur J Cardiothorac Surg*. 2015;47(3):556-562.
22. Sonobe M, Yamada T, Sato M, et al. Identification of subsets of patients with favorable prognosis after recurrence in completely resected non-small cell lung cancer. *Ann Surg Oncol*. 2014;21(8):2546-2554.
23. Takenaka T, Takenoyama M, Yamaguchi M, et al. Impact of the epidermal growth factor receptor mutation status on the post-recurrence survival of patients with surgically resected non-small-cell lung cancer. *Eur J Cardiothorac Surg*. 2015;47(3):550-555.
24. Jiang H. Overview of gefitinib in non-small cell lung cancer: an Asian perspective. *Jpn J Clin Oncol*. 2009;39(3):137-150.
25. Izar B, Sequist L, Lee M, et al. The impact of EGFR mutation status on outcomes in patients with resected stage I non-small cell lung cancers. *Ann Thorac Surg*. 2013;96(3):962-968.
26. Kosaka T, Yatabe Y, Onozato R, Kuwano H, Mitsudomi T. Prognostic implication of EGFR, KRAS, and TP53 gene mutations in a large cohort of Japanese patients with surgically treated lung adenocarcinoma. *J Thorac Oncol*. 2009;4(1):22-29.
27. Marks JL, Broderick S, Zhou Q, et al. Prognostic and therapeutic implications of EGFR and KRAS mutations in resected lung adenocarcinoma. *J Thorac Oncol*. 2008;3(2):111-116.
28. Kim YT, Seong YW, Jung YJ, et al. The presence of mutations in epidermal growth factor receptor gene is not a prognostic factor for long-term outcome after surgical resection of non-small-cell lung cancer. *J Thorac Oncol*. 2013;8(2):171-178.
29. Yang JC, Wu YL, Schuler M, et al. Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials. *Lancet Oncol*. 2015;16(2):141-151.

### OncoTargets and Therapy

## Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress