

RESEARCH ARTICLE

Increased risk for developing gambling disorder under the treatment with pramipexole, ropinirole, and aripiprazole: A nationwide register study in Sweden

Mirjam Wolfschlag^{1,2*}, Anders Håkansson^{1,2}

1 Department of Clinical Sciences Lund, Psychiatry, Faculty of Medicine, Lund University, Lund, Sweden, **2** Malmö Addiction Center, Clinical Research Unit, Malmö, Sweden

* mirjam.wolfschlag@kabelmail.de

Abstract

Gambling Disorder (GD) has recently been reclassified from an impulse-control disorder to a behavioural addiction and, as in other addictive disorders, the dopaminergic reward system is involved. According to neuroimaging studies, alterations within the striatal dopaminergic signalling can occur in GD. However, the findings to date are controversial and there has been no agreement yet on how the reward system is affected on a molecular basis. Within the last 20 years, there has been growing evidence for a higher risk to develop GD in response to certain dopaminergic medication. Especially the dopamine agonists pramipexole and ropinirole, and the dopamine modulator aripiprazole seem to increase the likelihood for GD. The goal of this study was to examine the association between a prescription for either of the three pharmaceuticals and a GD diagnosis in a large cross-sectional study of the Swedish population. Compared to patients with any other dopaminergic drug prescription (38.7% with GD), the diagnosis was more common in patients with a dopamine agonist prescription (69.8% with GD), resulting in an odds ratio of 3.2. A similar association was found between aripiprazole prescriptions and GD diagnoses, which were analysed within the subgroup of all patients with schizophrenia or a schizotypal, delusional, or another non-mood psychotic disorder. An aripiprazole prescription increased the likelihood of GD (88.8%) in comparison to patients without an aripiprazole prescription (71.2%) with an odds ratio of 3.4. This study contributes to the increasingly reliable evidence for an association between several dopaminergic drugs and a higher risk for developing GD. Therefore, one future research goal should be a better understanding of the neurobiology in GD to be able to design more selective dopaminergic medication with less severe side effects. Additionally, this knowledge could enable the development of pharmacotherapy in GD and other addictive disorders.

OPEN ACCESS

Citation: Wolfschlag M, Håkansson A (2021) Increased risk for developing gambling disorder under the treatment with pramipexole, ropinirole, and aripiprazole: A nationwide register study in Sweden. *PLoS ONE* 16(6): e0252516. <https://doi.org/10.1371/journal.pone.0252516>

Editor: Kenji Hashimoto, Chiba Daigaku, JAPAN

Received: April 1, 2021

Accepted: May 17, 2021

Published: June 1, 2021

Copyright: © 2021 Wolfschlag, Håkansson. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be shared publicly because they contain confidential patient information, and the authors hold no permission to share them. Data are available from the Swedish National Board of Health and Welfare, SE-106 30 Stockholm, Sweden (contact via +46 (0) 75 247 30 00 or socialstyrelsen@socialstyrelsen.se) for researchers who meet the criteria for access to confidential data.

Funding: The authors received no specific funding for this work.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: Mirjam Wolfschlag declares no competing interests. Anders Håkansson holds a position at Lund University which is supported by the state-owned Swedish gambling operator AB Svenska Spel as part of its responsible gambling policies, and also has research funding from the research councils of that body and from the research council of the Swedish alcohol monopoly, Systembolaget. Håkansson also has been the national sub-investigator of a pharmaco-epidemiological survey, carried out by the independent research institute Triangle Research Institute, United States, and financed by the pharmaceutical company Shire. No direct fees or compensations were paid to the individual sub-investigator. Also, a clinical follow-up study has been planned in collaboration with the company Kontigo care, which provides follow-up devices free of charge for a clinical treatment trial in gambling disorder, but without any further financial support to the research group or its members. None of the bodies mentioned above had any role in or influence on the present research and this does not alter our adherence to PLOS ONE policies on sharing data and materials.

Introduction

Gambling disorder and the dopaminergic reward system

Gambling disorder (GD) is the permanent and severe manifestation of the gambling activity that many in the general population pursue sporadically as a hobby. Common forms are casino games, slot-machines, and lotteries, but in the last years internet gambling has grown in popularity [1]. The prevalence of GD worldwide ranges between 0.12–5.80% and under the special circumstances of the recent COVID-19 pandemic, GD has become an increasing problem for vulnerable individuals [2, 3]. Possible negative consequences of GD include a detrimental impact on close relationships, and higher mortality and suicidality [4–7]. Furthermore, GD is associated with different psychiatric comorbidities such as substance-use, depression, and anxiety [8].

Formerly seen as an impulse-control disorder, GD has been reclassified as a behavioural addiction and been renamed from “Pathological Gambling” in the latest revisions of the International Classification of Diseases (ICD 11) and the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) [9, 10]. This decision was based on broad evidence for GD sharing many characteristics with substance-use disorders rather than impulse-control disorders, clinically and on a molecular basis [11–13]. Regarding the pathophysiology in GD, many different transmitter and hormone systems have been shown to be involved, including dopamine, opioids, serotonin, cortisol, and adrenaline [14]. Nevertheless, no clear understanding of the neurochemistry in GD could be obtained yet and there is no pharmacotherapy available specific for GD. As in other addictive disorders, opioid antagonists such as naltrexone have shown some beneficial effect but are insufficient as a sole approach.

One central element in the neurobiology of GD is the dopamine-based reward and reinforcement system [14]. Its main part are neurons within the mesolimbic dopamine pathway, that project from the ventral tegmental area in the brain stem into the nucleus accumbens (NAc) in the ventral striatum, located in the basal ganglia. Physiologically, this system enables the judgement of external stimuli, creates an evaluated memory for them, links them to each other, and conducts adequate behaviour [15]. In addiction, a certain substance or behaviour possibly over-activates the route of being given a positive value, even if the outcome is detrimental, for instance negative health effects. This leads to an impaired balance in the rating of stimuli and the subject of addiction receives pathologically increased attention, while other goals lose importance [16]. One important part in reward-related learning is Pavlovian conditioning, by which a neutral cue is linked to a secondary stimulus with a positive or a negative value [17]. The former unrelated cue becomes a conditioned stimulus which can from now on predict the secondary, unconditioned stimulus. On a molecular basis, this linkage is communicated by a shift from the basal tonic dopamine signalling at the NAc to phasic spikes. Whenever a hitherto neutral stimulus yields a more positive effect than expected, phasic signalling is the reaction to this reward prediction error (RPE). The cue is given a more positive value and causes no more changes from the tonic dopamine signalling in the future. Zack et al. recently published a comprehensive review on the uncertainty in gambling as a key factor for the behaviour to become an addiction [17]. In gambling there are no reliable cues for reward prediction and winning can always be seen as “better than expected”. Therefore, the idea was proposed that this uncertainty causes an extraordinary change in dopamine signalling based on constant RPEs, which leads to a pathological sensitisation of the reward system and gambling becomes addictive. This theory is supported by experimental findings, where increased dopamine release has been found to code the uncertainty in gambling [18]. Nevertheless, whether the addictive character of gambling is directly translated by changes in dopamine signalling at the NAc remains subject to controversies.

Alterations in striatal dopamine signalling in GD

Many contradictory studies have been published on the alterations within striatal dopamine signalling in GD, often based on positron emission tomography (PET) measurements [19]. On the one hand, there is some evidence for an elevation in the dopamine transmission in connection to typical characteristics of gambling behaviour. Increased binding of dopamine to inhibitory striatal D₂ and D₃ receptors has been found to positively correlate with symptom severity in GD patients but not in healthy controls, for instance higher impulsivity, increased excitement and alertness, or impaired decision making in a reward context [20–23]. Furthermore, the dopamine synthesis capacity has been found to be elevated in GD patients compared to a control group [24]. Amphetamine, which decreases the dopamine reuptake into the presynapse and hence increases the signalling, has been found to prime gambling motivation and induces a higher striatal dopamine release only in patients with GD, not in healthy controls [25, 26]. Modafinil, another dopamine reuptake inhibitor, was able to raise the tendency to be drawn toward rewards in GD patients in contrast to a placebo [27]. The D₂ antagonist haloperidol, which acts indirectly activating on dopamine signalling through inhibition blockage, can enhance the rewarding effects and priming caused by gambling within GD patients despite no observed effect in the control group [28].

On the other hand, several measurements have shown no difference between GD patients and healthy controls regarding striatal dopamine signalling. The striatal D₂ and D₃ receptor availability has been found to be the same in GD patients and controls [20, 29]. Additionally, the treatment of GD with the proposed D₂ antagonist olanzapine has shown no effect in comparison to a placebo [30, 31]. Hence, the findings regarding the striatal dopamine signalling in GD are inconclusive to date. There seems to be evidence for some alterations, but replicable data is rare.

The association between dopaminergic medication and GD

Within the last 20 years, the evidence for an association between developing GD and the intake of dopaminergic medication has grown. One of the first case reports was published in 2000, where 10 patients with Parkinson's disease (PD) under levodopa treatment had developed GD and the authors speculated on the altered dopaminergic tone being the neurobiological explanation [32]. This theory was specified by another research group, proposing the idea that the overstimulation of the mesolimbic dopamine system could be the mechanism behind dopamine agonists causing GD [33]. More case reports were published and a dose-response relationship between dopamine agonists and impulse-control disorders was observed [34–37]. In 2010, the DOMINION study received a lot of attention, which investigated impulse-control disorders in 3090 PD patients in a cross-sectional and multicenter design in Northern America. Within all patients, 5.0% had developed GD and the risk was higher under dopamine agonist treatment (6.9% with GD) [38]. Multiple other studies were published on the subject within the last decade, suggesting a connection between dopamine agonists, respectively dopamine modulators, and the diagnosis GD [39–44].

According to previous research, certain dopaminergic medication entails a higher risk for impaired impulse control and GD development than other substances. In comparison to other dopamine agonists, especially pramipexole and ropinirole seem to correlate with increased GD rates. Both pharmaceuticals were developed as agonists for the D₂-like receptor family, thus activating the inhibitory pathways of dopamine signalling, and are prescribed as part of the dopamine replacement therapy in PD but also against restless legs syndrome [45–48]. Pramipexole and ropinirole have a higher affinity for the D₃ receptor compared with the D₂ receptor, which has been proposed as one possible explanation for their increased property to cause addictive and impulse-control disorders [49]. Seeman et al. could show that the proportion of

patients that develop an impulse-control disorder directly correlates with D₃-selectivity over D₂ in dopamine agonists.

Another substance, that has been associated with an increased likelihood for developing GD, is the atypical antipsychotic aripiprazole, mainly prescribed against schizophrenia and bipolar disorder [50]. The partial dopamine agonism of aripiprazole results in its stabilising effect on dopamine levels, acting as a modulator [51]. Similar to pramipexole and ropinirole, aripiprazole shows highest affinity for D₂ and D₃ receptors, which could explain its impact on the development of GD.

Even if the connection between dopamine agonists, respectively aripiprazole, and an impaired impulse control could not be revealed on a molecular basis yet, the association has been shown and recognised increasingly within in the last years [52–55]. Nevertheless, more evidence based on large study populations is needed to raise more attention and find solutions for the issues with certain dopaminergic medication regarding adverse psychiatric effects. Therefore, this cross-sectional study aimed to investigate the prescribed drugs within all diagnosed GD patients in the Swedish population. The association between a pramipexole /ropinirole or an aripiprazole prescription and a GD diagnosis was analysed statistically, while certain other factors that could contribute to a higher likelihood of a GD diagnosis were eliminated by choosing adequate reference groups.

Materials and methods

Register material

Patient data was provided by the Swedish National Board of Health and Welfare and obtained from the Swedish National Patient Registry, which is based on in-patient and specialised out-patient health care records, and the Swedish Prescribed Drug Register. The study was approved by the Swedish Ethical Review Authority under the file number 2019–01559. Since the data was analysed anonymously, consent from the participants was not obtained. For the study population, all patients in Sweden with the diagnosis pathological gambling (F63.0 according to ICD-10) between 2005 and 2019 were selected (n = 3689). Each case was paired with two age- and gender-matched controls from the total population register by Statistics Sweden with the goal to analyse the association between the exposure to the medication and the development of gambling disorder independently of the variables age and gender. This resulted in a total study population of 11 067 individuals.

The register material from the Swedish National Patient Registry included data on diagnoses according to ICD-10, age at treatment, gender, and the date of treatment. These data were combined with information from the Swedish Prescribed Drug Register containing the prescription date and the full anatomical therapeutic chemical (ATC) codes for the drug classes N02, N03, N04, N05, N06, N07B, R06AD01, R06AD02 and R06AD52.

Statistical analysis

The aim of this study was to explore a possible association between the prescription for a certain medication and a GD diagnosis using the software IBM SPSS Statistics 26 to process and analyse the data. In preparation for the analysis, the different files received from the Swedish National Board of Health and Welfare were merged into one main file, containing all relevant information for each patient. Based on the ATC codes, binary variables were created for each medication group or pharmaceutical in question, for instance “pramipexole prescription/no pramipexole prescription” or “psychotropic drug prescription/no psychotropic drug prescription”. The same method was used to create binary variables for certain diagnoses, such as “GD/no GD” or “PD/no PD”, from the full list of diagnoses made in the in-patient and specialised out-patient health care.

In a next step, adequate subgroups within the total study population were identified. The goal was to select patients with a similar severity of disease to make individuals with a prescription for the pharmaceuticals in question as comparable as possible to the whole reference group. Since the counts of PD and restless legs syndrome patients were very low (32, respectively 27 patients in total), the subgroup for analysing dopamine agonists (pramipexole and/or ropinirole) was chosen to be all patients with any dopaminergic drug prescription (ATC code N04B, $n = 180$), which includes dopamine agonists but also for instance levodopa. Within this group, patients resembled each other in age, gender, and a possible PD diagnosis, not influenced by whether they had a prescription for a dopamine agonist ($n = 149$) or only another dopaminergic drug ($n = 31$) (see [S1 Appendix](#)). As a reference group for an aripiprazole prescription all patients with schizophrenia, schizotypal, delusional, and other non-mood psychotic disorders were selected (F20-29 diagnoses according to ICD-10, $n = 389$). Even within this group the age and gender distribution were similar in between patients with an aripiprazole prescription ($n = 170$) and without an aripiprazole prescription ($n = 219$) (see [S2 Appendix](#)) and the general mental health status could be expected to be comparable.

To determine the association between a dopamine agonist or an aripiprazole prescription with a GD diagnosis, chi-square tests were performed within the chosen subgroups. The tests were based on the binary variables “dopamine agonist prescription/no dopamine agonist prescription”, respectively “aripiprazole prescription/no aripiprazole prescription”, and “GD diagnosis/no GD diagnosis”. In both cases, all expected counts were higher than 5 per cell, making a chi-square test applicable.

Additionally, a binary logistic regression with the dependent variable “GD diagnosis/no GD diagnosis” was performed for each group to examine the effect size. Covariates were gender, age, dopamine agonist prescription, and psychotropic drug prescription, respectively gender, age, and aripiprazole prescription. The covariate psychotropic drug prescription was applied in the group with dopaminergic drug prescriptions to distinguish a compromised mental health status in general from GD as a specific psychiatric diagnosis. The variable included all prescriptions within psycholeptics, psychoanaleptics, drugs used in addictive disorders and antihistamines used as tranquillisers (ATC codes N05, N06, N07B, R06AD01, R06AD02, and R06AD52). Since the subgroup for analysing aripiprazole prescriptions was chosen according to the psychiatric diagnosis of the patients, the use of the covariate psychotropic drug prescription was assumed to be redundant in this case. The number of patients with the outcome GD diagnosis was large enough to apply four, respectively three covariates, with 116 GD patients in the group with dopaminergic prescriptions and 307 GD patients in the subgroup chosen to analyse aripiprazole prescriptions. For the chi-square tests and the logistic regressions, a significance level of 0.05 was applied and all hypothesis testing was performed two-tailed on independent samples.

Considering a possible causal relation, the time that passed in between the first dopamine agonist or aripiprazole prescription and the GD diagnosis was calculated based on the earliest date of prescription and the earliest date of the GD diagnosis for each patient. This analysis was performed on all 104 patients with a dopamine agonist prescription and a GD diagnosis, and all 151 patients with an aripiprazole prescription and a GD diagnosis.

Results

The association between a dopamine agonist prescription and GD in patients with any dopaminergic drug prescription

The chi-square test showed a significant association between a dopamine agonist prescription and a GD diagnosis with a P-value (P) of 0.001. Of all patients with any dopaminergic drug prescription ($n = 180$), including dopamine agonists but also for instance levodopa, 116

Table 1. Frequency counts and row percentages for dopamine agonist prescriptions and GD diagnoses in patients with any dopaminergic prescription.

		No GD diagnosis	GD diagnosis	Total
No dopamine agonist prescription	Count	19	12	31
	Expected Count	11.0	20.0	31.0
	Row percentage	61.3%	38.7%	100.0%
Dopamine agonist prescription	Count	45	104	149
	Expected Count	53.0	96.0	149.0
	Row percentage	30.2%	69.8%	100.0%
Total	Count	64	116	180
	Expected Count	64.0	116.0	180.0
	Row percentage	35.6%	64.4%	100.0%

<https://doi.org/10.1371/journal.pone.0252516.t001>

(64.4%) had been diagnosed with GD. While 104 of 149 patients (69.8%) with a dopamine agonist prescription had a GD diagnosis, only 12 out of 31 patients (38.7%) without a dopamine agonist prescription had a GD diagnosis (see [Table 1](#)).

In the logistic regression, the covariates gender and age had no significant influence on the outcome GD diagnosis. The odds ratio (OR) for having a GD diagnosis was 3.2 (95% confidence interval (CI) = 1.4–7.6; $P = 0.008$) for patients who had received a dopamine agonist prescription compared to patients without a dopamine agonist prescription. In the same regression model, the OR for having a GD diagnosis when having a psychotropic drug prescription was 5.8 (95% CI = 1.9–17.5; $P = 0.002$) in comparison to patients without any psychotropic drug prescription. The complete characterisation of the applied regression model can be found in the [S3 Appendix](#).

When the date of the first dopamine agonist prescription was compared with the date of the GD diagnosis, a majority of the patients had received the prescription first (81 of 104). The time that passed between prescription and diagnosis where 3.5 years as a median with an interquartile range (IQR) between 0.6 and 6.8 years, including all 104 patients regardless if prescription or diagnosis occurred first (see [S5 Appendix](#)).

The association between an aripiprazole prescription and GD in F20-29 patients

Even for an aripiprazole prescription the association with a GD diagnosis could be confirmed by a chi-square test with a significance of $P < 0.001$. Within the 389 patients with a F20-29 diagnosis, 307 (78.9%) had a GD diagnosis. Out of 170 patients with an aripiprazole prescription a total of 151 (88.8%) had a GD diagnosis, while only 156 out of 219 patients (71.2%) without the prescription had been diagnosed with GD (see [Table 2](#)).

Table 2. Frequency counts and row percentages for aripiprazole prescriptions and GD diagnoses in patients with F20-29 diagnoses.

		No GD diagnosis	GD diagnosis	Total
No aripiprazole prescription	Count	63	156	219
	Expected Count	46.2	172.8	219.0
	Row percentage	28.8%	71.2%	100.0%
Aripiprazole prescription	Count	19	151	170
	Expected Count	35.8	134.2	170.0
	Row percentage	11.2%	88.8%	100.0%
Total	Count	82	307	389
	Expected Count	82.0	307.0	389.0
	Row percentage	21.1%	78.9%	100.0%

<https://doi.org/10.1371/journal.pone.0252516.t002>

The covariates gender and age had no significant effect on the outcome GD diagnosis in the logistic regression performed. However, the OR for having a GD diagnosis when having an aripiprazole prescription was 3.4 (95% CI = 1.9–6.1) with a significance of $P < 0.001$ in comparison to patients without an aripiprazole prescription. A full description of the regression model is included in the [S4 Appendix](#).

Most of the patients had received their first aripiprazole prescription before the GD diagnosis (101 of 151). The median time difference including all 151 patients was 1.5 years between prescription and diagnosis with an IQR between -1.3 years and 4.9 years (see [S5 Appendix](#)).

Discussion

The results of the statistical analysis were convincing and coherent. Dopamine agonists as well as aripiprazole had an association with an increased risk for GD. Given the study design, these findings can be expected to have a high reliability and validity. The study population was large with more than 11 000 individuals, of which 3689 were diagnosed with GD. Since the data was obtained from a nationwide public institution, it can be assumed to represent the situation in Sweden accurately. Furthermore, the reference groups for the statistical analysis were selected to reach the highest comparability possible. The controls were age- and gender-matched and the overall health state was expected to be homogenous within the reference groups. Thus, a GD diagnosis is presumably linked to the intake of medication rather than a poor health condition in general. The OR for dopamine agonists was calculated while using a psychotropic drug prescription as another covariate. Both covariates showed significant ORs, and it can therefore be assumed that GD is not only associated with a compromised mental health state but also with dopamine agonist intake independently. For analysing the effects of aripiprazole, a subgroup with similar psychiatric diagnoses was chosen. Thus, GD presumably is connected to aripiprazole intake and is not only a comorbidity to other psychiatric diagnoses.

Since a GD diagnosis was treated as a binary variable, the course of disease could not be taken into consideration in this study. Future study designs should focus on the progression of the disorder under certain medication in more detail, for instance by using patient questionnaires, which could also take subclinical problematic gambling into account. Due to the cross-sectional study design, a causal relationship between the medication and GD can only be a speculation, given that most patients received their prescriptions before their diagnosis. For exploring causality, a longitudinal study would be more adequate, in which follow-up screenings for GD and impaired impulse control could be performed on patients who received the medication in question. Another limitation of this study is the missing information about the actual medication intake since the data was obtained from the prescription register. To what extent the patients made use of the prescribed medication is uncertain. Even this issue could be avoided in the future by following up on certain patients instead of using register data.

In contrast to the well-recognised DOMINION study on impulse-control disorders as a reaction to dopamine agonists [38], this study did not focus on PD patients. It was an important step to further support the results found in PD patients with findings in non-PD patients, since PD is characterised by a pathological dopamine transmission, which might even interfere with the dopaminergic reward system. The results of this study are in line with the DOMINION study, as with many other publications on the subject. While the association between increased GD rates and dopamine agonists and modulators starts to be well established raising assumptions about a causal relationship, the understanding on a molecular level is very limited so far. Only few studies have addressed the direct connection between the pharmacodynamics of dopaminergic drugs and the neurobiology of GD. Activating the striatal D_3 receptor could

be the possible mode of action for dopamine agonists, which would explain a dose-dependent relationship and pramipexole and ropinirole as agonists that increase the risk for GD the most. The pharmacodynamics of aripiprazole on the other hand, are more complex and still subject to debate [51]. It acts as a partial agonist on D2-like receptors, thus yielding an antagonistic effect under high endogenous dopamine levels and functioning as a modulator. Aripiprazole additionally interacts with serotonergic receptors and has been observed to show functional selectivity on postsynaptic D2 receptors, activating a different intracellular signalling pattern than the natural ligand dopamine [56]. Therefore, the neurochemical link between aripiprazole and developing GD is even less understood to date than the influence of dopamine agonists on the disorder.

In summary, our study puts additional emphasis on the possible effects of dopamine agonists and modulators on GD development. Even if institutions like the U.S. Food and Drug Administration (FDA) have recognised the problem, the knowledge needs to be established broadly in the primary care centres. One beneficial measure could be to assess a patient's background and risk factors for GD and impulse-control disorders, before prescribing dopaminergic medication. If, for instance, the general mental health state indicates a higher likelihood to develop GD, it could be advisable to choose another pharmaceutical if possible, or to closely monitor the patient for side effects related to impaired impulse-control.

Conclusions

The association between the intake of certain dopaminergic pharmaceuticals and the development of GD has started to become an international consensus over the last years. This study provides further evidence for GD being a possible response to dopamine agonists or aripiprazole. A causal relationship is likely, and the dopaminergic reward system seems to be the linkage between the medication and the addictive disorder. Since the molecular background of GD is only partly understood to date, it remains unclear how dopamine agonists or aripiprazole can influence its pathophysiology. The possible association with GD should be taken into consideration when prescribing dopamine agonists and aripiprazole, and for patients at risk for a compromised impulse control other medication seems more suitable. Furthermore, an extended knowledge on the matter is necessary to develop dopaminergic medication that can act more specifically on its intended target without causing impulse-control disorders and addictions. Revealing the molecular mechanisms behind GD would in addition contribute to a better understanding of addictive disorders in general and would open new possibilities for pharmacotherapy in this field.

Supporting information

S1 Appendix. Characterisation of the subgroup “Patients with any dopaminergic drug prescription”.

(DOCX)

S2 Appendix. Characterisation of the subgroup “Patients with a F20-29 diagnosis”.

(DOCX)

S3 Appendix. Logistic regression in the subgroup “Patients with any dopaminergic drug prescription”.

(DOCX)

S4 Appendix. Logistic regression in the subgroup “Patients with a F20-29 diagnosis”.

(DOCX)

S5 Appendix. Comparison between the date of GD diagnosis and the first prescription date for DAs and aripiprazole.

(DOCX)

Author Contributions

Conceptualization: Anders Håkansson.

Formal analysis: Mirjam Wolfschlag.

Funding acquisition: Anders Håkansson.

Investigation: Mirjam Wolfschlag.

Methodology: Mirjam Wolfschlag, Anders Håkansson.

Project administration: Anders Håkansson.

Resources: Anders Håkansson.

Supervision: Anders Håkansson.

Validation: Mirjam Wolfschlag.

Visualization: Mirjam Wolfschlag.

Writing – original draft: Mirjam Wolfschlag.

Writing – review & editing: Mirjam Wolfschlag, Anders Håkansson.

References

1. Potenza MN, Balodis IM, Derevensky J, Grant JE, Petry NM, Verdejo-Garcia A, et al. Gambling disorder. *Nature reviews Disease primers*. 2019; 5(1):51. Epub 2019/07/28. <https://doi.org/10.1038/s41572-019-0099-7> PMID: 31346179.
2. Calado F, Griffiths MD. Problem gambling worldwide: An update and systematic review of empirical research (2000–2015). *Journal of behavioral addictions*. 2016; 5(4):592–613. Epub 2016/10/28. <https://doi.org/10.1556/2006.5.2016.073> PMID: 27784180; PubMed Central PMCID: PMC5370365.
3. Håkansson A, Fernández-Aranda F, Menchón JM, Potenza MN, Jiménez-Murcia S. Gambling During the COVID-19 Crisis—A Cause for Concern. *Journal of addiction medicine*. 2020; 14(4):e10–e2. Epub 2020/05/21. <https://doi.org/10.1097/ADM.0000000000000690> PMID: 32433365; PubMed Central PMCID: PMC7273946 present work specifically. The authors declare that they have no conflicts of interest with respect to the content of this manuscript.
4. Dowling NA, Suomi A, Jackson AC, Lavis T. Problem Gambling Family Impacts: Development of the Problem Gambling Family Impact Scale. *Journal of gambling studies*. 2016; 32(3):935–55. Epub 2015/11/04. <https://doi.org/10.1007/s10899-015-9582-6> PMID: 26527482.
5. Karlsson A, Håkansson A. Gambling disorder, increased mortality, suicidality, and associated comorbidity: A longitudinal nationwide register study. *Journal of behavioral addictions*. 2018; 7(4):1091–9. Epub 2018/11/15. <https://doi.org/10.1556/2006.7.2018.112> PMID: 30427214; PubMed Central PMCID: PMC6376387.
6. Moghaddam JF, Yoon G, Dickerson DL, Kim SW, Westermeyer J. Suicidal ideation and suicide attempts in five groups with different severities of gambling: Findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *The American journal on addictions*. 2015; 24(4):292–8. Epub 2015/03/27. <https://doi.org/10.1111/ajad.12197> PMID: 25808267.
7. Jolly T, Trivedi C, Adnan M, Mansuri Z, Agarwal V. Gambling in patients with major depressive disorder is associated with an elevated risk of Suicide: Insights from 12-years of Nationwide inpatient sample data. *Addictive behaviors*. 2021; 118:106872. Epub 2021/03/07. <https://doi.org/10.1016/j.addbeh.2021.106872> PMID: 33676161.
8. Kessler RC, Hwang I, LaBrie R, Petukhova M, Sampson NA, Winters KC, et al. DSM-IV pathological gambling in the National Comorbidity Survey Replication. *Psychological medicine*. 2008; 38(9):1351–60. Epub 2008/02/09. <https://doi.org/10.1017/S0033291708002900> PMID: 18257941; PubMed Central PMCID: PMC2293303.

9. World Health Organization. ICD-11 for Mortality and Morbidity Statistics (Version: 09/2020) 2020 [05.01.2021]. Available from: <https://icd.who.int/browse11/l-m/en#/http%3a%2f%2fid.who.int%2fid%2fentity%2f1041487064>.
10. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders 5th edn. 2013.
11. Balodis IM, Potenza MN. Common neurobiological and psychological underpinnings of gambling and substance-use disorders. *Progress in neuro-psychopharmacology & biological psychiatry*. 2020; 99:109847. Epub 2019/12/22. <https://doi.org/10.1016/j.pnpbp.2019.109847> PMID: 31862419.
12. Petruccioli F, Diotaiuti P, Verrastro V, Petruccioli I, Carenti ML, De Berardis D, et al. Obsessive-compulsive aspects and pathological gambling in an Italian sample. *BioMed research international*. 2014; 2014:167438. Epub 2014/07/25. <https://doi.org/10.1155/2014/167438> PMID: 25057479; PubMed Central PMCID: PMC4095731.
13. Leeman RF, Potenza MN. Similarities and differences between pathological gambling and substance use disorders: a focus on impulsivity and compulsivity. *Psychopharmacology*. 2012; 219(2):469–90. Epub 2011/11/08. <https://doi.org/10.1007/s00213-011-2550-7> PMID: 22057662; PubMed Central PMCID: PMC3249521.
14. Potenza MN. Neurobiology of gambling behaviors. *Current opinion in neurobiology*. 2013; 23(4):660–7. Epub 2013/04/02. <https://doi.org/10.1016/j.conb.2013.03.004> PMID: 23541597; PubMed Central PMCID: PMC3803105.
15. Leeman RF, Potenza MN. A targeted review of the neurobiology and genetics of behavioural addictions: an emerging area of research. *Canadian journal of psychiatry Revue canadienne de psychiatrie*. 2013; 58(5):260–73. Epub 2013/06/13. <https://doi.org/10.1177/070674371305800503> PMID: 23756286; PubMed Central PMCID: PMC3762982.
16. Hyman SE, Malenka RC, Nestler EJ. Neural mechanisms of addiction: the role of reward-related learning and memory. *Annual review of neuroscience*. 2006; 29:565–98. Epub 2006/06/17. <https://doi.org/10.1146/annurev.neuro.29.051605.113009> PMID: 16776597.
17. Zack M, St George R, Clark L. Dopaminergic signaling of uncertainty and the aetiology of gambling addiction. *Progress in neuro-psychopharmacology & biological psychiatry*. 2020; 99:109853. Epub 2019/12/25. <https://doi.org/10.1016/j.pnpbp.2019.109853> PMID: 31870708.
18. Linnet J, Mouridsen K, Peterson E, Møller A, Doudet DJ, Gjedde A. Striatal dopamine release codes uncertainty in pathological gambling. *Psychiatry research*. 2012; 204(1):55–60. Epub 2012/08/15. <https://doi.org/10.1016/j.psychresns.2012.04.012> PMID: 22889563.
19. Clark L, Boileau I, Zack M. Neuroimaging of reward mechanisms in Gambling disorder: an integrative review. *Molecular psychiatry*. 2019; 24(5):674–93. Epub 2018/09/15. <https://doi.org/10.1038/s41380-018-0230-2> PMID: 30214041.
20. Clark L, Stokes PR, Wu K, Michalczuk R, Benecke A, Watson BJ, et al. Striatal dopamine D₂/D₃ receptor binding in pathological gambling is correlated with mood-related impulsivity. *NeuroImage*. 2012; 63(1):40–6. Epub 2012/07/11. <https://doi.org/10.1016/j.neuroimage.2012.06.067> PMID: 22776462; PubMed Central PMCID: PMC3438449.
21. Linnet J, Møller A, Peterson E, Gjedde A, Doudet D. Dopamine release in ventral striatum during Iowa Gambling Task performance is associated with increased excitement levels in pathological gambling. *Addiction (Abingdon, England)*. 2011; 106(2):383–90. Epub 2010/10/05. <https://doi.org/10.1111/j.1360-0443.2010.03126.x> PMID: 20883460.
22. Joutsa J, Johansson J, Niemelä S, Ollikainen A, Hirvonen MM, Piepponen P, et al. Mesolimbic dopamine release is linked to symptom severity in pathological gambling. *NeuroImage*. 2012; 60(4):1992–9. Epub 2012/02/22. <https://doi.org/10.1016/j.neuroimage.2012.02.006> PMID: 22348881.
23. Linnet J, Møller A, Peterson E, Gjedde A, Doudet D. Inverse association between dopaminergic neurotransmission and Iowa Gambling Task performance in pathological gamblers and healthy controls. *Scandinavian journal of psychology*. 2011; 52(1):28–34. Epub 2010/08/14. <https://doi.org/10.1111/j.1467-9450.2010.00837.x> PMID: 20704689.
24. van Holst RJ, Sescousse G, Janssen LK, Janssen M, Berry AS, Jagust WJ, et al. Increased Striatal Dopamine Synthesis Capacity in Gambling Addiction. *Biological psychiatry*. 2018; 83(12):1036–43. Epub 2017/07/22. <https://doi.org/10.1016/j.biopsych.2017.06.010> PMID: 28728675; PubMed Central PMCID: PMC6698370.
25. Zack M, Poulos CX. Amphetamine primes motivation to gamble and gambling-related semantic networks in problem gamblers. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*. 2004; 29(1):195–207. Epub 2003/10/23. <https://doi.org/10.1038/sj.npp.1300333> PMID: 14571257.
26. Boileau I, Payer D, Chugani B, Lobo DS, Houle S, Wilson AA, et al. In vivo evidence for greater amphetamine-induced dopamine release in pathological gambling: a positron emission tomography study with

- [(11)C]-(+)-PHNO. *Molecular psychiatry*. 2014; 19(12):1305–13. Epub 2013/12/11. <https://doi.org/10.1038/mp.2013.163> PMID: 24322203.
27. Smart K, Desmond RC, Poulos CX, Zack M. Modafinil increases reward salience in a slot machine game in low and high impulsivity pathological gamblers. *Neuropharmacology*. 2013; 73:66–74. Epub 2013/05/29. <https://doi.org/10.1016/j.neuropharm.2013.05.015> PMID: 23711549.
 28. Zack M, Poulos CX. A D2 antagonist enhances the rewarding and priming effects of a gambling episode in pathological gamblers. *Neuropsychopharmacology: official publication of the American College of Neuropsychopharmacology*. 2007; 32(8):1678–86. Epub 2007/01/05. <https://doi.org/10.1038/sj.npp.1301295> PMID: 17203013.
 29. Boileau I, Payer D, Chugani B, Lobo D, Behzadi A, Rusjan PM, et al. The D2/3 dopamine receptor in pathological gambling: a positron emission tomography study with [11C]-(+)-propyl-hexahydro-naphtho-oxazin and [11C]raclopride. *Addiction (Abingdon, England)*. 2013; 108(5):953–63. Epub 2012/11/22. <https://doi.org/10.1111/add.12066> PMID: 23167711.
 30. Fong T, Kalechstein A, Bernhard B, Rosenthal R, Rugle L. A double-blind, placebo-controlled trial of olanzapine for the treatment of video poker pathological gamblers. *Pharmacology, biochemistry, and behavior*. 2008; 89(3):298–303. Epub 2008/02/12. <https://doi.org/10.1016/j.pbb.2007.12.025> PMID: 18261787.
 31. McElroy SL, Nelson EB, Welge JA, Kaehler L, Keck PE Jr. Olanzapine in the treatment of pathological gambling: a negative randomized placebo-controlled trial. *The Journal of clinical psychiatry*. 2008; 69(3):433–40. Epub 2008/02/07. <https://doi.org/10.4088/jcp.v69n0314> PMID: 18251624.
 32. Molina JA, Sáinz-Artiga MJ, Fraile A, Jiménez-Jiménez FJ, Villanueva C, Ortí-Pareja M, et al. Pathologic gambling in Parkinson's disease: a behavioral manifestation of pharmacologic treatment? *Movement disorders: official journal of the Movement Disorder Society*. 2000; 15(5):869–72. Epub 2000/09/29. [https://doi.org/10.1002/1531-8257\(200009\)15:5<869::aid-mds1016>3.0.co;2-i](https://doi.org/10.1002/1531-8257(200009)15:5<869::aid-mds1016>3.0.co;2-i) PMID: 11009192.
 33. Gschwandtner U, Aston J, Renaud S, Fuhr P. Pathologic gambling in patients with Parkinson's disease. *Clinical neuropharmacology*. 2001; 24(3):170–2. Epub 2001/06/08. <https://doi.org/10.1097/00002826-200105000-00009> PMID: 11391129.
 34. Driver-Dunckley E, Samanta J, Stacy M. Pathological gambling associated with dopamine agonist therapy in Parkinson's disease. *Neurology*. 2003; 61(3):422–3. Epub 2003/08/13. <https://doi.org/10.1212/01.wnl.0000076478.45005.ec> PMID: 12913220.
 35. Ondo WG, Lai D. Predictors of impulsivity and reward seeking behavior with dopamine agonists. *Parkinsonism & related disorders*. 2008; 14(1):28–32. Epub 2007/08/19. <https://doi.org/10.1016/j.parkreldis.2007.05.006> PMID: 17702628.
 36. Lee JY, Kim JM, Kim JW, Cho J, Lee WY, Kim HJ, et al. Association between the dose of dopaminergic medication and the behavioral disturbances in Parkinson disease. *Parkinsonism & related disorders*. 2010; 16(3):202–7. Epub 2009/12/19. <https://doi.org/10.1016/j.parkreldis.2009.12.002> PMID: 20018549.
 37. Perez-Lloret S, Rey MV, Fabre N, Ory F, Spampinato U, Brefel-Courbon C, et al. Prevalence and pharmacological factors associated with impulse-control disorder symptoms in patients with Parkinson disease. *Clinical neuropharmacology*. 2012; 35(6):261–5. Epub 2012/11/06. <https://doi.org/10.1097/WNF.0b013e31826e6e6d> PMID: 23123689.
 38. Weintraub D, Koester J, Potenza MN, Siderowf AD, Stacy M, Voon V, et al. Impulse control disorders in Parkinson disease: a cross-sectional study of 3090 patients. *Archives of neurology*. 2010; 67(5):589–95. Epub 2010/05/12. <https://doi.org/10.1001/archneurol.2010.65> PMID: 20457959.
 39. Rizos A, Sauerbier A, Antonini A, Weintraub D, Martinez-Martin P, Kessel B, et al. A European multicentre survey of impulse control behaviours in Parkinson's disease patients treated with short- and long-acting dopamine agonists. *European journal of neurology*. 2016; 23(8):1255–61. Epub 2016/05/14. <https://doi.org/10.1111/ene.13034> PMID: 27170229.
 40. Grall-Bronnec M, Sauvaget A, Perrouin F, Leboucher J, Etcheverrigaray F, Challet-Bouju G, et al. Pathological Gambling Associated With Aripiprazole or Dopamine Replacement Therapy: Do Patients Share the Same Features? A Review. *Journal of clinical psychopharmacology*. 2016; 36(1):63–70. Epub 2015/12/15. <https://doi.org/10.1097/JCP.0000000000000444> PMID: 26658263; PubMed Central PMCID: PMC4700874.
 41. Scavone C, Stelitano B, Rafaniello C, Rossi F, Sportiello L, Capuano A. Drugs-Induced Pathological Gambling: An Analysis of Italian Spontaneous Reporting System. *Journal of gambling studies*. 2020; 36(1):85–96. Epub 2019/01/24. <https://doi.org/10.1007/s10899-019-09828-1> PMID: 30671764; PubMed Central PMCID: PMC7026297.
 42. Poletti M, Logi C, Lucetti C, Del Dotto P, Baldacci F, Vergallo A, et al. A single-center, cross-sectional prevalence study of impulse control disorders in Parkinson disease: association with dopaminergic

- drugs. *Journal of clinical psychopharmacology*. 2013; 33(5):691–4. Epub 2013/07/17. <https://doi.org/10.1097/JCP.0b013e3182979830> PMID: 23857310.
43. Garcia-Ruiz PJ, Martinez Castrillo JC, Alonso-Canovas A, Herranz Barcenos A, Vela L, Sanchez Alonso P, et al. Impulse control disorder in patients with Parkinson's disease under dopamine agonist therapy: a multicentre study. *Journal of neurology, neurosurgery, and psychiatry*. 2014; 85(8):840–4. Epub 2014/01/18. <https://doi.org/10.1136/jnnp-2013-306787> PMID: 24434037.
 44. Lanteri PF, Leguia A, Doladé NG, García GC, Figueras A. Drug-induced gambling disorder: A not so rare but underreported condition. *Psychiatry research*. 2018; 269:593–5. Epub 2018/09/12. <https://doi.org/10.1016/j.psychres.2018.09.008> PMID: 30205352.
 45. Piercey MF. Pharmacology of pramipexole, a dopamine D3-preferring agonist useful in treating Parkinson's disease. *Clinical neuropharmacology*. 1998; 21(3):141–51. Epub 1998/06/09. PMID: 9617505.
 46. Jost WH, Angersbach D. Ropinirole, a Non-Ergoline Dopamine Agonist. *CNS Drug Reviews*. 2005; 11(3):253–72. <https://doi.org/10.1111/j.1527-3458.2005.tb00046.x> PMID: 16389293
 47. Alonso Cánovas A, Luquin Piudo R, García Ruiz-Espiga P, Burguera JA, Campos Arillo V, Castro A, et al. Dopaminergic agonists in Parkinson's disease. *Neurologia (Barcelona, Spain)*. 2014; 29(4):230–41. Epub 2011/07/05. <https://doi.org/10.1016/j.nrl.2011.04.012> PMID: 21724302.
 48. Trenkwalder C, Allen R, Högl B, Clemens S, Patton S, Schormair B, et al. Comorbidities, treatment, and pathophysiology in restless legs syndrome. *The Lancet Neurology*. 2018; 17(11):994–1005. Epub 2018/09/25. [https://doi.org/10.1016/S1474-4422\(18\)30311-9](https://doi.org/10.1016/S1474-4422(18)30311-9) PMID: 30244828.
 49. Seeman P. Parkinson's disease treatment may cause impulse-control disorder via dopamine D3 receptors. *Synapse (New York, NY)*. 2015; 69(4):183–9. Epub 2015/02/04. <https://doi.org/10.1002/syn.21805> PMID: 25645960.
 50. Marder SR, McQuade RD, Stock E, Kaplita S, Marcus R, Safferman AZ, et al. Aripiprazole in the treatment of schizophrenia: safety and tolerability in short-term, placebo-controlled trials. *Schizophrenia research*. 2003; 61(2–3):123–36. Epub 2003/05/06. [https://doi.org/10.1016/s0920-9964\(03\)00050-1](https://doi.org/10.1016/s0920-9964(03)00050-1) PMID: 12729864.
 51. Tuplin EW, Holahan MR. Aripiprazole, A Drug that Displays Partial Agonism and Functional Selectivity. *Current neuropharmacology*. 2017; 15(8):1192–207. Epub 2017/04/18. <https://doi.org/10.2174/1570159X15666170413115754> PMID: 28412910; PubMed Central PMCID: PMC5725548.
 52. Stocchi F, Radicati FG, Torti M. Drug safety evaluation of ropinirole prolonged release. *Expert opinion on drug safety*. 2014; 13(3):383–9. Epub 2014/02/05. <https://doi.org/10.1517/14740338.2014.870152> PMID: 24490799.
 53. Torti M, Bravi D, Vacca L, Stocchi F. Are All Dopamine Agonists Essentially the Same? *Drugs*. 2019; 79(7):693–703. Epub 2019/04/11. <https://doi.org/10.1007/s40265-019-01103-2> PMID: 30968290.
 54. U.S. Food and Drug Administration. FDA Drug Safety Communication: FDA warns about new impulse-control problems associated with mental health drug aripiprazole (Abilify, Abilify Maintena, Aristada) 2016 [11.01.2021]. Available from: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-drug-safety-communication-fda-warns-about-new-impulse-control-problems-associated-mental-health>.
 55. Etminan M, Sodhi M, Samii A, Procyshyn RM, Guo M, Carleton BC. Risk of Gambling Disorder and Impulse Control Disorder With Aripiprazole, Pramipexole, and Ropinirole: A Pharmacoepidemiologic Study. *Journal of clinical psychopharmacology*. 2017; 37(1):102–4. Epub 2016/12/09. <https://doi.org/10.1097/JCP.0000000000000634> PMID: 27930495.
 56. Pan B, Chen J, Lian J, Huang XF, Deng C. Unique Effects of Acute Aripiprazole Treatment on the Dopamine D2 Receptor Downstream cAMP-PKA and Akt-GSK3 β Signalling Pathways in Rats. *PloS one*. 2015; 10(7):e0132722. Epub 2015/07/15. <https://doi.org/10.1371/journal.pone.0132722> PMID: 26162083; PubMed Central PMCID: PMC4498891.