

Zinc status and dietary quality of type 2 diabetic patients: implication of physical activity level*

Jin-Sook Yoon[§]

Department of Food and Nutrition, Keimyung University, Daegu 704-701, Korea

Received February 19, 2008; Revised March 10, 2008; Accepted March 24, 2008

Abstract

The purpose of this study was to analyze the relationships among zinc status, diet quality, glycemic control and self-rated physical activity level of type 2 diabetic patients. Dietary intakes for two non-consecutive days were measured by 24-hour recall method for seventy-six diabetic patients. Fasting blood glucose and HbA1c were measured for the assessment of glycemic control. We evaluated the extent of dietary adequacy by the percentage of subjects with a dietary intake of a nutrient less than the estimated average requirement(EAR), the dietary diversity score(DDS) and the dietary variety score(DVS). Zinc status was assessed from serum levels and urinary excretion. Dietary inadequacy was serious for five nutrients: riboflavin, calcium, thiamin, zinc and vitamin C. Dietary intakes from the meat, fish, and egg food groups and the milk food group were below the recommended level. We found that subjects with high levels of physical activities had significantly higher DVS and serum zinc levels compared to others ($p < 0.05$). Fasting blood glucose levels and HbA1c were not significantly different across self-reported physical activity levels. Therefore, we suggest that maintaining physical activity at or above a moderate level is beneficial to improving dietary quality and zinc status.

Key Words: Diabetes, zinc status, diet quality, physical activity level, glycemic control

Introduction

The incidence of diabetes among the Korean population has continuously increased in the recent twenty years (Kim *et al.*, 1998). Thus, the high prevalence and death rate from diabetes in Korea now places the country in the highest quartile group among the OECD countries (Ministry of Health & Welfare, 2007). The National Health and Nutrition Examination Survey for Koreans conducted in 2005 showed that the prevalence of diabetes was 9.0% for males and 7.2% for female adults (Ministry of Health & Welfare 2006).

Type 2 diabetic patients are characterized by reduced secretion of insulin and increased insulin resistance, resulting in elevated blood glucose levels (Salgueiro *et al.*, 2001). Zinc is known to be an essential micronutrient involved in the physiological action of insulin (Roth & Kirchgessner, 1981). Animal studies suggested that glucose tolerance in conjunction with insulin resistance aggravated after zinc deprivation (Kinlaw *et al.*, 1992; Salgueiro *et al.*, 2001). However, the relationship of zinc deficiency and glycemic control in diabetic patients has not been well established. Inadequate zinc intake induces higher absorption throughout the intestine resulting in less urinary excretion by homeostatic compensation mechanism in healthy subjects. Meanwhile, increased urinary excretion of zinc has been observed

in several studies (Chausmer 1998; Cunningham *et al.*, 1994; Salgueiro *et al.*, 2001).

Previous studies indicated that the zinc status of Korean diabetic patients is at the marginal level (Lee *et al.*, 2005; Yoon & Lee, 2007). Zinc intake has a positive relationship with the protein and fat intake, but is negatively correlated with the carbohydrate intake. Foods with high amounts of phytate were the major source of zinc, and therefore, did not contribute to increasing zinc densities of frequently consumed foods (Yoon & Lee, 2007).

It is known that regular physical activity lowers the risk of type 2 diabetes mellitus by 20-60% in dose-related manner. The size of the effect is greatest in subjects with high risk of non-insulin dependent diabetes (Vuori, 2001). Therefore, maintaining a moderate physical activity pattern along with dietary management is regarded as an essential element for glycemic control and weight reduction of diabetic patients. However, it is not yet explicated whether physical activity level is related to improving zinc status as well. Therefore, in this study, we assessed the quality of diet, glycemic control, and zinc status of diabetic patients and normal adults by using self-reported physical activity levels to figure out the association among physical activity level, glycemic control, and zinc nutrition. In particular, we aimed to explicate whether the diabetic patients

* This research was supported by a 2006 sabbatical research grant of Keimyung University.

[§] Corresponding Author: Jin-Sook Yoon, Tel. 82-53-580-5873, Fax. 82-53-580-5885, Email. jsook@kmu.ac.kr

with proper physical activity levels could also maintain better dietary practices, thereby improving their zinc status.

Subjects and Methods

Selection of subjects

Among adults visiting community public health centers or local clinics in the Daegu area, seventy-six type 2 diabetic patients with no known medical problems voluntarily participated in this study. In order to collect general characteristics such as age, education, physical activity level, medication, diabetic duration and complication, patients were interviewed by a trained dietitian.

Anthropometric assessment

While the subjects were wearing light indoor clothes and no shoes, body weight was measured to the nearest 0.1 kg, and height was measured to the nearest 0.5 cm. Circumferences of waist and hip were measured to the nearest 0.1 cm. Body mass index (BMI) was calculated from measured height and weight for each subject. Body fat content and percentage body fat were assessed using the Bio-electrical Impedance Fatness Analyzer (GIF-891).

Assessment of dietary intake

Information on the dietary intake of each subject was collected for two non-consecutive days during the week by the 24-hour dietary recall method. Food pictures were provided for the subjects to improve the accuracy of estimation of actual intake (Korean Food Research Institute, 1998).

Food intakes were converted to the amount of daily nutrients intake by CAN-Pro 3.0 (developed by Korean Nutrition Society). Dietary inadequacy of each nutrient was estimated from the proportion of subjects with intakes below the Estimated Average Requirement (EAR). Dietary diversity score (DDS) was assessed by the six-food group system described in the Dietary Reference Intakes for Koreans (KDRIs) (Korean Nutrition Society, 2005). Dietary variety score (DVS) was calculated as the average total number of food items consumed in a day by each subject from the 24-hour recall data.

Measurement of physical activity level

Three levels of physical activity (i.e. light, moderate, and heavy) were constructed as follows. Participants with a sedentary life style were classified into the "light physical activity" group. Those who walk or cycle regularly more than once a week were placed in the "moderate physical activity" group. Subjects engaging in regular exercise such as jogging or swimming at least 4-5 times a week were sorted into the "heavy physical activity" group. After explaining the typical activity patterns

representing light, moderate, heavy physical activities, each subject was asked to rate his or her physical activity level into one of the three levels: light, moderate or heavy.

Biochemical assessment

As for biochemical markers of individual zinc status, we measured the zinc levels in serum and urine. A fasting blood of 10 ml was collected from each subject and prepared serum samples were stored at -70°C . Serum glucose levels and HbA1c were measured for the assessment of blood glucose control. Urine samples for 24 hours were collected for one day from each subject. Urinary creatinine was measured using Hawk's method (Hawk, 1954). Urinary zinc was analyzed by Inductively Coupled Plasma Emission Spectroscopy (ICP) (Jobin Yvon, 38 plus, France). Serum level of zinc was assessed by atomic absorption spectrometry.

Statistical analysis

The Statistical Analysis System (SAS) Package was used to analyze the relationship among glycemic control status, dietary factors, and physical activity levels. Data were presented as mean \pm SD. The statistical significance of differences in dietary quality, zinc status, and glycemic control among different physical activity level groups was tested by using one-way analysis of variance (ANOVA) and Duncan's multiple range test.

Results

Subject characteristics

The general characteristics of the subjects are described in

Table 1. General characteristics of subjects

Variables	Male (n=34)	Female (n=42)
Age (years)	n (%)	n (%)
< 50	10 (29.4)	5 (11.9)
50-60	14 (41.2)	10 (23.8)
60 <	10 (29.4)	27 (64.3)
Education		
None	1 (3.3)	12 (31.6)
Primary	5 (16.7)	12 (31.6)
Middle	4 (13.3)	6 (15.8)
High	14 (46.7)	8 (21.1)
College <	6 (20.0)	0 (0.0)
Diabetic duration (years)		
< 1 yrs	12 (35.3)	16 (38.1)
1-10 yrs	16 (47.1)	17 (40.5)
10 yrs <	6 (17.7)	9 (21.4)
Diabetic complications		
Have	4 (13.3)	13 (37.1)
None	26 (86.7)	22 (62.9)

Table 2. Anthropometric assessment of the subjects¹⁾²⁾

Variables	Male (n=34)	Female (n=42)	p-value
Diabetic duration (yrs)	5.0 ± 6.2	6.5 ± 4.6	NS
Height (cm)	167.2 ± 7.6	157.3 ± 6.6	< 0.0001
Weight (kg)	66.1 ± 10.4	61.2 ± 8.7	< 0.05
Waist (cm)	84.5 ± 12.0	84.5 ± 8.7	NS
Hip (cm)	96.9 ± 5.9	96.8 ± 6.0	NS
BMI (kg/m ²)	23.6 ± 3.1	24.7 ± 2.8	NS
WHR	0.87 ± 0.11	0.87 ± 0.05	NS
Body fat ratio (%)	23.8 ± 4.7	30.8 ± 5.1	< 0.0001

¹⁾ Values are mean ± SD.

²⁾ NS : statistically not significant

Table 3. Percentage of subjects with the nutrients intake less than EAR

Nutrients	N (%)
Protein	10 (13.2)
Vitamin C	40 (52.6)
Thiamin	48 (63.2)
Riboflavin	63 (82.9)
Niacin	36 (47.4)
Ca	58 (76.3)
P	12 (15.8)
Fe	13 (17.1)
Zn	43 (56.6)

Table 4. Mean intake of each food group in diabetic subjects¹⁾

	Servings
Grain & starch	3.2 ± 1.1
Meat, poultry	3.1 ± 1.9
Vegetables	5.7 ± 2.5
Fruit	1.5 ± 1.5
Milk	0.4 ± 0.6
Oils & sugars	3.6 ± 3.8

¹⁾ Values are mean ± SD.

Table 1. The age distribution indicated that female diabetic patients who participated in this study were older than their male counterparts. As indicated in Table 2, there were no significant differences in body mass index (BMI), waist circumference, hip circumference, and waist-hip ratio (WHR) between males and females. Diabetic duration was 5.0 ± 6.2 years for men, 6.5 ± 4.6 for women.

Dietary quality and nutrient intake

Table 3 shows the profile of nutrition inadequacy estimated by the percentage of subjects whose nutrients intake was less than the EAR. More than half of the subjects showed dietary inadequacy for five nutrients such as riboflavin (= 82.9%), calcium (= 76.3%), thiamin (= 63.2%), zinc (= 56.6%) and vitamin C (=52.6%).

The average daily intake of each food group classified by KDRIs is presented in Table 4. Dietary intakes from the meat, poultry, fish, egg and bean (MPFEB) food groups and the milk food group were lower than the recommended level suggested

Table 5. Dietary zinc intake, serum and urinary zinc status by physical activity levels¹⁾²⁾

Activity Level	Zinc intake (mg)	Zn intake (%)	Serum Zn (mg/L)	Urinary Zn (mg/L)
Light (n=45)	6.7 ± 1.9 ^a	61.0 ± 15.0 ^a	0.77 ± 0.28 ^b	0.59 ± 0.36 ^a
Moderate (n=19)	7.0 ± 2.2 ^a	66.9 ± 20.3 ^a	0.90 ± 0.26 ^{ab}	0.45 ± 0.36 ^a
Heavy (n=9)	8.1 ± 2.2 ^a	69.7 ± 17.6 ^a	1.12 ± 0.19 ^a	0.71 ± 0.42 ^a
Total (n=73)	6.9 ± 2.0	63.3 ± 16.9	0.85 ± 0.29	0.57 ± 0.36

¹⁾ Values are mean ± SD.

²⁾ Values with the same superscript in the column were not significantly different at p < 0.05.

Table 6. Fasting blood glucose and HbA1c by physical activity levels¹⁾²⁾

Activity Levels	HbA1c (%)	Glucos (mg/dL)
Light (n=45)	7.9 ± 1.5 ^a	147.7 ± 63.2 ^a
Moderate (n=19)	7.5 ± 0.9 ^a	136.9 ± 33.3 ^a
Heavy (n=9)	7.2 ± 1.3 ^a	137.6 ± 67.1 ^a
Total (n=73)	7.7 ± 1.3	145.6 ± 56.2

¹⁾ Values are mean ± SD.

²⁾ Values with the same superscript in the column were not significantly different at p < 0.05.

Table 7. Diet diversity score and diet variety score by physical activity levels¹⁾²⁾

Activity Level	DDS ³⁾	DVS ⁴⁾
Light(n=45)	4.1 ± 0.6 ^a	18.2 ± 5.6 ^b
Moderate(n=19)	3.8 ± 0.7 ^a	20.4 ± 6.9 ^{ab}
Heavy(n=9)	4.0 ± 1.0 ^a	25.7 ± 2.5 ^a
Total(n=73)	4.1 ± 0.6	19.7 ± 6.3

¹⁾ Values are mean ± SD.

²⁾ Values with the same superscript in the column were not significantly different at p < 0.05.

³⁾ DDS : Diet diversity score

⁴⁾ DVS : Diet variety score

for 1800 kcal of energy intake level.

Zinc status, glycemic control and dietary quality by physical activity level

Results for dietary zinc intakes, as indicated in Table 5, show that zinc intake of the participants was not significantly different across groups classified by self-reported physical activity levels. However, we observed a tendency for higher zinc intake among those who reported “heavy” level of physical activity. Serum zinc level of diabetic patients was significantly lower for the “light” physical activity group, whereas no significant difference was found in urinary zinc excretion.

Table 6 compares the results of HbA1c and fasting blood glucose, which are possible indicators of glycemic control, by the physical activity level. Both fasting blood glucose and HbA1c were slightly higher in the “light” physical activity group, compared to those in the “moderate” or “heavy” physical activity group. However, these differences were not significant.

Table 7 shows the DDS and the DVS of the participants by physical activity levels. While the DDS was not significantly different across physical activity levels, the DVS was significantly lower among those engaging in light physical activities.

Discussion

In this study, we compared the zinc status, glycemic control indices, and the dietary quality of diabetic patients by their self-reported level of physical activity.

In general, diabetic patients showed a poorer intake of nutrients than the average Korean adult reported by National Health and Nutrition Examination Survey in 2005. In particular, a higher prevalence of riboflavin and calcium inadequacy was observed for diabetic patients and analyzed by the percentage of subjects with dietary intakes less than the EAR. This is possibly attributable to the insufficient consumption of milk. Our data demonstrated that the dietary intake of the meat, poultry, fish, egg, and bean food groups, and the milk food group, was lower than the daily-recommended level of 1800 kcal (Korean Nutrition Society, 2005).

There seems to be no general consensus on the protein intake for diabetes. Eating less protein has been recommended to avoid accelerating renal complications. Human trials, in which meat was withdrawn from the diet, have been reported to successfully reduce albuminuria in type 2 diabetes with macroalbuminuria (de Mello *et al.*, 2006). However, other investigators suggested that there is no reason to restrict protein in diabetes without clear renal indications to do so (American Dietetic Association, 2002; Eckel, 2006). Considering the fact that the bioavailability of zinc from animal sources is higher than from grain or vegetables, less intake from the animal food group could lead to inadequate zinc status in diabetic patients. Concerns on marginal zinc deficiency have also been reported by other investigators (Chausmer 1998; Kwun & Kwon, 2000; Lee *et al.*, 2005; Lonnerdal, 2000; Salgueiro *et al.*, 2001).

Most strategies of diabetes management encourage physical activity as well as healthy food choice to improve blood glucose control and weight loss for type 2 diabetes as increased physical activity can lead to improved glycemia and decreased insulin resistance (American Dietetic Association, 2002; Vuori, 2001). However, we did not find any statistically significant difference of glycemic control indicators across different physical activity levels. A decrease in blood glucose levels and HbA1c along with higher level of physical activity was observed in diabetic patients. Since the physical activity level was solely based on self-rating without any objective measurement for the detailed activity pattern of each subject, the classification of activity levels was not accurate enough to detect small differences in indicators across groups. It has been already known that diabetic patients benefit from regular physical activity in many ways such as increased insulin sensitivity, decreased need for insulin, and lower blood peak glucose levels (Vuori, 2001; Wallenberg-Henriksson *et al.*, 1998).

For the evaluation of dietary quality, the dietary variety score is generally accepted as one of the convenient measures (Choi *et al.*, 2000; Drewnoski *et al.*, 1998; Kant, 1996), since consumption of different food items is expected to improve the

overall quality of diet and the nutritional status. Our study showed that maintaining physical activity above moderate levels contributes to consumption of more diverse food items. It seems that the diabetic patients with proper physical activity levels would pay more attention to health and pursue health-promoting behaviors including dietary practices.

Our findings of higher urinary zinc excretion in diabetic patients are consistent with the previous studies (Cunningham *et al.*, 1994; Lee *et al.*, 2005; Pidduck *et al.*, 1970). A recent study also showed higher urinary levels of trace elements in diabetic patients compared to their age-matched healthy controls ($p < 0.001$). They reported that mean values of zinc levels in blood was reduced in diabetic patients compared with control subjects. However, these differences were not significant (Kazi *et al.*, 2008).

Although it has been speculated that the continued zinc loss through urinary excretion may result in zinc deficiency in diabetes, the correlation between urinary zinc and serum levels of zinc was not well demonstrated in previous studies (Levine *et al.*, 1983; Mateo *et al.*, 1978; Mooradian & Morley, 1987). In this study, higher serum zinc levels were shown in diabetic patients with heavy physical activities. It is beyond the scope of this study to draw any definite conclusions on the relationship between physical activity and zinc status. However, we believe that physical activity may potentially contribute to improving zinc status by improving the overall dietary status.

In conclusion, our study indicated that poorer dietary quality and zinc status were observed in diabetic patients with less physical activity. Therefore, more attention is needed to improve the zinc status among Korean diabetic patients, whose activity patterns tend to be sedentary. Appropriate nutrition education is urgently required for diabetic patients as well as the general adult population to attain dietary adequacy and to increase the amount of their physical activities.

Literature cited

- American Diabetes Association (2002). Evidence-based nutrition principles and recommendations for the treatment and prevention of diabetes and related complications. *J Am Diet Assoc* 102:109-118.
- Chausmer AB (1998). Zinc, Insulin and Diabetes. *J Am Coll Nutr* 17:109-115.
- Choi YJ, Kim SY, Jung KA & Chang YK (2000). An assessment of diet quality in the postmenopausal women. *The Korean Journal of Nutrition* 33:304-313.
- Cunningham JJ, Fu A, Mearkle PL & Brown RG (1994). Hyperzincuria in individuals with insulin-dependent diabetes mellitus: concurrent zinc status and the effect of high-dose zinc supplementation. *Metabolism* 43:1558-1562.
- De Mello VD, Zelmanovitz T, Perassolo MS, Azevedo MJ & Gross JL (2006). Withdrawal of red meat from the usual diet reduces albuminuria and improves serum fatty acid profile in type 2 diabetes patients with no albuminuria. *Am J Clin Nutr* 83:1032-1038.

- Drewnoski A, Henderson SA, Shore AB, Fishler C, Preziosi P & Hercberg S (1996). Dietary quality and dietary diversity in France implications for the French paradox. *J Am Diet Assoc* 96:663-669.
- Eckel RH (2006). A new look at dietary protein in diabetes. *Am J Clin Nutr* 78:671-672.
- Hawk PB, Oser BL & Summerson WH (1954). *Practical Physiology Chemistry*. 13th ed. 899 Blakston Co., Toronto, Canada
- Kant AK (1996). Indexes of overall dietary quality: a review. *J Am Diet Assoc* 96:785-791.
- Kazi TG, Afridi HI, Kazi N, Jamali MK, Arain MB, Jalbani N & Kandhro GA(2008). Copper, chromium, manganese, iron, nickel, and zinc levels in biological samples of diabetes mellitus patients. *Biol Trace Elem Res* 122:1-18.
- Kim YI, Choi CS, Kim SW, Lee JS, Kim HH, Lee MS, Lee SI, Park JY, Hong SK & Lee KU (1998). Prevalence of Diabetes Mellitus and impaired glucose tolerance in Korean adults living in Jungup District, South Korea. *Journal of Korean Diabetes Association* 22:363-371.
- Kinlaw WB, Levine AS, Morley JE, Silivis SE & McClain CJ (1983). Abnormal zinc metabolism in type II diabetes mellitus. *Am J Med* 75:273-277.
- Korean Food Research Institute (1998). Handbook of portion size for food intake research. Gyeonggi. Republic of Korea
- Korean Nutrition Society (2005). *Dietary Reference intakes for Koreans*, Hanareum Printing, Seoul. Republic of Korea
- Kwon IS & Kwon CS (2000). Dietary molar ratios of phytate: zinc and millimolar ratios of phytate x calcium: zinc in South Koreans. *Biol Trace Elem Res* 75:29-41.
- Lee JH, Lee HJ, Lee IK & Yoon JS (2005). Zinc and copper status of middle- and old-aged woman in type 2 diabetes. *The Korean Journal of Nutrition* 38:56-66.
- Levine AS, McClain CJ, Handwerger BS, Brown DM, Morley JE.(1983). Tissue zinc status of genetically diabetic and streptozotocin-induced diabetic mice. *Am J Clin Nutr* 37:382-386.
- LÖnnerdal B (2000). Dietary factors influencing zinc absorption *J Nutr* 130:1378-1383.
- Ministry of Health & Welfare (2006). Report on 2005 National health and nutritional survey. Seoul. Republic of Korea
- Ministry of Health & Welfare (2007). Symposium on In-Depth Analysis on 2005 National health and nutritional survey. Seoul. Republic of Korea
- Mooradian AD & Morley JE (1987). Micronutrient status in diabetes mellitus. *Am J Clin Nutr* 45:877-895.
- Pidduck HG, Wren PJJ & Price Evans DA (1970). The hyperzincuria of diabetes mellitus and possible genetic implications of this observation. *Diabetes* 19:240-247.
- Roth H & Kirchgessner M (1981). Zinc and insulin metabolism. *Biol Trace Elem Res* 3:13-32.
- Salgueiro MJ, Krebs N, Zubillaga MB, Weill R, Postaire E, Lysionek AE, Caro RA, Paoli TD, Hager A & Boccio J (2001). Zinc and Diabetes Mellitus. Is there a need of zinc supplementation in diabetes mellitus patients? *Biol Trace Elem Res* 81:215-228.
- Vuori IM (2001). Health benefits of physical activity with special reference to interaction with diet. *Public Health Nutr* 4:517-528.
- Wallenberg-Henriksson H, Rincon J & Zierath J (1998). Exercise in the management of non-insulin-dependent diabetes mellitus. *Sports Med* 25:25-35.
- Yoon JS & Lee JH (2007). A suggestion to improve zinc status of type 2 diabetic women: relation among zinc, protein and phytate intake. *Journal of Korean Dietetic Association* 13:301-310.