

Research article

Open Access

Analysis of the *Pythium ultimum* transcriptome using Sanger and Pyrosequencing approaches

Foo Cheung¹, Joe Win², Jillian M Lang³, John Hamilton⁴, Hue Vuong¹, Jan E Leach³, Sophien Kamoun², C André Lévesque⁵, Ned Tisserat³ and C Robin Buell*⁴

Address: ¹The J. Craig Venter Institute, 9704 Medical Center Dr, Rockville, MD 20850 USA, ²The Sainsbury Laboratory, Colney Lane, Norwich, NR4 7UH, UK, ³Colorado State University, Department of Bioagricultural Sciences and Pest Management, C129 Plant Sciences, Ft. Collins CO 80523, USA, ⁴Michigan State University, Department of Plant Biology, East Lansing MI 48824, USA and ⁵Agriculture and Agri-Food Canada, Ottawa, ON, K1A 0C6, Canada

Email: Foo Cheung - fcheung@jcvl.org; Joe Win - joe.win@bbsrc.ac.uk; Jillian M Lang - Jillian.Lang@ColoState.EDU; John Hamilton - hamilton@plantbiology.msu.edu; Hue Vuong - hvuong@jcvl.org; Jan E Leach - jan.leach@colostate.edu; Sophien Kamoun - sophien.kamoun@tsl.ac.uk; C André Lévesque - levesqueca@agr.gc.ca; Ned Tisserat - ned.tisserat@colostate.edu; C Robin Buell* - buell@msu.edu

* Corresponding author

Published: 15 November 2008

Received: 8 May 2008

BMC Genomics 2008, 9:542 doi:10.1186/1471-2164-9-542

Accepted: 15 November 2008

This article is available from: <http://www.biomedcentral.com/1471-2164/9/542>

© 2008 Cheung et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Pythium* species are an agriculturally important genus of plant pathogens, yet are not understood well at the molecular, genetic, or genomic level. They are closely related to other oomycete plant pathogens such as *Phytophthora* species and are ubiquitous in their geographic distribution and host range. To gain a better understanding of its gene complement, we generated Expressed Sequence Tags (ESTs) from the transcriptome of *Pythium ultimum* DAOM BR144 (= ATCC 200006 = CBS 805.95) using two high throughput sequencing methods, Sanger-based chain termination sequencing and pyrosequencing-based sequencing-by-synthesis.

Results: A single half-plate pyrosequencing (454 FLX) run on adapter-ligated cDNA from a normalized cDNA population generated 90,664 reads with an average read length of 190 nucleotides following cleaning and removal of sequences shorter than 100 base pairs. After clustering and assembly, a total of 35,507 unique sequences were generated. In parallel, 9,578 reads were generated from a library constructed from the same normalized cDNA population using dideoxy chain termination Sanger sequencing, which upon clustering and assembly generated 4,689 unique sequences. A hybrid assembly of both Sanger- and pyrosequencing-derived ESTs resulted in 34,495 unique sequences with 1,110 sequences (3.2%) that were solely derived from Sanger sequencing alone. A high degree of similarity was seen between *P. ultimum* sequences and other sequenced plant pathogenic oomycetes with 91% of the hybrid assembly derived sequences > 500 bp having similarity to sequences from plant pathogenic *Phytophthora* species. An analysis of Gene Ontology assignments revealed a similar representation of molecular function ontologies in the hybrid assembly in comparison to the predicted proteomes of three *Phytophthora* species, suggesting a broad representation of the *P. ultimum* transcriptome was present in the normalized cDNA population. *P. ultimum* sequences with similarity to oomycete RXLR and Crinkler effectors, Kazal-like and cystatin-like protease inhibitors, and elicitors were identified. Sequences with

similarity to thiamine biosynthesis enzymes that are lacking in the genome sequences of three *Phytophthora* species and one downy mildew were identified and could serve as useful phylogenetic markers. Furthermore, we identified 179 candidate simple sequence repeats that can be used for genotyping strains of *P. ultimum*.

Conclusion: Through these two technologies, we were able to generate a robust set (~10 Mb) of transcribed sequences for *P. ultimum*. We were able to identify known sequences present in oomycetes as well as identify novel sequences. An ample number of candidate polymorphic markers were identified in the dataset providing resources for phylogenetic and diagnostic marker development for this species. On a technical level, in spite of the depth possible with 454 FLX platform, the Sanger and pyro-based sequencing methodologies were complementary as each method generated sequences unique to each platform.

Background

Pythium species are ubiquitous fungal-like organisms in the Kingdom Straminipila. They are related to the oomycete plant pathogens *Phytophthora* and downy mildews (e.g. *Bremia*, *Peronospora*, and *Plasmopora*) in the Peronosporales and to *Saprolegnia* and *Aphanomyces* in the Saprolegniales. They differ from the true fungi (Eumycota) in that they are diploid, have coenocytic hyphae containing β -1–3 glucans and cellulose in their cell walls, reproduce sexually by fertilization of oogonia by antheridia and, in many species, form motile, biflagellate zoospores. Nevertheless, *Pythium* species have many similarities to true fungi in the way they obtain nutrients from the environment or attack plants [1].

The genus *Pythium* is biologically and ecologically diverse. Although approximately 250 *Pythium* species have been described, only half that number has been recognized as valid descriptions [2-4]. *Pythium* species have been divided into 11 phylogenetic clades based on rDNA ITS and other sequence data [3]. Most are soil inhabitants although some reside in salt water estuaries and other aquatic environments. Most *Pythium* species are saprophytic or facultative plant pathogens [4]. They are among the most important plant pathogens and cause a variety of diseases including seed rots and damping-off, root, stem and fruit rots, foliar blights and postharvest decay [5-8]. A few non-phytopathogenic species show promise as biological control agents [9]. Other *Pythium* species are parasites of insects [10] and fish [4] and at least one species (*P. insidiosum*) can infect animals and causes skin and bone lesions in pigs, dogs, and humans [11].

Pythium ultimum is a cosmopolitan plant pathogen with a broad host range. It is one of the most pathogenic *Pythium* species on corn, soybean, wheat, ornamentals, and many other crops [9]. It is homothallic, and in most cases, self fertile, although some outcrossing may occur [12]. *P. ultimum* is actually a species complex with three main morphological types. *P. ultimum* var. *ultimum*, the most common type, produces oospores but very rarely spor-

angia and zoospores, and then only at cool temperatures [4]. It is part of a large uniform clade based on rDNA ITS sequence comparisons which includes the neotype strain [3] as well as the isolate used to generate the Expressed Sequence Tags (ESTs) in this study. *P. ultimum* var. *sporangiferum* produces oospores and sporangia at room temperature [4]. In Barr et al. [13], the four *P. ultimum* var. *sporangiferum* isolates formed a unique genotype (U6) based on isozyme analysis whereas in Francis et al. [12] the two isolates of this variety were in two distinct genotypes, one belonging to the most common genotype of *P. ultimum* var. *ultimum* strains. It is unclear at this point whether or not the more infrequent but unique genotypes of *P. ultimum* such as the genotype U6 should be split into different species, reducing the observed genetic diversity within this species. Having more information on Single Nucleotide Polymorphisms would help to delineate gene flow and species boundaries in this ubiquitous species complex. Isolates in the third morphological type do not produce oospores in culture but can be identified based on other morphological characteristics, molecular studies, and the fact that they can be crossed with oospore producing strains of *P. ultimum* [3,12-14].

ESTs are a robust method for gene discovery and for identifying transcripts involved in specific biological processes. ESTs have been developed for over 40 oomycete and fungal pathogens [15]. For the oomycetes, EST projects have provided not only the first insight into the gene complement for these plant pathogens but also sets of genes involved in a number of biological processes [16-24]. With the advent of reduced costs and higher throughput sequencing methods, ESTs can be economically generated for a wider range of organisms and a greater breadth of conditions, thereby providing a more comprehensive assessment of an organism's transcriptome. One emerging platform for *de novo* sequencing of transcriptomes or genomes is pyrosequencing [25]. However, the pyrosequencing-derived read lengths on the GS20 and FLX platform are substantially shorter than conventional Sanger-based sequencing which is problematic for assembly into

a non-redundant set of consensus sequences. This limitation can be addressed through increased sequencing coverage of the transcriptome and/or through hybrid approaches in which conventional Sanger sequences are co-assembled [26] with pyrosequencing-derived sequences. Thus, for *de novo* sequencing of a transcriptome, both approaches are advantageous with pyrosequencing providing a deep representation of the transcriptome and Sanger-derived sequences providing seed sequences for assembly purposes. In this study, we report the first set of ESTs for this agriculturally important plant pathogen and provide a direct comparison of gene discovery that can be obtained with two high throughput sequencing methods, Sanger-based chain termination and 454 FLX pyrosequencing.

Results and Discussion

Sanger-based sequencing of a normalized cDNA library

A normalized cDNA library was generated from RNA isolated from hyphae growing on two contrasting media conditions, nutrient-rich and nutrient-starved conditions. A total of 9,578 reads from Sanger-based sequencing was generated; these assembled into 4,689 unique sequences (assemblies plus singletons) with an average length of 596 nucleotides and a total length of 2.8 Mb (Table 1, Figure 1). The majority of assemblies contain two component reads while very few assemblies contain six or more component reads (Figure 2). Overall, clustering and assembly of the reads resulted in a two-fold reduction in the number of unique sequences with few EST components per assembly consistent with an effective normalization treatment of the cDNA prior to library construction. We




Figure 1
Size distribution of assemblies and reads. Length distribution (in base pairs) of reads and assemblies from the Sanger, 454, and hybrid assemblies.




Figure 2
Component size distribution. The number of reads in each assembly is plotted for each of the three assemblies (Sanger reads, 454 reads, and hybrid assembly).

confirmed the origin of the ESTs by aligning the assemblies to a draft *P. ultimum* BR144 genome sequence [27]. For the Sanger-only assembly, 4,127 (88%) sequences (1,408 assemblies, 2,719 singletons) aligned to the draft genome. Of the 4,689 unique sequences, 2,345 (50.0%) had sequence similarity with an entry in the UniRef100 database [28]. Within the set of 4,689 unique sequences (assemblies and singletons), 4,444 open reading frames (ORFs) could be predicted. Of these, 3,676 ORFs contained an ATG translational start codon while 2,413 had both an ATG start codon and a stop codon. Of these 2,413 ORFs, 1,421 (58.6%) have sequence similarity with a protein sequence in the UniRef100 database that contains the corresponding start codon and are thus potentially full-length. The length of top UniRef100 matches ranged between 50 to >500 amino acids in length, indicating that the sequences and therefore the cDNAs generated from this library were not biased towards either long or short sequences.

454 Pyrosequencing sequencing and assembly

A single half-plate pyrosequencing run on the same normalized cDNA used in the Sanger-based sequencing generated 139,839 reads, which after trimming and cleaning, yielded 90,664 usable reads (>100 base pairs) with an average length of 190 nucleotides and a total length of 17.3 Mb (Table 1). After clustering and assembly using the TGICL clustering utilities [29], 66,312 sequence reads were incorporated into 11,155 assemblies and 24,352 singletons for a total of 35,507 unique sequences. Assemblies had an average length of 233 bp with a maximum length of 1,385 bp (Table 1, Figure 1). For the 454 assembly,

Table 1: Sequence and assembly statistics

	Pyrosequencing ESTs	Sanger ESTs	Pyrosequencing + Sanger ESTs	Pyrosequencing assembly	Sanger assembly	Hybrid Assembly
No. sequences	90,664	9,578	90,664 + 9,578 = 100,242	35,507 (11,155 assemblies)	4,689 (1,584 assemblies)	34,495 (10,811 assemblies)
Total bases (bp)	17,315,041	4,811,851	22,126,892	8,274,073	2,796,219	9,552,701
Minimum length (bp)	100	100	100	100	100	100
Average length (bp)	190	502	220	233	596	276
Maximum length (bp)	425	1,002	1,002	1,385	4,283	4,471
%GC	50.3	51.1	50.5	51.3	53.3	51.7
%N	0.04	0	0.03	0.05	0	0.04

25,285 (71%) sequences (9,226 assemblies, 16,059 singletons) aligned to the draft *P. ultimum* BR144 genome sequence. Of these 35,507 unique sequences, 7,991 (22.5%) had a protein match with UniRef100 [28], significantly less than what was observed for the Sanger-derived ESTs. If only sequences > 500 bp were considered, 71% had a match with an entry in UniRef100 suggesting that the short length of the 454 reads impact the ability to detect similar sequences. As the pyrosequencing library preparation involves random shearing of the normalized (but uncloned) cDNA population [25], the pyrosequencing reads originate from random locations within each cDNA and may have either orientation. The majority of assemblies contain 2–5 reads, while in contrast only a few of the assemblies contained over 100 pyrosequencing EST reads (Figure 1). This is presumably due to both the length of the individual reads, the normalization of the cDNA population, and the coverage of the transcriptome represented in this dataset.

Hybrid assembly of Sanger and Pyrosequencing-based reads

A hybrid assembly was constructed using the pyrosequencing- and Sanger-based EST reads which were clustered and assembled using the TGICL clustering utilities [29]; of these, 10,811 were assemblies and 23,684 were singletons for a total of 34,495 unique sequences (Table 1). Since the reads used in the hybrid assembly were predominantly derived from the pyrosequencing dataset, with the exception of the maximum assembly length, the assembly length and EST component statistics are similar to that for the pyrosequencing-only reads (Figure 1). In the hybrid assembly, a majority of assemblies are short with an average length of 276 base pairs (maximum length of 4,471 base pairs (Table 1)) and contain few component reads (Figure 2). The majority of assemblies contain 2–5 EST reads while only a limited number of assemblies contain over 100 EST reads. For the hybrid assembly, 24,140 (70%) sequences (8,922 assemblies, 15,218 singletons) aligned to the draft *P. ultimum* BR144 genome sequence. Within the hybrid assembly, 8,618 had a protein match using UniRef100 [28] significantly less

than what was observed for the Sanger-derived ESTs yet consistent with the abundance of 454 reads in the overall hybrid assembly. If only sequences > 500 bp within the hybrid assembly were considered, 69% had a match with an entry in UniRef100. Pyrosequencing using the 454 platform is reported to have reduced accuracy in homopolymer regions [25]. Examination of homopolymer regions (>4 nucleotides) detected between the Sanger- and the 454-derived reads revealed differences in homopolymer length between the Sanger and 454-reads in 837 instances, primarily in A or T homopolymer regions (407 A, 323 T, 65 G, 42 C).

A total of 1,110 unique sequences in the hybrid assembly were derived from Sanger sequencing alone. Although the majority of these had no similarity with entries in the UniRef100 database (Additional Data Files 1 and 2), 907 (81.7%) of them mapped to the *Pythium* genome at very high stringencies (95% identity, 95% coverage with a minimum of 100 bp of the Sanger read; Cheung and Buell, unpubl.), suggesting that these are not contaminants or artifacts within the cDNA library. Of the 1,110 unique sequences derived solely from Sanger reads, 576 had no match even at a very low threshold (BLASTN E-10 [30] or BLAT [31], default settings) with the pyrosequencing-derived sequences. Further analysis of these 576 Sanger-only reads showed that 461 (80%) could be mapped to the *Pythium* genome with high stringency ($\geq 95\%$ identity, $\geq 95\%$ coverage, of a minimum of 100 bp of the Sanger read). An analysis of average read length revealed that these Sanger sequences were slightly shorter on average (99 bp) than the complete set of Sanger sequences which would reduce to the probability of co-assembling with 454 reads during the hybrid assembly process (average length of 497 nucleotides in Sanger-only singletons and assemblies vs 596 nucleotides in the complete set of Sanger singletons and assemblies). Although the GC content did not differ substantially between the Sanger-only sequence set and the complete Sanger sequence set (51% vs 53%), other sources of possible bias could be the cloning and replication in *E. coli* which may amplify select sequences in the library compared to the

454 sequences. These results suggest that a small proportion of ESTs generated by Sanger sequencing are not present in the dataset generated by pyrosequencing.

Biological features of the *P. ultimum* transcriptome

The top 50 assemblies containing the most ESTs from all three builds (pyrosequencing reads only, Sanger reads only, hybrid assembly) ranged from 10 to 2,654 ESTs (Additional Data Files 3, 4, 5). Most of the protein matches to the top 50 assemblies from all three builds were to *Phytophthora* species. In general, the most common annotations based on similarity to UniRef100 entries were housekeeping genes although 17 of the top 50 deepest assemblies from the hybrid assembly had no match to an entry in UniRef100 (Additional Data File 3).

To determine how similar the *P. ultimum* transcriptome is to other plant pathogenic oomycetes, we compared our EST sequences with the predicted proteomes and transcriptomes available for four *Phytophthora* species: *P. infestans* (transcriptome and predicted proteome), *P. parasitica* (transcriptome), *P. ramorum* (predicted proteome), and *P. sojae* (transcriptome and predicted proteome) [32,33]. Using BLASTX with the predicted proteome data sets, 16,139 (46.8%) of the *P. ultimum* hybrid assembly sequences matched a sequence present within the *P. infestans*, *P. ramorum*, or *P. sojae* predicted proteomes. Using TBLASTX with a cutoff criterion of $E = 1e-10$, 14,021 (40.6%) of the *P. ultimum* hybrid sequences matched a sequence within the three *Phytophthora* transcriptomes with only 945 (2.7%) of the *P. ultimum* sequences that align with a predicted *Phytophthora* protein not matching a *Phytophthora* transcript. Using BLASTX with a cutoff criterion of $E = 1e-5$ against the UniRef100 database, 8,618 (25.0%) of the *P. ultimum* hybrid sequences matched the UniRef database, of which 673 (2.0%) did not also match a *Phytophthora* transcript or predicted protein. Collectively, from all three datasets (*Phytophthora* spp. predicted proteomes, *Phytophthora* spp. transcriptomes, and the UniRef database), 16,738 (48.5%) of the *P. ultimum* hybrid sequences do not have a match. This is most likely attributed to the short nature of the majority of the hybrid sequences. Indeed, if only sequences ≥ 500 bp within the hybrid assembly are considered (3,515 total sequences), the percentage of sequences with an alignment to the three data types significantly increases; 3,201 (91.1%) align to the *Phytophthora* spp. predicted proteomes, 2,959 (84.2%) align to the *Phytophthora* spp. transcriptomes, and 2,427 (69%) align to the UniRef database, with only 252 (7.2%) without a match to any sequence within these data sets.

We also performed a comparative analysis of Gene Ontologies [34] with *P. ultimum* and the three *Phytophthora* proteomes. As shown in Figure 3, the normalized cDNA

obtained from hyphae grown in rich and nutrient-starved conditions encoded a broad set of transcripts represented within the molecular function gene ontologies. Furthermore, the representation is very similar to that of the complete proteomes of the three plant pathogenic *Phytophthora* species suggesting that our approach of cDNA normalization, coupled with deep sequencing, provided a near complete representation of the *P. ultimum* transcriptome.

It recently emerged that *Phytophthora* and downy mildew species secrete a vast repertoire of effector proteins that modulate host defenses and enable pathogenicity [35,36]. The extent to which *Pythium* species also rely on secreted effectors to colonize host tissue is unclear. We therefore examined the *P. ultimum* ESTs for similarity to known oomycete effectors. We first scanned the *P. ultimum* EST hybrid assembly for candidate cytoplasmic effectors of the RXLR and Crinkler (CRN) families [36,37] using a combination of the PexFinder algorithm [23] to identify putative secreted protein genes and sequence similarity searches. An hmm profile based on an alignment of known RXLR-EER effectors [38] revealed one *P. ultimum* assembly (asmb1_7845) as a putative positive. *P. ultimum* asmb1_7845 encodes an ORF with a signal peptide (SignalP HMM score = 0.921) followed by the RLLRSAGD-VESSAVDDAAR sequence with similarity to the RXLR-DEER motif (Additional Data File 6A-B). The identification of only a single putative RXLR effector is surprising and contrasts to the common occurrence of RXLR effectors in similar sets of *Phytophthora* ESTs [21]. Apparently, RXLR effectors are not as widely present or expressed in *P. ultimum* as noted for *Phytophthora* species. There are several possible explanations. It is possible that *P. ultimum* does not have RXLR effector genes or has a highly reduced set compared to *Phytophthora*. This would be consistent with observations [39] that suggested that RXLR effectors are delivered through haustoria, specialized infection structures that are not produced by *Pythium*. The RXLR motif is similar in sequence, position and function to the *Plasmodium* Pexel/Host translocation motif [40]. The possible absence of RXLR effectors in *P. ultimum* indicates that although the motif is conserved across divergent parasitic eukaryotes it may not be ubiquitous in oomycetes. Four Crinkler-like sequences were identified among the *P. ultimum* hybrid assembly. In this case, the similarity to *Phytophthora* Crinklers was more convincing than for the single RXLR effector candidate with BLASTX E values as low as $e-48$. Clearly, these sequences displayed the consensus LXLFLAK instead of the LXLFLAXR motif that defines canonical *Phytophthora* and downy mildew Crinklers [37,38] (Additional Data File 6C). In summary, we detected one potential candidate RXLR and several Crinkler effectors in *P. ultimum*, however, they are not as abundantly represented among the examined *P. ultimum* ESTs




Figure 3
Distribution of GOSlim assignments in the *P. ultimum* hybrid EST sequences compared to the predicted proteomes of three *Phytophthora* species. The percentage of assignments of the total GOSlim assignments are shown. Only GOSlim terms present at level of 1% or greater in any of the four species are shown.

as they are in *Phytophthora* ESTs from similar developmental stages [21].

We also searched the *P. ultimum* assemblies for similarities to other oomycete effectors. We detected three assemblies with similarity to oomycete Kazal-like serine protease inhibitors [41,42] and another three with similarity to cystatin-like protease inhibitors [43] (Additional Data File 7). In *Phytophthora*, these apoplastic effectors are known to inhibit defense related proteases of plants [41-43]. In addition, at least 13 assemblies with similarity to elicitors were identified (Additional Data File 7). Elicitors are secreted lipid-binding oomycete proteins that trigger defense responses in plants [36]. These elicitors showed significant similarity to previously described *Phytophthora* and *Pythium* elicitors with their characteristic cysteine-rich domain [44,45]. Six assemblies were most similar to sylvaticin, a secreted elicitor of *Pythium sylvaticum* of

unknown function [45]. The same assemblies showed significant similarity with the elicitor-like protein of *Pythium oligandrum* [46] but the homology was much lower than that of *P. sylvaticum* (E value e^{-11}). This is consistent with the taxonomy and phylogeny of these species. Both *P. ultimum* (clade I) and *P. sylvaticum* (clade F) belong to the globose sporangia group of *Pythium* whereas *P. oligandrum* (clade D) belongs to a different group with contiguous sporangia [47].

Unlike *Phytophthora* spp., *Pythium* and other oomycetes like *Saprolegnia* spp. are not thiamine auxotrophs. Torto et al. [48] reported sequences with similarity to thiamine biosynthesis enzymes among ESTs of the fish pathogen *Saprolegnia parasitica* that are missing in the genome sequences of *P. sojae*, *P. ramorum*, *P. infestans*, and *Hyaloperonospora parasitica*. We identified four *P. ultimum* sequences within the hybrid assembly (asembl_312;

asmb1_353; asmb1_1697, and 334590_1440_2761) with similarity to the *S. parasitica* thiamine biosynthesis enzyme. This finding is consistent with the knowledge that *P. ultimum* can synthesize thiamine. The thiamine biosynthesis gene was apparently lost during the evolution of the *Phytophthora*/downy mildews lineage and could serve as a potential phylogenetic marker among Saprolegniales and the genus *Pythium*.

Oomycetes are often characterized by the ultrastructure of their flagellar apparatus. Although the strain we sequenced and *P. ultimum* var. *ultimum* are generally not known to produce zoospores *in vitro*, it might be possible that some flagellar associated proteins are expressed as was observed in *Phytophthora* grown under conditions that do not produce zoospores [21]. Some of the flagellar associated proteins from *Chlamydomonas reinhardtii* (177 in total, Additional Data File 8) that we identified in our ESTs are predicted to be commonly expressed in other structures/tissues than zoospores (e.g. calmodulin or elongation factor). As shown by Randall *et al*[21], we also found evidence of expression of dynein related to the flagellar apparatus (E values between e-6 and e-38) but we also identified several flagellar basal body proteins (E values between e-8 and e-21) that were expressed.

Random candidate markers for population genetic studies can also be derived from simple sequence repeats (SSRs). Within the hybrid assembly, a total of 179 SSRs were identified within 164 sequences. Among the SSRs, monucleotides (45) were the most abundant followed by dinucleotides (44), trinucleotides (33), pentanucleotides (27), tetranucleotides (26), and hexanucleotides (4) (Additional Data File 9). Lee and Moorman [49] developed SSR markers for *Pythium aphanidermatum*, *P. cryptosporangium* and *P. irregulare* from an SSR enriched library. Primers P18CCA1-41, P18TTC1-42, P18CAT1-74 amplified SSRs in *P. ultimum* but were not found in this EST library. It is possible that the genes in which these SSRs were located were not expressed, that these SSRs were in non-coding regions, or that the primers designed from other species worked on *P. ultimum* even if they had mismatches.

Conclusion

In this study, we report the integration of data from Sanger-based chain termination and 454 FLX pyrosequencing technologies. The two technologies were highly complementary although the shorter read length in the pyrosequencing-derived ESTs is problematic in that they limit biological interpretations due to the reduced information content in the predicted protein sequence. Thus, while the pyrosequencing did provide depth of coverage, we were able to generate a more robust set of transcribed

sequences for *Pythium* by co-assembly of Sanger and pyrosequencing derived ESTs. Furthermore, even with a greater depth of ESTs provided through the 454 FLX platform, the two sequencing methodologies generated sequences unique to each platform. The results presented contain an ample number of candidate polymorphic markers providing resources for potential phylogenetic and diagnostic and strain marker development for this agriculturally important group of plant pathogens. We expect that a similar analysis using other species and integration of data from 454 FLX pyrosequencing technologies would work synergistically with existing or new EST data and identify new genes/transcripts at a very cost effective and efficient manner.

Methods

Materials and Organismal Methods

Yeast extract broth (30 g/L sucrose, 1 g/L KH_2PO_4 , 0.5 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.5 g/L KCl, 10 mg/L $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 1 g/L yeast extract) was inoculated with several crushed plugs of 3 day old *P. ultimum* strain DAOM BR144 (= ATCC 200006 = CBS 805.95) grown on 10% V8 agar and incubated at room temperature on a shaker (Lab-Line Instruments, Melrose Park, IL) at 200 rpm for 2 days. Plugs of the same isolate were covered with modified Plich medium (Kamoun, S., unpub.) in a Petri dish and incubated in the dark at room temperature for 10 days. Mycelia from both media were rinsed with 50 ml of sterile water and harvested by straining through sterile cheesecloth, weighed and then ground in liquid nitrogen using a mortar and pestle.

Molecular and Bioinformatics Methods

RNA was extracted from this tissue using TRIzol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. Equal masses of total RNA from the two growth conditions were used for construction of a normalized cDNA population and subsequently, a normalized cDNA library. Using the services of Evrogen [50], full length cDNA was constructed using the Smart cDNA cloning methodology [51] and normalized using a duplex-specific nuclease [50]. Two methods were used to generate sequences from the normalized cDNA: Sanger-based dideoxy chain termination and Roche 454-FLX pyrosequencing [25]. Sanger-based sequences were generated from end sequencing of cDNA clones using standard high throughput sequencing methods on an ABI 3730xl sequencer. Sequences were trimmed to remove vector, low-quality, and *E. coli* sequences using Lucy [52] and iterative runs of the TIGR Seqclean Tool [53]. Accessions are available in GenBank dbEST ([EL774305-EL783216](#) and [ES607153-ES608087](#)). Pyrosequencing was performed on a Roche 454 FLX sequencer [25] using a half-plate. Pyrosequencing reads were trimmed to remove adaptors

using TIGR Seqclean Tool [14]. Short sequences (< 100 nucleotides) were removed from the build process as the addition of extremely short sequences will lead to false joining of ESTs, chimeras that were sequenced, and reduced quality of unique sequences. Sequences were deposited in GenBank dbEST under accession numbers [FE969956–FF060620](#).

Pyrosequencing and Sanger cDNA reads were assembled using the TGICL package [29] as was the hybrid assembly using the default parameters as described in Childs et al. [54] with the exception that sequences < 100 nucleotides were excluded. Sequences were searched against the UniRef100 database (Release 12.0) using BLASTX with a cutoff criterion of $E = 1e-5$. The EST assemblies were aligned with a draft assembly of the *P. ultimum* BR144 genome [27] using BLAT [31]. *Pythium* EST assemblies were downloaded from The Comprehensive Phytopathogen Resource Transcript Assemblies website [15] and *Phytophthora* genome sequences were downloaded from GenBank. Transcriptomes of *P. ultimum* version 1.0, *P. infestans* version 3.0, *P. sojae* version 1.0, and *P. parasitica* version 2.0 were aligned using TBLASTX with cutoff criteria of $E = 1e-5$. *P. ultimum* ESTs were aligned with the *P. ultimum* [55], *P. infestans* [56], *P. ramorum* [33] and *P. sojae* [33] genomes with TBLASTX requiring a minimum of 40% length matching. For the Gene ontology associations, the three *Phytophthora* proteomes were searched against the current UniRef100 database using BLASTP and a cutoff of $1e-5$. GO ids were assigned using the EBI UniProt Knowledgebase GOA file [57] using the top UniRef100 hit. The assigned GO identifiers were then mapped to the Generic GOSlim ontology using the map2slim tool [58]. Using BLASTX and a cutoff of $1e-5$, the hybrid EST set was searched against 892 *Chlamydomonas reinhardtii* proteins flagged in GenBank as being associated with the flagellar apparatus to identify transcripts associated with the flagellar mechanism in *Pythium*. Perfect mononucleotide to hexanucleotide simple sequence repeats were identified using MISA [59]. Perl scripts were used to specify the minimum number of the following repeats for microsatellites (unit size/minimum number of repeats): (1/20) (2/10) (3/7) (4/5) (5/5) (6/5).

Authors' contributions

FC was responsible for analysis of the EST data. JML was responsible for isolating mRNA and data analysis. JW and SK were responsible for analysis of effectors. NT, JEL, and CAL were responsible for strain selection, experimental design, data analysis, and discussion. JH was responsible for data analysis. CRB was responsible for experimental design, sequencing, data analysis, and discussion.

Additional material

Additional file 1

Hybrid assembly singleton reads derived solely from Sanger reads.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S1.xls>]

Additional file 2

Hybrid assemblies derived solely from Sanger reads.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S2.xls>]

Additional file 3

Top 50 largest assemblies from the hybrid assembly.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S3.xls>]

Additional file 4

Top 50 largest assemblies from 454-pyrosequencing only assembly.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S4.xls>]

Additional file 5

Top 50 largest assemblies from Sanger only sequence assembly.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S5.xls>]

Additional file 6

Alignments of P. ultimum sequences to effectors and Crinkler proteins.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S6.pdf>]

Additional file 7

Assemblies with similarity to protease inhibitors and elicitors.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S7.xls>]

Additional file 8

Chlamydomonas flagellar proteins identified in the hybrid EST assembly.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S8.xls>]

Additional file 9

Simple Sequence Repeats identified in the hybrid assembly.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-542-S9.xls>]

Acknowledgements

Funding for the work is provided by grants from the USDA National Research Initiative Cooperative State Research Extension Education Service (2006-55605-16645 (C.R.B., J.E.L., N.T.) and 2007-35600-18886 (C.R.B., N. T.)). S.K. and J.W. are funded by the Gatsby Charitable Foundation. C.A.L. is supported with funding to the Canadian Barcode of Life Network from Genome Canada through the Ontario Genomics Institute and NSERC.

References

- Latijnhouwers M, de Wit PJGM, Govers F: **Oomycetes and fungi: similar weaponry to attack plants.** *Trends in Microbiology* 2003, **11(10)**:462-469.
- Index Fungorum** [<http://www.indexfungorum.org>]
- Lévesque CA, De Cock WAM: **Molecular phylogeny and taxonomy of the genus *Pythium*.** *Mycol Res* 2004, **108**:1363-1383.
- Plaats-Niterink AJ van der: **Monograph of the genus *Pythium*.** *Studies in Mycology* 1981, **21**:1-242.
- Cook RJ, Sitton JW, Waldher JT: **Evidence for *Pythium* as a pathogen of direct-drilled wheat in the Pacific Northwest.** *Plant Dis* 1980, **64**:102-103.
- Larkin RP, English JT, Mihail JD: **Effects of infection by *Pythium* spp. on the root system morphology of alfalfa seedlings.** *Phytopathology* 1995, **85**:430-435.
- Sumner DR, Gascho GJ, Johnson AW, Hook JE, Threadgill ED: **Root diseases, populations of soil fungi, and yield decline in continuous double-crop corn.** *Plant Disease* 1990, **74(9)**:704-710.
- Snowdon AL: **A Color Atlas of Post-Harvest Diseases & Disorders of Fruits and Vegetables.** Volume 1. CRC Press; 1990:302.
- Martin FN, Loper JE: **Soilborne plant diseases caused by *Pythium* spp.: ecology, epidemiology, and prospects for biological control.** *Critical Reviews in Plant Sciences* 1999, **18(2)**:111-181.
- Saunders GA, Washburn JO, Egerter DE, Anderson JR: **Pathogenicity of fungi isolated from field-collected larvae of the western treehole mosquito, *Aedes sierrensis* (Diptera: Culicidae).** *Journal of Invertebrate Pathology* 1988, **52(2)**:360-363.
- Mendoza L, Ajello L, McGinnis MR: **Infections caused by the oomycetous pathogen *Pythium insidiosum*.** *Journal de Mycologie Medicale* 1996, **6(4)**:151-164.
- Francis DM, Gehlen MF, St Clair DA: **Genetic variation in homothallic and hyphal swelling isolates of *Pythium ultimum* var. *ultimum* and *P. ultimum* var. *sporangiferum*.** *Molecular Plant-Microbe Interactions* 1994, **7(6)**:766-775.
- Barr DJS, Warwick SI, Desaulniers NL: **Isozyme variation, morphology, and growth response to temperature in *Pythium ultimum*.** *Canadian Journal of Botany* 1996, **74(5)**:753-761.
- Huang HC, Morrison RJ, Muendel H-H, Barr DJS, Klassen GR, Bochko J: ***Pythium* sp. "group G", a form of *Pythium ultimum* causing damping-off of safflower.** *Canadian Journal of Phytopathology* 1992, **14**:229-232.
- The Comprehensive Phytopathogen Genome Resource Transcript Assemblies** [http://cpgr.plantbiology.msu.edu/cpgr_ta.shtml]
- Gaulin E, Madoui MA, Bottin A, Jacquet C, Mathe C, Couloux A, Wincker P, Dumas B: **Transcriptome of *Aphanomyces euteiches*: New oomycete putative pathogenicity factors and metabolic pathways.** *PLoS ONE* 2008, **3(3)**:e1723.
- Kamoun S, Hraber P, Sobral B, Nuss D, Govers F: **Initial assessment of gene diversity for the oomycete pathogen *Phytophthora infestans* based on expressed sequences.** *Fungal Genet Biol* 1999, **28(2)**:94-106.
- Le Berre JY, Engler G, Panabieres F: **Exploration of the late stages of the tomato-*Phytophthora parasitica* interactions through histological analysis and generation of expressed sequence tags.** *New Phytol* 2008, **177(2)**:480-492.
- Panabieres F, Amselem J, Galiana E, Le Berre JY: **Gene identification in the oomycete pathogen *Phytophthora parasitica* during in vitro vegetative growth through expressed sequence tags.** *Fungal Genet Biol* 2005, **42(7)**:611-623.
- Qutob D, Hraber PT, Sobral BW, Gijzen M: **Comparative analysis of expressed sequences in *Phytophthora sojae*.** *Plant Physiol* 2000, **123(1)**:243-254.
- Randall TA, Dwyer RA, Huitema E, Beyer K, Cvitanich C, Kelkar H, Fong AM, Gates K, Roberts S, Yatzkan E, Gaffney T, Law M, Testa A, Torto-Alalibo T, Zhang M, Zheng L, Mueller E, Windass J, Binder A, Birch PR, Gisi U, Govers F, Gow NA, Mauch F, van West P, Waugh ME, Yu J, Boller T, Kamoun S, Lam ST, et al.: **Large-scale gene discovery in the oomycete *Phytophthora infestans* reveals likely components of phytopathogenicity shared with true fungi.** *Mol Plant Microbe Interact* 2005, **18(3)**:229-243.
- Skalamera D, Wasson AP, Hardham AR: **Genes expressed in zoospores of *Phytophthora nicotianae*.** *Mol Genet Genomics* 2004, **270(6)**:549-557.
- Torto TA, Li S, Styer A, Huitema E, Testa A, Gow NA, van West P, Kamoun S: **EST mining and functional expression assays identify extracellular effector proteins from the plant pathogen *Phytophthora*.** *Genome Res* 2003, **13(7)**:1675-1685.
- Torto-Alalibo TA, Tripathy S, Smith BM, Arredondo FD, Zhou L, Li H, Chibucos MC, Qutob D, Gijzen M, Mao C, Sobral BW, Waugh ME, Mitchell TK, Dean RA, Tyler BM: **Expressed sequence tags from *Phytophthora sojae* reveal genes specific to development and infection.** *Mol Plant Microbe Interact* 2007, **20(7)**:781-793.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer ML, Jarvie TP, Jirage KB, Kim JB, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, et al.: **Genome sequencing in microfabricated high-density picolitre reactors.** *Nature* 2005, **437(7057)**:376-380.
- Goldberg SM, Johnson J, Busam D, Feldblyum T, Ferreira S, Friedman R, Halpern A, Khouri H, Kravitz SA, Lauro FM, Li K, Rogers YH, Strausberg R, Sutton G, Tallon L, Thomas T, Venter E, Frazier M, Venter JC: **A Sanger/pyrosequencing hybrid approach for the generation of high-quality draft assemblies of marine microbial genomes.** *Proc Natl Acad Sci USA* 2006, **103(30)**:11240-11245.
- Pythium ultimum* Genome Database** [<http://pythium.plantbiology.msu.edu>]
- Suzek BE, Huang H, McGarvey P, Mazumder R, Wu CH: **UniRef: comprehensive and non-redundant UniProt reference clusters.** *Bioinformatics* 2007, **23(10)**:1282-1288.
- Pertea G, Huang X, Liang F, Antonescu V, Sultana R, Karamycheva S, Lee Y, White J, Cheung F, Parvizli B, Tsai J, Quackenbush J: **TIGR Gene Indices clustering tools (TGICL): a software system for fast clustering of large EST datasets.** *Bioinformatics* 2003, **19(5)**:651-652.
- Altschul SF, Gish W, Miller W, Myers EV, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215(3)**:403-410.
- Kent WJ: **BLAT – the BLAST-like alignment tool.** *Genome Res* 2002, **12(4)**:656-664.
- The *Phytophthora infestans* Genome Database** [http://www.broad.mit.edu/annotation/genome/phytophthora_infestans/Home.html]
- Tyler BM, Tripathy S, Zhang X, Dehal P, Jiang RH, Aerts A, Arredondo FD, Baxter L, Bensasson D, Beynon JL, Chapman J, Damasceno CM, Dorrance AE, Dou D, Dickerman AW, Dubchak IL, Garbelotto M, Gijzen M, Gordon SG, Govers F, Grunwald NJ, Huang W, Ivors KL, Jones RW, Kamoun S, Krampis K, Lamour KH, Lee MK, McDonald WH, Medina M, et al.: ***Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis.** *Science* 2006, **313(5791)**:1261-1266.
- Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G: **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium.** *Nat Genet* 2000, **25(1)**:25-29.
- Birch PR, Rehmany AP, Pritchard L, Kamoun S, Beynon JL: **Trafficking arms: oomycete effectors enter host plant cells.** *Trends Microbiol* 2006, **14(1)**:8-11.
- Kamoun S: **A catalogue of the effector secretome of plant pathogenic oomycetes.** *Annu Rev Phytopathol* 2006, **44**:41-60.
- Kamoun S: **Groovy times: filamentous pathogen effectors revealed.** *Curr Opin Plant Biol* 2007, **10(4)**:358-365.
- Win J, Morgan W, Bos J, Krasileva KV, Cano LM, Chaparro-Garcia A, Ammar R, Staskawicz BJ, Kamoun S: **Adaptive evolution has targeted the C-terminal domain of the RXLR effectors of plant pathogenic oomycetes.** *Plant Cell* 2007, **19(8)**:2349-2369.

39. Whisson SC, Boevink PC, Moleleki L, Avrova AO, Morales JG, Gilroy EM, Armstrong MR, Grouffaud S, van West P, Chapman S, Hein I, Toth IK, Pritchard L, Birch PR: **A translocation signal for delivery of oomycete effector proteins into host plant cells.** *Nature* 2007, **450(7166)**:115-118.
40. Bhattacharjee S, Hiller NL, Liolios K, Win J, Kanneganti TD, Young C, Kamoun S, Haldar K: **The malarial host-targeting signal is conserved in the Irish potato famine pathogen.** *PLoS Pathog* 2006, **2(5)**:e50.
41. Tian M, Benedetti B, Kamoun S: **A second Kazal-like protease inhibitor from *Phytophthora infestans* inhibits and interacts with the apoplastic pathogenesis-related protease P69B of tomato.** *Plant Physiol* 2005, **138(3)**:1785-1793.
42. Tian M, Huitema E, Da Cunha L, Torto-Alalibo T, Kamoun S: **A Kazal-like extracellular serine protease inhibitor from *Phytophthora infestans* targets the tomato pathogenesis-related protease P69B.** *J Biol Chem* 2004, **279(25)**:26370-26377.
43. Tian M, Win J, Song J, Hoorn R van der, Knaap E van der, Kamoun S: **A *Phytophthora infestans* cystatin-like protein targets a novel tomato papain-like apoplastic protease.** *Plant Physiol* 2007, **143(1)**:364-377.
44. Jiang RH, Tyler BM, Whisson SC, Hardham AR, Govers F: **Ancient origin of elicitor gene clusters in *Phytophthora* genomes.** *Mol Biol Evol* 2006, **23(2)**:338-351.
45. Lascombe MB, Retailleau P, Ponchet M, Industri B, Blein JP, Prange T: **Structure of sylvaticin, a new alpha-elicitor-like protein from *Pythium sylvaticum*.** *Acta Crystallogr D Biol Crystallogr* 2007, **63(Pt 10)**:1102-1108.
46. Takenaka S, Nakamura Y, Kono T, Sekiguchi H, Masunaka A, Takahashi H: **Novel elicitor-like proteins isolated from the cell wall of the biocontrol agent *Pythium oligandrum* induce defence-related genes in sugar beet.** *Molecular Plant Pathology* 2006, **7**:325-339.
47. Lévesque CA, de Cock AWAM: **Molecular phylogeny and taxonomy of the genus *Pythium*.** *Mycological Research* 2004, **108**:1363-1383.
48. Torto-Alalibo T, Tian M, Gajendran K, Waugh ME, van West P, Kamoun S: **Expressed sequence tags from the oomycete fish pathogen *Saprolegnia parasitica* reveal putative virulence factors.** *BMC Microbiol* 2005, **5**:46.
49. Lee S, Moorman GW: **Identification and characterization of simple sequence repeat markers for *Pythium aphanidermatum*, *P. cryptoirregulare*, and *P. irregulare* and the potential use in *Pythium* population genetics.** *Curr Genet* 2008, **53(2)**:81-93.
50. **Evrogen** [<http://www.evrogen.com>]
51. Zhu YY, Machleder EM, Chenchik A, Li R, Siebert PD: **Reverse transcriptase template switching: a SMART approach for full-length cDNA library construction.** *Biotechniques* 2001, **30(4)**:892-897.
52. Chou HH, Holmes MH: **DNA sequence quality trimming and vector removal.** *Bioinformatics* 2001, **17(12)**:1093-1104.
53. **TIGR Seqclean Tool** [<http://www.tigr.org/tdb/tgi/software>]
54. Childs KL, Hamilton JP, Zhu W, Ly E, Cheung F, Wu H, Rabinowicz PD, Town CD, Buell CR, Chan AP: **The TIGR Plant Transcript Assemblies database.** *Nucleic Acids Res* 2007:D846-851.
55. ***Pythium ultimum* Genome Database** [<http://pythium.plantbiology.msu.edu>]
56. **The *Phytophthora infestans* Genome Database** [http://www.broad.mit.edu/annotation/genome/phytophthora_infestans/Home.html]
57. **EBI Uniprot Knowledgebase** [ftp://ftp.ebi.ac.uk/pub/databases/GO/goa/UNIPROT/gene_association.goa_uniprot.gz]
58. **Map2slim Tool** [<http://www.geneontology.org/GO.slims.shtml>]
59. Thiel T, Michalek W, Varshney RK, Graner A: **Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.).** *Theor Appl Genet* 2003, **106(3)**:411-422.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

