

Modulation of Host Immunity by Helminths: The Expanding Repertoire of Parasite Effector Molecules

Rick M. Maizels,^{1,*} Hermelijn H. Smits,^{2,*} and Henry J. McSorley^{3,*}

¹Wellcome Centre for Molecular Parasitology, Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK

²Leiden University Medical Center, Leiden, the Netherlands

³Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

*Correspondence: rick.maizels@glasgow.ac.uk (R.M.M.), h.h.smits@lumc.nl (H.H.S.), henry.mcsorley@ed.ac.uk (H.J.M.)

<https://doi.org/10.1016/j.immuni.2018.10.016>

Helminths are extraordinarily successful parasites due to their ability to modulate the host immune response. They have evolved a spectrum of immunomodulatory molecules that are now beginning to be defined, heralding a molecular revolution in parasite immunology. These discoveries have the potential both to transform our understanding of parasite adaptation to the host and to develop possible therapies for immune-mediated disease. In this review we will summarize the current state of the art in parasite immunomodulation and discuss perspectives on future areas for research and discovery.

Introduction

Helminths are extremely successful parasites, affecting around a quarter of the world's population (Bethony et al., 2006). They establish chronic infections and if untreated can persist for the lifetime of the host. While the immune system is capable of parasite expulsion (often incurring collateral damage), in natural infections there is frequently a muted immune response in which the host accommodates, and tolerates, the invader. Such re-setting of host immunity results from both host reparative responses to physical damage from tissue-migrating parasites and/or through active immunomodulation by their molecular products. The ability of parasites to defy host immunity reflects their masterful manipulation of the immune system, which as detailed below, is achieved through the release of a spectrum of finely tuned and highly evolved immuno-modulatory factors (Table 1).

Recent advances in helminth genomics and proteomics have uncovered a wealth of such immunomodulatory products. These act on every phase of the immune response, which for the purposes of this review we have divided into: (1) initiation, (2) antigen recognition and processing, (3) adaptive responses, (4) effector cell responses, and (5) coagulation, healing, and remodeling. Parasite immunomodulators active at each phase can be found that either share ancestral homology with host genes or have developed *de novo*, sharing no identifiable homology to the host (Figure 1). Here, we summarize the molecular actors known to interact at each step and their evolutionary and structural provenance, and we discuss future prospects for this exciting field.

Parasites Neutralize Initiating Alarmin Signals

Immune responses are initiated by danger signals, through detection of damage-associated molecular patterns (DAMPs) or introduction of pathogen-associated molecular patterns (PAMPs). Parasite migration induces DAMPs and introduces PAMPs from either the parasite itself or bacteria entering tissues through compromised barriers. The imperative to mute barrier

surface reactions has driven the evolution of potent mechanisms to suppress innate immune responses (Figure 2).

Activation of epithelial cells results in the release of “alarmin” cytokines such as thymic stromal lymphopoietin (TSLP), interleukin-25 (IL-25), and IL-33. IL-33 is tightly spatiotemporally controlled: it is released on epithelial necrosis and then rapidly oxidized and inactivated (Cohen et al., 2015). The murine intestinal nematode *Heligmosomoides polygyrus* suppresses IL-33 release through HpARI (*H. polygyrus* alarmin release inhibitor) (McSorley et al., 2014; Osbourn et al., 2017) within its excretory-secretory (ES) products. HpARI has a selective mode of action, binding DNA via its first domain, while the second and third domains bind to reduced (active) but not oxidized (inactive) IL-33. Binding obstructs interaction of the complex with the IL-33 receptor, ST2, while interaction with DNA tethers IL-33 within the nucleus of necrotic cells, preventing IL-33 release. HpARI administration ablates type 2 cell-mediated inflammation and improves lung function in an *Alternaria* allergen-dependent asthma model, while in *Nippostrongylus brasiliensis* infection, HpARI administration suppresses type 2 responses and increased worm burden. Crucially, these effects translate to the human setting, as HpARI prevents IL-33 release from human lung explants and blocks human IL-33 release in a transgenic mouse model (Osbourn et al., 2017).

H. polygyrus suppresses the IL-33 pathway at multiple levels additionally to HpARI, with an undefined *H. polygyrus* product further downregulating IL-33 production through induction of IL-1 β (Zaiss et al., 2013) and through the release of small RNA-containing extracellular vesicles that suppress transcription of the IL-33 receptor (Buck et al., 2014; Coakley et al., 2017).

IL-33 release can be provoked by signals of cell stress or death, in particular extracellular ATP, which induces epithelial and mast cell IL-33 production (Cekic and Linden, 2016). Many parasite secretions contain apyrases (Da'dara et al., 2014; Gounaris et al., 2004), which degrade ATP to non-inflammatory AMP, reducing inflammatory DAMP signals, and hence these could inhibit this important arm of the damage detection response.



Table 1. Different Classes of Helminth-Derived Immunomodulatory Molecules

	Abbreviation	Helminth Species	Action	References
Cytokine & Innate Defense Homologs and Growth Factors				
Alarmin release inhibitor	HpARI	<i>Heligmosomoides polygyrus</i>	Blocks human and mouse IL-33	Osbourn et al., 2017
Asparaginyl-tRNA synthetase	AsnRS	<i>Brugia malayi</i>	Structural homology to IL-8, binds IL-8 receptors CXCR1 and CXCR2; chemotactic for neutrophils and eosinophils; induced regulatory responses and IL-10 in a T cell transfer model of colitis	Kron et al., 2013
Chemokine binding protein	SmCKBP	<i>Schistosoma mansoni</i>	Neutralizes chemokine activity (CXCL8, CCL3, CX3CL1, CCL2, CCL5); inhibits neutrophil migration but not eosinophil migration	Smith et al., 2005
Granulin-like growth factor-1	OvGRN-1	<i>Opisthorchis viverrini</i>	Induces angiogenesis and wound healing (mice); proliferation in human cholangiocytes and pathways associated with wound healing and cancer	Smout et al., 2009, 2015
Helminth defense molecule-1	FhHDM-1	<i>Fasciola hepatica</i>	Molecular mimicry of antimicrobial peptide CAP18/LL-37, binds to LPS and reduces its activity; prevents acidification of the endolysosomal compartments and antigen processing; prevents NLRP3 inflammasome activation	Robinson et al., 2011, 2012
IL-4 inducing principle from <i>S. mansoni</i> eggs	IPSE	<i>S. mansoni</i> ; <i>S. haematobium</i>	Induces IL-4 secretion in basophils via binding to IgE; induces IL-10 in B cells and enhances their capacity to induce Treg cells; translocates to nucleus and reduces bladder hemorrhage <i>in vivo</i>	Haeberlein et al., 2017 ; Kaur et al., 2011 ; Mbanefo et al., 2018
Macrophage Migration Inhibitory Factor homolog-1	MIF-1	<i>B. malayi</i> , <i>Trichinella spiralis</i> , <i>Anisakis simplex</i>	Induces IL-8 release from monocytes; synergizes with IL-4 to induce alternatively activated macrophages; inhibits experimental arthritis, colitis & allergic airway inflammation via induction of IL-10 & Treg cells	Cho et al., 2015 ; Park et al., 2009 ; Tan et al., 2001 ; Zang et al., 2002
Metalloproteinases	–	<i>Necator americanus</i>	Causes proteolysis of eotaxin, but not of IL-8 or eotaxin-2	Culley et al., 2000
Neutrophil inhibitory factor	AcNIF	<i>Ancylostoma caninum</i>	Binds $\beta 2$ integrin CD11b/CD18, inhibit LPS-induced neutrophil migration and release of reactive oxygen species (ROS)	Anbu and Joshi, 2008 ; Moyle et al., 1994 ; Zhou et al., 1998
55 kDa glycoprotein	Hcgp55	<i>Haemonchus contortus</i>		
T cell immunomodulatory protein	EmTIP	<i>Echinococcus multilocularis</i>	Induces release of IFN- γ from CD4 ⁺ T cells <i>in vitro</i>	Nono et al., 2014
TGF- β homolog-2	TGH-2	<i>B. malayi</i>	Ligates mammalian TGF- β receptor and suppresses T cell responses	Gomez-Escobar et al., 2000
TGF- β mimic	TGM	<i>H. polygyrus</i>	Ligation of TGF- β receptor on T cells leading to induction of Treg cells	Grainger et al., 2010 ; Johnston et al., 2017
TGF-like molecule	FhTLM	<i>F. hepatica</i>	Ligates mammalian TGF- β receptor (albeit with a lower affinity) and induces IL-10 and Arginase in macrophages	Sulaiman et al., 2016

(Continued on next page)

Table 1. Continued

	Abbreviation	Helminth Species	Action	References
TLR Signaling				
ES-62	ES-62	<i>Acanthocheilonema viteae</i>	Modulates dendritic cell responses by inducing the selective autophagolysosomal degradation of TLR-transducers (e.g., TLR4) and interaction with MyD88; synthetic variants block inflammation in various disease models, i.e., allergy, rheumatoid arthritis, and colitis	Ball et al., 2018 ; Goodridge et al., 2007 ; Pineda et al., 2014 ; Rzepecka et al., 2014
Fatty acid binding protein	Fh12/15	<i>F. hepatica</i>	Suppresses LPS-induced activation via binding and blocking of CD14; induction of alternatively activated macrophages	Martin et al., 2015 ; Ramos-Benítez et al., 2017
Lysophosphatidylserine	Lyso-PS	<i>S. mansoni</i>	Ligation of TLR2 on dendritic cells; licenses DCs to develop IL-10-producing Treg cells	van der Kleij et al., 2002
Mucin-like polypeptide	Fhmuc	<i>F. hepatica</i>	Promotes TLR4 activation of DCs and Th1 cell induction	Noya et al., 2016, 2017
Intracellular Signaling and Gene Expression				
Abundant Larval Transcript	ALT	<i>B. malayi</i>	Upregulates SOCS1, the inhibitor of IFN- γ signaling	Gomez-Escobar et al., 2005
Acetylcholinesterase	AChE	<i>N. brasiliensis</i>	Degrades acetylcholine, reduces neural signaling; induces proinflammatory cytokines with diminished type 2 cytokines in transgenic AChE-expressing trypanosome infection	Vaux et al., 2016
<i>Ancylostoma</i> secreted protein-2	NaASP-2	<i>N. americanus</i>	Binding to CD79A on B cells, downregulation of lyn, PI3K, and BCR signaling	Tribolet et al., 2015
ATP diphosphohydrolase	SmATPDase1	<i>S. mansoni</i>	Degradation of the exogenous pro-inflammatory and pro-thrombotic nucleotides ATP and ADP: inhibition of blood coagulation	Da'dara et al., 2014
Calpain-1 & -2	SmCalp-1, SmCalp-2	<i>S. mansoni</i>	SmCalp1 and -2 cleave the blood clotting protein fibronectin and high-molecular-weight kininogen (HK)	Wang et al., 2017
Cathepsin L peptidases	FhCL-1,-2,-3	<i>F. hepatica</i>	Degrades fibrinogen and fibrin	Mebius et al., 2018
Ectonucleotide pyrophosphatase/ phosphodiesterase	SmNPP-5	<i>S. mansoni</i>	Ectonucleotide pyrophosphatase/phosphodiesterase homolog, which inhibits platelet aggregation by degradation of ADP	Elzoheiry et al., 2018
α -enolase	OvENO	<i>Onchocerca volvulus</i>	Binds to plasminogen and supports plasmin-mediated proteolysis including degradation of host's extracellular matrix	Jolodar et al., 2003
Inhibitor of potassium channel	AcK1	<i>A. caninum</i>	Inhibitor of the voltage-gated channel Kv1.3 in lymphocytes, similar to the polypeptide inhibitor of the sea anemone, Skh; suppressed delayed type hypersensitivity and the proliferation of memory T cells	Chhabra et al., 2014
	BmK1	<i>B. malayi</i>		

(Continued on next page)

Table 1. Continued

	Abbreviation	Helminth Species	Action	References
Omega-1	ω -1	<i>S. mansoni</i>	T2 ribonuclease binds to MR and DC-SIGN, after uptake degrades ribosomal and mRNA; primes DCs for enhances Th2 development; enhances IL-1 β secretion in macrophages; improves insulin sensitivity	Everts et al., 2009, 2012; Ferguson et al., 2015; Hams et al., 2016; Ke et al., 2017; Steinfeldt et al., 2009; Wilbers et al., 2017
	CP1412	<i>S. japonicum</i>		
Serine protease-2	SmSP-2	<i>S. mansoni</i>	Manipulation host vasodilatation and fibrinolysis: activates tissue plasminogen activator and plasminogen	Leontovyč et al., 2018
16 kDa polypeptide	Sj16	<i>S. japonicum</i>	Translocates to the nucleus, induces IL-10 in BM-derived DCs	Sun et al., 2016
Enzymes and Inhibitors				
Anti-inflammatory protein-1	AIP-1	<i>N. americanus</i>	Family member of tissue inhibitor of metalloprotease (TIMP)-like proteins; suppression in TNBS colitis model: promotes expression of colon IL-10, TGF- β , and TSLP and the accumulation of Treg cells in the colon	Ferreira et al., 2017
Anti-inflammatory protein-2	AIP-2	<i>N. americanus</i>	Family member of TIMP-like proteins; suppression in model of allergic airway inflammation via Treg cell induction and suppression of T cell proliferation in cells from house dust mite (HDM)-allergic patients	Navarro et al., 2016
Cystatins	AvCystatin	<i>A. viteae</i>	Induction of regulatory IL-10-producing macrophages and hyporesponsive T cells; compromised APC function; reduced airway allergy and intestinal colitic inflammation in mouse models	Daniłowicz-Luebert et al., 2013; Ziegler et al., 2015 Schönemeyer et al., 2001 Pfaff et al., 2002 Sun et al., 2013 Manoury et al., 2001 Dainichi et al., 2001 Wang et al., 2016 Coronado et al., 2017 Jang et al., 2011
	Onchocystatin	<i>O. volvulus</i>		
	LsCystatin	<i>Litomosoides sigmodontis</i>		
	HpCPI	<i>H. polygyrus</i>		
	BmCPI-2	<i>B. malayi</i>		
	Nippocystatin	<i>N. brasiliensis</i>		
	rSjCystatin	<i>S. japonicum</i>		
	rAi-CPI	<i>Acaris lumbricoides</i>		
	rCsStefin-1	<i>Clonorchis sinensis</i>		
Kunitz-type serine protease inhibitor	SjKT-1	<i>S. japonicum</i>	Inhibition of trypsin and chymotrypsin, neutrophil elastase, FXa, and plasma kallikrein: both anti-coagulant and anti-inflammatory properties. FhKTM reduces inflammatory cytokine production in DC	Falcón et al., 2014; Ranasinghe et al., 2015
	FhKTM	<i>F. hepatica</i>		
Peroxiredoxin	FhPrx	<i>F. hepatica</i>	Anti-oxidant enzyme: inactivation of ROS and induction of AA-MF in mouse models	Donnelly et al., 2008
Thioredoxin peroxidase	TPx	<i>F. hepatica</i>	Anti-oxidant enzyme, induction of AA-MF with increased IL-10 and PGE2 responses	Donnelly et al., 2005
Tissue inhibitor of metalloprotease	Ace-MTP-2	<i>A. ceylanicum</i>	Reduces MHC-I and MHCII molecules on dendritic cells and those DC induce CD4 and CD8 Treg cells	Baska et al., 2013

(Continued on next page)

Table 1. Continued

	Abbreviation	Helminth Species	Action	References
Lipid or Lipid-Binding				
Protein of <i>Ascaris suum</i> -1	PAS-1	<i>Ascaris suum</i>	Similar to ABA-1, a nematode tandemly repeated polyprotein with lipid binding properties; inhibitory in an experimental airway allergy model, dependent on IL-10-producing Treg cells and IFN- γ -producing CD8 γ δ TCR T cells	Araújo et al., 2008 ; de Araújo et al., 2010
Prostaglandin-2	PGE2	<i>Trichuris suis</i>	<i>T. suis</i> secretes large amounts of PGE2 in a COX-independent pathway; secreted PGE2 is effective in modulating DC responses	Kaisar et al., 2018 ; Laan et al., 2017 ; Liu et al., 1992
		<i>S. mansoni</i>	Ligation to Dectin1 and Dectin 2 on DC by SEA leads to autocrine production of PGE2 in DC, which lead to increased Th2 polarization	
Extracellular Vesicles				
Extracellular vesicles	EV	<i>H. polygyrus</i>	EVs contain microRNAs that target epithelial cells and macrophages leading to loss of IL-33 receptor (IL-33R, also termed ST2) expression, and downregulate other immune genes; inhibit eosinophilia in <i>Alternaria</i> allergic airway inflammation model	Buck et al., 2014 ; Coakley et al., 2017
Extracellular vesicles	EV	<i>S. japonicum</i>	Induce M1 macrophage differentiation <i>in vitro</i>	Wang et al., 2015




Figure 1. Helminth Modulators Act at All Phases of the Immune Response

Modulatory proteins can be homologous to host genes (shared evolutionary ancestry, green) or show no homology (developed *de novo*, red). The immune response is divided into five phases: (1) initiation, innate recognition of damage and danger; (2) innate cell activation and antigen presentation; (3) adaptive immune responses; (4) effector immune responses; and (5) resolution and healing.

further inhibitory functions (Martin et al., 2015). Furthermore, cysteine proteases from the flukes *F. hepatica* (FhCL1) and *S. mansoni* (SmCB1) both directly suppress myeloid cell TLR signaling by interfering with TLR-driven inflammation *in vivo*, resulting in the intracellular degradation of TLR3 and TLR4 (Donnelly et al., 2010).

Conversely, some helminth molecules stimulate, rather than inhibit, TLRs.

Helminths Target Dendritic Cell and Macrophage Functions

A pivotal point in host immunity is recognition of and reaction to pathogen molecules, typically by pathogen- or damage-associated molecules patterns (PAMPs and DAMPs) ligating pattern recognition receptors (PRRs) on myeloid cells, such as toll-like receptors (TLRs) and C-type lectin receptors (CLRs). These reactions are intensively targeted by helminth molecules (Figure 2), which block TLR ligand-induced dendritic cell and macrophage activation, interfering with receptors and their signaling, as well as antigen presentation and downstream effector mechanisms.

The MyD88 adaptor protein is required for signaling via all TLRs except TLR3, and also IL-1 family cytokine receptors (including the IL-33 receptor). ES-62, a multifunctional glycoprotein secreted by the rodent filarial nematode *Acanthocheilonema viteae* (Pineda et al., 2014), protects against pathology in mouse models of rheumatoid arthritis (RA) (Doonan et al., 2018), asthma (Rzepecka et al., 2014), and lung fibrosis (Suckling et al., 2018). ES-62 induces sequestration of the MyD88 signaling protein, leading to suppression of TLR and IL-33 signaling (Ball et al., 2018; Pineda et al., 2014). The immunomodulatory principle of ES-62 is phosphorylcholine (PC) side groups carried on N-linked glycan moieties (Goodridge et al., 2007), and synthetic small molecule variants of PC can reproduce many anti-inflammatory effects of the parent molecule (Al-Riyami et al., 2013).

A different mechanism is deployed by the immunomodulatory ES protein Fh12 (and its recombinant form, Fh15) from the liver fluke *Fasciola hepatica*. The Fh12 and Fh15 proteins induce alternative activation in human monocyte-derived macrophages (Mo-Macs), suppress macrophage activation by TLR2, TLR4, TLR5, and TLR8 ligands, and inhibit inflammatory cytokine production in septic shock (Ramos-Benítez et al., 2017). They bind to the TLR4 cofactor CD14 and reduce its expression, mediating suppression of TLR4 responses. The Fh12 and Fh15 proteins also suppress responses of CD14-deficient cells, suggesting

(lyso-PS) is enriched in the tegument of *S. mansoni* adult worms (Retra et al., 2015). LysoPS is a TLR2 ligand that increases the capacity of human DCs to drive IL-10-producing T cells, suppressing T cell proliferation (van der Kleij et al., 2002). In contrast, the 66-amino acid mucin-like polypeptide from *F. hepatica* (Fhmuc), while not itself a TLR4 ligand, promotes LPS-induced TLR4 activation of DCs, enhancing T helper 1 (Th1) cell responses, both *in vitro* and *in vivo* (Noya et al., 2017).

CLR signaling is important in responses to helminths, and parasite-specific glycans are a rich source of ligands (Hokke and van Diepen, 2017). The schistosome-specific GalNAc-di-GalNAc (LDN) motif is bound by soluble galectin-3 (van den Berg et al., 2004), while fucosylated (LDN-F) derivatives and fucose-containing Lewis X (Le^x) structures are bound by dendritic cell-specific ICAM3-grabbing non-integrin (DC-SIGN). Mannose-containing structures are recognized by mannose receptor (MR) and macrophage galactose-type lectin (MGL) (van Die et al., 2003). Similar glycan ligands for these receptors are also present in many other parasites including *F. hepatica*. During *F. hepatica* infection, glycan-CLR signaling induces a mixed type 2 and regulatory phenotype with IL-10 and transforming growth factor- β (TGF β) production (Rodríguez et al., 2017).

Schistosome glycans are strongly immunomodulatory (Meevissen et al., 2012) with a predominant role in Th2 cell polarization through DCs: *S. mansoni* soluble egg antigen (SEA) induces Th2 cell responses *in vivo* in a glycan-dependent manner (Okano et al., 1999). The LNFPIII (lactose-N-fucopentose III) motif, present in both on *S. mansoni* and in human milk, has similar DC-modulating effects as total SEA (Thomas et al., 2005) and can act as an adjuvant for a type 2 immune response (Okano et al., 2001). LNFPIII also stimulates murine B cells to produce the immunoregulatory cytokine IL-10 (Velupillai and Harn, 1994). SEA and human milk also contain LNNt (lacto-N-neotetraose), while intraperitoneally injected milk-derived LNNt induces the influx of a macrophage population that produces IL-10 and

turn prevents the acidification of the endolysosomal compartment, impairing antigen processing (Robinson et al., 2012) and preventing NLRP3 inflammasome activation and IL-1 β secretion (Alvarado et al., 2017). In addition to its immune-modulatory effects, FhHDM-1 may have parasite-intrinsic functions, as it also binds and detoxifies heme (Martínez-Sernández et al., 2017).

More broadly, and beyond signaling pathways and antigen processing, myeloid cell protein expression can be profoundly affected by helminth molecules. For example, ω -1 (Omega-1), a glycoprotein released by *S. mansoni* eggs, is a T2 ribonuclease (RNase). It is taken up by DCs through recognition of its glycan moiety, Lewis-X, by the mannose receptor, and extinguishes protein synthesis through the degradation of ribosomal RNA and mRNA within the cell. Omega-1 does not degrade specific transcripts, but rather targets the global pool of RNA. In this way omega-1, and its close homolog from *Schistosoma japonicum*, CP1412, suppress DC activation and maturation, preventing upregulation of CD86 and MHCII and synthesis of IL-12 in response to CD40 ligation, favoring both Th2 cell (Everts et al., 2012; Everts et al., 2009; Steinfeldt et al., 2009; Wilbers et al., 2017) and regulatory T (Treg) cell (Ke et al., 2017; Zacccone et al., 2011) induction. T cell priming in the absence of IL-12 and with diminished co-stimulation may favor Th2 cell polarization.

The Th2 cell-mediated priming capacity of omega-1 has also been confirmed *in vivo* using lentivirus-based transduction to silence gene expression in schistosome eggs, as intravenous injections of those eggs shows reduced lung granuloma formation compared to sham-transduced eggs (Hagen et al., 2014). Lastly, omega-1 induces inflammasome-dependent IL-1 β in TLR2-stimulated peritoneal macrophages, which is abrogated when blocking Dectin-1, suggesting that omega-1 can modulate multiple PRR pathways (Ferguson et al., 2015).

The modulatory effects of omega-1 were recently extended to changes in host metabolism. A series of recent studies have demonstrated the beneficial effects of helminth infection on host metabolism (Crowe et al., 2017; Wiria et al., 2014), which can be replicated by helminth products such as SEA (Hussaarts et al., 2015; Wolfs et al., 2014) or recombinant omega-1 (Hams et al., 2016). Injection of omega-1 into obese mice improves insulin sensitivity via IL-33 release from white adipose tissue, which in turn induces ILC2 activation and alternatively activated M2 macrophage differentiation. Mechanistically, the release of IL-33 is mediated by the RNase activity of omega-1, while the improved insulin sensitivity was dependent on the ligation of the mannose receptor (Hams et al., 2016).

Another example showing myeloid cell signaling manipulation is from *B. malayi* abundant larval transcript (ALT). The mosquito-borne *B. malayi* larvae stockpile quantities of ALT proteins prior to entry into the mammalian host. Investigation of gene function by a transfection approach within macrophages identified ALT-induced upregulation of SOCS1, the inhibitor of IFN- γ signaling in inflammation (Gomez-Escobar et al., 2005).

Finally, some helminth proteins contain a nuclear localization signal (NLS) for rapid translocation to the nucleus, including the *S. japonicum*-derived molecule Sj16, which induces IL-10 and inhibits DC maturation following nuclear translocation (Sun et al., 2016). IL-4-inducing principle from schistosome eggs (IPSE) is also rapidly translocated to the nucleus of DCs upon

uptake (Kaur et al., 2011), while the IPSE homolog from *S. haematobium* (H-IPSE) translocates to the nucleus of bladder epithelial cells (Pennington et al., 2017). Here, H-IPSE reduces bladder hemorrhage and accelerates urothelial repair in a chemotherapy-induced hemorrhagic cystitis (CHC) model (Mbanefo et al., 2018). The therapeutic effects are IL-4 dependent and may relate to its IL-4-inducing capacity described in basophils (detailed below). The effects of H-IPSE are, however, superior to IL-4 alone, implying that additional nuclear functions are involved (Mbanefo et al., 2018).

The end result of parasite modulation of dendritic cell and macrophage modulation is often aberrant, skewed, or tolerized T cell responses. In the next section we will deal with helminth products that act directly on adaptive immune responses.

Adaptive Response Modulation Is Targeted at Multiple Levels

On activation by APCs, naive T cells can differentiate into Th1, Th2, Th17, or Treg cell subsets. Modulation of adaptive immune responses has the potential for lasting effects in the host: if the parasite can appear as harmless or commensal, a memory response can be raised that treats the parasite as “self.” Parasitic helminths have developed multiple suppressive and modulatory pathways that suppress effector and/or induce regulatory T helper and B cell responses (Figure 3).

An example of both homologous and *de novo*-generated immunomodulators from helminth parasites can be found in parasite agonists of the TGF- β receptor. The effects of host immune TGF- β signaling are diverse, but are largely immunosuppressive, with generation of induced regulatory T cells, suppression of dendritic cell, macrophage, and T cell activation, and skewing of antibody isotypes to IgA (Chen and Ten Dijke, 2016). Thus, stimulation of the TGF- β pathway would be advantageous to an immunity-shy parasite.

The ancestral function of the TGF- β pathway, across the animal kingdom, is in development: the immunomodulatory role of TGF- β is a more recent adaptive branch within vertebrates (David and Massague, 2018). Notably, ligands and receptors show a high degree of conservation allowing, for example, *S. mansoni* and *Echinococcus multilocularis* TGF- β receptors to ligate human TGF- β family ligands (Beall and Pearce, 2001; Zavala-Góngora et al., 2006). The question is whether TGF- β family members from helminths activate immunosuppressive mechanisms in mammalian cells; indeed, *B. malayi* TGH-2 can signal through the TGF- β R (Gomez-Escobar et al., 2000) while extracts from *L. sigmodontis* also ligate the TGF- β R in a manner that cannot be inhibited with antibodies to TGF- β itself (Hartmann et al., 2015). Finally, a TGF- β homolog from *F. hepatica*, FhTLM, binds host receptors, but with lower affinity and this was inhibited by polyclonal anti-TGF- β antibodies. Immunologically, FhTLM induces IL-10 production in macrophages, indicative of a pro-regulatory outcome (Sulaiman et al., 2016).

In contrast to these instances of evolutionarily conserved members of the TGF- β family activating host receptors, a striking example of convergent evolution has been found in *H. polygyrus*. Secreted products from this intestinal nematode have been shown to activate the TGF- β signaling pathway to induce regulatory T cells *in vitro*, even in the presence of pro-inflammatory cytokines and to a similar degree as TGF- β itself




Figure 3. Helminth Modulators that Act on the Adaptive and Effector Immune Responses

Immune responses are controlled by helper T cell responses (Th1, Th2, or Th17) and B cell antibody responses. These responses can counter-regulate each other and are suppressed by regulatory T cell or regulatory B cell responses. Effector responses are mediated by T cell-derived cytokines which act on innate effector cells such as macrophages, eosinophils, basophils, and neutrophils. Helminth immunomodulators are shown in blue.

production of IL-10 and TGF- β (Kim et al., 2010). While the mechanism is assumed to parallel that of human galectin-9, this has not been verified for the helminth homolog.

Effector T cells are also targeted by certain products. The schistosome glutathione-S-transferase protein (P28GST) appears to promote a sufficiently strong eosinophil-rich Th2 cell response to prevent Th1 and Th17 cell-mediated inflammation in murine colitis in an eosinophil-dependent manner (Driss et al., 2016).

(Grainger et al., 2010). Subsequently, through biochemical fractionation, the factor responsible has been identified as an unrelated structure named TGM (TGF- β Mimic), a 5-domain protein around three times larger than TGF- β and able to bind with high affinity to both TGF- β RI and II, in distinction to mammalian TGF- β which ligates only to RII (Johnston et al., 2017). TGM is also active *in vivo*, for example in prolonging survival of fully allogeneic skin grafts (Johnston et al., 2017). Similarly to HpARI, HpTGM is also effective in cultures of human T cells, despite *H. polygyrus* lacking infectivity in humans.

While the TGF- β pathway directly induces regulatory T cells, other helminth mediators exploit indirect pathways to modulate host T cells. Hookworm-derived homologs of the tissue inhibitor of metalloproteases (TIMPs) modulate DC function, resulting in tolerogenic T cell responses (Cuéllar et al., 2009) and more recently these proteins (now called AIP-1 and AIP-2, for anti-inflammatory proteins) have been demonstrated to be effective quenchers of pathology in mouse models of airway allergy (Navarro et al., 2016) and TNBS-induced colitis (Ferreira et al., 2017). Protection is associated with increased Treg cell and IL-10 responses, alongside abated inflammatory cytokines and modified DC function. Effects are also seen in cultured human DCs and T cells, indicating potential as therapeutics for immune disorders (Navarro et al., 2016).

Treg cell induction *in vivo* is also seen with PAS-1, a 200-kDa native protein purified from *Ascaris suum*, which is reported to inhibit airway allergy in ovalbumin-alum primed mice (Araújo et al., 2008). PAS-1 induces an expands CD25⁺ Treg cell populations, which can suppress allergic responses on adoptive transfer (de Araújo et al., 2010). PAS-1 is similar to ABA-1, a previously described nematode tandemly repeated polypeptide with lipid-binding properties (Antunes et al., 2015; Meenan et al., 2011).

Host galectins promote Treg cell functionality and stability, and a homolog of human galectin-9 from *Toxascaris leonina* reduces DSS-induced colonic pathology in mice with increased

production of IL-10 and TGF- β (Kim et al., 2010). While the mechanism is assumed to parallel that of human galectin-9, this has not been verified for the helminth homolog.

Effector T cells are also targeted by certain products. The schistosome glutathione-S-transferase protein (P28GST) appears to promote a sufficiently strong eosinophil-rich Th2 cell response to prevent Th1 and Th17 cell-mediated inflammation in murine colitis in an eosinophil-dependent manner (Driss et al., 2016). In contrast, the *Echinococcus multilocularis* homolog of TIP (EmTIP) induces IFN- γ release from CD4⁺ T cells *in vitro*, which may explain how this parasite, unlike most other helminths, induces an initial Th1 cell response *in vivo* (Nono et al., 2014).

Potassium channels in lymphocytes, such as voltage-gated channel Kv1.3, are crucial for the activation and proliferation of T cells. Kv1.3 is considered an interesting therapeutic target in autoimmune conditions and inhibitors include ShK, a polypeptide toxin expressed in the sea anemone *Stichodactyla helianthus* (Norton et al., 2004). ShK homologs have been conserved and adapted in helminth T cell inhibitors, e.g., AcK1, found in the hookworms *Ancylostoma caninum* and *Ancylostoma ceylanicum*, and BmK1, identified in the filarial worm *B. malayi*. These peptides suppress a delayed-type hypersensitivity (DTH) reaction in rats and inhibit primarily the proliferation of memory T cells without affecting naive or central memory subsets (Chhabra et al., 2014).

An equally important target for helminth parasites is the host B cell response, which is known to be crucial for parasite ejection (Harris and Gause, 2011). Na-ASP-2 (*Ancylostoma* secreted protein 2 expressed by *N. americanus*) binds to CD79A, a component of the B cell receptor (BCR) complex, downregulating mRNA expression of targets involved in signaling and trans-endothelial migration (Tribolet et al., 2015) while ES-62 from *A. viteae* uncouples the BCR signaling machinery, rendering host cells unresponsive (Deehan et al., 2001). Other types of regulation of B cell responses in helminth infection are associated with a switch from the IgE to the IgG4 isotype, and a regulatory B cell (Breg) phenotype characterized by production of IL-10 (Hussaarts et al., 2011), similar to changes in B cell phenotypes found in allergen tolerance (Akdis and Akdis, 2014).

The secreted products of *S. mansoni* eggs can induce a regulatory B (Breg) cell phenotype in murine and human cells, through

or independently of the action of IPSE (Haeberlein et al., 2017). Uptake of IPSE is necessary to induce B cell IL-10 and Treg cell development *in vitro* (Haeberlein et al., 2017). IPSE also supports type 2 cell responses, through induction of IL-4 release from basophils that is IgE dependent but antigen independent and does not require uptake of IPSE by basophils, so this single protein has multiple functions and activation routes (Schramm et al., 2007).

By modulating CD4⁺ T cell differentiation, Treg cell induction, B cell isotype switching, and Breg cell induction, parasites can produce an environment most suited to them. These adaptive responses then determine effector cell responses and whether the outcome of infection is inflammation, resolution, parasite ejection, or tolerance.

Helminth Molecules Defeat Effector Cell Responses

The full spectrum of cell populations are involved in defense against helminths, from “professional” immune leukocytes to stromal and epithelial cells. In particular, granulocytes (eosinophils and neutrophils) are often the final effector cells in allergic and inflammatory responses, able to target the site of infection and kill pathogens by a variety of means including delivery of toxic compounds, sequestering vital nutrients, and destabilizing the parasite environment. Therefore, parasite survival can be maintained through secretions that block effector cell recruitment and function.

Chemokines are crucial in organizing the immune response through migration of immune cells to sites of inflammation. A key chemokine required for immune cell recruitment and activation in humans is IL-8. *B. malayi* produces asparaginyl-tRNA synthetase (AsnRS) which shows structural (but not sequence) homology to IL-8 and binds the IL-8 receptors CXCR1 and CXCR2, recruiting eosinophils and neutrophils (Kron et al., 2013). However, despite this apparently pro-inflammatory mechanism of action, administration of AsnRS in the mouse T cell transfer model of colitis induces regulatory responses and induction of IL-10 (Kron et al., 2013).

Conversely, human hookworms suppress chemokine responses through production of metalloproteinases which cleave eotaxin-1 (CCL11), but not IL-8 or eotaxin-2 (CCL24), specifically inhibiting the eotaxin-mediated recruitment of eosinophils (Culley et al., 2000). *S. mansoni* secretes a chemokine binding protein (SmCKBP) which binds and blocks IL-8, CCL2, CCL3, CCL5, and CX3CL1—this action inhibits neutrophil (but not eosinophil) migration and has suppressive activity in the DNFB contact hypersensitivity model, while blocking SmCKBP *in vivo* resulting in increased neutrophilia in *S. mansoni* egg-induced granulomas (Smith et al., 2005).

In concert with chemokines, cell migration also requires interaction with integrins such as CD11b/CD18 to facilitate extravasation. One hookworm protein, NIF from *Ancylostoma caninum*, binds this integrin together with fibrinogen and prevents binding of human neutrophils to vascular endothelium (Moyle et al., 1994). NIF is also effective in *in vitro* models of LPS-driven pulmonary inflammation (Zhou et al., 1998), and a NIF-IgE Fc fusion protein (NIF-IGHE-CH4) suppresses human neutrophil migration and blister formation in an *ex vivo* human model of subepidermal autoimmune blistering disease (Kemmer et al., 2015). Remarkably, NIF is a member of a multi-gene family (Pfam00188, vari-

ously named CAP, SCP, TAPS, or VAL) that is prominent across a wide range of helminths, and which includes a wide variety of immune response modulators in both animals and plants (Wilbers et al., 2018).

After responding to an infection, the host must heal after epithelial damage and inflammation. This can be mediated by immune pathways and helped (or hindered) by parasite products.

Helminths Can Promote Healing and Remodeling

As parasites enter the host, they must digest connective tissue to allow their migration, followed by rapid healing to maintain health of their host. They achieve this through secretion of proteases and other “chemical scissors” essential for the invasive process (Ranasinghe and McManus, 2017). Their entry is also facilitated by inhibiting coagulation, particularly prominent in blood-dwelling helminths like schistosomes with multiple strategies to manipulate host vasodilation, fibrinolysis, and coagulation (Mebius et al., 2013). Calpain family proteases SmCalp-1 and SmCalp-2 are expressed in the tegument of *S. mansoni* adult worms, where they appear to cleave and inactivate key coagulation proteins fibronectin and high-molecular-weight kininogen (HK) (Wang et al., 2018).

As discussed above, ATP-degrading enzymes (apyrases) are commonly secreted by helminths, suppressing DAMP-induced inflammatory responses. However, these enzymes have further roles in the control of thrombosis, as extracellular ATP and ADP are potent activators of platelets. Schistosome teguments contain ATP diphosphohydrolase (SmATPase1) (Da'dara et al., 2014) and the ectonucleotide pyrophosphatase/phosphodiesterase homolog SmNPP5 (Elzoheiry et al., 2018), both of which degrade ATP and ADP to non-thrombotic AMP.

Conversely, a serine protease-2 (SmSP-2) has been found to activate tissue plasminogen activator and plasminogen, both key components of the fibrinolytic system (Leontovyč et al., 2018). Similar activities have been described for other parasite secretions, such as the cathepsin L peptidases FhCL-1, -2, and -3 from *F. hepatica*, which degrade fibrinogen and fibrin (Mebius et al., 2018), and a plasminogen-binding alpha-enolase from *O. volvulus* (OvENO), which may promote proteolysis and degradation of the host extracellular matrix for migration of larvae through host tissues (Jolodar et al., 2003).

Parasite secretions can also encourage the healing of the epithelium. An extreme example of this is the bile duct-dwelling trematode *Opisthorcis viverrini*. This trematode feeds on bile duct epithelial cells and causes extensive inflammation *in situ*. It secretes a granulin homolog (Ov-GRN-1) which causes epithelial cell proliferation: this has potential as a therapeutic agent for non-healing wounds (Smout et al., 2015), but during infection can also result in epithelial transformation and contributes to mortality in *O. viverrini* infection due to cholangiocarcinoma (Smout et al., 2009).

Thus, the secretions of parasitic helminths can modulate every phase of immunity, from the most proximal events in immune initiation to terminal effector responses. In the next section we will discuss how these recent research findings could lead to the discovery of further parasite immunomodulators and how these could be developed for use in human immune-mediated diseases.




Figure 4. The Evolution and Diversity of Helminth Immunomodulation

(A) The TGF- β family is evolutionarily ancient, and nearly all helminth genomes (including free-living helminths) encode TGF- β family members. TGF- β homologs from *Brugia malayi* (Bm-TGH-2) and *Fasciola hepatica* (FhTLM) have co-evolved with their host to bind to the mammalian TGF- β receptor.

(B) In *Heligmosomoides polygyrus*, the CCP domain-containing family is over-represented in the genome, and HpTGM (a 5 CCP domain protein) has undergone convergent evolution to bind the TGF- β receptor, despite bearing no homology to the host cytokine. The CCP domain family in *H. polygyrus* appears highly adaptable, and the IL-33-blocking protein HpARI also consists of 3 CCP domains.

(C) T2 ribonucleases are involved in a number of homeostatic processes including RNA recycling, and most helminth genomes contain a member of this family. In the schistosomes, however, the T2 ribonuclease family has undergone expansion, and the T2 ribonuclease omega-1 (ω -1) is secreted into the host, where it gains entry to dendritic cells (DCs) through its Lewis X motifs which are bound by glycan receptors. Once inside the DC, omega-1 degrades host messenger and ribosomal RNA, suppressing DC activation and downstream T cell responses.

Blue arrows indicate evolutionary processes, black arrows indicate binding. Graphs show number of members of families in each group of genomes (WormBase ParaSite), represented as box and whiskers representing minimum to maximum values.

Future Horizons in Helminth Immune Modulation

The field of helminth immunomodulators has, until now, been dominated by protein actors, a tendency likely to continue with the torrent of sequence information extending to more than 100 helminth genomes, each with >10,000 protein-coding genes (International Helminth Genomes Consortium, 2018). Protein sequence analyses also provide insights into the molecular evolution of the different families of helminth immunomodulators (Figure 4). However, this should not overshadow the critical role of non-protein components (alone or in concert with protein moieties), which play many critical roles in helminth immunology.

A relatively neglected aspect of helminth biology has been their production of small molecules with potential for immune modulation, such as prostaglandins and short-chain fatty acids (SCFAs). Some helminths, for example *Ascaris*, release acetate and other metabolites likely to impact responses of the epithe-

lium (Tielens et al., 2010). In addition, helminths can modify the commensal microbiota in a fashion that raises SCFA concentrations, indirectly promoting regulatory mechanisms in the host (Zaiss et al., 2015). With metabolomics now being applied to helminths and genomics piecing together parasite biosynthetic pathways, the suite of such small molecules is likely to now expand rapidly and uncover many more critical interactions in host recognition and immune control. In a recent demonstration of this, small molecules from *A. caninum* suppressed pathology in a mouse model of TNBS-induced colitis (Wangchuk et al., 2018).

Lipid mediators are among the most potent small molecules in the host immune system, both activating and suppressing the immune response. They are <1 kDa, intensely active within their milieu, and have conserved chemical structures across multiple hosts. They represent a target for parasite neutralization or

molecular mimicry where helminths would benefit from their effects. PGE2 is an arachadonic acid metabolite, with roles in ILC2, T helper cell, macrophage, dendritic cell, and neutrophil responses (Martínez-Colón and Moore, 2018). PGE2 is strongly linked to Th2 cell and ILC2 responses, for example inducing IL-22-mediated atopic dermatitis (Robb et al., 2018), although it can also directly inhibit ILC2s (Maric et al., 2018). *T. suis*, *S. mansoni*, and filarial nematode secretions contain PGE2 (Laan et al., 2017; Liu et al., 1992), and *S. mansoni* SEA has the ability to induce further PGE2 from host cells (Kaisar et al., 2018), in this case mediating Th2 cell induction.

In the context of the pivotal role of lipid mediators, more attention is likely to be focused on lipid-binding proteins that are numerous among helminth products (Franchini et al., 2015). For example, *F. hepatica* Fh12 (also known as fatty acid binding protein) mentioned above induces alternative activation in human Mo-Macs and suppresses LPS-induced activation via binding and blocking of CD14.

A paradigm-shifting discovery has been that helminths release extracellular vesicles (EVs), often termed exosomes, which are able to fuse and enter host cells, carrying with them a cargo of miRNAs (Buck et al., 2014; Coakley et al., 2017). In *H. polygyrus* infections, these EVs target epithelial cells and macrophages, resulting in their loss of expression of the IL-33 receptor (IL-33R, also termed ST2) and downregulation of other immune genes. These effects result in suppression of both alternative and inflammatory macrophage activation (Coakley et al., 2017). These *in vitro* findings have been corroborated *in vivo*, as *N. brasiliensis* EVs protect against TNBS-induced colitis (Eichenberger et al., 2018), and *H. polygyrus* EVs suppress *Alternaria* allergen-induced allergic airway responses (Buck et al., 2014). In contrast, EVs from *S. japonicum* actively drive M1 macrophage differentiation *in vitro* (Wang et al., 2015), joining a range of helminth-secreted proteins that counter-intuitively induce inflammatory responses (Kron et al., 2013; Noya et al., 2017). Hence EVs are an adaptable part of the armory of helminth immunomodulators, allowing extracellular pathogens to manipulate events within host cells in a receptor-independent manner (Coakley et al., 2017). Further research is required in this area, to investigate the level of specificity of particular EV populations for specific host cells, to determine the relative importance of EV protein and nucleic acid cargoes, and to develop these findings toward potential therapeutic agents for human disease.

Identification, Characterization, and Clinical Development of the Next Generation of Parasite Immunomodulators

As the identification of defined immunomodulators accelerates, the scale of what the future may hold is also becoming apparent. Each large and complex helminth genome can encode 10,000–50,000 genes (Zarowiecki and Berriman, 2015) (compared to 20,000 in the human genome), and as a result many million previously unannotated genes are now being cataloged. Moreover, parasite secretions are extremely complex, containing hundreds to thousands of protein, carbohydrate, lipid, nucleotide, and vesicular moieties, each of which may be finely tuned to a particular host range.

Parasitism has arisen on multiple occasions within the helminth phyla, and each lineage has evolved specific molecular

strategies to defeat the host immune system. As helminths are metazoan members of the animal kingdom, they share many gene families with vertebrates that might provide a template for the molecular evolution of modulators interactive with counterparts in the host. Thus, as the host-adapted genes in the evolution of the immune system, parasites may have followed suit in parallel. Some cases, described below, appear to fulfil this prediction. In contrast, however, some of the most striking parasite immunomodulators such as HpARI, HpTGM, omega-1, and IPSE show no appreciable homology to host immune genes or previously known parasite immunomodulatory factors.

Immunomodulators derived from an ancient shared ancestor of both parasites and their hosts include the TGF- β and MIF families. In the case of TGF- β , a gene family originally required for developmental processes expanded to encompass members with cytokine functions. Thus, as the host developed an immunological role for TGF- β , some parasites have been able to adapt pre-existing developmental homologs toward an immune-modulatory role. Therefore, the TGF- β homologs encoded in parasite genomes do not necessarily indicate immune modulatory functions, while conversely a single TGF- β homolog may fulfil both developmental and immunomodulatory roles (Figure 4A).

It was recently discovered that, rather than adapt a pre-existing TGF- β homolog to interact with host receptors, *H. polygyrus* has convergently evolved an unrelated structure in the form of Hp-TGM, a protein consisting of five atypical complement control protein (CCP) domains with no homology to TGF- β , which ligates host TGF- β receptor (Johnston et al., 2017; Smyth et al., 2018). Furthermore, the same parasite blocks IL-33 through HpARI, another CCP domain protein (Osborn et al., 2017). Although CCP domains are present in all helminths studied, there has been a preferential expansion of these genes in *H. polygyrus* (Figure 4B). CCP domains occur in a wide range of organisms and serve diverse functions in neuronal signaling, cytokine receptors, hemoglobin binding, and of course, complement control. CCP domain proteins are highly stable and soluble (each tight, globular CCP domain contains two disulfide bonds and conserved glycosylation), structurally simple but functionally sophisticated (Makou et al., 2015) with independent activities of multiple domains within a single protein. This adaptable scaffold appears to have been exploited by *H. polygyrus* in particular.

Gene family expansion in helminths hints at functional adaptation to the parasitic life style. Among the most notable is Pfam00188, which in helminths include venom allergen-like (VAL) and *Ancylostoma* secreted (ASP) proteins. This gene family is dramatically expanded in many parasitic nematodes (especially the hookworms and strongylides) and have been implicated in lipid binding, immune modulation, and pathogen resistance, so they may, like the CCP domain family, represent an adaptable scaffold upon which a range of functions can be developed (Wilbers et al., 2018).

Conversely, expansion of the T2 ribonuclease family (containing omega-1) (Everts et al., 2012) appears specific to schistosomes (Figure 4C), implying that this functionality may have evolved within the trematode family relatively recently. T2 ribonucleases are broadly expressed across many parasites, plants, and animals and are implicated in recycling of RNA and in RNA

degradation during the stress response. T2 ribonucleases typically enter the secretory pathway and are often found in vacuoles and autophagosomes within cells. However, some T2 ribonucleases (for instance E^{ms} from pestiviruses) can also be secreted (Luhtala and Parker, 2010), for degradation of RNA in extracellular spaces or in target cells, and it is this latter function which appears to have been utilized in the schistosomes, degrading RNA in host DCs and suppressing their maturation (Everts et al., 2012). Thus, schistosomes have “weaponized” a normal metabolic pathway to suppress DC maturation.

Similarly the apyrase family, which is expanded in many parasitic nematodes (International Helminth Genomes Consortium, 2018), may have originally been a metabolic enzyme but have subsequently used to suppress of host damage detection and inhibit blood coagulation (Da'dara et al., 2014; Gounaris et al., 2004).

These examples of helminth modulation exemplify the techniques used by helminths to develop immunomodulatory products: convergent evolution of shared protein families, *de novo* production of cytokine mimics or antagonists from adaptable modular domains, and hi-jacking of pre-existing protein functions. The agenda for the future is to test whether administration of defined helminth-derived molecular entities (with known functions and in a specific context) can lead us to clinically effective therapeutic molecules.

Molecular Therapies from Helminths—Where Are We?

In 2005, the first helminth therapy trials (using live *T. suis* infection) showed impressive effects in IBD (Weinstock and Elliott, 2013) and generated a great deal of interest in the use of live helminths as therapeutic modalities. However, subsequent trials have shown smaller or absent effect sizes (Feary et al., 2010; Fleming et al., 2017; Schölmerich et al., 2017), and a lack of understanding of the mechanistic underpinning of helminth immunomodulation has hampered progression. With the recent advances in understanding of helminth immunomodulation at the molecular level, enormous potential exists for helminth-derived products for clinical use.

ES-62 was one of the original immunomodulatory molecules identified from parasites and with its activity through the small molecule phosphorylcholine (PC) moiety, it has great potential for therapeutic drug-like molecules. Two small molecule analogs (SMAs) of PC, termed 11a and 12b, can replicate many of the suppressive effects of ES-62, and due to their small size, stability, and lack of immunogenicity and efficacy are now being developed as therapeutic agents (Al-Riyami et al., 2013). Similarly, many of the recently identified immunomodulatory proteins are likely to be pursued as potential anti-inflammatory therapeutics for clinical use. These proteins have evolved to function in the mammalian milieu and represent natural biologicals that may offer unique mechanisms of action not available with existing drugs; in this context, the coming years of translational research promise to be very engaging.

We summarize here a remarkable panoply of helminth molecules which, when tested in isolation *in vitro* or *in vivo*, target distinct host populations and pathways. Little is yet known, however, of how products from parasites in specific niches (such as the intestinal lumen) are able to engage the broader host immune system, whether they exploit uptake mechanisms for systemic

dissemination or rely on localized effects in their preferred site. Certainly, in species such as *H. polygyrus*, secretion products differ markedly between immature tissue-dwelling larvae and the intestinal lumen-dwelling adults (Hewitson et al., 2013), arguing for developmental adaptability for interaction with the immune system. In other cases, blood-dwelling parasites such as schistosomes or lymphatic-dwelling filarial worms have more direct access to immune cells and may more readily exert a systemic effect.

Although parasite immunomodulators may act potently in laboratory tests, one obstacle to clinical development has been potential immunogenicity as foreign proteins *in vivo*. However, immunogenicity of helminth immunomodulators would represent an adverse outcome for the parasite, and thus mechanisms may be in place (including the innately immunosuppressive nature of these molecules) for avoidance of this: the fully foreign hookworm product AIP-2 is minimally immunogenic *in vivo* (Navarro et al., 2016).

There are also good precedents for the development of pathogen-derived proteins that have been successfully developed for human use, despite concerns of immunogenicity. Streptokinase (a *Streptococcus*-derived thrombolytic) has been used for decades in thrombotic disease such as myocardial infarction. Although highly immunogenic, as long as the drug is not repeatedly administered, it is effective and its immunogenicity does not present a barrier to use (Sikri and Bardia, 2007). Likewise, the myxoma viral protein Serp1 is an anti-inflammatory inhibitor of serine proteases; administration of Serp1 causes a non-neutralizing antibody response and subsequently the functional activity of the protein has been extended (and potential immunogenicity decreased) through development of small peptide mimetics that replicate the function of the full protein (Ambadipadi et al., 2016). Therefore, formulations of helminth immunomodulators are being developed in which immunogenicity is not present, or is not a barrier to use, for future therapeutic applications.

Conclusions

Helminth immunomodulation is coming of age. Through advances in -omics analysis of parasitic helminths, ever more information about helminth products is becoming available. However, only by careful analysis of defined recombinant products can the potential of helminth therapeutic agents be realized. In recent years this has led to the characterization of a handful of helminth immunomodulators that could be developed for use in the clinic. Considering the size and complexity of helminth secretions, the complex interaction with their host, and the intricacy of the immune response directed against them, it appears almost certain that we have only scratched the surface of the full gamut of helminth immunomodulators, and that many more will be identified in the next decade.

ACKNOWLEDGMENTS

R.M.M. is supported by the Wellcome Trust through an Investigator Award (106122), by the Kenneth Rainin Foundation (2016-3067), and by core funding to the Wellcome Centre for Molecular Parasitology (104111). H.H.S. is supported by a ZonMW-VIDI grant (91714352) from the Netherlands Organisation for Scientific Research and a Consortium Grant (5115015) from the Netherlands Lung Foundation. R.M.M., H.H.S., and H.J.M. are supported by

a Consortium grant (12017001) from the Netherlands Lung Foundation Accelerate Programme.

REFERENCES

- Akdis, M., and Akdis, C.A. (2014). Mechanisms of allergen-specific immunotherapy: multiple suppressor factors at work in immune tolerance to allergens. *J. Allergy Clin. Immunol.* **133**, 621–631.
- Al-Riyami, L., Pineda, M.A., Rzepecka, J., Huggan, J.K., Khalaf, A.I., Suckling, C.J., Scott, F.J., Rodgers, D.T., Harnett, M.M., and Harnett, W. (2013). Designing anti-inflammatory drugs from parasitic worms: a synthetic small molecule analogue of the *Acanthocheilonema viteae* product ES-62 prevents development of collagen-induced arthritis. *J. Med. Chem.* **56**, 9982–10002.
- Alvarado, R., To, J., Lund, M.E., Pinar, A., Mansell, A., Robinson, M.W., O'Brien, B.A., Dalton, J.P., and Donnelly, S. (2017). The immune modulatory peptide FhHDM-1 secreted by the helminth *Fasciola hepatica* prevents NLRP3 inflammasome activation by inhibiting endolysosomal acidification in macrophages. *FASEB J.* **31**, 85–95.
- Ambadapadi, S., Munuswamy-Ramanujam, G., Zheng, D., Sullivan, C., Dai, E., Morshed, S., McFadden, B., Feldman, E., Pinard, M., McKenna, R., et al. (2016). Reactive center loop (RCL) peptides derived from serpins display independent coagulation and immune modulating activities. *J. Biol. Chem.* **291**, 2874–2887.
- Anbu, K.A., and Joshi, P. (2008). Identification of a 55 kDa *Haemonchus contortus* excretory/secretory glycoprotein as a neutrophil inhibitory factor. *Parasite Immunol.* **30**, 23–30.
- Antunes, M.F., Titz, T.O., Batista, I.F., Marques-Porto, R., Oliveira, C.F., Alves de Araujo, C.A., and Macedo-Soares, M.F. (2015). Immunosuppressive PAS-1 is an excretory/secretory protein released by larval and adult worms of the ascarid nematode *Ascaris suum*. *J. Helminthol.* **89**, 367–374.
- Araújo, C.A., Perini, A., Martins, M.A., Macedo, M.S., and Macedo-Soares, M.F. (2008). PAS-1, a protein from *Ascaris suum*, modulates allergic inflammation via IL-10 and IFN- γ , but not IL-12. *Cytokine* **44**, 335–341.
- Ball, D.H., Al-Riyami, L., Harnett, W., and Harnett, M.M. (2018). IL-33/ST2 signalling and crosstalk with Fc ϵ R1 and TLR4 is targeted by the parasitic worm product, ES-62. *Sci. Rep.* **8**, 4497.
- Bąska, P., Wiśniewski, M., Krzyżowska, M., Długosz, E., Zygmier, W., Górski, P., and Wędrychowicz, H. (2013). Molecular cloning and characterisation of in vitro immune response against astacin-like metalloprotease Ace-MTP-2 from *Ancylostoma ceylanicum*. *Exp. Parasitol.* **133**, 472–482.
- Beall, M.J., and Pearce, E.J. (2001). Human transforming growth factor- β activates a receptor serine/threonine kinase from the intravascular parasite *Schistosoma mansoni*. *J. Biol. Chem.* **276**, 31613–31619.
- Bethony, J., Brooker, S., Albonico, M., Geiger, S.M., Loukas, A., Diemert, D., and Hotez, P.J. (2006). Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* **367**, 1521–1532.
- Buck, A.H., Coakley, G., Simbari, F., McSorley, H.J., Quintana, J.F., Le Bihan, T., Kumar, S., Abreu-Goodger, C., Lear, M., Harcus, Y., et al. (2014). Exosomes secreted by nematode parasites transfer small RNAs to mammalian cells and modulate innate immunity. *Nat. Commun.* **5**, 5488.
- Cekic, C., and Linden, J. (2016). Purinergic regulation of the immune system. *Nat. Rev. Immunol.* **16**, 177–192.
- Chen, W., and Ten Dijke, P. (2016). Immunoregulation by members of the TGF β superfamily. *Nat. Rev. Immunol.* **16**, 723–740.
- Chhabra, S., Chang, S.C., Nguyen, H.M., Huq, R., Tanner, M.R., Londono, L.M., Estrada, R., Dhawan, V., Chauhan, S., Upadhyay, S.K., et al. (2014). Kv1.3 channel-blocking immunomodulatory peptides from parasitic worms: implications for autoimmune diseases. *FASEB J.* **28**, 3952–3964.
- Cho, M.K., Park, M.K., Kang, S.A., Park, S.K., Lyu, J.H., Kim, D.H., Park, H.K., and Yu, H.S. (2015). TLR2-dependent amelioration of allergic airway inflammation by parasitic nematode type II MIF in mice. *Parasite Immunol.* **37**, 180–191.
- Coakley, G., McCaskill, J.L., Borger, J.G., Simbari, F., Robertson, E., Millar, M., Harcus, Y., McSorley, H.J., Maizels, R.M., and Buck, A.H. (2017). Extracellular vesicles from a helminth parasite suppress macrophage activation and constitute an effective vaccine for protective immunity. *Cell Rep.* **19**, 1545–1557.
- Cohen, E.S., Scott, I.C., Majithiya, J.B., Rapley, L., Kemp, B.P., England, E., Rees, D.G., Overed-Sayer, C.L., Woods, J., Bond, N.J., et al. (2015). Oxidation of the alarmin IL-33 regulates ST2-dependent inflammation. *Nat. Commun.* **6**, 8327.
- Coronado, S., Barrios, L., Zakzuk, J., Regino, R., Ahumada, V., Franco, L., Ocampo, Y., and Caraballo, L. (2017). A recombinant cystatin from *Ascaris lumbricoides* attenuates inflammation of DSS-induced colitis. *Parasite Immunol.* **39**, e12425.
- Crowe, J., Lumb, F.E., Harnett, M.M., and Harnett, W. (2017). Parasite excretory-secretory products and their effects on metabolic syndrome. *Parasite Immunol.* **39**, e12410.
- Cuéllar, C., Wu, W., and Mendez, S. (2009). The hookworm tissue inhibitor of metalloproteases (Ac-TMP-1) modifies dendritic cell function and induces generation of CD4 and CD8 suppressor T cells. *PLoS Negl. Trop. Dis.* **3**, e439.
- Culley, F.J., Brown, A., Conroy, D.M., Sabroe, I., Pritchard, D.I., and Williams, T.J. (2000). Eotaxin is specifically cleaved by hookworm metalloproteases preventing its action in vitro and in vivo. *J. Immunol.* **165**, 6447–6453.
- Da'dara, A.A., Bhardwaj, R., Ali, Y.B., and Skelly, P.J. (2014). Schistosome tegumental ecto-ATPase (SmATPase1) degrades exogenous pro-inflammatory and pro-thrombotic nucleotides. *PeerJ* **2**, e316.
- Dainichi, T., Maekawa, Y., Ishii, K., Zhang, T., Nashed, B.F., Sakai, T., Takashima, M., and Himeno, K. (2001). Nippocystatin, a cysteine protease inhibitor from *Nippostrongylus brasiliensis*, inhibits antigen processing and modulates antigen-specific immune response. *Infect. Immun.* **69**, 7380–7386.
- Daniłowicz-Luebert, E., Steinfeldt, S., Kühl, A.A., Drozdenko, G., Lucius, R., Worm, M., Hamelmann, E., and Hartmann, S. (2013). A nematode immunomodulator suppresses grass pollen-specific allergic responses by controlling excessive Th2 inflammation. *Int. J. Parasitol.* **43**, 201–210.
- David, C.J., and Massague, J. (2018). Contextual determinants of TGF β action in development, immunity and cancer. *Nat. Rev. Mol. Cell Biol.* **19**, 419–435.
- de Araújo, C.A.A., Perini, A., Martins, M.A., Macedo, M.S., and Macedo-Soares, M.F. (2010). PAS-1, an *Ascaris suum* protein, modulates allergic airway inflammation via CD8 $^+$ $\gamma\delta$ TCR $^+$ and CD4 $^+$ CD25 $^+$ FoxP3 $^+$ T cells. *Scand. J. Immunol.* **72**, 491–503.
- Deehan, M.R., Harnett, W., and Harnett, M.M. (2001). A filarial nematode-secreted phosphorylcholine-containing glycoprotein uncouples the B cell antigen receptor from extracellular signal-regulated kinase-mitogen-activated protein kinase by promoting the surface Ig-mediated recruitment of Src homology 2 domain-containing tyrosine phosphatase-1 and Pac-1 mitogen-activated kinase-phosphatase. *J. Immunol.* **166**, 7462–7468.
- Donnelly, S., O'Neill, S.M., Sekiya, M., Mulcahy, G., and Dalton, J.P. (2005). Thioredoxin peroxidase secreted by *Fasciola hepatica* induces the alternative activation of macrophages. *Infect. Immun.* **73**, 166–173.
- Donnelly, S., Stack, C.M., O'Neill, S.M., Sayed, A.A., Williams, D.L., and Dalton, J.P. (2008). Helminth 2-Cys peroxiredoxin drives Th2 responses through a mechanism involving alternatively activated macrophages. *FASEB J.* **22**, 4022–4032.
- Donnelly, S., O'Neill, S.M., Stack, C.M., Robinson, M.W., Turnbull, L., Whitchurch, C., and Dalton, J.P. (2010). Helminth cysteine proteases inhibit TRIF-dependent activation of macrophages via degradation of TLR3. *J. Biol. Chem.* **285**, 3383–3392.
- Doonan, J., Lumb, F.E., Pineda, M.A., Tarafdar, A., Crowe, J., Khan, A.M., Suckling, C.J., Harnett, M.M., and Harnett, W. (2018). Protection against arthritis by the parasitic worm product ES-62, and its drug-like small molecule analogues, is associated with inhibition of osteoclastogenesis. *Front. Immunol.* **9**, 1016.
- Driss, V., El Nady, M., Delbeke, M., Rousseaux, C., Dubuquoy, C., Sarazin, A., Gatault, S., Dendooven, A., Riveau, G., Colombel, J.F., et al. (2016). The schistosome glutathione S-transferase P28GST, a unique helminth protein, prevents intestinal inflammation in experimental colitis through a Th2-type response with mucosal eosinophils. *Mucosal Immunol.* **9**, 322–335.
- Eichenberger, R.M., Ryan, S., Jones, L., Buitrago, G., Polster, R., Montes de Oca, M., Zuvelek, J., Giacomini, P.R., Dent, L.A., Engwerda, C.R., et al.

- (2018). Hookworm secreted extracellular vesicles interact with host cells and prevent inducible colitis in mice. *Front. Immunol.* **9**, 850.
- Elzoheiry, M., Da'dara, A.A., deLaforcade, A.M., El-Beshbishi, S.N., and Skelly, P.J. (2018). The essential ectoenzyme SmNPP5 from the human intravascular parasite *Schistosoma mansoni* is an ADPase and a potent inhibitor of platelet aggregation. *Thromb. Haemost.* **118**, 979–989.
- Everts, B., Perona-Wright, G., Smits, H.H., Hokke, C.H., van der Ham, A.J., Fitzsimmons, C.M., Doenhoff, M.J., van der Bosch, J., Mohrs, K., Haas, H., et al. (2009). Omega-1, a glycoprotein secreted by *Schistosoma mansoni* eggs, drives Th2 responses. *J. Exp. Med.* **206**, 1673–1680.
- Everts, B., Hussaarts, L., Driessen, N.N., Meevissen, M.H., Schramm, G., van der Ham, A.J., van der Hoeven, B., Scholzen, T., Burgdorf, S., Mohrs, M., et al. (2012). Schistosome-derived omega-1 drives Th2 polarization by suppressing protein synthesis following internalization by the mannose receptor. *J. Exp. Med.* **209**, 1753–1767, S1.
- Falcón, C.R., Masih, D., Gatti, G., Sanchez, M.C., Motrán, C.C., and Cervi, L. (2014). *Fasciola hepatica* Kunitz type molecule decreases dendritic cell activation and their ability to induce inflammatory responses. *PLoS ONE* **9**, e114505.
- Faveeuw, C., Mallevaey, T., Paschinger, K., Wilson, I.B.H., Fontaine, J., Mollicone, R., Oriol, R., Altmann, F., Lerouge, P., Capron, M., and Trottein, F. (2003). Schistosome N-glycans containing core α 3-fucose and core β 2-xylose epitopes are strong inducers of Th2 responses in mice. *Eur. J. Immunol.* **33**, 1271–1281.
- Feary, J.R., Venn, A.J., Mortimer, K., Brown, A.P., Hooi, D., Falcone, F.H., Pritchard, D.I., and Britton, J.R. (2010). Experimental hookworm infection: a randomized placebo-controlled trial in asthma. *Clin. Exp. Allergy* **40**, 299–306.
- Ferguson, B.J., Newland, S.A., Gibbs, S.E., Tourlomousis, P., Fernandes dos Santos, P., Patel, M.N., Hall, S.W., Walczak, H., Schramm, G., Haas, H., et al. (2015). The *Schistosoma mansoni* T2 ribonuclease omega-1 modulates inflammation-dependent IL-1 β secretion in macrophages. *Int. J. Parasitol.* **45**, 809–813.
- Ferreira, I.B., Pickering, D.A., Troy, S., Croese, J., Loukas, A., and Navarro, S. (2017). Suppression of inflammation and tissue damage by a hookworm recombinant protein in experimental colitis. *Clin. Transl. Immunology* **6**, e157.
- Fleming, J., Hernandez, G., Hartman, L., Maksimovic, J., Nace, S., Lawler, B., Risa, T., Cook, T., Agni, R., Reichelderfer, M., et al. (2017). Safety and efficacy of helminth treatment in relapsing-remitting multiple sclerosis: Results of the HINT 2 clinical trial. *Mult. Scler.* Published online October 1, 2017. <https://doi.org/10.1177/1352458517736377>.
- Franchini, G.R., Pórfido, J.L., Ibáñez Shimabukuro, M., Rey Burusco, M.F., Bégamo, J.A., Smith, B.O., Kennedy, M.W., and Córscio, B. (2015). The unusual lipid binding proteins of parasitic helminths and their potential roles in parasitism and as therapeutic targets. *Prostaglandins Leukot. Essent. Fatty Acids* **93**, 31–36.
- Gomez-Escobar, N., Gregory, W.F., and Maizels, R.M. (2000). Identification of *tgh-2*, a filarial nematode homolog of *Caenorhabditis elegans daf-7* and human transforming growth factor β , expressed in microfilarial and adult stages of *Brugia malayi*. *Infect. Immun.* **68**, 6402–6410.
- Gomez-Escobar, N., Bennett, C., Prieto-Lafuente, L., Aebischer, T., Blackburn, C.C., and Maizels, R.M. (2005). Heterologous expression of the filarial nematode *alt* gene products reveals their potential to inhibit immune function. *BMC Biol.* **3**, 8.
- Goodridge, H.S., McGuinness, S., Houston, K.M., Egan, C.A., Al-Riyami, L., Alcocer, M.J., Harnett, M.M., and Harnett, W. (2007). Phosphorylcholine mimics the effects of ES-62 on macrophages and dendritic cells. *Parasite Immunol.* **29**, 127–137.
- Gounaris, K., Selkirk, M.E., and Sadeghi, S.J. (2004). A nucleotidase with unique catalytic properties is secreted by *Trichinella spiralis*. *Mol. Biochem. Parasitol.* **136**, 257–264.
- Grainger, J.R., Smith, K.A., Hewitson, J.P., McSorley, H.J., Harcus, Y., Filbey, K.J., Finney, C.A., Greenwood, E.J., Knox, D.P., Wilson, M.S., et al. (2010). Helminth secretions induce de novo T cell Foxp3 expression and regulatory function through the TGF- β pathway. *J. Exp. Med.* **207**, 2331–2341.
- Haeberlein, S., Obieglo, K., Ozir-Fazalalikhani, A., Chayé, M.A.M., Veninga, H., van der Vlugt, L.E.P.M., Voskamp, A., Boon, L., den Haan, J.M.M., Westerhof, L.B., et al. (2017). Schistosome egg antigens, including the glycoprotein IPSE/alpha-1, trigger the development of regulatory B cells. *PLoS Pathog.* **13**, e1006539.
- Hagen, J., Young, N.D., Every, A.L., Pagel, C.N., Schnoeller, C., Scheerlinck, J.P., Gasser, R.B., and Kalinna, B.H. (2014). Omega-1 knockdown in *Schistosoma mansoni* eggs by lentivirus transduction reduces granuloma size in vivo. *Nat. Commun.* **5**, 5375.
- Hams, E., Bermingham, R., Wurlod, F.A., Hogan, A.E., O'Shea, D., Preston, R.J., Rodewald, H.R., McKenzie, A.N.J., and Fallon, P.G. (2016). The helminth T2 RNase ω 1 promotes metabolic homeostasis in an IL-33- and group 2 innate lymphoid cell-dependent mechanism. *FASEB J.* **30**, 824–835.
- Harris, N., and Gause, W.C. (2011). To B or not to B: B cells and the Th2-type immune response to helminths. *Trends Immunol.* **32**, 80–88.
- Hartmann, W., Schramm, C., and Breloer, M. (2015). *Litomosoides sigmodontis* induces TGF- β receptor responsive, IL-10-producing T cells that suppress bystander T-cell proliferation in mice. *Eur. J. Immunol.* **45**, 2568–2581.
- Hewitson, J.P., Ivens, A.C., Harcus, Y., Filbey, K.J., McSorley, H.J., Murray, J., Bridgett, S., Ashford, D., Dowle, A.A., and Maizels, R.M. (2013). Secretion of protective antigens by tissue-stage nematode larvae revealed by proteomic analysis and vaccination-induced sterile immunity. *PLoS Pathog.* **9**, e1003492.
- Hokke, C.H., and van Diepen, A. (2017). Helminth glycomics - glycan repertoires and host-parasite interactions. *Mol. Biochem. Parasitol.* **215**, 47–57.
- Hussaarts, L., van der Vlugt, L.E., Yazdanbakhsh, M., and Smits, H.H. (2011). Regulatory B-cell induction by helminths: implications for allergic disease. *J. Allergy Clin. Immunol.* **128**, 733–739.
- Hussaarts, L., García-Tardón, N., van Beek, L., Heemskerck, M.M., Haeberlein, S., van der Zon, G.C., Ozir-Fazalalikhani, A., Berbé, J.F., Willems van Dijk, K., van Harmelen, V., et al. (2015). Chronic helminth infection and helminth-derived egg antigens promote adipose tissue M2 macrophages and improve insulin sensitivity in obese mice. *FASEB J.* **29**, 3027–3039.
- International Helminth Genomes Consortium (2018). Comparative genomics of the major parasitic worms. *Nat. Genet.* Published online November 5, 2018. <https://doi.org/10.1038/s41588-018-0262-1>.
- Jang, S.W., Cho, M.K., Park, M.K., Kang, S.A., Na, B.-K., Ahn, S.C., Kim, D.-H., and Yu, H.S. (2011). Parasitic helminth cystatin inhibits DSS-induced intestinal inflammation via IL-10⁺F4/80⁺ macrophage recruitment. *Korean J. Parasitol.* **49**, 245–254.
- Johnston, C.J.C., Smyth, D.J., Kodali, R.B., White, M.P.J., Harcus, Y., Filbey, K.J., Hewitson, J.P., Hinck, C.S., Ivens, A., Kemter, A.M., et al. (2017). A structurally distinct TGF- β mimic from an intestinal helminth parasite potently induces regulatory T cells. *Nat. Commun.* **8**, 1741.
- Jolodar, A., Fischer, P., Bergmann, S., Büttner, D.W., Hammerschmidt, S., and Brattig, N.W. (2003). Molecular cloning of an α -enolase from the human filarial parasite *Onchocerca volvulus* that binds human plasminogen. *Biochim. Biophys. Acta* **1627**, 111–120.
- Kaiser, M.M.M., Ritter, M., Del Fresno, C., Jónasdóttir, H.S., van der Ham, A.J., Pelgrom, L.R., Schramm, G., Layland, L.E., Sancho, D., Prazeres da Costa, C., et al. (2018). Dectin-1/2-induced autocrine PGE2 signaling licenses dendritic cells to prime Th2 responses. *PLoS Biol.* **16**, e2005504.
- Kaur, I., Schramm, G., Everts, B., Scholzen, T., Kindle, K.B., Beetz, C., Montiel-Duarte, C., Blindow, S., Jones, A.T., Haas, H., et al. (2011). Interleukin-4-inducing principle from *Schistosoma mansoni* eggs contains a functional C-terminal nuclear localization signal necessary for nuclear translocation in mammalian cells but not for its uptake. *Infect. Immun.* **79**, 1779–1788.
- Ke, X.D., Shen, S., Song, L.J., Yu, C.X., Kikuchi, M., Hirayama, K., Gao, H., Wang, J., Yin, X., Yao, Y., et al. (2017). Characterization of *Schistosoma japonicum* CP1412 protein as a novel member of the ribonuclease T2 molecule family with immune regulatory function. *Parasit. Vectors* **10**, 89.
- Kemmer, A., Bieber, K., Abadpour, A., Yu, X., Mitschker, N., Roth, S., Kauderer, C., Ludwig, R.J., Seeger, K., Köhl, J., et al. (2015). A recombinant fusion protein derived from dog hookworm inhibits autoantibody-induced dermal-epidermal separation ex vivo. *Exp. Dermatol.* **24**, 872–878.
- Kim, J.Y., Cho, M.K., Choi, S.H., Lee, K.H., Ahn, S.C., Kim, D.-H., and Yu, H.S. (2010). Inhibition of dextran sulfate sodium (DSS)-induced intestinal inflammation via enhanced IL-10 and TGF- β production by galectin-9 homologues isolated from intestinal parasites. *Mol. Biochem. Parasitol.* **174**, 53–61.

- Kron, M.A., Metwali, A., Vodanovic-Jankovic, S., and Elliott, D. (2013). Nematode asparaginyl-tRNA synthetase resolves intestinal inflammation in mice with T-cell transfer colitis. *Clin. Vaccine Immunol.* **20**, 276–281.
- Laan, L.C., Williams, A.R., Stavenhagen, K., Giera, M., Kooij, G., Vlasakov, I., Kalay, H., Kringle, H., Nejsun, P., Thamsborg, S.M., et al. (2017). The whipworm (*Trichuris suis*) secretes prostaglandin E2 to suppress proinflammatory properties in human dendritic cells. *FASEB J.* **31**, 719–731.
- Leontovyc, A., Ulrychová, L., O'Donoghue, A.J., Vondrášek, J., Marešová, L., Hubálek, M., Fajtová, P., Chanová, M., Jiang, Z., Craik, C.S., et al. (2018). SmSP2: A serine protease secreted by the blood fluke pathogen *Schistosoma mansoni* with anti-hemostatic properties. *PLoS Negl. Trop. Dis.* **12**, e0006446.
- Liu, L.X., Buhlmann, J.E., and Weller, P.F. (1992). Release of prostaglandin E2 by microfilariae of *Wuchereria bancrofti* and *Brugia malayi*. *Am. J. Trop. Med. Hyg.* **46**, 520–523.
- Luhtala, N., and Parker, R. (2010). T2 Family ribonucleases: ancient enzymes with diverse roles. *Trends Biochem. Sci.* **35**, 253–259.
- Makou, E., Herbert, A.P., and Barlow, P.N. (2015). Creating functional sophistication from simple protein building blocks, exemplified by factor H and the regulators of complement activation. *Biochem. Soc. Trans.* **43**, 812–818.
- Manoury, B., Gregory, W.F., Maizels, R.M., and Watts, C. (2001). *Bm*-CPI-2, a cystatin homolog secreted by the filarial parasite *Brugia malayi*, inhibits class II MHC-restricted antigen processing. *Curr. Biol.* **11**, 447–451.
- Maric, J., Ravindran, A., Mazzurana, L., Björklund, A.K., Van Acker, A., Rao, A., Friberg, D., Dahlén, S.E., Heinemann, A., Konya, V., and Mjösberg, J. (2018). Prostaglandin E₂ suppresses human group 2 innate lymphoid cell function. *J. Allergy Clin. Immunol.* **141**, 1761–1773.e6.
- Martin, I., Cabán-Hernández, K., Figueroa-Santiago, O., and Espino, A.M. (2015). *Fasciola hepatica* fatty acid binding protein inhibits TLR4 activation and suppresses the inflammatory cytokines induced by lipopolysaccharide in vitro and in vivo. *J. Immunol.* **194**, 3924–3936.
- Martínez-Colón, G.J., and Moore, B.B. (2018). Prostaglandin E2 as a regulator of immunity to pathogens. *Pharmacol. Ther.* **185**, 135–146.
- Martínez-Sernández, V., Mezo, M., González-Warleta, M., Perteguer, M.J., Gárate, T., Romarís, F., and Ubeira, F.M. (2017). Delineating distinct heme-scavenging and -binding functions of domains in MF6p/helminth defense molecule (HDM) proteins from parasitic flatworms. *J. Biol. Chem.* **292**, 8667–8682.
- Mbanefo, E.C., Le, L., Pennington, L.F., Odegaard, J.I., Jardetzky, T.S., Alouffi, A., Falcone, F.H., and Hsieh, M.H. (2018). Therapeutic exploitation of IPSE, a urogenital parasite-derived host modulatory protein, for chemotherapy-induced hemorrhagic cystitis. *FASEB J.* **32**, 4408–4419.
- McSorley, H.J., Blair, N.F., Smith, K.A., McKenzie, A.N.J., and Maizels, R.M. (2014). Blockade of IL-33 release and suppression of type 2 innate lymphoid cell responses by helminth secreted products in airway allergy. *Mucosal Immunol.* **7**, 1068–1078.
- Mebius, M.M., van Genderen, P.J., Urbanus, R.T., Tielens, A.G., de Groot, P.G., and van Hellemond, J.J. (2013). Interference with the host haemostatic system by schistosomes. *PLoS Pathog.* **9**, e1003781.
- Mebius, M.M., Op Heij, J.M.J., Tielens, A.G.M., de Groot, P.G., Urbanus, R.T., and van Hellemond, J.J. (2018). Fibrinogen and fibrin are novel substrates for *Fasciola hepatica* cathepsin L peptidases. *Mol. Biochem. Parasitol.* **221**, 10–13.
- Meenan, N.A., Ball, G., Bromek, K., Uhrin, D., Cooper, A., Kennedy, M.W., and Smith, B.O. (2011). Solution structure of a repeated unit of the ABA-1 nematode polyprotein allergen of *Ascaris* reveals a novel fold and two discrete lipid-binding sites. *PLoS Negl. Trop. Dis.* **5**, e1040.
- Meevissen, M.H., Yazdanbakhsh, M., and Hokke, C.H. (2012). *Schistosoma mansoni* egg glycoproteins and C-type lectins of host immune cells: molecular partners that shape immune responses. *Exp. Parasitol.* **132**, 14–21.
- Moyle, M., Foster, D.L., McGrath, D.E., Brown, S.M., Laroche, Y., De Meutter, J., Stanssens, P., Bogowitz, C.A., Fried, V.A., Ely, J.A., et al. (1994). A hookworm glycoprotein that inhibits neutrophil function is a ligand of the integrin CD11b/CD18. *J. Biol. Chem.* **269**, 10008–10015.
- Navarro, S., Pickering, D.A., Ferreira, I.B., Jones, L., Ryan, S., Troy, S., Leech, A., Hotez, P.J., Zhan, B., Laha, T., et al. (2016). Hookworm recombinant protein promotes regulatory T cell responses that suppress experimental asthma. *Sci. Transl. Med.* **8**, 362ra143.
- Nono, J.K., Lutz, M.B., and Brehm, K. (2014). EmTIP, a T-Cell immunomodulatory protein secreted by the tapeworm *Echinococcus multilocularis* is important for early metacystode development. *PLoS Negl. Trop. Dis.* **8**, e2632.
- Norton, R.S., Pennington, M.W., and Wulff, H. (2004). Potassium channel blockade by the sea anemone toxin ShK for the treatment of multiple sclerosis and other autoimmune diseases. *Curr. Med. Chem.* **11**, 3041–3052.
- Noya, V., Brossard, N., Berasaín, P., Rodríguez, E., Chiale, C., Mazal, D., Carmona, C., and Freire, T. (2016). A mucin-like peptide from *Fasciola hepatica* induces parasite-specific Th1-type cell immunity. *Parasitol. Res.* **115**, 1053–1063.
- Noya, V., Brossard, N., Rodríguez, E., Dergan-Dylon, L.S., Carmona, C., Rabinovich, G.A., and Freire, T. (2017). A mucin-like peptide from *Fasciola hepatica* instructs dendritic cells with parasite specific Th1-polarizing activity. *Sci. Rep.* **7**, 40615.
- Okano, M., Satoskar, A.R., Nishizaki, K., Abe, M., and Harn, D.A., Jr. (1999). Induction of Th2 responses and IgE is largely due to carbohydrates functioning as adjuvants on *Schistosoma mansoni* egg antigens. *J. Immunol.* **163**, 6712–6717.
- Okano, M., Satoskar, A.R., Nishizaki, K., and Harn, D.A., Jr. (2001). Lacto-N-fucopentaose III found on *Schistosoma mansoni* egg antigens functions as adjuvant for proteins by inducing Th2-type response. *J. Immunol.* **167**, 442–450.
- Osborn, M., Soares, D.C., Vacca, F., Cohen, E.S., Scott, I.C., Gregory, W.F., Smyth, D.J., Toivakka, M., Kemter, A.M., le Bihan, T., et al. (2017). HpARI protein secreted by a helminth parasite suppresses interleukin-33. *Immunity* **47**, 739–751.e5.
- Park, S.K., Cho, M.K., Park, H.K., Lee, K.H., Lee, S.J., Choi, S.H., Ock, M.S., Jeong, H.J., Lee, M.H., and Yu, H.S. (2009). Macrophage migration inhibitory factor homologs of *anisakis simplex* suppress Th2 response in allergic airway inflammation model via CD4+CD25+Foxp3+ T cell recruitment. *J. Immunol.* **182**, 6907–6914.
- Pennington, L.F., Alouffi, A., Mbanefo, E.C., Ray, D., Heery, D.M., Jardetzky, T.S., Hsieh, M.H., and Falcone, F.H. (2017). H-IPSE is a pathogen-secreted host nucleus infiltrating protein (infiltrin) expressed exclusively by the *Schistosoma haematobium* egg stage. *Infect. Immun., IAI*.00301-17.
- Pfaff, A.W., Schulz-Key, H., Soboslay, P.T., Taylor, D.W., MacLennan, K., and Hoffmann, W.H. (2002). *Litomosoides sigmodontis* cystatin acts as an immunomodulator during experimental filariasis. *Int. J. Parasitol.* **32**, 171–178.
- Pineda, M.A., Lumb, F., Harnett, M.M., and Harnett, W. (2014). ES-62, a therapeutic anti-inflammatory agent evolved by the filarial nematode *Acanthocheilonema viteae*. *Mol. Biochem. Parasitol.* **194**, 1–8.
- Prieto-Lafuente, L., Gregory, W.F., Allen, J.E., and Maizels, R.M. (2009). MIF homologues from a filarial nematode parasite synergize with IL-4 to induce alternative activation of host macrophages. *J. Leukoc. Biol.* **85**, 844–854.
- Ramos-Benítez, M.J., Ruiz-Jiménez, C., Aguayo, V., and Espino, A.M. (2017). Recombinant *Fasciola hepatica* fatty acid binding protein suppresses toll-like receptor stimulation in response to multiple bacterial ligands. *Sci. Rep.* **7**, 5455.
- Ranasinghe, S.L., and McManus, D.P. (2017). Protease inhibitors of parasitic flukes: emerging roles in parasite survival and immune defence. *Trends Parasitol.* **33**, 400–413.
- Ranasinghe, S.L., Fischer, K., Gobert, G.N., and McManus, D.P. (2015). Functional expression of a novel Kunitz type protease inhibitor from the human blood fluke *Schistosoma mansoni*. *Parasit. Vectors* **8**, 408.
- Retra, K., deWalick, S., Schmitz, M., Yazdanbakhsh, M., Tielens, A.G., Brouwers, J.F., and van Hellemond, J.J. (2015). The tegumental surface membranes of *Schistosoma mansoni* are enriched in parasite-specific phospholipid species. *Int. J. Parasitol.* **45**, 629–636.
- Ritter, M., Gross, O., Kays, S., Ruland, J., Nimmerjahn, F., Saijo, S., Tschopp, J., Layland, L.E., and Prazeres da Costa, C. (2010). *Schistosoma mansoni* triggers Dectin-2, which activates the Nlrp3 inflammasome and alters adaptive immune responses. *Proc. Natl. Acad. Sci. USA* **107**, 20459–20464.

- Robb, C.T., McSorley, H.J., Lee, J., Aoki, T., Yu, C., Crittenden, S., Astier, A., Felton, J.M., Parkinson, N., Ayele, A., et al. (2018). Prostaglandin E₂ stimulates adaptive IL-22 production and promotes allergic contact dermatitis. *J. Allergy Clin. Immunol.* *141*, 152–162.
- Robinson, M.W., Donnelly, S., Hutchinson, A.T., To, J., Taylor, N.L., Norton, R.S., Perugini, M.A., and Dalton, J.P. (2011). A family of helminth molecules that modulate innate cell responses via molecular mimicry of host antimicrobial peptides. *PLoS Pathog.* *7*, e1002042.
- Robinson, M.W., Alvarado, R., To, J., Hutchinson, A.T., Dowdell, S.N., Lund, M., Turnbull, L., Whitchurch, C.B., O'Brien, B.A., Dalton, J.P., and Donnelly, S. (2012). A helminth cathelicidin-like protein suppresses antigen processing and presentation in macrophages via inhibition of lysosomal vATPase. *FASEB J.* *26*, 4614–4627.
- Rodríguez, E., Carasi, P., Frigerio, S., da Costa, V., van Vliet, S., Noya, V., Brosard, N., van Kooyk, Y., García-Vallejo, J.J., and Freire, T. (2017). *Fasciola hepatica* immune regulates CD11c⁺ cells by interacting with the macrophage Gal/GalNAc lectin. *Front. Immunol.* *8*, 264.
- Rzepecka, J., Coates, M.L., Saggari, M., Al-Riyami, L., Coltherd, J., Tay, H.K., Huggan, J.K., Janicova, L., Khalaf, A.I., Siebecke, I., et al. (2014). Small molecule analogues of the immunomodulatory parasitic helminth product ES-62 have anti-allergy properties. *Int. J. Parasitol.* *44*, 669–674.
- Schölmerich, J., Fellermann, K., Seibold, F.W., Rogler, G., Langhorst, J., Howaldt, S., Novacek, G., Petersen, A.M., Bachmann, O., Matthes, H., et al.; International TRUST-2 Study Group (2017). A randomised, double-blind, placebo-controlled trial of *Trichuris suis* ova in active Crohn's disease. *J. Crohn's Colitis* *11*, 390–399.
- Schönemeyer, A., Lucius, R., Sonnenburg, B., Brattig, N., Sabat, R., Schilling, K., Bradley, J., and Hartmann, S. (2001). Modulation of human T cell responses and macrophage functions by onchocystatin, a secreted protein of the filarial nematode *Onchocerca volvulus*. *J. Immunol.* *167*, 3207–3215.
- Schramm, G., Mohrs, K., Wodrich, M., Doenhoff, M.J., Pearce, E.J., Haas, H., and Mohrs, M. (2007). Cutting edge: IPSE/alpha-1, a glycoprotein from *Schistosoma mansoni* eggs, induces IgE-dependent, antigen-independent IL-4 production by murine basophils in vivo. *J. Immunol.* *178*, 6023–6027.
- Sikri, N., and Bardia, A. (2007). A history of streptokinase use in acute myocardial infarction. *Tex. Heart Inst. J.* *34*, 318–327.
- Smith, P., Fallon, R.E., Mangan, N.E., Walsh, C.M., Saraiva, M., Sayers, J.R., McKenzie, A.N.J., Alcamí, A., and Fallon, P.G. (2005). *Schistosoma mansoni* secretes a chemokine binding protein with antiinflammatory activity. *J. Exp. Med.* *202*, 1319–1325.
- Smout, M.J., Laha, T., Mulvenna, J., Srija, B., Suttiprapa, S., Jones, A., Brindley, P.J., and Loukas, A. (2009). A granulins-like growth factor secreted by the carcinogenic liver fluke, *Opisthorchis viverrini*, promotes proliferation of host cells. *PLoS Pathog.* *5*, e1000611.
- Smout, M.J., Sotillo, J., Laha, T., Papatpremsiri, A., Rinaldi, G., Pimenta, R.N., Chan, L.Y., Johnson, M.S., Turnbull, L., Whitchurch, C.B., et al. (2015). Carcinogenic parasite secretes growth factor that accelerates wound healing and potentially promotes neoplasia. *PLoS Pathog.* *11*, e1005209.
- Smyth, D.J., Harcus, Y., White, M.P.J., Gregory, W.F., Nahler, J., Stephens, I., Toke-Bjølgerud, E., Hewitson, J.P., Ivens, A., McSorley, H.J., and Maizels, R.M. (2018). TGF-β mimic proteins form an extending gene family in the murine parasite *Heligmosomoides polygyrus*. *Int. J. Parasitol.* *48*, 379–385.
- Steinfelder, S., Andersen, J.F., Cannons, J.L., Feng, C.G., Joshi, M., Dwyer, D., Caspar, P., Schwartzberg, P.L., Sher, A., and Jankovic, D. (2009). The major component in schistosoma eggs responsible for conditioning dendritic cells for Th2 polarization is a T2 ribonuclease (omega-1). *J. Exp. Med.* *206*, 1681–1690.
- Suckling, C.J., Mukherjee, S., Khalaf, A.I., Narayan, A., Scott, F.J., Khare, S., Dhakshinamoorthy, S., Harnett, M.M., and Harnett, W. (2018). Synthetic analogues of the parasitic worm product ES-62 reduce disease development in in vivo models of lung fibrosis. *Acta Trop.* *185*, 212–218.
- Sulaiman, A.A., Zolnierczyk, K., Japa, O., Owen, J.P., Maddison, B.C., Emes, R.D., Hodgkinson, J.E., Gough, K.C., and Flynn, R.J. (2016). A trematode parasite derived growth factor binds and exerts influences on host immune functions via host cytokine receptor complexes. *PLoS Pathog.* *12*, e1005991.
- Sun, Y., Liu, G., Li, Z., Chen, Y., Liu, Y., Liu, B., and Su, Z. (2013). Modulation of dendritic cell function and immune response by cysteine protease inhibitor from murine nematode parasite *Heligmosomoides polygyrus*. *Immunology* *138*, 370–381.
- Sun, X., Yang, F., Shen, J., Liu, Z., Liang, J., Zheng, H., Fung, M., and Wu, Z. (2016). Recombinant Sj16 from *Schistosoma japonicum* contains a functional N-terminal nuclear localization signal necessary for nuclear translocation in dendritic cells and interleukin-10 production. *Parasitol. Res.* *115*, 4559–4571.
- Tan, T.H., Edgerton, S.A., Kumari, R., McAlister, M.S., Roe, S.M., Nagl, S., Pearl, L.H., Selkirk, M.E., Bianco, A.E., Totty, N.F., et al. (2001). Macrophage migration inhibitory factor of the parasitic nematode *Trichinella spiralis*. *Biochem. J.* *357*, 373–383.
- Teclé, T., Tripathi, S., and Hartshorn, K.L. (2010). Review: Defensins and cathelicidins in lung immunity. *Innate Immun.* *16*, 151–159.
- Terrazas, L.I., Walsh, K.L., Piskorska, D., McGuire, E., and Harn, D.A., Jr. (2001). The schistosome oligosaccharide lacto-N-neotetraose expands Gr1⁺ cells that secrete anti-inflammatory cytokines and inhibit proliferation of naive CD4⁺ cells: a potential mechanism for immune polarization in helminth infections. *J. Immunol.* *167*, 5294–5303.
- Thomas, P.G., Carter, M.R., Da'dara, A.A., DeSimone, T.M., and Harn, D.A. (2005). A helminth glycan induces APC maturation via alternative NF-κB activation independent of IκBα degradation. *J. Immunol.* *175*, 2082–2090.
- Tielens, A.G.M., van Grinsven, K.W.A., Henze, K., van Hellemond, J.J., and Martin, W. (2010). Acetate formation in the energy metabolism of parasitic helminths and protists. *Int. J. Parasitol.* *40*, 387–397.
- Tribolet, L., Cantacessi, C., Pickering, D.A., Navarro, S., Doolan, D.L., Trieu, A., Fei, H., Chao, Y., Hofmann, A., Gasser, R.B., et al. (2015). Probing of a human proteome microarray with a recombinant pathogen protein reveals a novel mechanism by which hookworms suppress B-cell receptor signaling. *J. Infect. Dis.* *211*, 416–425.
- van den Berg, T.K., Honing, H., Franke, N., van Remoortere, A., Schiphorst, W.E.C.M., Liu, F.-T., Deelder, A.M., Cummings, R.D., Hokke, C.H., and van Die, I. (2004). LacdiNAc-glycans constitute a parasite pattern for galectin-3-mediated immune recognition. *J. Immunol.* *173*, 1902–1907.
- van der Kleij, D., Latz, E., Brouwers, J.F., Kruize, Y.C., Schmitz, M., Kurt-Jones, E.A., Espevik, T., de Jong, E.C., Kapsenberg, M.L., Golenbock, D.T., et al. (2002). A novel host-parasite lipid cross-talk. *Schistosoma* lysophosphatidylserine activates toll-like receptor 2 and affects immune polarization. *J. Biol. Chem.* *277*, 48122–48129.
- van Die, I., van Vliet, S.J., Nyame, A.K., Cummings, R.D., Bank, C.M., Appel-melk, B., Geijtenbeek, T.B., and van Kooyk, Y. (2003). The dendritic cell-specific C-type lectin DC-SIGN is a receptor for *Schistosoma mansoni* egg antigens and recognizes the glycan antigen Lewis x. *Glycobiology* *13*, 471–478.
- Van Dyken, S.J., Mohapatra, A., Nussbaum, J.C., Molofsky, A.B., Thornton, E.E., Ziegler, S.F., McKenzie, A.N.J., Krummel, M.F., Liang, H.E., and Locksley, R.M. (2014). Chitin activates parallel immune modules that direct distinct inflammatory responses via innate lymphoid type 2 and γδ T cells. *Immunity* *40*, 414–424.
- Vaux, R., Schnoeller, C., Berkachy, R., Roberts, L.B., Hagen, J., Gounaris, K., and Selkirk, M.E. (2016). Modulation of the immune response by nematode secreted acetylcholinesterase revealed by heterologous expression in *Trypanosoma muscili*. *PLoS Pathog.* *12*, e1005998.
- Velupillai, P., and Harn, D.A. (1994). Oligosaccharide-specific induction of interleukin 10 production by B220⁺ cells from schistosoma-infected mice: a mechanism for regulation of CD4⁺ T-cell subsets. *Proc. Natl. Acad. Sci. USA* *91*, 18–22.
- Vermeire, J.J., Cho, Y., Lolis, E., Bucala, R., and Cappello, M. (2008). Orthologs of macrophage migration inhibitory factor from parasitic nematodes. *Trends Parasitol.* *24*, 355–363.
- Wang, L., Li, Z., Shen, J., Liu, Z., Liang, J., Wu, X., Sun, X., and Wu, Z. (2015). Exosome-like vesicles derived by *Schistosoma japonicum* adult worms mediates M1 type immune- activity of macrophage. *Parasitol. Res.* *114*, 1865–1873.
- Wang, S., Xie, Y., Yang, X., Wang, X., Yan, K., Zhong, Z., Wang, X., Xu, Y., Zhang, Y., Liu, F., and Shen, J. (2016). Therapeutic potential of recombinant

- cystatin from *Schistosoma japonicum* in TNBS-induced experimental colitis of mice. *Parasit. Vectors* 9, 6.
- Wang, Q., Da'dara, A.A., and Skelly, P.J. (2017). The human blood parasite *Schistosoma mansoni* expresses extracellular tegumental calpains that cleave the blood clotting protein fibrinogen. *Sci. Rep.* 7, 12912.
- Wang, Q., Da'dara, A.A., and Skelly, P.J. (2018). The blood fluke *Schistosoma mansoni* cleaves the coagulation protein high molecular weight kininogen (HK) but does not generate the vasodilator bradykinin. *Parasit. Vectors* 11, 182.
- Wangchuk, P., Constantinoiu, C., Kouremenos, K.A., Becker, L., Jones, L., Shepherd, C., Buitrago, G., Giacomini, P., Daly, N., McConville, M.J., et al. (2018). Hookworm-derived small molecule extracts suppress pathology in a mouse model of colitis and inhibit secretion of key inflammatory cytokines in primary human leukocytes. *bioRxiv* <https://doi.org/10.1101/316885>.
- Weinstock, J.V., and Elliott, D.E. (2013). Translatability of helminth therapy in inflammatory bowel diseases. *Int. J. Parasitol.* 43, 245–251.
- Wilbers, R.H., Westerhof, L.B., van Noort, K., Obieglo, K., Driessen, N.N., Everts, B., Gringhuis, S.I., Schramm, G., Goverse, A., Smant, G., et al. (2017). Production and glyco-engineering of immunomodulatory helminth glycoproteins in plants. *Sci. Rep.* 7, 45910.
- Wilbers, R.H.P., Schneiter, R., Holterman, M.H.M., Drurey, C., Smant, G., Asojo, O.A., Maizels, R.M., and Lozano-Torres, J.L. (2018). Secreted venom allergen-like proteins of helminths: Conserved modulators of host responses in animals and plants. *PLoS Pathog.* 14, e1007300.
- Wiria, A.E., Sartono, E., Supali, T., and Yazdanbakhsh, M. (2014). Helminth infections, type-2 immune response, and metabolic syndrome. *PLoS Pathog.* 10, e1004140.
- Wolfs, I.M., Stöger, J.L., Goossens, P., Pöttgens, C., Gijbels, M.J., Wijnands, E., van der Vorst, E.P., van Gorp, P., Beckers, L., Engel, D., et al. (2014). Reprogramming macrophages to an anti-inflammatory phenotype by helminth antigens reduces murine atherosclerosis. *FASEB J.* 28, 288–299.
- Zaccone, P., Burton, O.T., Gibbs, S.E., Miller, N., Jones, F.M., Schramm, G., Haas, H., Doenhoff, M.J., Dunne, D.W., and Cooke, A. (2011). The *S. mansoni* glycoprotein ω -1 induces Foxp3 expression in NOD mouse CD4⁺ T cells. *Eur. J. Immunol.* 41, 2709–2718.
- Zaiss, M.M., Maslowski, K.M., Mosconi, I., Guenat, N., Marsland, B.J., and Harris, N.L. (2013). IL-1 β suppresses innate IL-25 and IL-33 production and maintains helminth chronicity. *PLoS Pathog.* 9, e1003531.
- Zaiss, M.M., Rapin, A., Lebon, L., Dubey, L.K., Mosconi, I., Sarter, K., Piersigilli, A., Menin, L., Walker, A.W., Rougemont, J., et al. (2015). The intestinal microbiota contributes to the ability of helminths to modulate allergic inflammation. *Immunity* 43, 998–1010.
- Zang, X., Taylor, P., Wang, J.M., Meyer, D.J., Scott, A.L., Walkinshaw, M.D., and Maizels, R.M. (2002). Homologues of human macrophage migration inhibitory factor from a parasitic nematode. Gene cloning, protein activity, and crystal structure. *J. Biol. Chem.* 277, 44261–44267.
- Zarowiecki, M., and Berriman, M. (2015). What helminth genomes have taught us about parasite evolution. *Parasitology* 142 (Suppl 1), S85–S97.
- Zavala-Góngora, R., Kroner, A., Bernthaler, P., Knaus, P., and Brehm, K. (2006). A member of the transforming growth factor- β receptor family from *Echinococcus multilocularis* is activated by human bone morphogenetic protein 2. *Mol. Biochem. Parasitol.* 146, 265–271.
- Zhou, M.Y., Lo, S.K., Bergenfeldt, M., Tiruppathi, C., Jaffe, A., Xu, N., and Malik, A.B. (1998). In vivo expression of neutrophil inhibitory factor via gene transfer prevents lipopolysaccharide-induced lung neutrophil infiltration and injury by a beta2 integrin-dependent mechanism. *J. Clin. Invest.* 101, 2427–2437.
- Ziegler, T., Rausch, S., Steinfeldt, S., Klotz, C., Hepworth, M.R., Kühl, A.A., Burda, P.C., Lucius, R., and Hartmann, S. (2015). A novel regulatory macrophage induced by a helminth molecule instructs IL-10 in CD4⁺ T cells and protects against mucosal inflammation. *J. Immunol.* 194, 1555–1564.