



## Original Article

# Effects of curcumin and metformin on oxidative stress and apoptosis in heart tissue of type 1 diabetic rats

Atefeh Naghdi<sup>1</sup>, Mohammad Taghi Goodarzi<sup>1,2</sup>, Jamshid Karimi<sup>1</sup>, Mohammad Hashemnia<sup>3</sup>, Iraj Khodadadi<sup>1\*</sup>

<sup>1</sup>Department of Clinical Biochemistry, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Iran

<sup>2</sup>Department of Biochemistry, Shahrood Branch, Islamic Azad University, Shahrood, Iran

<sup>3</sup>Department of Pathobiology, Faculty of Veterinary Medicine, Razi University, Kermanshah, Iran

### Article info

#### Article History:

Received: 23 January 2022

Accepted: 01 June 2022

published: 28 June 2022

#### Keywords:

Apoptosis

Curcumin

Diabetes Mellitus

Metformin

Oxidative Stress

### Abstract

**Introduction:** Hyperglycemia enhances oxidative stress and apoptosis and induces damages in heart tissue. Based on antioxidant properties of curcumin and metformin, we hypothesized that these agents may exhibit cardioprotective effects by attenuating oxidative stress and modulating expression of the genes involved in apoptosis in type-1 diabetes.

**Methods:** Thirty-six male rats were randomly divided into six groups; (N): control; (D): streptozotocin-induced diabetic rats; (D+Cur50) and (D+Cur150): diabetic rats treated with 50 and 150 milligram of curcumin per kilogram of body weight (mg/kg.bw), respectively; (D+Met300) and (D+Met500): diabetic rats received 300 and 500 mg/kg.bw of metformin, respectively. Heart tissues were dissected and gene expression levels of Bax, Bcl-2, and caspase-3 were analyzed. Total anti-oxidant capacity (TAC), total oxidant status (TOS), and malondialdehyde (MDA) level, and activities of catalase (CAT), superoxide dismutase (SOD) and glutathione peroxidase (GPx) were measured.

**Results:** Enhancement in TOS, OSI, and MDA levels as well as increased in the activity of CAT and reduction in SOD and GPx activities were observed in diabetic group (D) compared with control rats. Treatment of diabetic animals with either curcumin or metformin normalized TOS, OSI, and MDA levels and restored CAT, SOD, and GPx activities. Diabetes caused extensive damages in heart tissue of rats (group D) and increased expression of caspase-3 and Bax genes and enhanced ratio of Bax/Bcl-2 expression compared with controls. Treatment with curcumin or metformin mitigated histopathological changes and dampened apoptosis by normalizing Bax and caspase-3 expression.

**Conclusion:** Curcumin and metformin modulated diabetes-induced cardiac damage probably by reducing oxidative stress.

### Introduction

Diabetes mellitus (DM), as one of the most common metabolic disorders, is characterized by hyperglycemia and comes from defects in the insulin secretion by the pancreas and insulin resistance in the end-organ tissues or both.<sup>1</sup> The prevalence of DM is growing in the world and it is estimated to reach up to 693 million by 2045.<sup>2</sup> Chronic hyperglycemia results in damage and disrupted functions of multiple organs especially, eyes, kidneys, nervous and cardiovascular systems. Cardiovascular disorders (CVD) remain the most important causes of death and disability in DM-affected patients.<sup>3</sup> Despite several lines of evidence suggesting CVD risks among diabetic patients, the pathophysiologic mechanisms which link these two diseases are yet to be clearly understood.<sup>2</sup>

DM-associated CVD includes a range of clinical

conditions, such as diabetic cardiomyopathy, coronary heart disease and congestive heart failure, where oxidative stress seems to be involved in pathogenesis of all these conditions.<sup>4</sup> Hyperglycemia-induced intracellular changes finally lead to generate reactive oxygen species (ROS) which promote apoptosis process.<sup>5</sup> Apoptosis is considered as a major player of CVD because it leads to loss of cardiomyocytes. Cardiac regenerative capacity is limited in adult mammalian, thus attempts to prevent losing of cardiomyocytes or stimulating formation of new muscles in damaged heart have received much attention.<sup>6</sup>

Besides common therapeutic strategies, administration of natural compounds is increasing as major medicinal alternatives, because of their safety, success in the clinical conditions and less side effects. Curcumin is a lipophilic polyphenol compound which is extracted as a major active



\*Corresponding Author: Iraj Khodadadi, Email: khodadadi@umsha.ac.ir

© 2022 The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

compound from dried rhizomes of turmeric (*Curcuma longa*).<sup>7, 8</sup> Available evidence suggests that curcumin protects organs against oxidative stress. In addition to the molecular structure-associated antioxidant nature of the curcumin, this compound increases the expression of antioxidant enzymes such as superoxide dismutase (SOD), glutathione peroxidase (GPx) and catalase (CAT) through upregulation of nuclear factor erythroid 2-related factor 2 (Nrf-2).<sup>9, 10</sup> Available evidence also suggests that metformin has antioxidant, anti-inflammatory, and anti-apoptotic effects and might be therapeutically beneficial in improving diseases with inflammatory, apoptotic, and oxidative stress basis in their pathogenesis.<sup>11</sup>

Recently, protective effect of curcumin on the heart in diabetic patients has attracted considerable attention. Curcumin beneficially modulates oxidative stress and improves metabolic abnormalities in cardiomyocytes. The therapeutic potential of curcumin in the treatment of diabetic cardiomyopathy is probably mediated through several mechanisms such as attenuation of oxidative stress and modulation of Sirt1-Foxo1 and PI3K-Akt pathways<sup>12</sup> and by regulation of calmodulin-dependent protein kinase II (CaMKII) and peroxisome proliferator-activated receptor gamma (PPAR- $\gamma$ ) expression.<sup>8</sup>

Since diabetes is associated with apoptosis in cardiomyocytes<sup>12</sup> and the role of oxidative stress in evoking apoptosis has already been reported,<sup>3</sup> we hypothesized that curcumin and metformin may ameliorate apoptotic process in cardiomyocytes of type-1 diabetic animal models due to their antioxidant properties. In addition, due to the involvement of caspase-3 enzyme in apoptotic process we postulated that curcumin and/or metformin may modulate cardiomyocyte apoptosis through regulating of caspase-3 gene expression. Therefore, in the present study, beneficial effects of curcumin and metformin were investigated on the oxide-redox status of cardiac tissue by determining total antioxidant capacity (TAC), total oxidant status (TOS), and lipid peroxidation marker malondialdehyde (MDA). In addition, the activities of enzymes involved in cellular antioxidant defense system including CAT, SOD, and GPx were measured and the expression levels of apoptosis-related genes such as Bax, Bcl-2, and caspase-3 in the heart tissue of streptozotocin-induced type 1 diabetic rats were determined.

## Material and Methods

### Experimental design

In this case-control study thirty-six 8 weeks old male Wistar rats (150-200 g) were procured from Animal House of Hamadan University of Medical Sciences (Hamadan-Iran). Animals were housed under standard laboratory conditions in a temperature-controlled room under natural light/dark cycle and unlimited access to water and standard chow diet was provided for the rats. All experiments were performed in accordance with the National Institutes of Health Guide for the Care and

Use of Laboratory Animals (NIH Publication No.80-23, revised 1996) and the study was approved by the Research Ethics Committee of Hamadan University of Medical Sciences (IR.UMSHA.REC.1395.563).

After acclimatization of rats with regular chow diet for one week, all animals were randomly divided into six groups (n=6) as following: (N): normal control group; (D): diabetic untreated group; (D+Cur50): diabetic group treated with 50 milligram of curcumin per kilogram of body weight (mg/kg.bw); (D+Cur150): diabetic group treated with 150 mg/kg.bw of curcumin; (D+Met300): diabetic group treated with 300 mg/kg.bw of metformin; and (D+Met500): diabetic group treated with 500 mg/kg.bw of metformin. Metformin was given to the rats at the concentrations of 300 and 500 mg/kg.bw while rats received curcumin at the 50 and 150 mg/kg.bw concentrations based on previous studies.<sup>13-15</sup> Metformin and curcumin were purchased from Sigma-Aldrich (Sigma-Aldrich, Ontario-Canada), dissolved and suspended in deionized water, and were orally administered each day for six weeks (42 days).

### Induction of diabetes

After acclimatization of rats with regular chow diet for one week and, streptozotocin (STZ) (Santa Cruz Biotechnology, Heidelberg-Germany) was dissolved in sodium citrate buffer (0.1 M, pH: 4.5) and a dose of 55 mg per kg of body weight was intraperitoneally injected to the rats to induce type 1 diabetes mellitus (T1DM) in diabetic groups (D, D+Cur50, D+Cur150, D+Met300, and D+Met500). Seventy-two hours later, the level of fasting blood sugar (FBS) was measured using a glucometer to confirm induction of diabetes. FBS level higher than 250 mg/dl was considered as diagnostic off point of diabetes.<sup>16</sup> After induction of diabetes (day zero; D0) treatment of the rats with curcumin or metformin was started and continued for 42 days.<sup>14</sup> Twenty-four hours after the last treatment, the rats were anesthetized with ketamine-xylazine (50mg/kg ketamine and 12mg/kg xylazine) and sacrificed. Blood samples were collected from jugular vein and sera were separated and stored at -20°C until analysis of biochemical parameters. Heart tissue samples were dissected and used to determine oxide-redox status and to evaluate gene expression levels. Portions of tissue samples were also fixed in 10% neutral-buffered formalin for histopathological evaluation.

### Analysis of serum biochemical parameters

Serum total creatine kinase (CK), creatine kinase MB isoform (CK-MB), lactate dehydrogenase (LDH), and aspartate transaminase (AST) activities were measured using commercially available kits (Parsazmoon, Iran) and a BS-480 Mindray Chemistry Analyzer (Shenzhen Mindray Bio-Medical Electronics Co., Guangdong-China).

### Determination of oxide-redox status in heart samples

About 100 mg of heart tissues were rinsed with ice-cold

saline, crushed in liquid nitrogen, and homogenized by 1 ml of lysis buffer containing DTT 0.5 mM, HEPES 10 mM, KCl 10 mM, MgCl<sub>2</sub> 1.5 mM and Triton X100 2%. Homogenates were then centrifuged at 14,000×g for 20 min (4°C). The supernatants were aliquoted and total protein contents of samples were measured by Bradford assay using bovine serum albumin (BSA) as standard.<sup>17</sup>

Total antioxidant capacity (TAC) was measured by ferric reducing antioxidant power (FRAP) method.<sup>18</sup> The principle of FRAP method was based on the reduction of ferric tripyridyl triazine (Fe<sup>+3</sup>-TPTZ) to ferropyridyl triazine (Fe<sup>+2</sup>-TPTZ) by the antioxidant content of the sample. The absorbance of the Fe<sup>+2</sup>-TPTZ complex (blue color) was assayed at 593 nm and the TAC levels were calculated based on a standard curve (FeSO<sub>4</sub>.7H<sub>2</sub>O) and expressed as nmol/mg of protein.

Total oxidant status (TOS) assay was carried out by ferrous ion oxidation xylenol orange (Fox1) method.<sup>19</sup> The Fox1 method was based on the oxidation of Fe<sup>+2</sup> ions to Fe<sup>+3</sup> in the presence of sample's oxidizing agents under the acidic medium. The absorbance of Fe<sup>+3</sup>-xylenol orange color complex was read at 546 nm and TOS levels were calculated against H<sub>2</sub>O<sub>2</sub> as a standard and expressed as μmol/mg of protein.

Malondialdehyde (MDA), as a lipid peroxidation marker was determined by spectrofluorometric method.<sup>20</sup> Briefly, the reaction of thiobarbituric acid with MDA led to the formation of a fluorescence complex with excitation and emission wavelengths at 530 and 550 nm, respectively. Serial dilution of tetraethoxypropane was used as standard and MDA levels were reported as μmol/mg of protein.

#### **Determination of catalase activity in heart tissue**

Catalase (CAT) activity was determined using spectrophotometric Hadwan method<sup>21</sup> based on the decomposition of H<sub>2</sub>O<sub>2</sub> by the catalase activity of the samples. Briefly, to 100 μl heart tissue lysate, 1 ml of H<sub>2</sub>O<sub>2</sub> or deionized water was added (test and control tubes, respectively) while in standard tube 1 ml deionized water was mixed with 100 μl of H<sub>2</sub>O<sub>2</sub>. After incubation of tubes in 37°C for 3 min, enzyme reaction was stopped by 4 ml of ammonium molybdate and absorbances of the tubes were measured at 374 nm. Finally, CAT activity was calculated by the following formula where t was time, S<sup>o</sup>, S, and M were absorbances of standard, test, and control tubes respectively, Vt was total volume of reagents in test tube, and Vs was volume of supernatant.

$$CAT\ activity = \frac{2.303}{t} \times \left( \log \frac{S^o}{S-M} \right) \times \frac{Vt}{Vs}$$

#### **Determination of glutathione peroxidase activity in heart tissue**

Glutathione peroxidase (GPx) activity in heart tissue samples was determined using Paglia and Valentine method, as previously described.<sup>22</sup> Briefly, 100 μl NADPH

(8 mM), 100 μl reduced glutathione, 20 μl glutathione reductase (30 U/ml) and 20 μl sodium azide (0.12 M) were added to 50 μl of heart lysate. Next, potassium phosphate buffer (50 mM, pH=7.5) was added to the tubes and mixtures were incubated at 37°C for 30 minute. Then, 100 μl H<sub>2</sub>O<sub>2</sub> was added and the reductions in the absorbance at 340 nm were measured. Finally, GPx activities were calculated as μmol of NADPH used to reduce glutathione.

#### **Determination of superoxide dismutase activity in heart tissue**

Superoxide dismutase (SOD) activity was measured by the Marklund method<sup>23</sup> in which one unit of enzyme activity was equal to the quantity of enzyme required to inhibit 50% of pyrogallol oxidation rate. Briefly, 2 ml Tris-HCl (50 mM, pH=8.2) and 20 μl pyrogallol solution (10 mM, pH=7.4) were added to 30 μl heart lysate and absorbances of test and control tubes were measured at 420 nm. Finally, the results of enzyme activity were calculated based on the inhibition of pyrogallol autoxidation percent [(ΔA test/ΔA control) × 100]/50% and expressed as U/mg of protein.

#### **Determination the expression levels of apoptosis-related genes by RT-qPCR**

Gene expression levels of Bax, Bcl-2, and caspase-3 in heart tissue samples were determined by RT-qPCR on a Roche Light Cycler<sup>®</sup> 96 System (Roche Life Science, Sandhofer-Germany) and qPCR master mix (QuantiFast SYBR Green PCR Kit). Briefly, total RNA was isolated using RNX-plus reagent (Cinnagen, Iran), according to the manufacturer's protocol'. The quantity and quality of the isolated RNA were evaluated by NanoDrop<sup>™</sup> (Thermo Fisher Scientific-USA) and 1% agarose gel electrophoresis, respectively. cDNA was synthesized by reverse transcription of 500 ng of total RNA using PrimeScript first Standard cDNA Synthesis Kit (Takara Biotechnology, Japan). Gene-specific primers (Table 1) were designed using AlleleID<sup>®</sup>6 software (Premier Biosoft Corporation, USA). Online primer blast software (<https://www.ncbi.nlm.nih.gov/tools/primer-blast/>) was used to confirm specificities of designed primers to the corresponding genes. β-actin was considered as housekeeping gene and relative mRNA expression levels (fold change) were calculated by 2<sup>-ΔΔCt</sup> formula based on Livak method<sup>24</sup> compared with the expression of β-actin.

#### **Histopathological evaluation**

For histopathological studies, heart tissue samples were fixed in 10% neutral-buffered formalin, processed, and embedded in paraffin. Tissue samples were then sectioned into 5 μm thickness slides and stained with hematoxylin and eosin (H&E). Finally, the slides were examined by a pathologist using a routine light microscope and characteristics of cardiomyocyte structure including uniformity, shape and location of nuclei and intercalated

**Table 1.** Sequences and characteristics of primers used in the study.

Gene name	Accession number	Sequences	Tm°C	Product size (bp)
β-Actin	NM_031144.3	F: 5'-ATCAGCAAGCAGGAGTACGAT-3' R: 5'-AAAGGGTGTAAAACGCAGCTC-3'	56	94
Bcl-2	NM_016993.1	F: 5'-TGTGGATGACTGAGTACCTGAACC-3' R: 5'-CAGCCAGGAGAAATCAAACAGAGG-3'	52	122
Bax	NM_017059.2	F: 5'-GCAGAGGATGATTGCTGATGTGG-3' R: 5'-AGGAAGTCCAGTCCAGCCCAT-3'	52	204
Caspase-3	NM_012922.9	F: 5'-ACAACAACGAAACCTCCGTG-3' R: 5'-CATTGCGAGCTGACATTCCA-3'	56	192

Gene sequences were obtained from online Gene Bank database (<https://www.ncbi.nlm.nih.gov/genbank/>). Gene-specific primers were designed using AlleleID® software (Premier Biosoft Corporation, USA). Online primer blast software (<https://www.ncbi.nlm.nih.gov/tools/primer-blast/>) was used to confirm specificities of designed primers to the corresponding genes.

discs of plasma membrane were evaluated. Presence of hypertrophic myofibers, vacuolation and congestion of cytoplasm, and deformation of nuclei were also investigated.

### Statistical analysis

Statistical analysis was carried out using the SPSS 16 (SPSS Inc., Chicago-USA). One-way ANOVA followed by post hoc Tukey test was used for comparison between groups. Data were presented as mean ± SD and *P* values less than 0.05 (*P* < 0.05) was considered as statistical significance.

## Results

### Effects of curcumin and metformin on serum biochemical parameters

As shown in Table 2, fasting blood sugar was significantly higher in all treated or untreated diabetic rats compared with control rats at the end of study (FBS D42). Although treatment with 150 mg/kg.bw of curcumin slightly lowered FBS compared with untreated diabetic animals (group D), the results confirmed that neither of low or high doses of curcumin or metformin show notable hypoglycemic property in type 1 diabetes. AST significantly (*P* < 0.001) increased in serum to a mean of 296.66 U/L after induction of diabetes (D group) but treatment of the rats with curcumin (*P* = 0.006 and *P* = 0.009 for low and high doses) or metformin (*P* = 0.004 and *P* = 0.026 for low and

high doses) successfully lowered AST level compared with group D. At the end of experiment (day 42) no significant difference was observed in AST level between treated rats and control animals. Despite of significant increase in the level of LDH in diabetic group (D) compared with the control group, treatment with curcumin and metformin almost completely restored LDH level to normal level (*P* < 0.05). As shown in Table 2, the activity of serum creatine kinase (CK) and its CK-MB isoform did not significantly differ between control and diabetic groups (N and D, respectively), however treated rats (D + Cur50, D + Cur150, D + Met300, and D + Met500) showed almost 50% lower CK and CK-MB enzyme activities compared with both control and untreated diabetic rats in D group. Since the pattern of alteration in total CK activity was similar to that of CK-MB, the CK-MB/CK ratio was calculated for further evaluation of creatin kinase activity in diabetes. Diabetic rats (group D) showed higher CK-MB/CK ratio than control group ( $3.72 \pm 0.15$  vs  $3.27 \pm 0.12$ , *P* < 0.01) while treatment with both doses of curcumin in groups D + Cur50 and D + Cur150, resulted in significant reduction in CK-MB/CK ratio compared with untreated diabetic rats (*P* = 0.026 and *P* = 0.015, respectively) and completely restored the ratio level to that of control rats. Treatment with metformin showed relatively mild effect and despite to reducing CK-MB/CK ratio to the normal range, it did not significantly differ from that of diabetic

**Table 2.** Effects of curcumin or metformin on biochemical parameters

Groups	FBS D42 (mg/dl)	AST (U/L)	LDH (U/L)	CK (U/L)	CK-MB (U/L)	CK-MB/CK
N	92.33 ± 7.52	163.33 ± 15.05	1268.33 ± 177.81	1511.66 ± 686.18	493.33 ± 215.19	3.27 ± 0.12
DM	586.67 ± 16.14 <sup>a</sup>	296.66 ± 52.40 <sup>a</sup>	1966.66 ± 186.19 <sup>a</sup>	1218.33 ± 431.15	403.33 ± 138.94	3.72 ± 0.15 <sup>a</sup>
D + Cur50	570.50 ± 21.03 <sup>a</sup>	208 ± 27.75 <sup>b</sup>	1356.66 ± 520.48 <sup>b</sup>	581.66 ± 239.28 <sup>a</sup>	228.57 ± 74.41 <sup>a</sup>	3.3 ± 0.26 <sup>b</sup>
D + Cur150	541.60 ± 27.72 <sup>a,b</sup>	212 ± 25.88 <sup>b</sup>	1340 ± 380.98 <sup>b</sup>	650 ± 195.32 <sup>a</sup>	244 ± 72.66 <sup>a</sup>	3.24 ± 0.34 <sup>b</sup>
D + Met300	590.50 ± 12.72 <sup>a,c</sup>	212.85 ± 24.30 <sup>b</sup>	1362.85 ± 252.50 <sup>b</sup>	630 ± 265.20 <sup>a</sup>	228.57 ± 81.94 <sup>a</sup>	3.35 ± 0.22
D + Met500	588.60 ± 11.97 <sup>a,c</sup>	222 ± 59.33 <sup>b</sup>	1396 ± 309.56	610 ± 306.35 <sup>a</sup>	228 ± 102.05 <sup>a</sup>	3.43 ± 0.11

N: Normal group; DM: Diabetic group, D + Cur50: Diabetic group treated with 50 mg/Kg curcumin, D + Cur150: Diabetic group treated with 150 mg/Kg curcumin, D + Met300: Diabetic group treated with 300 mg/Kg metformin; D + Met500: Diabetic group treated with 500 mg/Kg metformin. FBS: Fasting blood sugar; AST: Aspartate transaminase; LDH: Lactate dehydrogenase; CK: Total creatine kinase; CK-MB: creatine kinase MB. Data is represented as mean ± SD and *P* < 0.05 is considered as statistically significance difference. In each column "a" represents significant difference with normal group (N), "b" indicates significant difference with diabetic group (DM), and "c" shows significance difference with D + Cur150.

rats (group D).

### **Antioxidant properties of curcumin and metformin in heart tissue**

Heart level of TAC showed no significant difference between studied groups (Table 3). On the other hand, TOS level was significantly higher in group D than the control group ( $P < 0.001$ ). Although TOS level in curcumin- and metformin-treated diabetic rats were relatively lower compared with untreated diabetic rats (group D), this difference was statistically significance only in D + Cur50 and D + Cur150 groups ( $P = 0.015$  and  $P = 0.007$ , respectively). As shown in Table 3, determination of oxidative stress index (OSI) as a ratio of TOS to TAC showed that induction of diabetes markedly increased OSI in D group ( $P = 0.013$ ) but treatment of the diabetic animals with both low and high doses of curcumin or metformin successfully lowered OSI compared with un-treated diabetic rats (group D) and normalized OSI to the level observed in control group. Likewise, lipid peroxidation marker (MDA) which was increased over 20% in D group ( $P = 0.005$ ) was markedly reduced in all D + Cur50, D + Cur150, D + Met300, and D + Met500 treated rats compared to group D ( $P = 0.008$ ,  $P = 0.007$ ,  $P = 0.001$ , and  $P = 0.029$  respectively) and completely restored to the normal level. Interestingly, no difference was observed in efficiency of curcumin and metformin in lowering lipid peroxidation level, as indicated in Table 3.

### **Curcumin or metformin favorably restored activities of antioxidant enzymes in heart**

Determination of activities of enzymes involved in antioxidant defense system showed that (Table 4) although induction of diabetes (D group) significantly increased CAT activity ( $P = 0.006$ ), treatment with curcumin and metformin prevented alteration in CAT activity where no significant difference was observed in the activity of CAT compared with normal group. Unlike catalase, both

SOD and GPx decreased by the induction of diabetes ( $P = 0.001$  for both enzymes), as observed in group D. However, reduction in the activity of both enzymes was completely prevented by the treatment of the rats using either low or high doses of curcumin and metformin in all treated animals (D + Cur50, D + Cur150, D + Met300, and D + Met500), as shown in Table 4. Interestingly, treatment with curcumin (both low and high doses) were more effective than metformin in preventing declines in SOD and GPx activities caused by the induction of diabetes in D + Cur50 and D + Cur150 groups.

### **Effects of curcumin or metformin on gene expression of Bax, Bcl-2, and Caspase-3 in heart**

The gene expression analysis revealed that expression of Bax was almost doubled ( $P < 0.001$ ) by the induction of diabetes in the rats (D group), as shown in Figure 1A, but treatment of animals with low or high doses of curcumin or metformin significantly suppressed evaluation of Bax gene expression in treated rats in D + Cur50, D + Cur150, D + Met300, and D + Met500 groups ( $P < 0.001$  for all). Unlike the increment in the expression of proapoptotic gene (Bax) by the induction of diabetes in D group, the expression of antiapoptotic Bcl-2 gene neither altered by inducing of diabetes (in group D) nor by the treatment of the diabetic rats with curcumin or metformin (Figure 1B). However, the ratio of Bax/Bcl-2 gene expressions which determines cell susceptibility to apoptosis showed higher susceptibility of heart tissues to apoptosis in diabetic rats from group D compared with control rats ( $P < 0.001$ ) whereas all curcumin- or metformin-treated animals (D + Cur50, D + Cur150, D + Met300, and D + Met500) showed almost statistically ( $P < 0.001$ ) similar lower apoptotic susceptibility (Figure 1C). Antiapoptotic properties of curcumin and metformin were further confirmed by determination of caspase-3 gene expression which showed over 50% elevation ( $P < 0.001$ ) in gene expression in diabetic animals (D group) while none of

**Table 3.** Effects of curcumin and metformin on antioxidant parameters


Groups	TAC (nmol/mg pr)	TOS (nmol/mg pro)	OSI (arbitrary)	MDA ( $\mu$ mol/mg pr)
N	0.52 $\pm$ 0.13	0.82 $\pm$ 0.05	1.62 $\pm$ 0.30	0.38 $\pm$ 0.02
DM	0.53 $\pm$ 0.07	1.03 $\pm$ 0.05 <sup>a</sup>	2.09 $\pm$ 0.31 <sup>a</sup>	0.46 $\pm$ 0.03 <sup>a</sup>
D+Cur50	0.53 $\pm$ 0.06	0.89 $\pm$ 0.10 <sup>b</sup>	1.67 $\pm$ 0.16 <sup>b</sup>	0.39 $\pm$ 0.03 <sup>b</sup>
D+Cur150	0.55 $\pm$ 0.12	0.87 $\pm$ 0.06 <sup>b</sup>	1.60 $\pm$ 0.19 <sup>b</sup>	0.38 $\pm$ 0.02 <sup>b</sup>
D+Met300	0.52 $\pm$ 0.07	0.93 $\pm$ 0.03	1.66 $\pm$ 0.08 <sup>b</sup>	0.38 $\pm$ 0.05 <sup>b</sup>
D+Met500	0.54 $\pm$ 0.05	0.92 $\pm$ 0.09	1.67 $\pm$ 0.17	0.39 $\pm$ 0.02 <sup>b</sup>

N: Normal group; DM: Diabetic group, D+Cur50: Diabetic group treated with 50 mg/Kg curcumin, D+Cur150: Diabetic group treated with 150 mg/Kg curcumin, D+Met300: Diabetic group treated with 300 mg/Kg metformin; D+Met500: Diabetic group treated with 500 mg/Kg metformin. TAC: Total antioxidant capacity; TOS: Total oxidant status; MDA: Malondialdehyde; OSI: oxidative stress index. Data is represented as mean  $\pm$  SD and  $P < 0.05$  is considered as statistically significance difference. In each column "a" represents significant difference with normal group (N) and "b" indicates significant difference with diabetic group (DM).

**Table 4.** Effects of curcumin and metformin on the activity of enzymes involved in antioxidant system

Groups	CAT (U/mg pr)	SOD (U/mg pr)	GPx (U/mg pr)
N	19.34 $\pm$ 0.94	37.64 $\pm$ 0.82	26.19 $\pm$ 1.00
DM	22.75 $\pm$ 1.72 <sup>a</sup>	35.74 $\pm$ 0.90 <sup>a</sup>	24.66 $\pm$ 0.49 <sup>a</sup>
D+Cur50	21.12 $\pm$ 1.42	37.55 $\pm$ 0.75 <sup>b</sup>	25.78 $\pm$ 0.15 <sup>b</sup>
D+Cur150	21.57 $\pm$ 1.31	37.52 $\pm$ 0.78 <sup>b</sup>	25.99 $\pm$ 0.10 <sup>b</sup>
D+Met300	20.78 $\pm$ 1.03	36.68 $\pm$ 0.31	25.35 $\pm$ 0.42
D+Met500	21.25 $\pm$ 1.96	36.92 $\pm$ 0.26	25.35 $\pm$ 0.58

N: Normal group; DM: Diabetic group, D+Cur50: Diabetic group treated with 50 mg/Kg curcumin, D+Cur150: Diabetic group treated with 150 mg/Kg curcumin, D+Met300: Diabetic group treated with 300 mg/Kg metformin; D+Met500: Diabetic group treated with 500 mg/Kg metformin. CAT: Catalase; SOD: Superoxide dismutase; GPx: Glutathione peroxidases. Data is represented as mean  $\pm$  SD and  $P < 0.05$  is considered as statistically significance difference. In each column "a" represents significant difference with normal group (N) and "b" indicates significant difference with diabetic group (DM).



**Figure 1.** Effects of curcumin and metformin on Bax, Bcl-2, and Caspase-3 gene expression in the heart tissue: (A) Bax expression, (B) Bcl-2 expression, (C) Caspase-3 expression and (D) Bax-/Bcl-2 expression ratio, as determined by RT-qPCR. N: normal group; D: diabetic group D+Cur50: diabetic group treated with curcumin (50 mg/kg), D+Cur150: diabetic group treated with curcumin (150 mg/kg), D+Met300: diabetic group treated with metformin (300 mg/kg); D+Met500: diabetic group treated with metformin (500 mg/kg). Data is represented as mean  $\pm$  SD. In each figure «a» represents significant difference ( $p < 0.001$ ) with normal group (N) and «b» indicates significant difference ( $p < 0.001$ ) with diabetic group (D)

curcumin- or metformin-treated rats showed significant changes in caspase-3 gene expression level (Figure 1D).

### Curcumin and metformin prevented diabetes-induced tissue damages

Histopathological examination of the H&E-stained heart sections of control rats revealed normal structure. The cardiomyocytes were uniform with single, oval and centrally located nuclei and the plasma membrane and intercalated discs were intact (Figure 2) whereas induction of diabetes led to extensive structural changes in heart tissue and the presence of hypertrophic myofibers with disarrayed pattern, a mild distortion of intercalated discs, degenerated cardiomyocytes with vacuolated cytoplasm and moderate congestion were the main histopathologic findings. Moreover, nuclei of the cardiomyocytes showed deformation in sizes and shapes (Figure 2).

Treatment with low doses of curcumin (D + Cur50) and metformin (D + Met300), though could not completely alleviate the lesions, showed milder degenerative changes such as nuclear alteration, vacuolated cytoplasm and congestion, when compared to untreated diabetic rats (Figure 2). Heart sections of diabetic rats treated with high doses of curcumin (D + Cur150) and metformin (D + Met500) showed no pathological changes and were comparable to those of normal rats except the presence of small cytoplasmic vacuoles and mild myocytes hypertrophy (Figure 2).


### Discussion

The major mechanisms involved in the pathogenesis of diabetes-induced heart disorders are not yet clearly understood. It is believed that oxidative stress and inflammation resulted from diabetes play critical role in linking diabetes and cardiac disorders through

induction of apoptosis.<sup>25</sup> In diabetes, hyperglycemia triggers intracellular changes leading to the generation of ROS which in turn promotes cellular apoptosis and loss of cardiomyocytes in CVD.<sup>5, 26</sup> Nowadays, besides common therapeutic strategies using natural compounds as alternative or adjuvant therapeutic agents has received much attention due to their beneficial therapeutic potency, lack of side effects and safety. Available evidence suggests that curcumin which is extracted as major lipophilic polyphenol from turmeric<sup>7-8</sup> exhibits antioxidant potential<sup>27</sup> and upregulates the expression of antioxidant enzymes of cellular defense system.<sup>9, 10</sup> Similarly, antioxidant, anti-inflammatory, and anti-apoptotic properties have also been reported for metformin in some experimental conditions.<sup>28, 29</sup> Thus, based on the antioxidant effects of curcumin and metformin, we hypothesized that these compounds may inhibit oxidative stress-induced apoptosis in heart tissue of diabetic rats. Therefore, in the present study beneficial effects of curcumin and metformin on heart tissue were investigated by determining TAC, TOS, OSI, and MDA levels and measuring the activities of enzymes involved in cellular antioxidant defense system including CAT, SOD, and GPx. In addition, the expression levels of apoptosis-related genes such as Bax, Bcl-2, and caspase-3 were determined by RT-qPCR.

We showed that induction of diabetes in D group led to the significant increase in serum AST and LDH levels compared with the control groups which is in line with previous reports.<sup>30, 31</sup>

Interestingly, treatment with both low and high doses of curcumin or metformin significantly prevented alteration of AST and LDH and restored them to the normal range. It is postulated that increased production of ROS, as the consequences of hyperglycemia, leads to damage



**Figure 2.** Curcumin and metformin protected cardiac tissue from diabetes-induced damages, as shown by histopathological examinations. In control rats, (A) histopathology of normal heart tissue with normal features of myocardial cells (H&E; Bar=75  $\mu$ m) and Normal myocardial cells with normal nuclei, longitudinal striation of fibers and intercalated discs, as indicated with arrows (B) were observed (H&E; Bar=20  $\mu$ m). In contrast, in untreated diabetic rats (group D), hypertrophic myofibers with disarrayed pattern (H&E; Bar=20  $\mu$ m) were observed (C) and cardiomyocytes degeneration with vacuolated cytoplasm was detected (D), as shown by arrows (H&E; Bar=20  $\mu$ m). In addition, degenerated cardiomyocytes with pyknotic nuclei and eosinophilic cytoplasm (H&E; Bar=20  $\mu$ m) as shown in (E) and congested blood vessels (F) were also observed in diabetic rats (H&E; Bar=75  $\mu$ m). After treatment of the diabetic rats with low dose of curcumin (group D+Cur50) a few degenerated myocardial cells with vacuolated cytoplasm and mild hypertrophy was observed (G) while higher dose of curcumin (group D+Cur150) resulted in mild myocardial hypertrophy with a minimal degree of cytoplasmic vacuolation (H). Similarly, in those diabetic rats received lower dose of metformin (D+Met300), degenerated myocardial cells with vacuolated cytoplasm (H&E; Bar=20  $\mu$ m) were seen (I) but treatment of the rats with 500 mg/kg of metformin (D500+Met) mostly prevented diabetes-induced damages (J) and a few myocardial cells with mild hypertrophy (H&E; Bar=20  $\mu$ m) was detected in heart tissue.

of the cardiomyocytes cell membrane through reaction with lipids, proteins and other cellular constituents, and ultimately results in the releasing of AST and LDH from damaged cardiac cells into the circulation.<sup>5, 32</sup> The beneficial effects of curcumin and metformin in restoring AST and LDH activities, as observed in this study, are probably attributed to the antioxidant nature of curcumin and metformin,<sup>28, 29</sup> which protected cardiomyocytes from oxidative damages.

Although in the present study none of total CK and its cardiac isoform (CK-MB) increased in diabetic rats (D group), as expected based on the previous reports in the literature,<sup>30, 33</sup> the ratio of CK-MB to total CK (defined as CK-MB/CK) was significantly increased by the induction of diabetes in the rats. Treatment of the rats with either curcumin or metformin normalized the CK-MB/CK ratio to that of control animals but curcumin was found more effective than metformin in restoring CK-MB/CK ratio. Unexpectedly, despite normalizing of CK-MB/CK ratio treatment of diabetic rats with curcumin or metformin lowered total CK and CK-MB levels to nearly 50% of their normal levels. At the moment underlying molecular mechanism of action of metformin or polyphenolic curcumin in greatly lowering CK and CK-MB activities is not clear and further studies are required to show whether reduction in enzyme activities is due to the downregulation at the transcriptional level, suppressing translational stage, or inhibition of post-translational modification processes. Nevertheless, it is supposed that antioxidant properties of these compounds may play critical roles in their wide-range and multifaced cardioprotective potentials.

In the present study although TAC level did not alter

by the induction of diabetes or even by the treatment of diabetic rats with curcumin or metformin, significant enhancements were observed in TOS, OSI, and MDA levels in rats from diabetic group (D). In contrast, treatment of diabetic rats (D + 50Cur, D + 150Cur, D + 300Met, and D + 500Met groups) completely returned these parameters of oxidative stress to their normal ranges, being curcumin more effective than metformin. Similar to our observation, a previously published report has stated increasing in TOS level in heart tissue without any significant changes in TAC level in diabetic rats.<sup>34</sup> Increased TOS level is an expected finding in diabetes mellitus because diabetes-induced hyperglycemia eventually leads to increased production of ROS<sup>5</sup> and ROS itself constitute a major part of TOS.<sup>35</sup> Taş et al also have shown that diabetes resulted in increased heart MDA level<sup>36</sup> through peroxidation of lipids<sup>37</sup> which occurs as a consequence of diabetes-induced enhancement in reactive oxygen species. Available evidence indicated that apart from the antioxidant properties of curcumin which is related to its chemical nature,<sup>9</sup> curcumin may prevent increasing of ROS through reducing of electron transfer chain and activities of enzymes which produce ROS.<sup>25, 38</sup> Metformin may also decrease ROS production through inhibition of mitochondrial respiratory complex-I and increasing of pentose phosphate pathway activity.<sup>39</sup>

Surprisingly, results of antioxidant enzymes activities were contradictory in the present study. In line with results of study by Mollazadeh et al,<sup>30</sup> we showed that diabetic rats had significantly reduced GPx and SOD activities in the heart, whereas showed markedly increased CAT activity compared with control rats. Although this discrepancy

in CAT, SOD, and GPx activities can be explained by the fact that Nrf-2 protein may slightly increase in the heart tissue of two-months old diabetic rats where as a transcription factor upregulates expression of genes involved in antioxidant defense system,<sup>40,41</sup> and therefore, the increased CAT activity in our study might be occurred as a compensatory response against hyperglycemia-induced H<sub>2</sub>O<sub>2</sub> production.<sup>42</sup> However, further studies are required to deeply investigate contradictory responses of CAT, SOD, and GPx in diabetes since upregulation of Nrf-2 expression in acute hyperglycemia as well as its downregulation in chronic hyperglycemia are both reported.<sup>40,41</sup> Treatment with curcumin and metformin completely prevented increasing of CAT activity in the treated groups. Given that increased CAT activity in diabetic rats (D group) is a compensatory response against diabetes-induced oxidative stress,<sup>42</sup> so treatment with curcumin and metformin probably modulated this response through their antioxidant capacities. Upregulation of Nrf-2 by curcumin and metformin has previously been observed,<sup>9,43</sup> thus Nrf-2-dependent increase in GPx and SOD transcription (and activity) might be a cause of increased SOD and GPx observed in treated diabetic rats (D + 50Cur, D + 150Cur, D + 300Met, and D + 500Met).

It has previously been known that diabetes induces apoptosis and cardiomyocyte damage through increased production of ROS.<sup>5,6,26</sup> Here, we showed that induction of diabetes in DM group up-regulated expression of caspase-3 and Bax gene expression and increased ratio of Bax/Bcl-2. Similar observations confirming of increased expression of caspase-3 and Bax genes and enhanced Bax/Bcl-2 ratio have previously been reported in diabetes.<sup>44</sup> It is supposed that increased ROS levels in hyperglycemic conditions reduces AKT activation through reduction of its phosphorylation. AKT inhibits transcription factor FOXO3 which participates in the expression of pro-apoptotic genes such as Bax, thus decreased phosphorylation (and activity) of AKT abolishes inhibitory effects on FOXO3 and induces the expression of Bax and increases the chance of formation Bax-Bax homodimers which is in favor of progressing the apoptosis process.<sup>45</sup> Therefore, the effects of curcumin and metformin in suppressing Bax gene expression might be explained by the fact that curcumin induces phosphorylation and activation of AKT through pro-survival kinase such as phosphoinositide 3-kinase<sup>12</sup> and metformin reduces pro-apoptotic genes through down-regulation of FOXO3.<sup>39</sup>

Induction of diabetes resulted in severe damage in heart tissue on diabetic rats in D group. Hypertrophic myofibers with disarrayed pattern, cardiomyocytes degeneration with vacuolated cytoplasm, and congested blood vessels were observed in histopathological evaluation of heart tissue in diabetic rats which are in line with previous studies.<sup>46,47</sup> Increased ROS level directly or indirectly activates some signal transduction pathways which foster

the expression or augment activation of the hypertrophic factors such as beta-myosin heavy chain.<sup>48</sup> Cardiomyocyte degeneration also may be occurred through autophagy as process which can be mediated through ROS.<sup>49</sup> In the other hand, histopathological evaluation indicated the presence of pyknotic nuclei in the heart tissue of diabetic rats, which are considered as a small condensed nuclei of apoptotic cells.<sup>50</sup> Treatment with curcumin and metformin especially in high doses prevented progressing of diabetes-induced tissue damages through their antioxidant and anti-apoptotic effects. Therefore, it can be mentioned that the findings of histopathological evaluation confirmed the results of gene expression and evaluation of oxidative stress in this present study.

Comparison of beneficial effects of low (50mg/kg) and high (150mg/kg) doses of curcumin showed that there was no statistically significant difference between these two doses in any of parameters measured in the present study including FBS, AST, LDH, CK, CK-MB, CK-MB/CK, TAC, TOS, OSI, MDA, CAT, SOD, GPx, and the expression of Bax, Bcl-2, Bax/Bcl-2, and caspase-3 genes. Similarly, no difference was observed in these parameters between the effects of low (300mg/kg) and high (500mg/kg) doses of metformin. Therefore, it is postulated that wider range of concentrations (higher doses) of curcumin and/or metformin might be needed to see different impacts. However, the probable toxic effects of consuming higher doses of these agents should be taken into consideration in long term treatments.

Moreover, the effects of curcumin and metformin on dependent variables did not differ between groups in exception to the FBS level, where curcumin was significantly effective than metformin. Although no significant difference was observed between curcumin and metformin in AST, LDH, CK, CK-MB, CK-MB/CK, TAC, TOS, OSI, MDA, CAT, SOD, GPx, and the expression of Bax, Bcl-2, Bax/Bcl-2, and caspase-3 genes, the mean values for almost all parameters in curcumin-treated animals were relatively closer than those of metformin to the normal values determined in control rats.

Collectively, it can be suggested that the favorable effects of curcumin and metformin in ameliorating oxidative stress and attenuating diabetes-induced damages in cardiomyocytes, as observed here, were more possibly mediated through their antioxidant properties and not because of their hypoglycemic potency, particularly for metformin. These results may be considered as our incipient understanding about effectiveness of curcumin in ameliorating diabetes-induced oxidative stress and apoptosis in the heart tissue since a deeper investigation of the cardioprotective mechanisms of curcumin and/or metformin requires measurement of ROS, evaluation of apoptosis using TUNEL assay or flow cytometry, measurement of caspase-3 enzyme activity, investigating synergistic effects of combined treatments (curcumin + metformin), and determination of Bax and



Bcl-2 at the protein level.

### Conclusion

Curcumin and metformin exhibit strong antioxidant properties and therefore might be beneficial in ameliorating diabetes-induced oxidative stress and modulating diabetes-induced cardiac damage in the heart tissue.

### Acknowledgments

The authors would like to thank Hamadan University of Medical Sciences for financial support (IR.UMSHA.9512037389).

### Funding

This study did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Competing of interests

The authors have declared no conflicts of interests.

### Ethical approval

The study was approved by the Research Ethics Committee of Hamadan University of Medical Sciences (IR.UMSHA.REC.1395.563).

### References

- Ernawati U, Wihastuti TA, Utami YW. Effectiveness of diabetes self-management education (DSME) in type 2 diabetes mellitus (T2DM) patients: systematic literature review. **J Public Health Res.** 2021;10(2):2240. doi:10.4081/jphr.2021.2240
- Cole JB, Florez JC. Genetics of diabetes mellitus and diabetes complications. **Nat Rev Nephrol.** 2020;16(7):377-390. doi:10.1038/s41581-020-0278-5
- Low Wang CC, Hess CN, Hiatt WR, Goldfine AB. Clinical update: cardiovascular disease in diabetes mellitus: atherosclerotic cardiovascular disease and heart failure in type 2 diabetes mellitus - mechanisms, management, and clinical considerations. **Circulation.** 2016;133(24):2459-2502. doi:10.1161/circulationaha.116.022194
- Faria A, Persaud SJ. Cardiac oxidative stress in diabetes: mechanisms and therapeutic potential. **Pharmacol Ther.** 2017;172:50-62. doi:10.1016/j.pharmthera.2016.11.013
- Yoshinaga A, Kajihara N, Kukidome D, Motoshima H, Matsumura T, Nishikawa T, et al. Hypoglycemia induces mitochondrial reactive oxygen species production through increased fatty acid oxidation and promotes retinal vascular permeability in diabetic mice. **Antioxid Redox Signal.** 2021;34(16):1245-1259. doi:10.1089/ars.2019.8008
- Lee Y, Gustafsson ÅB. Role of apoptosis in cardiovascular disease. **Apoptosis.** 2009;14(4):536-548. doi:10.1007/s10495-008-0302-x
- Pourbagher-Shahri AM, Farkhondeh T, Ashrafzadeh M, Talebi M, Samargahndian S. Curcumin and cardiovascular diseases: focus on cellular targets and cascades. **Biomed Pharmacother.** 2021;136:111214. doi:10.1016/j.biopha.2020.111214
- Gbr AA, Abdel Baky NA, Mohamed EA, Zaky HS. Cardioprotective effect of pioglitazone and curcumin against diabetic cardiomyopathy in type 1 diabetes mellitus: impact on CaMKII/NF-κB/TGF-β1 and PPAR-γ signaling pathway. **Naunyn Schmiedebergs Arch Pharmacol.** 2021;394(2):349-360. doi:10.1007/s00210-020-01979-y
- Kotha RR, Luthria DL. Curcumin: biological, pharmaceutical, nutraceutical, and analytical aspects. **Molecules.** 2019;24(16):2930. doi:10.3390/molecules24162930
- Bahmani M, Ziamajidi N, Hashemnia M, Abbasalipourkabir R, Salehzadeh A. Human umbilical cord blood mesenchymal stem cell conditioned medium (hMSC-CM) improves antioxidant status in carbon tetrachloride-induced oxidative damage in rat. **Iran J Sci Technol Trans A Sci.** 2020;44(5):1327-1335. doi:10.1007/s40995-020-00944-x
- Hasanpour Dehkordi A, Abbaszadeh A, Mir S, Hasanvand A. Metformin and its anti-inflammatory and anti-oxidative effects; new concepts. **J Renal Inj Prev.** 2019;8(1):54-61. doi:10.15171/jrip.2019.11
- Ren BC, Zhang YF, Liu SS, Cheng XJ, Yang X, Cui XG, et al. Curcumin alleviates oxidative stress and inhibits apoptosis in diabetic cardiomyopathy via Sirt1-Foxo1 and PI3K-Akt signalling pathways. **J Cell Mol Med.** 2020;24(21):12355-12367. doi:10.1111/jcmm.15725
- Zhai L, Gu J, Yang D, Hu W, Wang W, Ye S. Metformin ameliorates podocyte damage by restoring renal tissue nephrin expression in type 2 diabetic rats. **J Diabetes.** 2017;9(5):510-517. doi:10.1111/1753-0407.12437
- Asadi S, Goodarzi MT, Karimi J, Hashemnia M, Khodadadi I. Does curcumin or metformin attenuate oxidative stress and diabetic nephropathy in rats? **J Nephropathol.** 2019;8(1):e08. doi:10.15171/jnp.2019.08
- Sharma S, Kulkarni SK, Chopra K. Curcumin, the active principle of turmeric (*Curcuma longa*), ameliorates diabetic nephropathy in rats. **Clin Exp Pharmacol Physiol.** 2006;33(10):940-945. doi:10.1111/j.1440-1681.2006.04468.x
- Jun JE, Lee SE, Lee YB, Ahn JY, Kim G, Hur KY, et al. Continuous glucose monitoring defined glucose variability is associated with cardiovascular autonomic neuropathy in type 1 diabetes. **Diabetes Metab Res Rev.** 2019;35(2):e3092. doi:10.1002/dmrr.3092
- Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. **Anal Biochem.** 1976;72:248-254. doi:10.1006/abio.1976.9999
- Benzie IF, Strain JJ. Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. **Methods Enzymol.** 1999;299:15-27. doi:10.1016/s0076-6879(99)99005-5
- Erel O. A new automated colorimetric method for measuring total oxidant status. **Clin Biochem.** 2005;38(12):1103-1111. doi:10.1016/j.clinbiochem.2005.08.008
- Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. **Anal Biochem.** 1979;95(2):351-358. doi:10.1016/0003-2697(79)90738-3
- Hadwan MH, Abed HN. Data supporting the spectrophotometric method for the estimation of catalase activity. **Data Brief.** 2016;6:194-199. doi:10.1016/j.dib.2015.12.012
- Paglia DE, Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. **J Lab Clin Med.** 1967;70(1):158-169. doi:10.5555/uri:pii:0022214367900765
- Marklund S, Marklund G. Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. **Eur J Biochem.** 1974;47(3):469-474. doi:10.1111/j.1432-1033.1974.tb03714.x
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>ΔΔCT method. **Methods.** 2001;25(4):402-408. doi:10.1006/meth.2001.1262
- Soetikno V, Suzuki K, Veeraveedu PT, Arumugam S, Lakshmanan AP, Sone H, et al. Molecular understanding of curcumin in diabetic nephropathy. **Drug Discov Today.** 2013;18(15-16):756-763. doi:10.1016/j.drudis.2013.04.009

26. Jang JH, Surh YJ. Potentiation of cellular antioxidant capacity by Bcl-2: implications for its antiapoptotic function. **Biochem Pharmacol.** 2003;66(8):1371-1379. doi:10.1016/s0006-2952(03)00487-8
27. Bhattacharyya S, Ghosh S, Sil PC. Amelioration of aspirin induced oxidative impairment and apoptotic cell death by a novel antioxidant protein molecule isolated from the herb *Phyllanthus niruri*. **PLoS One.** 2014;9(2):e89026. doi:10.1371/journal.pone.0089026
28. Kelly B, Tannahill GM, Murphy MP, O'Neill LA. Metformin inhibits the production of reactive oxygen species from NADH: ubiquinone oxidoreductase to limit induction of interleukin-1 $\beta$  (IL-1 $\beta$ ) and boosts interleukin-10 (IL-10) in lipopolysaccharide (LPS)-activated macrophages. **J Biol Chem.** 2015;290(33):20348-20359. doi:10.1074/jbc.M115.662114
29. Pandey A, Kumar VL. Protective effect of metformin against acute inflammation and oxidative stress in rat. **Drug Dev Res.** 2016;77(6):278-284. doi:10.1002/ddr.21322
30. Mollazadeh H, Sadeghnia HR, Hoseini A, Farzadnia M, Boroushaki MT. Effects of pomegranate seed oil on oxidative stress markers, serum biochemical parameters and pathological findings in kidney and heart of streptozotocin-induced diabetic rats. **Ren Fail.** 2016;38(8):1256-1266. doi:10.1080/0886022x.2016.1207053
31. Qin Q, Lin N, Huang H, Zhang X, Cao X, Wang Y, et al. Ginsenoside Rg1 ameliorates cardiac oxidative stress and inflammation in streptozotocin-induced diabetic rats. **Diabetes Metab Syndr Obes.** 2019;12:1091-1103. doi:10.2147/dms.s208989
32. Boghdady NA. Antioxidant and antiapoptotic effects of proanthocyanidin and ginkgo biloba extract against doxorubicin-induced cardiac injury in rats. **Cell Biochem Funct.** 2013;31(4):344-351. doi:10.1002/cbf.2907
33. Ebrahimi E, Shirali S, Talaei R. The protective effect of marigold hydroalcoholic extract in STZ-induced diabetic rats: evaluation of cardiac and pancreatic biomarkers in the serum. **J Bot.** 2016;2016:9803928. doi:10.1155/2016/9803928
34. Akat F, Fıçıcılar H, Durak A, Tuncay E, Dursun AD, Topal Çelikkın F, et al. Intermittent hypoxia induces beneficial cardiovascular remodeling in left ventricular function of type 1 diabetic rat. **Anatol J Cardiol.** 2018;19(4):259-266. doi:10.14744/AnatolJCardiol.2018.00236
35. Suresh DR, Kumaran SE, Annam V, Veena H. Age related changes in malondialdehyde: total antioxidant capacity ratio—a novel marker of oxidative stress. **Int J Pharma Bio Sci.** 2010;1(2):1-6.
36. Taş S, Sarandöl E, Dirican M. Vitamin B6 supplementation improves oxidative stress and enhances serum paraoxonase/arylesterase activities in streptozotocin-induced diabetic rats. **ScientificWorldJournal.** 2014;2014:351598. doi:10.1155/2014/351598
37. Janero DR. Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. **Free Radic Biol Med.** 1990;9(6):515-540. doi:10.1016/0891-5849(90)90131-2
38. Kumar S, Prasad S, Sitasawad SL. Multiple antioxidants improve cardiac complications and inhibit cardiac cell death in streptozotocin-induced diabetic rats. **PLoS One.** 2013;8(7):e67009. doi:10.1371/journal.pone.0067009
39. Nesti L, Natali A. Metformin effects on the heart and the cardiovascular system: a review of experimental and clinical data. **Nutr Metab Cardiovasc Dis.** 2017;27(8):657-669. doi:10.1016/j.numecd.2017.04.009
40. Tan Y, Ichikawa T, Li J, Si Q, Yang H, Chen X, et al. Diabetic downregulation of Nrf2 activity via ERK contributes to oxidative stress-induced insulin resistance in cardiac cells in vitro and in vivo. **Diabetes.** 2011;60(2):625-633. doi:10.2337/db10-1164
41. Wang G, Song X, Zhao L, Li Z, Liu B. Resveratrol prevents diabetic cardiomyopathy by increasing Nrf2 expression and transcriptional activity. **Biomed Res Int.** 2018;2018:2150218. doi:10.1155/2018/2150218
42. El-Bahr SM. Curcumin regulates gene expression of insulin like growth factor, B-cell CLL/lymphoma 2 and antioxidant enzymes in streptozotocin induced diabetic rats. **BMC Complement Altern Med.** 2013;13:368. doi:10.1186/1472-6882-13-368
43. Onken B, Driscoll M. Metformin induces a dietary restriction-like state and the oxidative stress response to extend *C. elegans* Healthspan via AMPK, LKB1, and SKN-1. **PLoS One.** 2010;5(1):e8758. doi:10.1371/journal.pone.0008758
44. Wang Y, Sun H, Zhang J, Xia Z, Chen W. Streptozotocin-induced diabetic cardiomyopathy in rats: ameliorative effect of PIPERINE via Bcl2, Bax/Bcl2, and caspase-3 pathways. **Biosci Biotechnol Biochem.** 2020;84(12):2533-2544. doi:10.1080/09168451.2020.1815170
45. Dong LY, Yao LP, Zhao J, Jin KK, Qiu XX. Captopril inhibits calpain-mediated apoptosis of myocardial cells in diabetic rats and improves cardiac function. **Mol Med Rep.** 2018;18(2):2300-2306. doi:10.3892/mmr.2018.9192
46. Jemai H, Sayadi S. Heart histopathology and oxidative features in diabetic rats and protective effects of oleuropein. **Adv Biosci Biotechnol.** 2015;6(6):383-389. doi:10.4236/abb.2015.66038
47. Al-Rasheed NM, Al-Rasheed NM, Hasan IH, Al-Amin MA, Al-Ajmi HN, Mohamad RA, et al. Simvastatin ameliorates diabetic cardiomyopathy by attenuating oxidative stress and inflammation in rats. **Oxid Med Cell Longev.** 2017;2017:1092015. doi:10.1155/2017/1092015
48. Moris D, Spartalis M, Spartalis E, Karachaliou GS, Karaolanis GI, Tsourouflis G, et al. The role of reactive oxygen species in the pathophysiology of cardiovascular diseases and the clinical significance of myocardial redox. **Ann Transl Med.** 2017;5(16):326. doi:10.21037/atm.2017.06.27
49. Takemura G, Kanamori H, Okada H, Miyazaki N, Watanabe T, Tsujimoto A, et al. Anti-apoptosis in nonmyocytes and pro-autophagy in cardiomyocytes: two strategies against postinfarction heart failure through regulation of cell death/degeneration. **Heart Fail Rev.** 2018;23(5):759-772. doi:10.1007/s10741-018-9708-x
50. Ortiz A, Lorz C, Catalan MP, Justo P, Egido J. Role and regulation of apoptotic cell death in the kidney. Y2K update. **Front Biosci.** 2000;5:D735-749. doi:10.2741/ortiz