



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

4. Liu Y, Yang Y, Zhang C, Huang F, Wang F, Yuan J, et al. Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury. *Sci China Life Sci* 2020;63:364-74.
5. Chan JF, Yuan S, Kok K-H, To KK, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* 2020;395:514-23.
6. Cai J, Xu J, Lin D, Yang Z, Xu L, Qu Z, et al. A case series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clin Infect Dis* 2020 [Epub ahead of print].
7. Cai JH, Wang XS, Ge YL, Xia AM, Chang HL, Tian H, et al. [First case of 2019 novel coronavirus infection in children in Shanghai]. *Zhonghua Er Ke Za Zhi* 2020;58:86-7.
8. Chen F, Liu Z, Zhang F, Xiong RH, Chen Y, Cheng XF, et al. [First case of severe childhood novel coronavirus pneumonia in China]. *Chin J Pediatr* 2020;58:179-82.
9. Feng K, Yun YX, Wang XF, Yang GD, Zheng YJ, Lin CM, et al. Analysis of CT features of 15 children with 2019 novel coronavirus infection. *Zhonghua Er Ke Za Zhi* 2020;58:007.
10. Kam K, Yung CF, Cui L, Lin Tzer Pin R, Mak TM, Maiwald M, et al. A well infant with coronavirus disease 2019 (COVID-19) with high viral load. *Clin Infect Dis* 2020 [Epub ahead of print].
11. Wang D, Ju XL, Xie F, Lu Y, Li FY, Huang HH, et al. Clinical analysis of 31 cases of 2019 novel coronavirus infection in children from six provinces (autonomous region) of northern China. *Zhonghua Er Ke Za Zhi* 2020;58:011.
12. Zhang YH, Lin DJ, Xiao MF, Wang JC, Wei Y, Lei ZX, et al. 2019 Novel coronavirus infection in a three-month-old baby. *Zhonghua Er Ke Za Zhi* 2020;58:182-4.
13. Zeng L, Tao X, Yuan W, Wang J, Liu X, Liu Z. China's first neonatal coronavirus pneumonia. *Chin J Pediatr* 2020;58.
14. Zhang G, Zjang A, Huang L, Cheng LY, Liu ZX, Peng XL, et al. Twin girls infected with SARS-CoV-2. *Chin J Contemp Pediatr* 2020;22:221-5.
15. Zhao R, Shen X, Yu K, Sheng. A case of children with 2019 novel Coronavirus Infection. *Zhejiang Med J*; 2020 [Epub ahead of print].
16. Lu X, Zhang L, Du H, Zhang J, Li YY, Qu J, et al. Chinese Pediatric Novel Coronavirus Study Team. SARS-CoV-2 infection in children. *N Engl J Med* 2020 [Epub ahead of print].
17. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. China Medical Treatment Expert Group for Covid-19. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020 [Epub ahead of print].
18. Husain EH, Mullah-Ali A, Al-Sharidah S, Azab AF, Adekile A. Infectious etiologies of transient neutropenia in previously healthy children. *Pediatr Infect Dis J* 2012;31:575-7.
19. Karavanaki K, Polychronopoulou S, Giannaki M, Haliotis F, Sider B, Brisimitzi M, et al. Transient and chronic neutropenias detected in children with different viral and bacterial infections. *Acta Paediatr* 2006;95:565-72.

## Missed or delayed diagnosis of Kawasaki disease during the 2019 novel coronavirus disease (COVID-19) pandemic



### To the Editor:

Due to “stay-at-home” orders and the risk of novel coronavirus disease 2019 (COVID-19), many parents now hesitate or fear seeking in-person consultations for their children. This has led to reductions in emergency department visits and hospital admissions for other critical illnesses. In addition, healthcare providers have focused on COVID-19 management during the pandemic. Because of Bayesian

thinking, other diseases may be underdiagnosed or undergo delayed treatment.

Because COVID-19 now leads as the probable diagnosis for first-line providers encountering febrile patients, the potential for missed or late diagnosis and treatment of Kawasaki disease in children is particularly concerning.<sup>1</sup> Prompt diagnosis of Kawasaki disease and treatment with intravenous immunoglobulin (IVIG) prevents coronary artery aneurysms (CAA).<sup>2,3</sup> Without timely treatment, CAAs could occur in up to 25% of children with Kawasaki disease.<sup>3</sup>

We respectfully remind caregivers of the following principles for the care of children with suspected or definite Kawasaki disease: (1) Keep a high suspicion for Kawasaki disease in all children with prolonged fever, but especially in those younger than 1 year of age. (2) Administer IVIG within 10 days, and ideally within 7 days, from onset of fever. (3) In the presence of ongoing systemic inflammation, children with Kawasaki disease presenting with greater than 10 days of fever and/or CAA may warrant IVIG treatment. (4) Continue to obtain recommended echocardiograms according to published guidelines.<sup>3</sup> (5) Watch for late manifestations of Kawasaki disease, review the clinical history, and seek pediatric cardiology consultation.<sup>4,5</sup> (6) In the case of delayed diagnosis, refer to the American Heart Association management guidelines or contact an expert in Kawasaki disease.<sup>3</sup> (7) Offer telemedicine services, remote echocardiogram, and direct-to-consumer visits that allow for nonverbal communication when evaluating children with confirmed or suspected Kawasaki disease.<sup>6-8</sup>

With this, we hope to avoid a future surge in the prevalence of CAAs in patients due to missed or delayed diagnosis of Kawasaki disease.

*We thank Angela J Doty, MD, and Maryam Harahsheh for their editorial assistance.*

**Ashraf S. Harahsheh, MD, FACC, FAAP**

Division of Pediatric Cardiology

Department of Pediatrics

Children's National Hospital

The George Washington University School of Medicine

Washington, District of Columbia

**Nagib Dahdah, MD**

Division of Pediatric Cardiology

Department of Pediatrics

CHU Sainte Justine

University of Montreal

Montreal, Quebec, Canada

**Jane W. Newburger, MD, MPH**

Department of Cardiology

Boston Children's Hospital

Department of Pediatrics

Harvard Medical School

Boston, Massachusetts

**Michael A. Portman, MD**  
Seattle Children's Hospital  
University of Washington  
Seattle, Washington

**Maryam Piram, MD, MPH, PhD**  
Division of Pediatric Dermatology  
Department of Pediatrics  
CHU Sainte Justine  
University of Montreal  
Montreal, Quebec, Canada  
  
CEREMAIA  
Université Paris Saclay  
Le Kremlin Bicêtre, France

**Robert Tulloh, DM, FRCPC**  
University of Bristol  
Bristol, United Kingdom

**Brian W. McCrindle, MD, MPH**  
The Hospital for Sick Children  
Toronto, Ontario, Canada

**Sarah D. de Ferranti, MD, MPH**  
Department of Cardiology  
Boston Children's Hospital  
Department of Pediatrics  
Harvard Medical School  
Boston, Massachusetts

**Rolando Cimaz, MD**  
University of Milano  
Milano, Italy

**Dongngan T. Truong, MD**  
University of Utah  
Primary Children's Hospital  
Salt Lake City, Utah

**Jane C. Burns, MD**  
Department of Pediatrics  
Rady Children's Hospital  
University of California San Diego  
San Diego, California

<https://doi.org/10.1016/j.jpeds.2020.04.052>

*J.N. serves on the Editorial Board for The Journal of Pediatrics. The other authors declare no conflicts of interest.*

## References

1. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020 Apr 30;382:1708-20.
2. Jones V, Mills M, Suarez D, Hogan C, Yeh D, Segal J, et al. COVID-19 and Kawasaki disease: novel virus and novel case. *Hosp Pediatr* 2020;10(6):537-40.
3. McCrindle BW, Rowley AH, Newburger JW, Burns JC, Bolger AF, Gewitz M, et al. Diagnosis, treatment, and long-term management of Ka-

wasaki disease: a scientific statement for health professionals from the American Heart Association. *Circulation* 2017;135:e927-99.

4. Wang S, Best BM, Burns JC. Periungual desquamation in patients with Kawasaki disease. *Pediatr Infect Dis J* 2009;28:538-9.
5. Lindsley CB. Nail-bed lines in Kawasaki disease. *Am J Dis Child* 1992;146:659-60.
6. Satou GM, Rheuban K, Alverson D, Lewin M, Mahnke C, Marcin J, et al. Telemedicine in pediatric cardiology: a scientific statement from the American Heart Association. *Circulation* 2017;135:e648-78.
7. Phillips A, Sable C, Waggaman C, Harahsheh AS. Direct-to consumer cardiology telemedicine: a single large academic pediatric center experience (abstract). Paper presented at the Nov. 16-18, 2019 American Heart Association Scientific Sessions; Philadelphia, PA.
8. Harahsheh AS, Ottolini M, Lewis K, Blatt B, Mitchell S, Greenberg L. An innovative pilot curriculum training pediatric residents in referral and communication skills on a cardiology rotation. *Acad Pediatr* 2016;16:700-2.

## Iron, ketone bodies, and brain development



### To the Editor:

Iron deficiency is deleterious in early life brain development and a risk for short- and long-term cognitive, motor, and socioemotional impairment.<sup>1,2</sup> Parkin et al recently reported that increasing serum ferritin values, up to a level of 17  $\mu\text{g/L}$ , were correlated with higher cognitive function in infants of 1-3 years of age.<sup>3</sup> Here, we address another aspect of the role of iron deficiency in brain development in relation to ketone bodies as an energy source for brain metabolism.

In the starved state, ketone bodies are synthesized from free fatty acids produced by breakdown of body fat, and cross the blood-brain barrier for use in brain metabolism. Ketone bodies may have neuroprotective effects and a ketogenic diet is used in treatment of neurological diseases such as refractory epilepsy, Parkinson disease, Alzheimer disease, and traumatic brain injury.<sup>4,5</sup>

Kuzawa and Blair hypothesized that body fat is a critical brain energy source, especially during infancy and early childhood.<sup>6</sup> In this period, the brain consumes about 40% of daily energy expenditure (20% for adults). Glucose alone cannot supply this energy, and ketone bodies are used. Thus, brain energy expenditure is inversely related to body fat gain (increased body mass index) during early childhood, and thus, decreases the risk of obesity.<sup>6,7</sup>

In an iron-deficient state, ketogenesis is impaired, citrate synthase and succinate dehydrogenase activities are decreased, and production of free fatty acids and ketone bodies is limited.<sup>8,9</sup> Therefore, iron deficiency may not only have a direct effect on brain function and development, but may also decrease the availability of ketone bodies as an energy source.<sup>3</sup> This possibility that the brain is deprived of ketone bodies as an energy source in iron deficiency indicates a need for further research into how iron is involved in brain development during early childhood.

**Osamu Arisaka, MD, PhD**  
Department of Pediatrics  
Nasu Red Cross Hospital  
Ohtawara, Tochigi-ken, Japan