



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Review article

# COVID-19 in solid organ transplant recipients: A systematic review and meta-analysis of current literature



Mohammed A. Raja <sup>a,1</sup>, Maria A. Mendoza <sup>b,\*</sup>, Aasith Villavicencio <sup>b</sup>, Shweta Anjan <sup>a,c</sup>, John M. Reynolds <sup>d</sup>, Veraprapas Kittipibul <sup>b</sup>, Anmary Fernandez <sup>a</sup>, Giselle Guerra <sup>c,e</sup>, Jose F. Camargo <sup>a</sup>, Jacques Simkins <sup>a,c</sup>, Michele I. Morris <sup>a</sup>, Lilian A. Abbo <sup>a,c</sup>, Yoichiro Natori <sup>a,c,\*</sup>

<sup>a</sup> Department of Medicine, Division of Infectious Disease, University of Miami Miller School of Medicine, Miami, FL, USA

<sup>b</sup> Department of Medicine, Division of Internal Medicine, University of Miami Miller School of Medicine, Miami, FL, USA

<sup>c</sup> Miami Transplant Institute, Jackson Health System, Miami, FL, USA

<sup>d</sup> Department of Health Informatics, Calder Memorial Library, University of Miami Miller School of Medicine Miami, FL, USA

<sup>e</sup> Department of Medicine, Division of Nephrology, University of Miami Miller School of Medicine Miami, FL, USA

## ARTICLE INFO

## Keywords:

Solid organ transplant  
COVID-19  
SARS-CoV-2  
Immunosuppression  
Systematic review  
Meta-analysis

## ABSTRACT

Severe acute respiratory virus syndrome 2 (SARS-CoV-2) has led to a worldwide pandemic. Early studies in solid organ transplant (SOT) recipients suggested a wide variety of presentations, however, there remains a paucity of robust data in this population. We conducted a systematic review and meta-analysis of SOT recipients with SARS-CoV-2 infection from January 1<sup>st</sup> to October 9<sup>th</sup>, 2020. Pooled incidence of symptoms, treatments and outcomes were assessed. Two hundred and fifteen studies were included for systematic review and 60 for meta-analysis. We identified 2,772 unique SOT recipients including 1,500 kidney, 505 liver, 141 heart and 97 lung. Most common presenting symptoms were fever and cough in 70.2% and 63.8% respectively. Majority (81%) required hospital admission. Immunosuppressive medications, especially antimetabolites, were decreased in 76.2%. Hydroxychloroquine and interleukin six antagonists were administered in 59.5% and 14.9% respectively, while only few patients received remdesivir and convalescent plasma. Intensive care unit admission was 29% from amongst hospitalized patients. Only few studies reported secondary infections. Overall mortality was 18.6%. Our analysis shows a high incidence of hospital admission in SOT recipients with SARS-CoV-2 infection. As management of SARS-CoV-2 continues to evolve, long-term outcomes among SOT recipients should be assessed in future studies.

© 2020 Published by Elsevier Inc.

## Contents

1. Introduction . . . . .	2
2. Materials and methods . . . . .	2
2.1. Search methods . . . . .	2
2.2. Inclusion and exclusion criteria . . . . .	3
2.3. Data analysis . . . . .	4
3. Results . . . . .	4
3.1. Patient background . . . . .	5
3.2. Symptoms with COVID-19 in solid organ transplant recipients . . . . .	5
3.3. Laboratory results and radiologic findings . . . . .	5

*Abbreviation:* AKI, Acute kidney injury; CI, Confidence interval; CNI, Calcineurin inhibitor; COVID-19, Coronavirus disease 2019; CRP, C-Reactive Protein; FEM, Fixed Effect Model; HSCT, Hematopoietic Stem Cell Transplant; ICU, Intensive Care Unit; IL-6, Interleukin Six; MERS, Middle East Respiratory Syndrome; NP, Nasopharyngeal; PRESS, Peer Review for Electronic Search Strategies; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; REM, Random Effect Model; RT-PCR, Reverse Transcriptase-Polymerase Chain Reaction; SARS, Severe Acute Respiratory Syndrome Coronavirus 1; SARS-CoV-2, Severe Acute Respiratory Syndrome Coronavirus 2; SOT, Solid organ transplant.

\* Corresponding author at: Division of Infectious Diseases, Miami Transplant Institute, University of Miami Miller School of Medicine, 1801 N.W. 9th Ave., Suite 733, Miami, FL 33136, USA.

E-mail address: [yxn138@med.miami.edu](mailto:yxn138@med.miami.edu) (Y. Natori).

<sup>1</sup> Co-first author.

3.4. Treatment method . . . . .	6
3.5. Outcomes . . . . .	7
3.6. Kidney transplant recipients . . . . .	9
3.7. Non-kidney solid organ transplant recipients. . . . .	9
4. Discussion. . . . .	10
Funding . . . . .	11
Disclosure . . . . .	11
Authorship . . . . .	11
Declaration of Competing Interest . . . . .	12
Acknowledgement . . . . .	12
Appendix A. Supplementary data . . . . .	12
References . . . . .	12

## 1. Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the virus responsible for the coronavirus disease 2019 (COVID-19), has been an unmitigated worldwide pandemic, since the first case was reported in December 2019 [1,2]. Due to the high rate of transmissibility in the asymptomatic phase, the virus has continued to spread globally with catastrophic outcomes and has been a great challenge for clinicians all over the world [2,3]. Even though it causes mild disease in certain populations, to date it has led to 228,998 and 1,178,475 deaths in the U.S and worldwide, respectively [3].

Solid organ transplant (SOT) recipients are known to be vulnerable to several respiratory virus infections, such as influenza [4] due to a weakened T-cell mediated immune response [5]. The Centers for Disease Control and Prevention include SOT recipients amongst patients at increased risk for severe illness from SARS-CoV-2 [6]. Due to the novelty of this virus, there is still a paucity of data regarding many aspects including its natural course within immunocompromised hosts, the utility of current treatment regimens employed in non-transplant individuals for SOT recipients, the effect of immunosuppression on the course of the disease, and much more [7,8]. Even though presence of comorbidities, such as obesity, have been reported as risk factors for severe disease [9], an immunocompromised state has not yet been proven to have a worse clinical outcome with SARS-CoV-2 infection. Furthermore, as Vishnevetsky correctly points out, viruses that belong to the same coronavirus family such as Middle East Respiratory Syndrome (MERS) and Severe Acute Respiratory Syndrome Coronavirus 1 (SARS) have not shown an increased risk of infection or worse outcome in immunocompromised patients [10–12]. Lai et. al. also published their systematic review comparing disease outcomes among the three major coronavirus diseases with preliminary data [13]. Furthermore, immunosuppression in the setting of hyperinflammation may even be beneficial, by blunting excessive cytokine release, which is still left to be established [14].

Since the first report of COVID-19 in SOT recipient from China was published in early 2020 [15], researchers from all over the world, have published their own experiences, especially from epidemic centers [16–18] to try and bridge the gap in knowledge, including several recently published registries [19–22]. COVID-19 presentation amongst SOT recipients has ranged from mild upper respiratory infection to severe acute respiratory distress and death [23,24]. Studies in the general population so far suggest a higher prevalence and severity of disease among middle-aged patients and in those with underlying comorbidities such as hypertension and obesity, however, data in SOT recipients is still lacking [25]. The wide variety in clinical presentation among SOT recipients is still not fully understood. Fung and Bekik recently published their narrative review that included SOT recipients and their manuscript broadly covered this area but did not assess present data quantitatively [24].

We performed a systematic review and meta-analysis of the existing literature to help expand our current limited knowledge base regarding

COVID-19 in SOT recipients, especially looking at clinical presentations, treatment modalities and outcomes, with the aim to provide transplant physicians worldwide with important decision-making tools to manage these critically ill patients during this time of crisis.

## 2. Materials and methods


Our review was conducted according to the Methodological Expectations for Cochrane Intervention Reviews [26] and reported following Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines [27].

### 2.1. Search methods

We conducted a comprehensive literature database search to find all relevant literature, published from January 1<sup>st</sup> to October 9<sup>th</sup>, 2020, on SARS-CoV-2 and SOT. The database search strategy was developed by an academic health science librarian (J.R.) in consultation with the project leader (Y.N.) and was reviewed using the Peer Review for Electronic Search Strategies (PRESS) tool through the PRESS forum [28].

We searched Ovid Medline (including Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily), Embase (Elsevier Embase.com), Scopus (Elsevier), Web of Science (Clarivate Analytics, using Science Citation Index Expanded, Social Sciences Citation Index, Arts & Humanities Citation Index, Conference Proceedings Citation Index- Science, Conference Proceedings Citation Index- Social Science & Humanities, Emerging Sources Citation Index, BIOSIS Citation Index, KCI-Korean Journal Database, Russian Science Citation Index, SciELO Citation Index, and Zoological Record), and LitCovid (National Center for Biotechnology Information, U.S. National Library of Medicine, without date restrictions). We searched clinical trial registries with Clinicaltrials.gov and the World Health Organization International Clinical Trials Registry Platform for clinical trials with reported results. In lieu of handsearching, we also searched the online tables of contents of seven journals: *American Journal of Transplantation*, *Journal of Heart and Lung Transplantation*, *Liver Transplantation*, *Pediatric Transplantation*, *Transplant Infectious Disease*, *Transplant International*, and *Transplantation*. Databases and registries were searched between June 11<sup>th</sup> and June 15<sup>th</sup>, 2020, and individual journals were searched on June 19<sup>th</sup>, 2020. The database and journal site searches were updated on July 13<sup>th</sup>, 15<sup>th</sup>, August 20<sup>th</sup>, October 5<sup>th</sup>, as well as October 8<sup>th</sup> and again on October 9<sup>th</sup> 2020. The research team also monitored social media for new studies up to August 23<sup>rd</sup>, 2020. The search strategy was initially based on terms from 36 articles found by the research team prior to the comprehensive database search, and by refining, expanding, and adding to COVID-19 specific search strategies by Ellen Aaronson at the Mayo Clinic [29], and Michelle Volesko Brewer and Pieter van der Houwen, at Wolters Kluwer/Ovid [30].

The search strategy was written for Ovid Medline and translated using each database's syntax, controlled vocabulary, and search fields. MeSH terms, Emtree terms, and text words were used for the search



**Fig. 1.** Study selection flow. Majority of the cases were excluded due to opinion commentaries for COVID-19 or related to other viruses. Abbreviations: COVID; Coronavirus disease 2019, SOT; Solid organ transplant, RT-PCR; Reverse transcriptase-polymerase chain reaction.

concepts of Covid-19/SARS-Cov-2 and solid organ transplantation and their synonyms. No language, study type, or publication type limits were applied at the search phase. Animal studies were excluded. Except for LitCovid, searches were limited by date from January 1, 2020 to the present to allow for broader search terms while still focusing on SARS-Cov-2.

For full search strategies, see Appendix 1. All database records were downloaded to EndNote X9, then uploaded to Covidence web-based software for deduplication, screening, and data extraction. We

contacted several authors whose cases potentially can be used several times in the different articles. Two reviewers (MAM and APV) independently assessed all studies for risk of bias. Any discrepancies in data extraction were resolved with discussion with the third and fourth reviewers (M.R. and Y.N.). After study selection, one author (J.R.) checked for retractions at The Retraction Watch Database and individual journal websites.

### 2.2. Inclusion and exclusion criteria

We included all randomized controlled trials, cohort studies, case reports, and letters to the editor that included targeted data in SOT and reverse transcriptase-polymerase chain reaction (RT-PCR) or antibody testing confirmed SARS-CoV-2 infection. As previously mentioned, animal studies and articles where the main topic was other than COVID-19 in SOT patients were excluded. We excluded studies dealing with patients with suspected infection (eg. chest imaging abnormality and symptoms), without positive RT-PCR or antibody testing. We also excluded other non-SARS-CoV-2 coronavirus related studies such as MERS and SARS. If the study subject was published more than once,

**Table 1**  
Summary of symptoms with COVID 19 in Solid Organ Transplant Recipients.

Symptoms	No. of Studies	Model	Pooled incidence (%) (95% CI)	Heterogeneity ( $I^2$ ), $p$ value
Fever	38	REM	70.2 (61.0-79.5)	96.2, $p < 0.01$
Cough	38	REM	63.8 (55.0-72.6)	94.4, $p < 0.01$
Shortness of breath	36	REM	49.1 (41.1-57.1)	92.1, $p < 0.01$
Diarrhea	36	REM	30.4 (23.6-37.2)	85.4, $p < 0.01$

Abbreviations: CI, confidence interval, REM, random effect model

we entered in the analysis only once using all available sources. For the meta-analysis, we only included articles with 5 or more patients to minimize small study effect. From included studies, the following variables were collected: number of subjects, organ transplant types, symptoms, laboratory data, treatment modalities used, type of immunosuppression, modification to the immunosuppression, graft loss, follow up period, duration from first positivity to negativity of nasopharyngeal (NP) swab RT-PCR, and all-cause mortality. After obtaining these results we excluded studies for hospital admission, intensive care unit (ICU) admission, mechanical ventilation, and all-cause mortality, for studies dealing only with critically ill patients, to avoid selection bias and avoid heterogeneity.

2.3. Data analysis

We conducted this data analysis using STATA/IC 14.0 (StataCorp LP, College Station, TX, USA). Dichotomous data were shown as relative risk ratio (RR, event/total) and their confidence interval (CI), where relevant. Pooled incidence was calculated with either random effect model or fixed-effect model, wherever appropriate. Heterogeneity among studies was graphically shown and checked with I<sup>2</sup> values,

which show the variation among studies that are not due to chance. I<sup>2</sup> values of 75%, 50%, and 25% correspond to a high, medium, and low level of heterogeneity. Fixed-effect model (FEM) was used if I<sup>2</sup> was less than 50%, otherwise random effect model (REM) was used. To visually inspect the publication bias, we created both forest and funnel plots.

3. Results

With this search strategy, we found 1,667 potential studies, after excluding the duplicates. Of these, 1,263 studies were excluded during the abstract and title review as they did not meet eligibility criteria. The most common reasons for exclusion were perspective or opinion commentaries regarding COVID-19 and articles reporting viruses other than COVID-19. As a result, we conducted a full-text review for 403 articles, of which 188 were excluded with reasons shown in Fig. 1. Finally, a total of 215 studies were included in the systematic review [8,15–24,31–235], and 60 studies were included in a meta-analysis [16,18,21,22,24,31,32,38,40,43,50,51,55,59,60,65,69,70,79,81,85,88,96,100,102,108,114,117,128,133,135,136,141–143,149,151,153,157,158,162,166,175,179,183,185,189,190,192,199,210,212,216,217,219,228,229,231].




Fig. 2A. Pooled incidence of fever showing the absolute risk and 95% confidence intervals of each study (n=38) [18,20,22,31,56,59,60,65,70,79,81,85,88,96,102,108,117,133,136,141–143,151,152,156,158,166,179,189,190,192,210,212,216,217,219,229,231]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \*=study period not defined.

3.1. Patient background

In total, 2,772 unique SOT recipients were identified including 1,500 kidney, 505 liver, 141 heart, 97 lung, 1 face and 43 unidentified combined transplants. One thousand seven hundred sixty four (63.63%) patients were male. Calcineurin inhibitor (CNI) and corticosteroids were used in the majority of the patients at the time of diagnosis, 91%, (95% Confidence Interval [CI], 87.6%-94.3%, REM, I<sup>2</sup> 86.4%) and 70.1%, (95% CI, 61.5%-78.7%, REM, I<sup>2</sup> 96.2%) respectively. Antimetabolite, such as azathioprine and mycophenolate mofetil, were used in 76.9% (95% CI, 71.3%-82.4%, REM, I<sup>2</sup> 87.6%). Time from transplant to diagnosis of SARS-CoV-2 widely varied from 4 days to 31 years. Of note, no donor derived SARS-2 COVID infection was reported.

3.2. Symptoms with COVID-19 in solid organ transplant recipients

At first, we assessed the most common presenting symptoms among SOT recipients with COVID-19. A summary is shown in table 1. Of note, pooled incidence of fever, shortness of breath, cough and diarrhea were 70.2% (95% CI, 61.0%-79.5%, REM, I<sup>2</sup> 96.2%), 49.1% (95% CI, 41.1%-57.1%, REM, I<sup>2</sup> 92.1%), 63.8% (95% CI, 55.0%-72.6%, REM, I<sup>2</sup> 94.4%) and 30.4%

(95% CI, 23.6%-37.2%, REM, I<sup>2</sup> 85.4%), respectively (Fig. 2A-2D) [16,18,20,22,24,31,38,40,43,51,55,56,59,60,65,69,70,79,81,85, 88,96, 100,102,108,117,133,136,141-143,149,151,152,156,158,162,166,179, 185,189,190,192,210,212,216,217,219,228,229,231].

3.3. Laboratory results and radiologic findings

We then analyzed laboratory results at and after diagnosis of COVID-19. Leukopenia (white blood cell count <4000/mm<sup>3</sup>) was noted in 19 studies with pooled incidence of 17.8% (95% CI 10.7%-24.8%, REM, I<sup>2</sup> 57.2%) [24,31,40,51,88,96,108,133,135,143,151,152,158,162,166,216, 217,219,229]. C-reactive protein (CRP) level was documented in 161 studies [8,14,16-19,21,23-25,31-38,40,42-48,50-52,55,56,59,60,62, 63,66-69,71-73,76-82,84-89,92-98,100-104,106,108-110,112,115, 116,118-121,124-130,133-139,141,143,146,147,151-156,158,159, 161-167,169,171,174,177-179,185,187-192,194,196-200,203,204, 206,207,210-214,216,218-221,224-226,228-232,234-240]. Cravedi et. al. reported a median CRP level of 41.0 (interquartile range 11.5-125.3) mg/L amongst 144 patients [19], while Pereira et. al. reported a median CRP level of 68.5 (interquartile range 15.4-126) mg/L from among 90 patients [18].




Fig. 2B. Pooled incidence of shortness of breath showing the absolute risk and 95% confidence intervals of each study (n=36) [18,20,22,31,55,60,65,70,79,81,85,88, 96,102,108,117,133,136, 142,143,151,152,156,158,166,179,189,190,192,210,212,216,217,219,229,231] Abbreviations. ES; Effect Size, CI; Confidence Interval, \*=study period not defined.

Radiological abnormalities on chest imaging, either radiography or computerized tomography, was found in 28 studies with pooled incidence of 79.7% (95% CI 69.5%-89.9%, REM, I<sup>2</sup> 96.0%) [22,31,59,65,79, 81,85,96,102,114,117,133,143,152,156,158,166,179,189,190,192,210, 212,216,217,219,229,231] (Fig. 3).

3.4. Treatment method

We also analyzed the different treatment modalities used in SOT patients with COVID 19 and findings are summarized in table 2 [18,20,22,24,31,55,59,60,65,70,79,81,85,88,96,102,108,114,117,133, 136,142, 143,151,156,158,166,179,183,189,192,210,212,216,219,231]. Immunosuppressive medications, especially antimetabolites, were decreased in 76.2% (95% CI, 68.4%-84.1%, REM, I<sup>2</sup> 91.9% ,Fig. 4A) [18,21,22,55,59,79,85,88, 108,114,117,133,135,143,151,156,166,179, 183,189,190,192,212,216,217,229,231]. On the other hand, reduction in CNI highly varied from 0%-100% with pooled incidence of 38.7% (95% CI, 29.0%-48.4%, REM, I<sup>2</sup> 93.0, Fig. 4B) [16,18,21,23,24,38,40,51, 55,69,81,85,88,100,114,117,128,133,136,143,151,166,179,183,189,

212,216,217,228,229,231]. Corticosteroids were very seldom reduced in 1.7% (95% CI, 0.1%-3.5%, FEM, I<sup>2</sup> 0.0%, Fig. 4C) [18,21,51,59,69,88, 100,114,128,133,151,166,192,212,216,217,229].

Majority of cases were treated with hydroxychloroquine (59.5% 95% CI, 45.0%-74.0%, REM, I<sup>2</sup> 98.7%, supplemental Fig. A) [18,20,22, 31,55, 59,60,65,70,81,85,88,96,102,108,114,117,133,143,151,158,166,183, 189,190, 192,210,212,216,219]. Forty eight percent of cases were also placed on azithromycin (95%CI, 35.2%-62.0%, REM, I<sup>2</sup> 97.4%, supplemental Fig. B) [18,20,22,55, 59,60,65,70,88, 96,102,114,158,166, 179,183, 189,190,210,216,219]. Interleukin Six (IL-6) antagonists were used in 14.9% (95% CI, 9.9%-19.9%, REM, I<sup>2</sup> 86.7% supplemental Fig. C) [18,20, 22,55,60,70,79,85,88,117,133,136,189,192,210]. Protease inhibitors were used in 28.8% (95% CI, 19.1%-38.5%, REM, I<sup>2</sup> 82.8% supplemental Fig. D) [20,24,31,56,70,108,143,151,189,210,212]. Use of high dose steroids, [defined as prednisone at least 1mg/kg, hydrocortisone at least 100mg/day, methylprednisolone at least 200mg/daily, dexamethasone (>2mg/day)], was reported in 19 studies with a pooled incidence of 38.9% (95% CI, 27.9%-49.8%, REM, I<sup>2</sup> 97.0% ,supplemental Fig. E) [18,20,22,55,60,70,81,108,117,133,142,156,158,179,183,189,192, 210,




Fig. 2C. Pooled incidence of cough showing the absolute risk and 95% confidence intervals of each study (n=38) [18,20,22,31,55,59,60,65,70,79,81,85,88,96,102,108,114,117,133,136, 141,143,151,152,156,158,166,179,189,190,192,210,212,216,217,219,229,231]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \* =study period not defined.

231]. Concurrent antibiotic use was seen in 69.4% (95% CI, 58.9%-79.9%, REM, I<sup>2</sup> 94.9%) [16,24,32,40,50,51,55,60,79, 81,102,108,136,151, 162,166,179,185,190,199,210,212,216,219,229].

Fourteen studies, excluding potential duplicates, documented use of remdesivir with a total of 36 unique patients [18,22,50,69,70,114, 127,133,136, 165,183,217,228,240], of which 9 studies documented mortality with 2 deaths amongst 14 patients [50,69,114,127,133,136, 165,217,228]. Thirteen studies, excluding potential duplicates, mentioned use of convalescent plasma with a total of 33 unique patients [24,33,46,69,90,110,114,133,136,165,185,199,228], of which 11 studies documented about mortality with a total of 4 deaths from among 31 patients [24,33,46,69,90,110,114,133,165,185,228].

3.5. Outcomes

Finally, we analyzed outcomes in SOT recipients post COVID-19 infection.

Twenty two studies showed 81.0% pooled incidence of hospital admission (95% CI 75.2%-86.7%, REM, I<sup>2</sup> 91.4%) [18,20,22, 55,60, 65,70, 79,81,85,88,96,102,108,114,136,141,156, 166,190, 212,216], while 30

studies showed a 29.3% pooled incidence (95% CI 22.6%-36.1%, REM, I<sup>2</sup> 89.2%) of admission to the ICU from amongst hospital admissions [18,20,22,31,59,60,70,79,81,85,88,96, 102,108,114, 117,133, 143,151, 152,158,166,179,183,189,190,210,212,216,229]. Overall, 25.9% of patients required mechanical ventilation (95% CI 20.1%-31.8%, REM, I<sup>2</sup> 74.4%) from among hospital admissions [18,20,22,31,59,60,65, 70,79,81,85,88,96,102,108,114,117,133,142,143, 151,152,158,166, 179, 183,190,192,212,216,217,219,229,231].

Thirty-four studies addressed secondary or co-infections [22,24, 50,51,89,91,92,96,105,106, 110,127,133,143,145,151,157,158, 162,163, 165,166,177,182,185,189-191,194,199,200,212,217,219]. The majority of reported cases were bacterial urinary tract infection in kidney transplant recipients [89,96,151,165,166,212]. Of note, Nair et. al documented that 3 recipients, out of 10 kidney transplant patients, developed positive urine culture (*Enterococcus*, *Klebsiella* and *E.coli*) [166] and while Goss et. al report a culture with Group B streptococcus [96]. Respiratory secondary bacterial infections due to gram negative bacteria, including *Escherichia coli*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Klebsiella pneumoniae*, *Morganella morganii*, *Stenotrophomonas maltophilia*, as well as gram positive bacteria, including *Enterococcus faecalis*,




Fig. 2D. Pooled incidence of diarrhea showing the absolute risk and 95% confidence intervals of each study (n=36) [16,18,24,31,38,40,43,51,55,65,69,79,85,96,100,117,133,136, 141,142,149,151,158,162, 166,179,185,189,190,192,212,216,217,219,228,231]. Abbreviations. ES: Effect Size, CI: Confidence Interval, \*=study period not defined.






Fig. 3. Pooled incidence of abnormal chest imaging (either radiography or computerized tomography) in solid organ transplant recipients and COVID-19, showing the absolute risk and 95% confidence intervals of each study (n=28)[22,31,59,65,79,81,85,96,102,114,117,133,143,152,156,158,166,179,189,190,192,210,212,216,217,219,229,231]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \* =study period not defined.

*Staphylococcus aureus*, *Streptococcus haemolyticus* were also noted [22,50,92,127,157,169,177,185,219][22,50]. Bacteremia was reported in 4 cases with different pathogens [50,91,133,143]: 2 patients with *Klebsiella pneumoniae*, *Staphylococcus epidermidis* and two patients with *Staphylococcus aureus* [143]. Clostridium difficile colitis was reported in three patients [177,185].

Regarding viral secondary infection we six reports of Cytomegalovirus infection [50,51,143,158,177,190]. Molaei et. al reports four patients having COVID-19 and CMV co-infection with a viremia of 412–592 IU/mL [158], on the other hand, Pereira et. al also reports five patients with CMV with a median of 1469 (1326–8994) IU/mL [177]

while Pierrotti et. al reports 9 patients with CMV viremia [179]. Only one case of Influenza A co-infection was reported [200]. Two other patients had co-infections with HHV-6 and EBV [177]. There is also a report on a patient with a recent history of slight BK viremia that largely increased after COVID-19 infection [145].

Fungal secondary infections were also reported. Candidemia was reported in 1 liver transplant recipient [106] and other solid organ transplant patients with *Candida glabrata* candidemia [177] and *Candida krusei* bacteremia [185] while Rinaldi et al. reported 3 invasive candidiasis as well as 12 pulmonary aspergillosis [189]. Morlacchi et. al. reported on 4 lung transplant recipients with a bronchoalveolar lavage from 1 patient with *Aspergillus fumigatus* [163] while Pereira et. al reported another solid organ transplant recipient with aspergillus infection [177]. Also, Roberts et. al. reported 2 fungal infections with *Aspergillus versicolor* and severe cutaneous candidiasis [191]. Finally, Kates et. al. reported a pneumonia due to cryptococcus species as well as a patient with Pneumocystis pneumonia [22].

There are 62 studies documented about the duration of viral shedding and time to negativity of RT-PCR on NP specimens [1,15,24, 35,36,39,45,47,55,60,61,66,67,70,72,75,80,91,92,96–98,105,106,110, 113,115,118–122,125,130,134,136–138,140,147,148, 161–164, 170, 174,176,180,182,197,198,200,207,213, 218,220,221,226, 229,230, 232, 241] Majority of patients achieved negativity around 3 to 5 weeks from the initial positive result. The NP swab RT-PCR methods varied from

Table 2 Summary of treatment modalities with COVID 19 in Solid Organ Transplant Recipients.

Treatment Modalities	No. of Studies	Model	Pooled incidence (%) (95% CI)	Heterogeneity (I <sup>2</sup> , p value)
Hydroxychloroquine	30	REM	59.5 (45.0–74.0)	98.7, p<0.01
Azithromycin	21	REM	48.6 (35.2–62.0)	97.4, p<0.01
IL-6 antagonist	15	REM	14.9 (9.9–19.9)	86.7, p<0.01
Protease inhibitors	11	REM	28.8 (19.1–38.5)	82.8, p<0.01
Steroids	19	REM	38.9 (27.9–49.8)	97.0, p<0.01

Abbreviations: CI, confidence interval, REM, random effects model, IL-6, interleukin six

study to study and the time from onset of symptoms to first positive RT-PCR result was not assessed. However, we identified that SOT recipients achieved RT-PCR negativity on day 5 at the earliest [170] and day 56 at the latest [182] in this systematic review. No studies were found documenting viral culture of SARS-CoV-2 among SOT recipients.

Also, 52 articles documented about rejection after COVID-19 [18,22-24,31,40,43,50,65,69,70, 81,84,88,92,98,101,103-105, 108,114, 116,118,120,123,125,130,133-135, 143,145,154,165,171,173,175,179, 183,186,191,192,199,201,205,207,210,220,224,230,231][18, 22-24,65, 81, 84,92,98,101,103-105,108,116,118,120,123, 125,130,133-135, 143,173,175,186,191,192,199,201,210,224,230,231]. During the follow up period, the pooled incidence of rejection was calculated only at 1.0% (95% CI, 0.2%-1.7%, FEM, I<sup>2</sup> 0.0%) [22,23,65,81,88,108,114,133,135, 143,179,183,192,210,231].

Finally, we assessed all-cause mortality after COVID-19 in SOT recipients. We identified 37 articles and 18.6% died from any cause (95% CI, 14.8%-22.3%, REM, I<sup>2</sup> 72.4%) during their follow-up period [18,20, 22,31,55,59,60,65,70,79,81,85,88,96, 102,108,114,117,133,141-143, 151,152,156,158,166,179,189,190,192,210,212,216,219,229,231]

### 3.6. Kidney transplant recipients

We conducted the same analysis for the subset of kidney transplant recipients [16,18,20-24,31,32,38,43, 51,55,60,65,79,81,85,88, 96,100, 102, 114,136,142,143,149,151,152,156,158, 166,179,185,189,192, 210, 212,228,229,231]. Fever, cough, and shortness of breath were seen in

70.3% (95% CI, 62.5%-78.2%, REM, I<sup>2</sup> 90.6%), 65.3% (95% CI, 58.9%-71.7%, REM, I<sup>2</sup> 74.3%), and 49.3% (95% CI, 40.5%-58.2%, REM, I<sup>2</sup> 87.8%), respectively. Of note, acute kidney injury (AKI) in this group was seen in 45.4% (95% CI, 35.7%-55.1%, REM, I<sup>2</sup> 85.5%) [21,24,40,43, 51,60,81,85, 100,114,136,149,156,166,179,189,192,210,229,231] during entire follow-up. Anti-metabolites were reduced in 85.5% (95% CI, 79.4%-91.6%, REM, I<sup>2</sup> 84.0%), while change in CNI were seen in 48.2% (95% CI, 31.4%-65.0%, REM, I<sup>2</sup> 95.5%), and steroids only among 1.8% 95% CI, 0.6%-4.1%, FEM, I<sup>2</sup> 0.0%). In regards to treatment modalities, majority of cases were treated with hydroxychloroquine in 75.6% (95% CI, 62.3%-88.8%, REM, I<sup>2</sup> 97.1%), with azithromycin use among 50.8% (95% CI, 38.7%-62.8%, REM, I<sup>2</sup> 89.0%). IL-6 antagonists were used in 17.3% (95% CI, 10.0%-24.6%, REM, I<sup>2</sup> 88.1%), protease inhibitors in 37.7% (95% CI, 23.2%-52.2%, REM, I<sup>2</sup> 89.1%), with high dose steroids in 44.4% (95% CI, 27.3%-61.4%, REM, I<sup>2</sup> 96.5%). All-cause mortality in the subset of kidney transplant recipients was noted to be 22.0% (95% CI, 19.5%-24.4%, FEM, I<sup>2</sup> 47.1%).

### 3.7. Non-kidney solid organ transplant recipients

We also looked at similar outcomes in non-kidney transplant recipients. However, due to small sample size, stratification per organ type was difficult. In terms of all-cause mortality, a pooled incidence of 11.8% (95% CI, 4.2%-19.3%, REM, I<sup>2</sup> 73.4%) among liver transplant recipients [43,50,70,96,128,141,212,219] and a pooled incidence of 15.6%




Fig. 4A. Pooled incidence of reduction in anti-metabolite showing the absolute risk and 95% confidence intervals of each study (n=27)[18,20-23,55,59,79,81,108,117,133,135,143,156, 166,175,189-192,212,216,229,231]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \* =study period not defined.

(95% CI, 8.5%-22.7, FEM, I<sup>2</sup> 48.5%) among heart transplant recipients was seen [40,65,96,108,117,133,190,199].

**4. Discussion**

This is the first full systematic review and meta-analysis regarding COVID-19 in SOT recipients during the first wave of the pandemic in terms of symptoms, treatment strategy (including reduction in immunosuppression), and short-term outcomes. First, one fifth of this vulnerable population died after COVID-19 in a short term. Next, fever was common and reported in 70.2% but respiratory symptoms (cough and shortness of breath) were found in 63.8% and 49.1% respectively. Non-respiratory symptoms such as diarrhea were also seen in 30.4%. Also, we found that in the majority of SOT recipients, antimetabolites were either reduced or stopped completely.

In the general population, risk factors for severe COVID-19 infection have been well documented, such as hypertension, older age, and diabetes mellitus. Immunosuppressive medications have been thought to be a risk factor for severe illness from COVID-19 [6]. Even our meta-analysis showed a significantly higher admission rate (81.0%) as compared to the general population, which may not entirely reflect disease severity but rather the management strategy as employed by treating physicians who may prefer closer inpatient monitoring. Our meta-analysis showed incidence of lower respiratory tract infection (based on abnormal imaging), ICU admission rate and mortality of

79.7%, 29%, and 18.6% respectively, which are all comparable to the large published data in non-SOT, general population [242]. Even though T-cell immunity against respiratory viruses in SOT recipients is thought to be lower than the general population, viral clearance for SARS-CoV-2 in SOT recipients was around a median of 3-5 weeks, which again, is almost comparable to general population [242].

Similarly, as compared to oncology patients, which were also recognized as immunocompromised and high risk for severe COVID-19, our data showed comparable outcomes especially for hospital admission rate and mortality [243]. Even though this meta-analysis shows high incidence of mortality in SOT recipients when compared with the general population [242], it is important to note that these studies are not entirely comparable given the differences in associated comorbidities within the cohorts, such as incidence of diabetes or obesity.

On the other hand, hematopoietic stem cell transplant (HSCT) recipients are treated with immunosuppressive medications as well, and sometimes even at higher doses as compared to SOT recipients. Few reports have been published regarding COVID-19 in HSCT population and Malard et. al. documented a mortality of around 40%[244], which is significantly higher than SOT recipients.

In general, secondary infection was reported and found to be rare. In our study, some authors reported bacterial infection, especially urine culture positivity in kidney transplant recipients [89,151,165,166,212]. We cannot conclude if these infections occurred due to COVID-19 or were simply hospital acquired. Six reports of Cytomegalovirus infection




Fig. 4B. Pooled incidence of reduction in calcineurin inhibitor showing the absolute risk and 95% confidence intervals of each study (n=31)[16,18,20,21,23,24,38,51,55,79,81,117,128,133,143,151,166,175,189,191,212,216,228,229,231]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \* =study period not defined.




Fig. 4C. Pooled incidence of reduction in steroid showing the absolute risk and 95% confidence intervals of each study (n=17)[18,21,23,51,81,100,128,151,166,191,192,212,216,229]. Abbreviations. ES; Effect Size, CI; Confidence Interval, \*=study period not defined.

were found in our study [50,51,143,158,177,190]. In general, *Aspergillus* co-infection was also reported, Rinaldi et. al reported twelve pulmonary aspergillosis [189], Pereira et.al reported also a patient with aspergillus infection [177], while other authors report a case of a lung transplant recipient with a positive bronchoalveolar lavage for *Aspergillus fumigatus* [163] and a case of *Aspergillus versicolor* [191]. Even though SOT recipients, especially lung transplant, are at highest risk for Aspergillosis, we found a relatively lower rate of Aspergillosis after COVID-19. This may be due to a short follow-up period or use of post-lung transplant antimold prophylaxis, even prior to COVID-19.

This systematic review and meta-analysis has several limitations. First, we found significant heterogeneity between studies. This may be in part due to a wide variety of different transplant types, immunosuppressive agents, and a lack of uniform treatment of COVID-19. Definition of certain clinical presentations, such as fever, also varied from study to study, giving rise to further heterogeneity. Even after factoring in heterogeneity into our analyses, we could not completely overcome it. Secondly, we could only perform analyses in the subset of kidney transplant recipients. Due to sample size limitations, we could not perform the same for other transplant types. Next, the follow-up period varied significantly from study to study and that prevented us from capturing certain outcomes, such as rate of rejection, allograft dysfunction, secondary infection, and even mortality. Finally, it is important to note that this systematic review and meta-analysis reflects our initial experience with this ever rapidly growing pandemic, and as treatment modalities and strategies evolve in the coming months, this is likely to change.

Moreover, differences in the available resources, epidemiology and demographics, from country to country, also play an important factor when it comes to the management of these highly immunosuppressed patients.

In conclusion, we performed a systematic review and meta-analysis of COVID-19 in SOT recipients regarding symptoms, treatment options, and outcomes. Based on this review and meta-analysis, we conclude that a higher admission rate was noted but overall outcome was similar to the general population. Further studies are still required to assess the long-term outcomes including allograft rejection, secondary infections and mortality in this highly immunocompromised patient population.

**Funding**

The authors declare no-funding for this study.

**Disclosure**

Authors of this manuscript do not have any conflicts of interest to disclose.

**Authorship**

M.R, MAM, APV, J.R. and Y.N. performed the literature search. M.R., MAM, V.K. and Y.N. performed the data analysis. All authors were responsible for the study design, data interpretation, and writing.

## Declaration of Competing Interest

The authors of this manuscript declare no competing conflicts of interest.

## Acknowledgement

The authors would like to thank Tara Brigham, MLIS, from the Mayo Clinic Libraries in Jacksonville, FL, for peer review of the Medline database search strategy.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.trre.2020.100588>.

## References

- [1] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- [2] Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. N Engl J Med 2020;382:727–33.
- [3] Coronavirus disease (COVID-19). Situation Report. World Health Organization; 2020.
- [4] Kumar D, Ferreira VH, Blumberg E, Silveira F, Cordero E, Perez-Romero P, et al. A 5-Year Prospective Multicenter Evaluation of Influenza Infection in Transplant Recipients. *Clin Infect Dis* 2018;67:1322–9.
- [5] L'Huillier AG, Ferreira VH, Hirzel C, Nellmarla S, Ku T, Natori Y, et al. T-cell responses following Natural Influenza Infection or Vaccination in Solid Organ Transplant Recipients. *Sci Rep* 2020;10:10104.
- [6] Coronavirus Disease 2019 - People of Any Age with Underlying Medical Conditions, Centers for Disease Control and Prevention.
- [7] Cao X. COVID-19: immunopathology and its implications for therapy. *Nat Rev Immunol* 2020;20:269–70.
- [8] Guillen E, Pineiro GJ, Revuelta I, Rodriguez D, Bodro M, Moreno A, et al. Case report of COVID-19 in a kidney transplant recipient: Does immunosuppression alter the clinical presentation? *Am J Transplant* 2020;20.
- [9] Singh AK, Gillies CL, Singh R, Singh A, Chudasama Y, Coles B, et al. Prevalence of comorbidities and their association with mortality in patients with COVID-19: A Systematic Review and Meta-analysis. *Diabetes Obes Metab* 2020 [Online ahead of print].
- [10] Chan JW, Ng CK, Chan YH, Mok TY, Lee S, Chu SY, et al. Short term outcome and risk factors for adverse clinical outcomes in adults with severe acute respiratory syndrome (SARS). *Thorax* 2003;58:686–9.
- [11] Park JE, Jung S, Kim A, Park JE. MERS transmission and risk factors: a systematic review. *BMC Public Health* 2018;18:574.
- [12] Vishnevsky A, Levy M. Rethinking high-risk groups in COVID-19. *Mult Scler Relat Disord* 2020;42:102139.
- [13] Lai Q, Spoletini G, Bianco G, Graceffa D, Agnes S, Rossi M, et al. SARS-CoV2 and immunosuppression: A double-edged sword. *Transpl Infect Dis* 2020:e13404.
- [14] Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395:1033–4.
- [15] Zhu L, Xu X, Ma K, Yang J, Guan H, Chen S, et al. Successful recovery of COVID-19 pneumonia in a renal transplant recipient with long-term immunosuppression. *Am J Transplant* 2020;17:17.
- [16] Chaudhry ZS, Williams JD, Vahia A, Fadel R, Acosta TP, Prashar R, et al. Clinical Characteristics and Outcomes of COVID-19 in Solid Organ Transplant Recipients: A Case-Control Study. *Am J Transplant* 2020;20:3051–60.
- [17] Gandolfini I, Delsante M, Fiaccadori E, Zaza G, Manenti L, Degli Antoni A, et al. COVID-19 in kidney transplant recipients. *Am J Transplant* 2020;31:31.
- [18] M.R. Pereira, S. Mohan, D.J. Cohen, S.A. Husain, G.K. Dube, L.E. Ratner, S. Arcasoy, M.M. Aversa, L.J. Benvenuto, D.M. Dadhania, S. Kapur, L.M. Dove, R.S. Brown Jr, R.E. Rosenblatt, B. Samstein, N. Uriel, M.A. Farr, M. Satlin, C.B. Small, T.J. Walsh, R.P. Kodiyannplakkal, B.A. Miko, J.G. Aaron, D.S. Tsapepas, J.C. Emond, E.C. Verna, COVID-19 in solid organ transplant recipients: Initial report from the US epicenter, *Am J Transplant*, (n/a).
- [19] P. Cravedi, S.M. Suraj, Y. Azzi, M. Haverly, S. Farouk, M.J. Pérez-Sáez, M.D. Redondo-Pachón, B. Murphy, S. Florman, L.G. Cyrino, M. Grafals, S. Venkataraman, X.S. Cheng, A.X. Wang, G. Zaza, A. Ranghino, L. Furian, J. Manrique, U. Maggiore, I. Gandolfini, N. Agrawal, H. Patel, E. Akalin, L.V. Riella, T.T.s. group, COVID-19 and Kidney Transplantation: Results from the TANGO International Transplant Consortium, *Am J Transplant*, (n/a).
- [20] M. Crespo, A. Mazuecos, E. Rodrigo, E. Gavela, F. Villanega, E. Sánchez-Alvarez, E. González-Monte, C. Jiménez-Martín, E. Melilli, F. Diekman, S. Zárraga, D. Hernández, J. Pascual, f.t.S.S.o.N.C.-. Group, Respiratory and Gastrointestinal COVID-19 Phenotypes in Kidney Transplant Recipients, *Transplantation*, (Online First (9000)).
- [21] A. Favà, D. Cucchiari, N. Montero, N. Toapanta, J. Centellas, A. Vila-Santandreu, A. Coloma, M. Meneghini, A. Manonelles, J. Sellarés, I. Torres, R. Gelpi, I. Lorenzo, P. Ventura-Aguar, F. Cofan, V. Torregrosa, M. Perelló, C. Facundo, D. Seron, F. Oppenheimer, O. Bestard, J. Cruzado, F. Moreso, E. Melilli, Clinical characteristics and risk factors for severe COVID-19 in hospitalized kidney transplant recipients: A multicentric cohort study, *Am J Transplant*, (n/a).
- [22] Kates OS, Haydel BM, Florman SS, Rana MM, Chaudhry ZS, Ramesh MS, et al. COVID-19 in solid organ transplant: A multi-center cohort study. *Clin Infect Dis* 2020.
- [23] Fernandez-Ruiz M, Andres A, Loinaz C, Delgado JF, Lopez-Medrano F, San Juan R, et al. COVID-19 in solid organ transplant recipients: A single-center case series from Spain. *Am J Transplant* 2020;16:16.
- [24] Fung M, Chiu CY, DeVoe C, Doernberg SB, Schwartz BS, Langelier C, et al. Clinical Outcomes and Serologic Response in Solid Organ Transplant Recipients with COVID-19: A Case Series from the United States. *Am J Transplant* 2020;01:01.
- [25] Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020;8:475–81.
- [26] TL JC Julian Higgins, Tovey David, Thomas James, Flemyng Ella, Churchill Rachel. Methodological Expectations of Cochrane Intervention Reviews (MECIR). *Cochrane Methods* 2020.
- [27] Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 2009;62:e1–34.
- [28] McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016;75:40–6.
- [29] L. Kristin, COVID-19 Literature Searches, in.
- [30] MEDLINE® 2020 Database Guide, in.
- [31] S. Abolghasemi, M. Mardani, S. Sali, N. Honarvar, M. Baziboroun, COVID-19 and kidney transplant recipients, *Transpl Infect Dis*, (n/a e13413).
- [32] Abrishami A, Samavat S, Behnam B, Arab-Ahmadi M, Nafar M, Sane'i Taheri M. Clinical Course, Imaging Features, and Outcomes of COVID-19 in Kidney Transplant Recipients. *Eur Urol* 2020;06:06.
- [33] Abuzeineh M, Muzaale AD, Crews DC, Avery RK, Brotman DJ, Brennan DC, et al. Telemedicine in the Care of Kidney Transplant Recipients With Coronavirus Disease 2019: Case Reports. *Transplant Proc* 2020;52(9):2620–5.
- [34] Adapa S, Konala VM, Naramala S, Daggubati SR, Koduri NM, Gayam V, et al. COVID-19 in Renal Transplant Patient Presenting With Active Typical Symptoms and Resolved Atypical Symptoms. *J Investig Med High Impact Case Rep* 2020;8(2324709620949307).
- [35] Adrogue AH, Mithani F, Ibrahim HN, Schwartz MR, Gaber L, Hebert SA, et al. A Kidney Transplant Recipient With Coronavirus Disease 2019: Utility of a Prognostication Score. *Transplant Proc* 2020;52:2620–5.
- [36] Aiello P, Johnson S, Ramos Mercado A, Hussein S. Pneumatosis intestinalis in a patient with COVID-19. *BMJ Case Rep* 2020;13.
- [37] Aigner C, Dittmer U, Kamler M, Collaud S, Taube C. COVID-19 in a lung transplant recipient. *J Heart Lung Transplant* 2020;39:610–1.
- [38] Akalin E, Azzi Y, Bartash R, Seethamraju H, Parides M, Hemmige V, et al. Covid-19 and Kidney Transplantation. *N Engl J Med* 2020;24:24.
- [39] Akdur A, Karakaya E, Ayvazoglu Soy EH, Alshalabi O, Kirnap M, Arslan H, et al. Coronavirus Disease (COVID-19) in Kidney and Liver Transplant Patients: A Single-Center Experience. *Exp Clin Transplant* 2020;18:270–4.
- [40] Al-Darzi W, Aurora L, Michaels A, Cowger J, Grafton G, Selektor Y, et al. Heart transplant recipients with confirmed 2019 novel coronavirus infection: The Detroit experience. *Clin Transplant* 2020:e14091.
- [41] Al-Makki A, Taber T. Neutropenic fever in COVID-19 in kidney transplant patient. *Rev Med Virol* 2020:e2156.
- [42] Alberici F, Delbarba E, Manenti C, Econimo L, Valerio F, Pola A, et al. A single center observational study of the clinical characteristics and short-term outcome of 20 kidney transplant patients admitted for SARS-CoV2 pneumonia. *Kidney Int* 2020;97:1083–8.
- [43] Ali T, Al-Ali A, Fajji L, Hammad E, Nazmi A, Alahmadi I, et al. Coronavirus Disease-19: Disease Severity and Outcomes of Solid Organ Transplant Recipients: Different Spectrum of Disease in Different Populations? *Transplantation* 2020.
- [44] Allam SR, Dao A, Madhira MM, Antiporta PB, Nair RR, Guiteau JJ, et al. Interleukin-6 receptor antagonist therapy to treat SARS-CoV-2 driven inflammatory syndrome in a kidney transplant recipient. *Transpl Infect Dis* 2020:e13326.
- [45] Ammirati E, Travi G, Orcese C, Sacco A, Auricchio S, Frigerio M, et al. Heart-Kidney Transplanted patient affected by COVID-19 pneumonia treated with tocilizumab on top of immunosuppressive maintenance therapy. *IJC Heart Vasc* 2020;29:100596.
- [46] Antony SJ, Singh J, de Jesus M, Lance J. Early use of tocilizumab in respiratory failure associated with acute COVID -19 pneumonia in recipients with solid organ transplantation. *IDCases* 2020;21:e00888.
- [47] Araten DJ, Belmont HM, Schaefer-Cuttillo J, Iyengar A, Mattoo A, Reddy R. Mild Clinical Course of COVID-19 in 3 Patients Receiving Therapeutic Monoclonal Antibodies Targeting C5 Complement for Hematologic Disorders. *Am J Case Rep* 2020;21:e927418.
- [48] Arenas MD, Crespo M, Perez-Saez MJ, Collado S, Redondo-Pachon D, Llinas-Mallol L, et al. Clinical Profiles in Renal Patients with COVID-19. *J Clin Med* 2020;9.
- [49] Arpali E, Akyollu B, Yelken B, Tekin S, Turkmen A, Kocak B. Case report: A kidney transplant patient with mild COVID-19. *Transpl Infect Dis* 2020:e13296.
- [50] M. Aversa, L. Benvenuto, M. Anderson, L. Shah, H. Robbins, M. Pereira, J. Scheffert, M. Carroll, J. Hum, M. Nolan, G. Reilly, P. Lemaître, B.P. Stanifer, F. D'Ovidio, J. Sonett, S. Arcasoy, T.C.U.L.T. Program, COVID-19 in lung transplant recipients: A single center case series from New York City, *Am J Transplant*, (n/a).
- [51] Banerjee D, Popoola J, Shah S, Ster IC, Quan V, Phanish M. COVID-19 infection in kidney transplant recipients. *Kidney Int* 2020;97:1076–82.

- [52] Bartiromo M, Borchi B, Botta A, Bagala A, Lugli G, Tilli M, et al. Threatening drug-drug interaction in a kidney transplant patient with Coronavirus Disease 2019 (COVID-19). *Transpl Infect Dis* 2020;12:12.
- [53] Becchetti C, Zambelli MF, Pasulo L, Donato MF, Invernizzi F, Detry O, et al. COVID-19 in an international European liver transplant recipient cohort. *Gut* 2020 (gutjnl-2020-321923).
- [54] Belli LS, Duvoux C, Karam V, Adam R, Cuervas-Mons V, Pasulo L, et al. COVID-19 in liver transplant recipients: preliminary data from the ELITA/ELTR registry. *Lancet Gastroenterol Hepatol* 2020;04:04.
- [55] I. Benotmane, G. Gautier Vargas, M.-J. Wendling, P. Perrin, A. Velay, X. Bassand, D. Bedo, C. Baldacini, M. Sagnard, D. Bozman, M. Della Chiesa, M. Solis, F. Gallais, N. Cognard, J. Olagne, H. Delagrèverie, L. Gontard, B. Panaget, D. Marx, F. Heibel, L. Braun, B. Moulin, S. Caillard, S. Fafi-Kremer, In-depth virological assessment of kidney transplant recipients with COVID-19, *Am J Transplant*, (n/a).
- [56] Benotmane I, Perrin P, Gautier Vargas G, Bassand X, Keller N, Lavaux T, et al. Biomarkers of Cytokine Release Syndrome Predict Disease Severity and Mortality From COVID-19 in Kidney Transplant Recipients. *Transplantation* 2020.
- [57] Bhoori S, Rossi RE, Citterio D, Mazzaferro V. COVID-19 in long-term liver transplant patients: preliminary experience from an Italian transplant centre in Lombardy. *Lancet Gastroenterol Hepatol* 2020;5:532-3.
- [58] Billah M, Santeusano A, Delaney V, Cravedi P, Farouk SS. A catabolic state in a kidney transplant recipient with COVID-19. *Transpl Int* 2020;01:01.
- [59] F. Bösch, N. Börner, S. Kemmer, C. Lampert, S. Jacob, D. Koliogiannis, M. Stangl, S. Michel, N. Kneidinger, C. Schneider, M. Fischeder, M. Irlbeck, G. Denk, J. Werner, M.K. Angele, M.O. Guba, Attenuated early inflammatory response in solid organ recipients with COVID-19, *Clin Transplant*, (n/a e14027).
- [60] N. Bossini, F. Alberici, E. Delbarba, F. Valerio, C. Manenti, S. Possenti, L. Econimo, C. Maffei, A. Pola, V. Terlizzi, C. Salviani, M. Moscato, S. Pasquali, N. Zambetti, M. Tonoli, S. Affatato, P. Pecchini, F. Battista Viola, F. Malberti, G. Depetri, M. Gaggiotti, F. Scolari, T.B.R.C.t. force, Kidney transplant patients with SARS-CoV-2 infection: the brescia renal COVID task force experience, *Am J Transplant*, (n/a).
- [61] Buscemi V, De Carlis R, Lauterio A, Merli M, Puoti M, De Carlis L. Does interval time between liver transplant and COVID-19 infection make the difference? *Dig Liver Dis* 2020.
- [62] Bush R, Johns F, Acharya R, Upadhyay K. Mild COVID-19 in a pediatric renal transplant recipient. *Am J Transplant* 2020;13:13.
- [63] Bussalino E, De Maria A, Russo R, Paoletti E. Immunosuppressive therapy maintenance in a kidney transplant recipient with SARS-CoV-2 pneumonia: A case report. *Am J Transplant* 2020;13:13.
- [64] Carey SA, Afzal A, Jamil A, Williams S, Gottlieb RL. Outpatient COVID-19 surveillance testing in orthotopic heart transplant recipients. *Clin Transplant* 2020; e14105.
- [65] Cavagna L, Seminari E, Zanframundo G, Gregorini M, Di Matteo A, Rampino T, et al. Calcineurin Inhibitor-Based Immunosuppression and COVID-19: Results from a Multidisciplinary Cohort of Patients in Northern Italy. *Microorganisms* 2020;8.
- [66] Chen S, Yin Q, Shi H, Du D, Chang S, Ni L, et al. A familial cluster, including a kidney transplant recipient, of Coronavirus Disease 2019 (COVID-19) in Wuhan, China. *Am J Transplant* 2020;03:03.
- [67] Cheng DR, Wen JQ, Liu ZZ, Lv TF, Chen JS. Coronavirus disease 2019 in renal transplant recipients: report of two cases. *Transpl Infect Dis* 2020;e13329.
- [68] Chenna A, Konala VM, Gayam V, Naramala S, Adapa S. Coronavirus Disease 2019 (COVID-19) in a Renal Transplant Patient. *Cureus* 2020;12.
- [69] Christensen J, Kumar D, Moinuddin I, Bryson A, Kashi Z, Kimball P, et al. Coronavirus Disease 2019 Viremia, Serologies, and Clinical Course in a Case Series of Transplant Recipients. *Transplant Proc* 2020;52:2637-41.
- [70] J. Colmenero, M. Rodríguez-Perálvarez, M. Salcedo, A. Arias-Milla, A. Muñoz-Serrano, J. Graus, J. Nuño, M. Gastaca, J. Bustamante-Schneider, A. Cacherro, L. Lladó, A. Caballero, A. Fernández-Yunquera, C. Loinaz, I. Fernández, C. Fondevila, M. Navasa, M. Iñarrairaegui, L. Castells, S. Pascual, P. Ramírez, C. Vinaixa, M.L. González-Díez, R. González-Grande, L. Hierro, F. Noguerras, A. Otero, J.M. Álamo, G. Blanco-Fernández, E. Fábrega, F. García-Pajares, J.L. Montero, S. Tomé, G. De la Rosa, J.A. Pons, Epidemiological pattern, incidence and outcomes of COVID-19 in liver transplant patients, *J Hepatol*.
- [71] P. Columbia University Kidney Transplant, Early Description of Coronavirus 2019 Disease in Kidney Transplant Recipients in New York, *J Am Soc Nephrol*, 31 (2020) 1150-1156.
- [72] Cozzi E, Faccioli E, Marinello S, Loy M, Congedi S, Calabrese F, et al. COVID-19 pneumonia in lung transplant recipients: report of two cases. *Am J Transplant* 2020;12:12.
- [73] Crespo M, Jose Perez-Saez M, Redondo-Pachon D, Llinas-Mallol L, Montero MM, Villar J, et al. COVID-19 in elderly kidney transplant recipients. *Am J Transplant* 2020;29:29.
- [74] Dahl Mathiasen V, Jensen-Fangel S, Skov K, Leth S. Uneventful case of COVID-19 in a kidney transplant recipient. *BMJ Case Rep* 2020;13:e237427.
- [75] Dajti E, Cristini F, Tamanini G, Cescon M, Bazzoli F, Tame M. COVID-19 in a young liver transplant recipient: caution for drug-drug interactions. *J Gastrointest Liver Dis* 2020;29:470.
- [76] De Francesco MA, Alberici F, Bossini N, Scolari F, Pascucci F, Tomasoni G, et al. Pneumocystis jirevicii and SARS-CoV-2 Co-Infection: A Common Feature in Transplant Recipients? *Vaccines (Basel)* 2020;8.
- [77] De Gottardi A, Fratila C, Bertoli R, Cerny A, Magenta L, Gianella P, et al. Clinical characteristics and management of a liver transplanted patient admitted with SARS-CoV-2 infection. *Clin Res Hepatol Gastroenterol* 2020.
- [78] Decker A, Welzel M, Laubner K, Grundmann S, Kochs G, Panning M, et al. Prolonged SARS-CoV-2 shedding and mild course of COVID-19 in a patient after recent heart transplantation. *Am J Transplant* 2020;09:09.
- [79] E. Demir, M. Uyar, E. Parmaksiz, A. Sinangil, B. Yelken, A.B. Dirim, O. Merhametsiz, S. Yadigar, Z. Atan Ucar, A.R. Ucar, M.E. Demir, M. Mese, E.B. Akin, N. Garayeva, S. Safak, O.A. Oto, H. Yazici, A. Turkmen, COVID-19 in kidney transplant recipients: A multicenter experience in Istanbul, *Transpl Infect Dis*, (n/a e13371).
- [80] N. Desmazes-Dufeu, B. Coltey, L. Amari, M. Gouitaa, C. Touzery, M. Reynaud-Gaubert, P. Chanez, N. Cassir, Discordant courses of COVID-19 in a cohabiting couple of lung transplant recipients, *Transpl Infect Dis*, (n/a e13410).
- [81] A. Devresse, L. Belkhir, B. Vo, B. Ghaye, A. Scohy, B. Kabamba, E. Goffin, J. De Greef, M. Mourad, M. De Meyer, J.-C. Yombi, N. Kanaan, COVID-19 Infection in Kidney Transplant Recipients: A Single-Center Case Series of 22 Cases From Belgium, *Kidney Medicine*
- [82] Dirim AB, Demir E, Ucar AR, Garayeva N, Safak S, Oto OA, et al. Fatal SARS-CoV-2 infection in a renal transplant recipient. *CEN Case Reports* 2020;409-12.
- [83] Doevelaar AAN, Holzer B, Seibert FS, Bauer F, Stervbo U, Rohn BJ, et al. Lessons for the clinical nephrologist: recurrence of nephrotic syndrome induced by SARS-CoV-2. *J Nephrol* 2020.
- [84] Dube GK, Husain SA, McCune KR, Sandoval PR, Ratner LE, Cohen DJ. COVID-19 infection in pancreas transplant recipients. *Transpl Infect Dis* 2020;e13359.
- [85] Elias M, Pievani D, Randoux C, Louis K, Denis B, Delion A, et al. COVID-19 Infection in Kidney Transplant Recipients: Disease Incidence and Clinical Outcomes. *J Am Soc Nephrol* 2020;31:2413-23.
- [86] Faguer S, Del Bello A, Abravanel F, Nicolau-Travers ML, Kamar N. Tocilizumab for Hemophagocytic Syndrome in a Kidney Transplant Recipient With COVID-19. *Ann Intern Med* 2020;18:18.
- [87] Farfour E, Picard C, Beaumont L, Lesprit P, Ballester MC, Ackermann F, et al. COVID-19 in lung-transplanted and cystic fibrosis patients: Be careful. *J Cyst Fibros* 2020; 10:10.
- [88] Felldin M, Softeland JM, Magnusson J, Ekberg J, Karason K, Schult A, et al. Initial Report From a Swedish High-Volume Transplant Center After The First Wave of The COVID-19 Pandemic. *Transplantation* 2020.
- [89] Fontana F, Alfano G, Mori G, Amurri A, Tei L, Ballestri M, et al. COVID-19 pneumonia in a kidney transplant recipient successfully treated with tocilizumab and hydroxychloroquine. *Am J Transplant* 2020;23:23.
- [90] Fung M, Nambiar A, Pandey S, Aldrich JM, Teraoka J, Freise C, et al. Treatment of Immunocompromised COVID-19 patients with Convalescent Plasma. *Transpl Infect Dis* 2020.
- [91] K. Gajurel, Persistently positive severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) nasopharyngeal PCR in a kidney transplant recipient, *Transpl Infect Dis*, (n/a e13408).
- [92] Gao F, Zheng KI, Gu JY, George J, Zheng MH. COVID-19 and liver transplantation: Lessons learned from three reported cases. *Transpl Infect Dis* 2020;e13335.
- [93] Gautier-Vargas G, Baldacini C, Benotmane I, Keller N, Perrin P, Moulin B, et al. Rapid resolution of cytokine release syndrome and favorable clinical course of severe COVID-19 in a kidney transplant recipient treated with tocilizumab. *Kidney Int* 2020;04:04.
- [94] E. Giorgakis, S.P. Zehetaban, A.E. Stevens, S. Bhusal, L. Burdine, COVID-19 in solid organ transplant recipients, *Transpl Infect Dis*, (n/a e13419).
- [95] Goodlet KJ, Bansal S, Arjuna A, Nailor MD, Buddhdev B, Abdelrazek H, et al. COVID-19 in a lung transplant recipient: Exploring the diagnostic role of circulating exosomes and the clinical impact of advanced immunosuppression. *Transpl Infect Dis* 2020;e13480.
- [96] Goss MB, Galvan NTN, Ruan W, Munoz FM, Brewer ED, O'Mahony CA, et al. The Pediatric Solid Organ Transplant Experience with COVID-19: An Initial Multi-Center, Multi-Organ Case Series. *Pediatr Transplant* 2020;e13868.
- [97] Guella A, Elfadil O, Abdulrazaq G, Osman S, Khan MM, Ahmed A, et al. Favorable Outcome of COVID-19 Pneumonia in a Kidney Transplant Recipient Treated with Tocilizumab. *Case Rep Infect Dis* 2020;2020:8830626.
- [98] Hammami MB, Garibaldi B, Shah P, Liu G, Jain T, Chen PH, et al. Clinical course of COVID-19 in a liver transplant recipient on hemodialysis and response to tocilizumab therapy: A case report. *Am J Transplant* 2020;02:02.
- [99] Hann A, Lembach H, McKay SC, Perrin M, Isaac J, Oo YH, et al. Controversies Regarding Shielding and Susceptibility to COVID-19 Disease in Liver transplant Recipients in the United Kingdom. *Transpl Infect Dis* 2020;e13352.
- [100] S. Hartzell, S. Bin, C. Benedetti, M. Haverly, L. Gallon, G. Zaza, L.V. Riella, M.C. Menon, S. Florman, A.H. Rahman, J.M. Leech, P.S. Heeger, P. Cravedi, Evidence of potent humoral immune activity in COVID-19-infected kidney transplant recipients, *Am J Transplant*, (n/a).
- [101] N. Heinz, A. Griesemer, J. Kinney, J. Vittorio, S.M. Lagana, D. Goldner, M. Velasco, T. Kato, S. Lobritto, M. Martinez, A case of an Infant with SARS-CoV-2 hepatitis early after liver transplantation, *Pediatr Transplant*, (n/a e13778).
- [102] Hoek RAS, Manintveld OC, Betjes MGH, Hellemons ME, Seghers L, van Kampen JAA, et al. Covid-19 in solid organ transplant recipients: A single center experience. *Transpl Int* 2020;27:27.
- [103] Holzhauser L, Lourenco L, Sarswat N, Kim G, Chung B, Nguyen AB. Early Experience of COVID-19 in Two Heart Transplant Recipients: Case Reports and Review of Treatment Options. *American journal of transplantation* 2020.
- [104] Hsu JJ, Gaynor P, Kamath M, Fan A, Al-Saffar F, Cruz D, et al. COVID-19 in a High-Risk Dual Heart and Kidney Transplant Recipient. *Am J Transplant* 2020;21:21.
- [105] Huang J, Lin H, Wu Y, Fang Y, Kumar R, Chen G, et al. COVID-19 in posttransplant patients-report of 2 cases. *Am J Transplant* 2020;03:03.
- [106] Huang JF, Zheng KI, George J, Gao HN, Wei RN, Yan HD, et al. Fatal outcome in a liver transplant recipient with COVID-19. *Am J Transplant* 2020;10:10.
- [107] Husain SA, Dube G, Morris H, Fernandez H, Chang JH, Paget K, et al. Early Outcomes of Outpatient Management of Kidney Transplant Recipients with Coronavirus Disease 2019. *Clin J Am Soc Nephrol* 2020;15:1174-8.

- [108] Iacovoni A, Boffini M, Pidello S, Simonato E, Barbero C, Sebastiani R, et al. A case series of Novel-Coronavirus infection in heart transplantation from two centers in the pandemic area in the North of Italy. *J Heart Lung Transplant* 2020;39:1081–8.
- [109] Jang K, Khatri A, Majure DT. COVID-19 Leading to Acute Encephalopathy in a Heart Transplant Patient. *J Heart Lung Transplant* 2020;39:853–5.
- [110] J, Jiang, Y. Miao, Y. Zhao, X. Lu, P. Zhou, X. Zhou, Z. Chen, D. Du, Convalescent plasma therapy: Helpful treatment of COVID-19 in a kidney transplant recipient presenting with severe clinical manifestation and complex complications, *Clin Transplant*, (n/a e14025).
- [111] Johnson KM, Belfer JJ, Peterson GR, Boelkins MR, Dumkow LE. Managing COVID-19 in Renal Transplant Recipients: A Review of Recent Literature and Case Supporting Corticosteroid-sparing Immunosuppression. *Pharmacotherapy* 2020;40:517–24.
- [112] Kadosh BS, Pavone J, Wu M, Reyentovich A, Gidea C. Collapsing Glomerulopathy Associated with COVID-19 Infection in a Heart Transplant Recipient. *J Heart Lung Transplant* 2020;39:855–7.
- [113] Kates OS, Fisher CE, Stankiewicz-Karita HC, Shepherd AK, Church EC, Kapnadak SG, et al. Earliest cases of coronavirus disease 2019 (COVID-19) identified in solid organ transplant recipients in the United States. *Am J Transplant* 2020;24:24.
- [114] Katz-Greenberg G, Yadav A, Gupta M, Martinez-Cantarin MP, Gulati R, Ackerman L, et al. Outcomes of COVID-19-positive kidney transplant recipients: A single-center experience. *Clin Nephrol* 2020;94.
- [115] Keller BC, Le A, Sobhanie M, Colburn N, Burcham P, Rosenheck J, et al. Early COVID-19 infection after lung transplantation. *Am J Transplant* 2020;29:29.
- [116] Kemmner S, Guba MO, Schonermarck U, Stangl M, Fischereder M. Cyclosporine as a preferred calcineurin inhibitor in renal allograft recipients with COVID-19 infection. *Kidney Int* 2020;98.
- [117] Ketcham SW, Adie SK, Malliett A, Abdul-Aziz AA, Bitar A, Grafton G, et al. Coronavirus Disease-2019 in Heart Transplant Recipients in Southeastern Michigan: A Case Series. *J Card Fail* 2020;14:14.
- [118] Kim Y, Kwon O, Paek JH, Park WY, Jin K, Hyun M, et al. Two distinct cases with COVID-19 in kidney transplant recipients. *Am J Transplant* 2020;26:26.
- [119] Kocak B, Arpali E, Akyollu B, Yelken B, Tekin S, Kanbay M, et al. A Case Report of Oligosymptomatic Kidney Transplant Patients with COVID-19: Do They Pose a Risk to Other Recipients? *Transplant Proc* 2020;15:15.
- [120] Koczulla RA, Szczepanski B, Kotecki A, Kuhnert S, Hecker M, Askevold I, et al. SARS-CoV-2 infection in two patients following recent lung transplantation. *Am J Transplant* 2020;12:12.
- [121] A. Kolonko, S. Dudzicz, A. Wiecek, R. Król, COVID-19 infection in solid organ transplant recipients: A single-center experience with patients immediately after transplantation, *Transpl Infect Dis*, (n/a e13381).
- [122] Kumar RN, Tanna SD, Shetty AA, Stosor V. COVID-19 in an HIV-positive Kidney Transplant Recipient. *Transpl Infect Dis* 2020:e13338.
- [123] Lagana SM, De Michele S, Lee MJ, Emond JC, Griesemer AD, Tulin-Silver SA, et al. COVID-19 Associated Hepatitis Complicating Recent Living Donor Liver Transplantation. *Arch Pathol Lab Med* 2020;17:17.
- [124] Latif F, Farr MA, Clerkin KJ, Habal MV, Takeda K, Naka Y, et al. Characteristics and Outcomes of Recipients of Heart Transplant With Coronavirus Disease 2019. *JAMA Cardiol* 2020;13:13.
- [125] Lauterio A, Valsecchi M, Santambrogio S, De Carlis R, Merli M, Calini A, et al. Successful recovery from severe COVID-19 pneumonia after kidney transplantation: The interplay between immunosuppression and novel therapy including tocilizumab. *Transpl Infect Dis* 2020:e13334.
- [126] H. Lazareth, H. Péré, Y. Binois, M. Chabannes, J. Schurder, T. Bruneau, A. Karras, E. Thervet, M. Rabant, D. Veyer, N. Pallet, COVID-19-related Collapsing Glomerulopathy in a Kidney Transplant Recipient, *Am J Kidney Dis*
- [127] Le MP, Le Hingrat Q, Jaquet P, Wicky PH, Bunel V, Massias L, et al. Removal of Remdesivir's Metabolite GS-441524 by Hemodialysis in a Double Lung Transplant Recipient with COVID-19. *Antimicrob Agents Chemother* 2020;64.
- [128] Lee BT, Perumalswami PV, Im GY, Florman S, Schiano TD. COVID-19 in Liver Transplant Recipients: An Initial Experience from the U.S. Epicenter. *Gastroenterology* 2020;19:19.
- [129] H. Lee, B.S. Mantell, M.E. Richmond, S.P. Law, W.A. Zuckerman, L.J. Addonizio, T.M. Lee, I.D. Lytrivi, Varying Presentations of COVID-19 in Young Heart Transplant Recipients: A Case Series, *Pediatr Transplant*, (n/a e13780).
- [130] Li F, Cai J, Dong N. First cases of COVID-19 in heart transplantation from China. *J Heart Lung Transplant* 2020;39:496–7.
- [131] Li H, Liu L, Zhang D, Xu J, Dai H, Tang N, et al. SARS-CoV-2 and viral sepsis: observations and hypotheses. *Lancet* 2020;395:1517–20.
- [132] Li QY, Cheng Q, Zhao ZL, Dai NN, Zeng L, Zhu L, et al. Severe acute respiratory syndrome coronavirus 2 infection in renal transplant recipients: A case report. *Health sciences* 2020;52:780–4.
- [133] B. Lima, G.T. Gibson, S. Vullaganti, K. Malhame, S. Maybaum, S.T. Hussain, S. Shah, D.T. Majure, F. Wallach, K. Jang, V. Bijol, M.J. Esposito, A.K. Williamson, R.M. Thomas, T.A. Bhuiya, H.A. Fernandez, G.R. Stevens, COVID-19 in recent heart transplant recipients: Clinicopathologic features and early outcomes, *Transpl Infect Dis*, (n/a e13382).
- [134] Liu B, Wang Y, Zhao Y, Shi H, Zeng F, Chen Z. Successful treatment of severe COVID-19 pneumonia in a liver transplant recipient. *Am J Transplant* 2020;03:03.
- [135] C. Loíñez, A. Marcauczo, M. Fernández-Ruiz, O. Caso, F. Cambra, R. San Juan, I. Justo, J. Calvo, A. García-Sesma, A. Manrique, M.A. Pérez-Jacoiste Asín, M.D. Folgueira, J.M. Aguado, C. Lumberras, Varied clinical presentation and outcome of SARS-CoV-2 infection in liver transplant recipients: Initial experience at a single center in Madrid, Spain, *Transpl Infect Dis*, (n/a e13372).
- [136] Lubetzky M, Aull MJ, Craig-Schapiro R, Lee JR, Marku-Podvorica J, Salinas T, et al. Kidney allograft recipients, immunosuppression, and coronavirus disease-2019: a report of consecutive cases from a New York City transplant center. *Nephrol Dial Transplant* 2020;35:1250–61.
- [137] Ma BM, HUNG IFN, Chan GCW, Tam AR, Chan SSK, Wong BCK, et al. Case of "relapsing" COVID-19 in a kidney transplant recipient. *Nephrology (Carlton)* 2020.
- [138] Machado DJB, Ianhez LE. COVID-19 pneumonia in kidney transplant recipients-Where we are? *Transpl Infect Dis* 2020:e13306.
- [139] Malekhosseini SA, Nikoupour H, Gholami S, Shamsaeefar A, Arasteh P, Kazemi K, et al. A Report of 85 Cases of COVID-19 and Abdominal Transplantation From a Single Center: What Are the Associated Factors With Death Among Organ Transplantation Patients. *Transplantation* 2020.
- [140] Man Z, Jing Z, Huibo S, Bin L, Fanjun Z. Viral shedding prolongation in a kidney transplant patient with COVID-19 pneumonia. *Am J Transplant* 2020;13:13.
- [141] Mansoor E, Perez A, Abou-Saleh M, Sclair SN, Cohen S, Cooper GS, et al. Clinical Characteristics, Hospitalization and Mortality Rates of COVID-19 Among Liver Transplant Patients in the United States: A Multi-Center Research Network Study. *Gastroenterology* 2020.
- [142] Marcault C, Fodil S, Dupont T, Darmon M, Azoulay E. In response to the letter of Montagud-Marrahi et al. *Am J Transplant* 2020;08:08.
- [143] F. Maritati, E. Cerutti, L. Zuccatosta, A. Fiorentini, C. Finale, M. Ficoeoco, F. Cristiano, A. Capestro, E. Balestra, D. Tarascia, M. Vivarelli, A. Donati, G.P. Perna, A. Giacometti, M. Tavio, M. Onesta, L. Di Sante, A. Ranghino, SARS-CoV-2 infection in kidney transplant recipients: experience of the Italian Marche region, *Transpl Infect Dis*, (n/a e13377).
- [144] Marx D, Moulin B, Fafi-Kremer S, Benotmane I, Gautier G, Perrin P, et al. First case of COVID-19 in a kidney transplant recipient treated with belatacept. *Am J Transplant* 2020;13:13.
- [145] Masset C, Ville S, Halary F, Gaborit B, Bressolette-Bodin C, Deltombe C, et al. Resurgence of BK virus following Covid-19 in kidney transplant recipients. *Transpl Infect Dis* 2020:e13465.
- [146] Massoumi H, Rocca J, Frager S, Kinkhabwala M. COVID-19 Infection in Early Post-Operative Period after Liver Transplantation. *Liver Transpl* 2020;05:05.
- [147] Mathies D, Rauschnig D, Wagner U, Mueller F, Maibaum M, Binnemann C, et al. A Case of SARS-CoV-2-pneumonia with successful antiviral therapy in a 77-year-old male with heart transplant. *Am J Transplant* 2020;21:21.
- [148] M. Mattioli, E. Fustini, S. Gennarini, Heart transplant recipient patient with COVID-19 treated with tocilizumab, *Transpl Infect Dis*, (n/a e13380).
- [149] S.A. Mehta, J. Leonard, P. Labella, K. Cartiera, I. Soomro, H. Neumann, R.A. Montgomery, N.M. Ali, Outpatient management of kidney transplant recipients with suspected COVID-19—Single-center experience during the New York City surge, *Transpl Infect Dis*, (n/a e13383).
- [150] Melgosa M, Madrid A, Alvarez O, Lumberras J, Nieto F, Parada E, et al. Spanish Pediatric Nephrology, SARS-CoV-2 infection in Spanish children with chronic kidney pathologies. *Pediatr Nephrol* 2020;20:20.
- [151] Mella A, Mingozzi S, Gallo E, Lavacca A, Rossetti MM, Clari R, et al. Case series of six kidney transplanted patients with COVID-19 pneumonia treated with tocilizumab. *Transpl Infect Dis* 2020:e13348.
- [152] S. Meziyerh, D. van der Helm, A.P.J. de Vries, Vulnerabilities in kidney transplant recipients with COVID-19: a single center experience, *Transpl Int*, (n/a).
- [153] Meziyerh S, Zwart TC, van Etten RW, Janson JA, van Gelder T, Alwayn IPJ, et al. Severe COVID-19 in a renal transplant recipient: A focus on pharmacokinetics. *Am J Transplant* 2020;26:26.
- [154] Miarons M, Larrosa-García M, García-García S, Los-Arcos I, Moreso F, Berastegui C, et al. COVID-19 in Solid Organ Transplantation: A Matched Retrospective Cohort Study and Evaluation of Immunosuppression Management. *Transplantation* 2020.
- [155] Modi AR, Koval CE, Taegle AJ, Modaresi Esfeh J, Eghtesad B, Menon KVN, et al. Coronavirus Disease 2019 in an Orthotopic Liver Transplant Recipient Living with Human Immunodeficiency Virus. *Transpl Infect Dis* 2020:e13351.
- [156] I.H. Mohamed, P.B. Chowdary, S. Shetty, C. Sammartino, R. Sivaprakasam, B. Lindsey, R. Thuraingham, M.M. Yaqoob, M.A. Khurram, Outcomes of Renal Transplant Recipients With SARS-CoV-2 Infection in the Eye of the Storm: A Comparative Study With Waitlisted Patients, *Transplantation*, (Online First (9000)).
- [157] Mohamed MA, Kaur J, Wani F, Kichloo A, Bhanot R. Renal Transplant Recipient with Concurrent COVID-19 and Stenotrophomonas maltophilia Pneumonia Treated with Trimethoprim/Sulfamethoxazole Leading to Acute Kidney Injury: A Therapeutic Dilemma. *Am J Case Rep* 2020;21:e926464.
- [158] Molaie H, Khedmat L, Nemati E, Rostami Z, Saadat SH. Iranian kidney transplant recipients with COVID-19 infection: Clinical outcomes and cytomegalovirus coinfection. *Transpl Infect Dis* 2020:e13455.
- [159] Molnar MZ, Bhalla A, Azhar A, Tsujita M, Talwar M, Balaraman V, et al. Outcomes of critically ill solid organ transplant patients with COVID-19 in the United States. *Am J Transplant* 2020;20:3061–71.
- [160] A. Monfared, S. Dashti-Khavidaki, R. Jafari, A. Jafari, E. Ramezanzade, M.-K. Lebad, Y. Haghdar-Saheli, P. Aghajanzadeh, M. Khosravi, A. Movassaghi, H. Taghvaye-Masoumi, A. Alavi Foumani, Clinical characteristics and outcome of COVID-19 pneumonia in kidney transplant recipients in Razi hospital, Rasht, Iran, *Transpl Infect Dis*, (n/a e13420).
- [161] Morand A, Roquelaure B, Colson P, Amrane S, Bosdure E, Raoult D, et al. Child with liver transplant recovers from COVID-19 infection. A case report. *Arch Pediatr* 2020;06:06.
- [162] Morillas JA, Marco Canosa F, Srinivas P, Asadi T, Calabrese C, Rajendram P, et al. Tocilizumab therapy in five solid and composite tissue transplant recipients with early ARDS due to SARS-CoV-2. *Am J Transplant* 2020;31:31.
- [163] Morlacchi LC, Rossetti V, Gigli L, Amati F, Rosso L, Aliberti S, et al. COVID-19 In Lung Transplant Recipients: A Case Series From Milan, Italy. *Transpl Infect Dis* 2020:e13356.

- [164] Muller H, Kniepeiss D, Stauber R, Schrem H, Rauter M, Krause R, et al. Recovery from COVID-19 following Hepatitis C, Human Immunodeficiency Virus Infection and Liver Transplantation. *Am J Transplant* 2020;03:03.
- [165] S. Naem, R. Gohh, G. Bayliss, C. Cosgrove, D. Farmakiotis, B. Merhi, P. Morrissey, A. Osband, J.A. Bailey, J. Sweeney, R. Rogers, Successful recovery from COVID-19 in three kidney transplant recipients who received convalescent plasma therapy. *Transpl Infect Dis*, (n/a e13451).
- [166] Nair V, Jandovitz N, Hirsch JS, Nair G, Abate M, Bhaskaran M, et al. COVID-19 in kidney transplant recipients. *Am J Transplant* 2020;29:29.
- [167] Namazee N, Mahmoudi H, Afzal P, Ghaffari S. Novel Corona Virus 2019 pneumonia in a kidney transplant recipient. *Am J Transplant* 2020;13:13.
- [168] Niknam R, Malek-Hosseini SA, Hashemeh SS, Dehghani M. COVID-19 in Liver Transplant Patients: Report of 2 Cases and Review of the Literature. *Int Med Case Rep J* 2020;13:317–21.
- [169] Nikoupour H, Arasteh P, Gholami S, Nikeghbalian S. Liver transplantation and COVID-19: a case report and cross comparison between two identical twins with COVID-19. *BMC Surg* 2020;20:181.
- [170] Ning L, Liu L, Li W, Liu H, Wang J, Yao Z, et al. Novel coronavirus (SARS-CoV-2) infection in a renal transplant recipient: Case report. *Am J Transplant* 2020;10:10.
- [171] Noble R, Tan MY, McCulloch T, Shantier M, Byrne C, Hall M, et al. Collapsing Glomerulopathy Affecting Native and Transplant Kidneys in Individuals with COVID-19. *Nephron* 2020:1–6.
- [172] Oniszczuk J, Moktefi A, Mausoleo A, Pallet N, Malard-Castagnet S, Fourati S, et al. De Novo Focal and Segmental Glomerulosclerosis After COVID-19 in a Patient With a Transplanted Kidney From a Donor With a High-risk APOL1 Variant. *Transplantation* 2020.
- [173] Ossami Sady RR, Globke B, Pratschke J, Schoening W, Eurich D. Successful implementation of preventive measures leads to low relevance of SARS-CoV-2 in liver transplant patients: Observations from a German outpatient department. *Transpl Infect Dis* 2020:e13363.
- [174] R. Papa-Gobbi, A. Bueno, J. Serradilla, P. Talayero, P. Stringa, B. Pascual-Miguel, A. Alcolea-Sánchez, R. González-Sacristan, A.M. Andrés, M. López-Santamaría, M. Rumbo, E. Ramos-Boluda, F. Hernández-Oliveros. Novel coronavirus (SARS-CoV-2) infection in a patient with multivisceral transplant, *Transpl Infect Dis*, (n/a e13430).
- [175] J. Pascual, E. Melilli, C. Jiménez-Martín, E. González-Monte, S. Zárraga, A. Gutiérrez-Dalmau, V. López-Jiménez, J. Juega, M. Muñoz-Cepeda, I. Lorenzo, C. Fucundo, M.D.C. Ruiz-Fuentes, A. Mazuecos, E. Sánchez-Álvarez, M. Crespo, C.-G. Spanish Society of Nephrology, COVID-19-related Mortality During the First 60 Days After Kidney Transplantation, *Eur Urol*, (2020) S0302-2838(0320)30465-30466.
- [176] Patrono D, Lupo F, Canta F, Mazza E, Mirabella S, Corcione S, et al. Outcome of COVID-19 in Liver Transplant Recipients: a Preliminary Report from Northwestern Italy. *Transpl Infect Dis* 2020:e13353.
- [177] Pereira MR, Aversa MM, Farr MA, Miko BA, Aaron JG, Mohan S, et al. Tocilizumab for severe COVID-19 in solid organ transplant recipients: a matched cohort study. *Am J Transplant* 2020;2:3198–205.
- [178] M.J. Pérez-Sáez, M. Blasco, D. Redondo-Pachón, P. Ventura Aguilar, T. Bada-Bosch, I. Pérez-Flores, E. Melilli, L.A. Sánchez-Cámara, M.O. López-Oliva, C. Canal, A. Shabaka, N. Garra Moncau, P.L. Martín-Moreno, V. López, R. Hernández-Gallego, O. Siverio, C. Galeano, J. Espí Reig, C.J. Cabezas, M.T. Rodrigo, L. Llinàs-Mallol, M.J. Fernández-Reyes, L. Cruzado Vega, L. Pérez-Tamajón, R. Santana-Estupiñán, M.C. Ruiz-Fuentes, G. Tabernerero, S. Zárraga, J. Carlos Ruiz, A. Gutiérrez-Dalmau, A. Mazuecos, E. Sánchez-Álvarez, M. Crespo, J. Pascual, f.T.S.S.o.N.C.- Group, I. García-Méndez, L. Pérez-Oller, M. Giorgi, Use of tocilizumab in kidney transplant recipients with COVID-19, *Am J Transplant*, (n/a).
- [179] Pierrotti LC, Reusing Junior JO, Freire MP, Barros Machado DJ, Megale Moreira R, Ventura CG, et al. COVID-19 among kidney transplant recipients requiring hospitalization: preliminary data and outcomes from a single-center in Brazil. *Transpl Int* 2020.
- [180] Prieto M, Gastaca M, Ruiz P, Ventoso A, Palomares I, Rodríguez-Álvarez RJ, et al. A case of COVID-19 immediately after liver transplantation: Not only bad news. *Ann Hepatobiliary Pancreat Surg* 2020;24:314–8.
- [181] T.C.U.K.T. Program, Early Description of Coronavirus 2019 Disease in Kidney Transplant Recipients in New York, *J Am Soc Nephrol*, 31 (2020) 1150.
- [182] Qin J, Wang H, Qin X, Zhang P, Zhu L, Cai J, et al. Perioperative Presentation of COVID-19 Disease in a Liver Transplant Recipient. *Hepatology* 2020;27:27.
- [183] Rabiee A, Sadowski B, Adeniji N, Perumalswami P, Nguyen V, Moghe A, et al. Liver Injury in Liver Transplant Recipients with Coronavirus Disease 2019 (COVID-19): US Multicenter Experience. *Hepatology* 2020.
- [184] Raëth J, Tomio A, Eugene A, Mouffak A, Durand M, Hamidfar R, et al. Immunosuppression in a lung transplant recipient with COVID-19? Lessons from an early case. *Respiratory Medicine and Research* 2020;78:100782.
- [185] Rahman F, Liu STH, Taimur S, Jacobs S, Sullivan T, Dunn D, et al. Treatment with convalescent plasma in solid organ transplant recipients with COVID-19: Experience at large transplant center in New York City. *Clin Transplant* 2020:e14089.
- [186] Ramos KJ, Kapnadak SG, Collins BF, Pottinger PS, Wall R, Mays JA, et al. Detection of SARS-CoV-2 by bronchoscopy after negative nasopharyngeal testing: Stay vigilant for COVID-19. *Respir Med Case Rep* 2020;30:101120.
- [187] Rasekhi A, Hoseinyazdi M, Esmaeilian S, Teimouri A, Safaei A, Rafiee F. COVID-19 pneumonia presenting as a single pulmonary nodule in a kidney transplant recipient: A case report and literature review. *Radiology Case Reports* 2020;15:1587–90.
- [188] Renaud-Picard B, Gallais F, Ohana M, Zeyons F, Kretz B, Andre J, et al. Bilateral Acute Cardioembolic Limb Ischemia After Coronavirus Disease 2019 Pneumonia in a Lung Transplant Recipient: A Case Report. *Transplant Proc* 2020;52:2715–8.
- [189] M. Rinaldi, M. Bartoletti, L. Bussini, L. Pancaldi, R. Pascale, G. Comai, M. Morelli, M. Ravaoli, M. Cescon, F. Cristini, P. Viale, M. Giannella, COVID-19 in solid organ transplant recipients: No difference in survival compared to general population, *Transpl Infect Dis*, (n/a e13421).
- [190] Rivinius R, Kaya Z, Schramm R, Boeken U, Provaznik Z, Heim C, et al. COVID-19 among heart transplant recipients in Germany: a multicenter survey. *Clin Res Cardiol* 2020.
- [191] M.B. Roberts, S. Izzy, Z. Tahir, A. Al Jarrah, J.A. Fishman, J. El Khoury, COVID-19 in solid organ transplant recipients: dynamics of disease progression and inflammatory markers in ICU and non-ICU admitted patients, *Transpl Infect Dis*, (n/a e13407).
- [192] Rodriguez-Cubillo B, Moreno de la Higuera MA, Lucena R, Valdes Franci E, Hurtado M, Calvo Romero N, et al. Should cyclosporine be useful in renal transplant recipients affected by SARS-CoV-2? *Am J Transplant* 2020.
- [193] Russell MR, Halnon NJ, Alejos JC, Salem MM, Reardon LC. COVID-19 in a pediatric heart transplant recipient: Emergence of donor-specific antibodies. *J Heart Lung Transplant* 2020;29:29.
- [194] P. Sakulkonkij, J. Bruminhent, C. Pankongngam, N. Chalermphunchai, A family cluster of diagnosed coronavirus disease 2019 (COVID-19) kidney transplant recipient in Thailand, *Immunity, Inflammation and Disease*, (n/a).
- [195] Sánchez Cadena AD, Negreira Caamaño M, Pérez Serrano R, Porras Leal ML. Inmunoglobulinas por vía intravenosa: una alternativa terapéutica a tener en cuenta en el paciente trasplantado renal con COVID-19. *Nefrología* 2020.
- [196] E. Seminari, M. Colaneri, M. Sambo, I. Gallazzi, A. Di Matteo, S. Roda, R. Bruno, C.I.S.M.P.T.F. the, SARS Cov-2 infection in a renal-transplanted patient: A case report, *Am J Transplant*, (n/a).
- [197] Serrano OK, Kutzler HL, Rochon C, Radojevic JA, Lawlor MT, Hammond JA, et al. Incidental COVID-19 in a Heart-Kidney Transplant Recipient with Malnutrition and Recurrent Infections: Implications for the SARS-CoV-2 Immune Response. *Transpl Infect Dis* 2020.
- [198] Shingare A, Bahadur MM, Raina S. COVID-19 in recent kidney transplant recipients. *Am J Transplant* 2020;08:08.
- [199] Singhvi A, Barghash M, Lala A, Mitter SS, Parikh A, Oliveros E, et al. Challenges in Heart Transplantation during COVID-19: A Single Center Experience. *J Heart Lung Transplant* 2020;39:894–903.
- [200] Song SH, Chen TL, Deng LP, Zhang YX, Mo PZ, Gao SC, et al. Clinical characteristics of four cancer patients with SARS-CoV-2 infection in Wuhan, China. *Infect Dis Poverty* 2020;9:82.
- [201] Soquet J, Rousse N, Moussa M, Goeminne C, Deblauwe D, Vuotto F, et al. Heart Retransplantation Following COVID-19 Illness in a Heart Transplant Recipient. *J Heart Lung Transplant* 2020;39:983–5.
- [202] Sperry BW, Khumri TM, Kao AC. Donor-derived cell-free DNA in a heart transplant patient with COVID-19. *Clin Transplant* 2020:e14070.
- [203] Stachel MW, Gidea CG, Reventovich A, Mehta SA, Moazami N. COVID-19 pneumonia in a dual heart-kidney recipient. *J Heart Lung Transplant* 2020;39:612–4.
- [204] Suwanwongse K, Shabarek N. Fatal Outcome in a Kidney-Pancreas Transplant Recipient With COVID-19. *Cureus* 2020;12.
- [205] Taha M, Sharma A, Taha M, Samavati L. Coronavirus Disease 2019 in Immunocompromised Organ Transplant Recipients: A Case Report and Review of the Literature. *Transplant Proc* 2020;52:2698–702.
- [206] E. Tatar, M. Karatas, I. Bozaci, A. Ari, T. Acar, C. Simsek, A.M. Yildirim, O. Yildirim, A. Uslu, Intravenous Immunoglobulin and Favipiravir treatment for A Kidney Transplant Patient with Severe Covid-19 Pneumonia, *Transfus Apher Sci*
- [207] D.R.B. Terrabuio, L. Haddad, L. Ducatti, L. Nunes Gouveia, V. Rocha-Santos, R.M.T. Ferreira, G.F. Darce, A.J.A. Cardoso, F.J. Carrilho, W. Andraus, E. Abdala, L.A.C. D'Albuquerque, Insights in the approach of long-term liver transplant recipients with COVID-19, *Transpl Infect Dis*, (n/a e13424).
- [208] T. Thammathiwat, S. Tungsanga, K. Tiankanon, P. Torvorapanit, W. Chumpangern, S. Udomkarnjananun, Y. Avihingsanon, T. Sriprasart, N. Srisawat, K. Jutivorakool, L. Paitoonpong, O. Puchareon, N. Townamchai, A case of successful treatment of severe COVID-19 pneumonia with favipiravir and tocilizumab in post-kidney transplant recipient, *Transpl Infect Dis*, (n/a e13388).
- [209] Tian S, Xiong Y, Liu H, Niu L, Guo J, Liao M, et al. Pathological study of the 2019 novel coronavirus disease (COVID-19) through postmortem core biopsies. *Mod Pathol* 2020;33:1007–14.
- [210] Trujillo H, Caravaca-Fontan F, Sevillano A, Gutierrez E, Caro J, Gutierrez E, et al. SARS-CoV-2 Infection in Hospitalized Patients with Kidney Disease. *KL Reports* 2020;01:01.
- [211] Trujillo H, Caravaca-Fontan F, Sevillano A, Gutierrez E, Fernandez-Ruiz M, Lopez-Medrano F, et al. Tocilizumab use in Kidney Transplant Patients with COVID-19. *Clin Transplant* 2020:e14072.
- [212] J. Tschopp, A.G. L'Huilier, M. Mombelli, N.J. Mueller, N. Khanna, C. Garzoni, D. Meloni, M. Papadimitriou-Oliverigs, D. Neofytos, H.H. Hirsch, M.M. Schuurmans, T. Müller, T. Berney, J. Steiger, M. Pascual, O. Manuel, C. van Delden, S. Swiss Transplant Cohort, First experience of SARS-CoV-2 infections in solid organ transplant recipients in the Swiss Transplant Cohort Study, *Am J Transplant*, (n/a).
- [213] Vaidya G, Czer LSC, Kobashigawa J, Kittleson M, Patel J, Chang D, et al. Successful Treatment of Severe COVID-19 Pneumonia With Clazakizumab in a Heart Transplant Recipient: A Case Report. *Transplant Proc* 2020;52:2711–4.
- [214] Vechorko V, Gordeev I, Gubareva E, Ryndyayeva E, Averkov O. COVID-19 infection after recent heart transplantation: a case report. *Russian Journal of Cardiology* 2020;25:3904.
- [215] A. Velioglu, S. Tuglular, Care of Asymptomatic SARS-CoV-2 positive Kidney Transplant Recipients, *Transpl Int*, (n/a).
- [216] G.M. Verleden, L. Godinas, N. Lorent, P. Van Bleyenbergh, L. Dupont, M. Delcroix, J. Yserbyt, C. Doms, R. Vos, COVID-19 in lung transplant patients: a case series, *Am J Transplant*, (n/a).



- [217] Verma A, Khorsandi SE, Dolcet A, Prachalias A, Suddle A, Heaton N, et al. Low prevalence and disease severity of COVID-19 in post liver transplant recipients - a single centre experience. *Liver Int* 2020;29:29.
- [218] Vilaro J, Al-Ani M, Manjarres DG, Lascano JE, Cherabuddi K, Elgendy AY, et al. Severe COVID-19 After Recent Heart Transplantation Complicated by Allograft Dysfunction. *JACC* 2020;2:1347–50.
- [219] D.R. Waisberg, E. Abdala, L.S. Nacif, L.B. Haddad, L. Ducatti, V. Rocha-Santos, L.N. Gouveia, C.S. Lazari, R.B. Martino, R.S. Pinheiro, R.M. Arantes, D.R. Terrabuio, L.M. Malbouisson, F.H. Galvao, W. Andraus, L.A. Carneiro-D'Albuquerque, Liver Transplant Recipients Infected with SARS-CoV-2 in the Early Postoperative Period: Lessons from a Single Center in the Epicenter of the Pandemic, *Transpl Infect Dis*, (n/a e13418).
- [220] A.X. Wang, O. Quintero Cardona, D.Y. Ho, S. Busque, C.R. Lenihan, Influence of immunosuppression on seroconversion against SARS-CoV-2 in two kidney transplant recipients, *Transpl Infect Dis*, (n/a e13423).
- [221] Wang J, Li X, Cao G, Wu X, Wang Z, Yan T. COVID-19 in a Kidney Transplant Patient. *Eur Urol* 2020;77:769–70.
- [222] Webb GJ, Marjot T, Cook JA, Aloman C, Armstrong MJ, Brenner EJ, et al. Outcomes following SARS-CoV-2 infection in liver transplant recipients: an international registry study. *Lancet Gastroenterol Hepatol* 2020;5:1008–16.
- [223] Webb GJ, Moon AM, Barnes E, Barritt AS, Marjot T. Determining risk factors for mortality in liver transplant patients with COVID-19. *Lancet Gastroenterol Hepatol* 2020;5:643–4.
- [224] T.H. Westhoff, F.S. Seibert, F. Bauer, U. Stervbo, M. Anft, A.A.N. Doevelaar, B.J. Rohn, G. Winnekendonk, U. Dittmer, P. Schenker, E. Vonbrunn, K. Amann, R. Viebahn, N. Babel, Allograft infiltration and meningoencephalitis by SARS-CoV-2 in a pancreas-kidney transplant recipient, *Am J Transplant*, (n/a).
- [225] Xia Z, Liu X, Hu X, Zhong Z, Wang Y, Peng G, et al. Failed antibody response in a renal transplant recipient with SARS-CoV-2 infected. *Transpl Infect Dis* 2020; e13349.
- [226] J.J. Xu, D. Samaha, S. Mondhe, D. Massicotte-Azarniouch, G. Knoll, M. Ruzicka, Renal Infarct in a COVID-19 Positive Kidney-Pancreas Transplant Recipient, *Am J Transplant*, (n/a).
- [227] Yamada M, Rastogi P, Ince D, Thayyil A, Adela Mansilla M, Smith RJH, et al. Minimal Change Disease With Nephrotic Syndrome Associated With Coronavirus Disease 2019 After Apolipoprotein L1 Risk Variant Kidney Transplant: A Case Report. *Transplant Proc* 2020;52:2693–7.
- [228] S.G. Yi, A.W. Rogers, A. Saharia, M. Aoun, R. Faour, M. Abdelrahim, R.J. Knight, K. Grimes, S. Bullock, M. Hobeika, R. McMillan, C. Mobley, M. Moaddab, H.J. Huang, A. Bhimaraj, R.M. Ghobrial, A.O. Gaber, Early Experience With COVID-19 and Solid Organ Transplantation at a US High-volume Transplant Center, *Transplantation*, Online First (9000).
- [229] Zhang H, Chen Y, Yuan Q, Xia QX, Zeng XP, Peng JT, et al. Identification of Kidney Transplant Recipients with Coronavirus Disease 2019. *Eur Urol* 2020;77:742–7.
- [230] Zhong Z, Zhang Q, Xia H, Wang A, Liang W, Zhou W, et al. Clinical characteristics and immunosuppressant management of coronavirus disease 2019 in solid organ transplant recipients. *Am J Transplant* 2020;13:13.
- [231] Zhu L, Gong N, Liu B, Lu X, Chen D, Chen S, et al. Coronavirus Disease 2019 Pneumonia in Immunosuppressed Renal Transplant Recipients: A Summary of 10 Confirmed Cases in Wuhan, China. *Eur Urol* 2020;77:748–54.
- [232] L. Zhu, H. Shu, H. Li, T. Qiu, J. Zhou, G. Chen, Slow Recovery from Critical Coronavirus Disease 2019 Pneumonia in an Immunosuppressed Renal Transplant Recipient with Early Acute Cardiorenal Syndrome, *Cardiorenal Med*, (2020) 1–6.
- [233] Chen D, Yang B, Zhang Y, Chen L, Wei L, Zhang W, et al. Withdrawing mycophenolate mofetil in treating a young kidney transplant recipient with COVID-19: A case report. *Medicine (Baltimore)* 2020;99:e20481.
- [234] K.T. Mahendraraj, Tsuyoshi; Nissen, Nicholas; Kosari, Kambiz; Brennan, Todd; Voidonikolas, Georgios; Klein, Andrew; Kim, Irene, First report of novel coronavirus infection following abdominal organ transplantation in North America during the COVID-19 pandemic, *Transplantation*, 104 (2020) S321.
- [235] Li J, Chen G, Zhang M, Tu S, Chen C. Different clinical presentations of two renal transplant recipients with coronavirus disease 2019: a case report. *BMC Infect Dis* 2020;20:707.
- [236] M. Ahluwalia, M.M. Givertz, M.R. Mehra, A Proposed Strategy for Management of Immunosuppression in Heart Transplant Patients with COVID-19, *Clin Transplant*, (n/a e14032).
- [237] Binda B, Picchi G, Carucci AC, Sinatti G, Di Norcia M, Grimaldi A, et al. Follow-up and Management of Kidney Transplant Recipients During the COVID-19 Lockdown: The Experience of an Italian Transplant Center, Including Two Cases of COVID-19 Pneumonia. *Transplant Proc* 2020;52:2614–9.
- [238] M. Ghaffari Rahbar, M. Nafar, A. Khoshdel, N. Dalili, A. Abrishami, A. Firouzan, F. Poorrezagholi, F. Samadian, S. Ziaie, S. Fatemizadeh, S. Samavat, Low Rate of COVID-19 pneumonia in Kidney Transplant Recipients - A Battle Between Infection and Immune Response?, *Transpl Infect Dis*, (n/a e13406).
- [239] F. Silva, A. Cipriano, H. Cruz, J. Tavares, J. Fragoso, J. Malheiro, M. Almeida, L.S. Martins, M. Abreu, S. Pedroso, L. Dias, A.C. Henriques, SARS-CoV-2 infection in kidney transplant recipients: Early report of five cases, *Transpl Infect Dis*, (n/a e13394).
- [240] Caillard S, Anglicheau D, Matignon M, Durrbach A, Greze C, Frimat L, et al. An initial report from the French SOT COVID Registry suggests high mortality due to Covid-19 in recipients of kidney transplants. *Kidney Int* 2020.
- [241] Ahmad SH, Smith R, Camilleri B. Belatacept, kidney transplantation and COVID-19: Successful management of the first reported case within the United Kingdom. *Clin Transplant* 2020:e14026.
- [242] Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. C. China Medical Treatment Expert Group for, Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med* 2020;382:1708–20.
- [243] Desai A, Sachdeva S, Parekh T, Desai R. COVID-19 and Cancer: Lessons From a Pooled Meta-Analysis. *JCO Glob Oncol* 2020;6:557–9.
- [244] Malard F, Genthon A, Brissot E, van de Wyngaert Z, Marjanovic Z, Ikhlef S, et al. COVID-19 outcomes in patients with hematologic disease. *Bone Marrow Transplant* 2020;50:2180–4.