

RESEARCH ARTICLE

Evaluation of reference genes for reverse transcription quantitative real-time PCR (RT-qPCR) studies in *Silene vulgaris* considering the method of cDNA preparation

Pavla Koloušková, James D. Stone, Helena Štorchová*

Plant Reproduction Laboratory, Institute of Experimental Botany v.v.i., Academy of Sciences of the Czech Republic, Prague, Czech Republic

* storchova@ueb.cas.cz



OPEN ACCESS

Citation: Koloušková P, Stone JD, Štorchová H (2017) Evaluation of reference genes for reverse transcription quantitative real-time PCR (RT-qPCR) studies in *Silene vulgaris* considering the method of cDNA preparation. PLoS ONE 12(8): e0183470. <https://doi.org/10.1371/journal.pone.0183470>

Editor: Serena Aceto, University of Naples Federico II, ITALY

Received: January 25, 2017

Accepted: August 4, 2017

Published: August 17, 2017

Copyright: © 2017 Koloušková et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by the Ministry of Education, Youth and Sports of the Czech Republic (MŠMT) (<http://www.msmt.cz/research-and-development-1>) grant: KONTAKT II LH15075 and the Grantová agentura České republiky (<https://gacr.cz/en/>) grant: 16-09220S. The funders had no role in study design, data collection and

Abstract

Accurate gene expression measurements are essential in studies of both crop and wild plants. Reverse transcription quantitative real-time PCR (RT-qPCR) has become a preferred tool for gene expression estimation. A selection of suitable reference genes for the normalization of transcript levels is an essential prerequisite of accurate RT-qPCR results. We evaluated the expression stability of eight candidate reference genes across roots, leaves, flower buds and pollen of *Silene vulgaris* (bladder campion), a model plant for the study of gynodioecy. As random priming of cDNA is recommended for the study of organellar transcripts and poly(A) selection is indicated for nuclear transcripts, we estimated gene expression with both random-primed and oligo(dT)-primed cDNA. Accordingly, we determined reference genes that perform well with oligo(dT)- and random-primed cDNA, making it possible to estimate levels of nucleus-derived transcripts in the same cDNA samples as used for organellar transcripts, a key benefit in studies of cyto-nuclear interactions. Gene expression variance was estimated by RefFinder, which integrates four different analytical tools. The *SvACT* and *SvGAPDH* genes were the most stable candidates across various organs of *S. vulgaris*, regardless of whether pollen was included or not.

Introduction

The quantification of gene expression based on RT-qPCR is an increasingly important method across diverse fields of plant biology. Initially developed in *Arabidopsis* [1] and crop plants [2, 3], it has been adapted for use in wild species investigated by ecologists and population geneticists [4, 5]. The accuracy of RT-qPCR results is influenced by numerous factors including the quality of RNA, method of cDNA preparation, PCR efficiency, and very importantly, by the choice of reference gene [6]. An ideal reference gene has to show invariant expression across a range of tissues, developmental stages and environmental conditions. However, expression of the vast majority of genes fluctuates in response to various stresses [7, 8, 9, 10, 11, 12, 13, 14, 15], circadian rhythms [12, 16], or by developmental stage [10, 17, 18, 19, 20, 21]. Furthermore,

analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

most genes show substantial differences between gametophyte and sporophyte [20, 22]. It is therefore necessary to confirm invariant expression of a candidate reference gene in relevant tissues, organs and under experimental treatments before it is used in an experiment. A reference gene successfully used in one species may serve as a starting point for identifying them in another, but cannot necessarily be applied in a different, albeit related, species. The choice of reference gene(s) should be always justified for the given species and conditions as recommended by MIQE guidelines (Minimum Information for Publication of Quantitative Real-time PCR Experiments) [6].

Although the majority of angiosperms are hermaphroditic, some species produce female, male or hermaphroditic flowers on the same individual (monoecy), or on distinct individuals (dioecy). In these species, gender-specific expression patterns, in addition to organ- and tissue-specificity, has to be considered, when evaluating reference genes [4, 23, 24].

Silene vulgaris has emerged as a model system for investigating gynodioecy, a plant breeding system characterized by the co-occurrence of females (F) and hermaphrodites (H) [25]. The female gender is attributable to the interaction of cytoplasmic male sterility (CMS) genes, located in the mitochondrial genome, and nuclear genes [26]. This system represents a textbook example of cyto-nuclear interaction [27].

Recent publications of the complete mitochondrial genomes [28] and transcriptomes of *S. vulgaris* [29, 30] pave the way for detailed investigations of gene expression and function in this species. RT-qPCR plays an essential role in gene expression studies, but no detailed examination of reference genes suitable for transcript level normalization across various tissues and organs of *S. vulgaris* has been published. This study aims to identify reference genes with invariant expression in F and H plants of *S. vulgaris*, across leaves, roots, flower buds and pollen.

To investigate the role of gene expression in cyto-nuclear interactions, it is important to estimate both organellar and nucleus-derived transcript levels in the same cDNA specimens. Unlike cytoplasmic mRNAs, organellar mRNAs cannot be efficiently reverse transcribed using oligo(dT)-primers, because they often lack poly(A) tails [31, 32]. To achieve balanced reverse transcription of both cytoplasmic and organellar RNAs, cDNA synthesis primed with random oligomers is necessary. On the other hand, most gene expression studies rely on oligo(dT)-priming, taking an advantage of selective enrichment of poly(A)-tailed mRNAs. To better study the transcription of genes involved in cyto-nuclear interactions in *S. vulgaris*, reference genes performing well with both of these cDNA priming methods will be identified. This is the first study that reported the comparison of two different cDNA synthesis method on the transcription stability of reference genes. Accordingly, we have measured the stability of expression of eight reference gene candidates in *S. vulgaris* with cDNAs generated by both oligo(dT)- and random-priming.

Finally, we demonstrated the utility of the best candidate reference genes by normalizing the expression of the nuclear *MutS HOMOLOGUE 1 (MSH1)* gene, encoding the mitochondrial- and plastid-targeted protein which influences recombination and replication in plant organelles [33, 34].

Material and methods

Plant material

S. vulgaris plants were grown from seedlings to mature plants in 1.5 L pots filled with a substrate of coconut coir, Vermiculite and Agropelrite (1:1:1 ratio) in the IEB greenhouse under a 16 hour light and 8 hour darkness period, temperature between 18 and 24°C and humidity around 80%. Plants were regularly watered and fertilized with a combination of Kristalon

fertilizers (Agro CS, Česká Skalice, Czech Republic) and $\text{Ca}(\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$ (0.66 g of Kristalon-Fruit and flower, 0.8 g of Kristalon-Autumn and 0.53 g of $\text{Ca}(\text{NO}_3)_2 \cdot 2\text{H}_2\text{O}$ per L). Temperature, humidity and light conditions varied slightly with outside conditions due to the greenhouse's semi-insulated design. We used 4 H plants for the dataset with pollen, and 3 F and 3 H plants for the other datasets.

RNA extraction and cDNA synthesis

Tissue samples were taken from mature plants (flower buds about 0.5 cm long, young upper leaves, mature pollen after anther dehiscence and apical roots). A maximum of 100 mg tissue sample was snap-frozen in liquid nitrogen and stored at -80°C until further use. Plant material was homogenized with mortar and pestle in liquid nitrogen, total RNA was extracted using the RNeasy Plant Mini kit (Qiagen, Valencia, USA) and treated by DNaseI (Ambion, Foster City, USA). To perform the non-RT control, treated RNA was used as a PCR template with specific *S. vulgaris* primers (2 min at 94°C initial denaturation followed by 36 cycles: 1 min at 93°C , 1 min at 58°C and 1 min s at 72°C). This produced no amplification, confirming the absence of DNA contamination.

RNA was visualised on 0.8% agarose gel under RNase-free conditions. The concentration of RNA was quantified using a NanoDrop ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). RNA samples with A260/A280 ratios from 1.9 to 2.1 and A260/A230 ratios between 2.0 and 2.2 were used for cDNA synthesis. An equal amount of total RNA (500 ng) from each tissue sample was reverse transcribed with Transcriptor High Fidelity cDNA synthesis kit (Roche Applied Science, Mannheim, Germany) using oligo(dT)- or random hexamer-primers, in a 20 μL reaction volume according to the manufacturer's instructions. The reverse transcription reaction was subsequently diluted to 20x with RNase-free water for RT-qPCR.

Selection of candidate reference genes and primer design

Twenty candidate reference genes were selected based on their previous applications in plant gene expression studies. They included *18S rRNA* [35, 36, 37, 38], *ACTIN (ACT)* [38, 39, 40], *ATP-BINDING CASSETTE 16 (ABC16)* [12], *CLATHRIN ADAPTOR PROTEIN-2 (AP2M)* [2], *CONSERVED OLIGOMERIC GOLGI COMPLEX (COG)* [41], *CYCLOPHILIN (CYP)* [42, 43, 44], *ELONGATION FACTOR 1 alpha (ELF1 α)* [38, 44, 45, 46, 47], *GLYCERALDEHYDE-3-PHOSPHATE DEHYDROGENASE (GAPDH)* [48, 49, 50], *S-ADENOSYLMETHIONINE DECARBOXYLASE (SAMDC)* [51], *SAND family protein (SAND)* [2, 50], *SKP1/ASK-INTERACTING PROTEIN 16 (SKIP16)* [52], *TIP41-like family protein (TIP41)* [52], *ALPHA TUBULIN (TUBA)* [9, 53], *BETA TUBULIN (TUBB)*, *UBIQUITIN-CONJUGATING ENZYME 18 (UBC18)* [51], *UBIQUITIN 4 (UBQ4)* [54], *UBIQUITIN 5 (UBQ5)* [55], *UBIQUITIN 6 (UBQ6)* [56], *POLYUBIQUITIN 14 (UBQ14)* [54], and *VACUOLAR MEMBRANE ATPASE 10 (VMA10)* [41]. All amino acid sequences were blasted against the *S. vulgaris* transcriptome database (<http://silene-genomics.biology.virginia.edu/>) using blastx with a cutoff of $e \leq 0.001$, identifying isotigs with nearly complete coding sequences and overall amino acid similarity between *Arabidopsis* and *S. vulgaris* $> 60\%$ (Table 1). These isotig sequences from *S. vulgaris* were selected for primer development. At least two primer pairs were designed for each isotig using Primer 3 (<http://simgene.com/Primer3>) as follows: length of primers 20–24 nt, size of amplicons 90–200 bp; the function penalty weight for primer pairs, of any complementarity, was set to 1.0. Similarly, the sequence of *SvMSH1* was retrieved from the *S. vulgaris* transcriptome database by a blastn search of sugar beet *MSH1* mRNA (XM010692765). The final

Table 1. Description of *Silene vulgaris* candidate reference genes.

<i>S. vulgaris</i>		Arabidopsis		
Gene Symbol	Isotig*	Ortholog Locus	Gene Product	Amino Acid Identity between Arabidopsis and <i>S. vulgaris</i> (%)
<i>SvELF</i>	05995	AT1G07920	Elongation factor 1A	94
<i>SvGAPDH</i>	26878	AT3G04120	Glyceraldehyde-3-phosphate dehydrogenase	91
<i>SvTUBA</i>	05561	AT1G04820	Alpha Tubulin	96
<i>SvTUBB</i>	07210	AT5G62700	Beta Tubulin	95
<i>SvACT</i>	09472	AT5G09810	Actin	96
<i>SvCOG</i>	34935	AT4G24840	Golgi (COG) Complex Component	63
<i>SvCYP</i>	11915	AT4G38740	Cyclophilin ROC1	83
<i>18S rRNA</i>	36580	AT3G41768	18S ribosomal RNA	89

* *S. vulgaris* transcriptomic database available at <http://sileneomics.biology.virginia.edu>

<https://doi.org/10.1371/journal.pone.0183470.t001>

SvMSH1 sequence (total length 4007 bp, ORF 3414 bp) was assembled from 5 isotigs, with gaps filled by PCR and Sanger sequencing.

Quantitative real-time PCR (qPCR)

qPCR was carried out in 96-well plates on a LightCycler[®] 480 Instrument II (Roche Molecular System, Germany) using LightCycler[®] 480 SYBR Green I Master (Roche Applied Science, Mannheim, Germany). One sample reaction was run in four replicates (two replicates per plate, to check run-to-run variation) in 10 µL total volume containing 5 µL of 2x SYBR Green I Master mix, 2.5 µL of diluted cDNA and gene-specific primers in proper concentrations (Table 2).

Cycling conditions: 5 min at 95°C for initial denaturation followed by 50 cycles: 10 s at 95°C, 10 s at 60°C and 15 s at 72°C, except for 18S rRNA primers where annealing was set to 10 s at 63°C. The PCR efficiency for each assay was calculated with LightCycler[®] 480 software from the calibration curve based on a serial dilution of cDNA (most often 20x, 40x, 80x, 160x, 320x). Specificity of qPCR in each sample was confirmed by melting curve analyses. The qPCR amplicons (Fig 1) were sequenced to confirm their identity.

Table 2. Details of qPCR primers and amplicons.

Gene Symbol	Primer Pair	Primer Sequence Forward / Reverse	Ann Temp (°C)	Optimized Primer Concentration (pM)	cDNA Amplicon Length (bp)	qPCR efficiency
<i>SvELF</i>	Sv_ELF 1	TCTCCCTGGTGACAATGTTGGT / GAACTGGGGCATATCCGTTACC	60	0.5	176	1.0
	Sv_ELF 2	GCATGCACTTCTTGCTTTCA / GACCTTGTCGGGATTGTACC	60	0.5	160	1.0
<i>SvGAPDH</i>	Sv_GAPDH 1	GGCCAAGGTTATCAATGACAG / CCTTCCACCTCTCCAGTCCT	60	0.4	119	1.0
	Sv_GAPDH 2	GATCGGAATCAACGGATTTG / TGGACCATGAACACTGTCTGT	60	0.5	151	0.98
<i>SvTUBA</i>	Sv_TUB A	ACATGGCTTGCTGTCTGATG / TGGGGCTGGTAGTTGATAC	60	0.4	146	1.0
<i>SvTUBB</i>	Sv_TUB B	ACAACCCATCGACTCGAAAC / GCAACATGAATGACCTCGTG	60	0.5	173	1.0
<i>SvACT</i>	Sv_ACT	GGGCTGTGATCTCTTTGCTC / ATTGTTCCGGTATGGAAGCTC	60	0.4	163	0.96
<i>SvCOG</i>	Sv_COG	CCTGTTCCGCATCTCTCTTA / CGACTTCAGATGCCAATTCA	60	0.5	166	1.0
<i>SvCYP</i>	Sv_CYP	TCGCAGTTCTTCATCTGCAC / AGCCTTCACAACATCCAACC	60	0.4	96	1.0
<i>18S rRNA</i>	Sv_18S rRNA	CCTCCAATGGATCCTCGTTA / AAACGGCTACCACATCCAAG	63	0.3	154	1.0
<i>SvMSH1</i>	Sv_MSH_1	CTGAGGATTTCTCCCA / ACCAACTGCTTCGTGTC	60	0.5	105	1.0

<https://doi.org/10.1371/journal.pone.0183470.t002>




Fig 1. qPCR amplicons generated with cDNA from leaf tissue by the indicated primer pairs. Agarose gel (1.5%) with specific qPCR products of expected sizes. **A:** oligo(dT)- primed amplicons; **B:** random hexamer-primed amplicons. Size standards are loaded on the left and right sides—GeneRuler 100 bp DNA Ladder.

<https://doi.org/10.1371/journal.pone.0183470.g001>

Validation of gene expression using RNA-seq data

RNA-seq data obtained by Illumina HiSeq (single reads, 50 cycles) from eight cDNA libraries constructed by oligo(dT)-priming from the flower buds and leaves of H and F plants were used to evaluate the expression variance of each reference gene candidate. Amplicon coverage was estimated as follows: each primer sequence, as well as the 30 nt immediately upstream and downstream were used for read alignment, ensuring complete coverage of the target primer sequence. The sequences between primer pairs would have attracted reads derived from alternative transcripts and, therefore, not unique to the focal primer pairs and were not used for read alignment. Thus, the reference sequence for each primer pair was 160 mappable bases (two 20 nt primer sequences and four flanking 30 nt regions). Occasionally, polymorphism in the flanking regions necessitated multiple reference sequences per primer pair to capture all the reads derived from RNA potentially amplified by a particular primer pair. Reads from individual transcriptomes were aligned to these reference sequences using Bowtie2 [57], allowing zero mismatches and requiring that the full read length aligned. The average coverage of each sequence was obtained by BEDTools genomecov tool [58]. The coverage values were multiplied by 10^9 , divided by sequence length adjusted for edge effects (50% average coverage in flanking regions gives an adjusted length of 100 nt rather than 160 nt), divided by 50 (read length), and lastly divided by the total number of reads from each sample. If multiple mapping sequences were needed for a particular primer pair, their coverage values were summed. The resulting number is equivalent to RPKM (reads per 1 thousand bp of sequence length and 1 million total reads). Coefficient of variation (CV) was calculated as the standard deviation across samples divided by the mean value and expressed as a percentage.

Statistical evaluation

The web tool RefFinder [59] (<http://fulxie.0fees.us/>) which integrates four widely used algorithms: geNorm [60], NormFinder [61], BestKeeper [62] and the comparative delta-Ct method [63] was used to estimate the stability of candidate reference genes, providing us comprehensive information for ranking the candidate reference genes. We used all four technical cycle threshold values (C_t) replicates as input when calculating stabilities. Expression stability was also tested using the stand-alone programs geNorm and NormFinder, where relative quantities (RQ) calculated with qPCR efficiencies, rather than raw C_t data, were used as inputs, as directed in the geNorm manual.

MSH1 expression data were analyzed by a two-way analysis of variance (ANOVA) with gender and organ modeled as fixed-effects. Differences between means were tested using Tukey's test and a critical value of $P < 0.05$. Analyses were performed using the software SPSS 15.0.

Results

Expression profiles of candidate reference genes

We selected eight *S. vulgaris* isotigs homologous to reference genes previously employed for RT-qPCR normalization in plants (Table 1): *SvELF1 α* [38], *SvGAPDH* [50], *SvTUBA*, *SvTUBB*, *SvACT* [52], *SvCOG* [41], *SvCYP* [44], and *18S rRNA* [64]. We performed several RT-qPCR assays for each gene and chose the primer pairs which worked with a PCR efficiency of at least 1.95 and produced only the specified PCR products (Table 2). We continued with two primer pairs targeting different genic regions of *SvELF1 α* and *SvGAPDH* and with one primer pair for the rest of the candidates.

As we aimed to find reliable reference genes to evaluate gene expression across various tissues of both F and H plants of *S. vulgaris*, we examined two groups of plants. The first group, H_BLRP, contained only H individuals (4 plants), which made it possible to measure gene expression in pollen (P) in addition to flower buds (B), leaves (L) and roots (R). The second group (named HF_BLR) consisted of 3 H and 3 F plants and was used to determine gene expression in flower buds (B), leaves (L) and roots (R).

We also tested whether the same reference genes could be used with cDNAs prepared by different priming methods. Accordingly, we performed RT-qPCR measurements of all the genes except 18S rRNA in parallel with cDNA primed either with oligo dT or with random hexamers. The 18S rRNA expression was estimated only in random hexamer-primed cDNA, because their lack of poly(A) tails precluded oligo(dT)-priming.

Fig 2 shows a comparison of C_t values of the candidate reference genes obtained with oligo dT primed or random hexamer-primed cDNA. Medians and percentiles of C_t values for the candidate genes are consistent between the two datasets. The medians from hexamer-primed cDNAs are several cycles higher than those obtained with oligo(dT)-primed cDNA, because hexamer-primed cDNA contains a much lower proportion of mRNA-derived reverse transcripts.

The two primer pairs designed for the *SvELF1 α* and *SvGAPDH* genes differed in median C_t values. This may be caused by the distinct PCR efficiencies of RT-qPCR assays (in the case of *SvGAPDH*) or by targeting multiple loci. The C_t values for two tissue sets with pollen (H_BLRP—buds, leaves, roots, pollen) and without pollen (HF_BLR—buds, leaves, roots) were concordant. The set with pollen exhibited larger variation.

Stability of expression of candidate reference genes

The two most stably expressed candidate reference genes of *S. vulgaris* across the oligo dT primed tissue set with pollen (H_BLRP), as determined by RefFinder, were *SvGAPDH* (primer pair Sv_GAPDH 1) and *SvACT* (Table 3). They were followed by *SvELF1 α* (primer pair Sv_elf 1) and *SvGAPDH* (primer pair Sv_GAPDH 2). Random-primed cDNA provided a very similar order of reference gene candidates for the complete set of tissues including pollen (H_BLRP). A minor difference in the oligo(dT)-primed ranking was the exchange of first and second positions, which placed *SvACT* on the top. *18S rRNA*, frequently used to normalize random primed cDNA, was not the best performer, ranking, on average, fifth.

When pollen was excluded in the oligo(dT)-dataset, the two most stable candidates remained, but the third most stable gene became *SvCOG*, which ranked lower in dataset including pollen (Table 3). Its overall transcript level was the lowest of the examined candidate genes (Fig 2).

Hexamer-primed cDNA gave results consistent with the oligo(dT)-dataset, with *SvACT* and *SvGAPDH* (primer pair Sv_GAPDH 1) as the most stable reference genes. The least stable




Fig 2. Box-plot of threshold cycle (C_t) values of the candidate reference genes. Flower buds, leaves, roots and pollen (BLRP) or in flower buds, leaves, roots (BLR) were analyzed. **A:** Oligo dT primed cDNAs. **B:** Random hexamer-primed cDNAs. Graph shows the median values as line across the box. Lower and upper box boundaries indicate the 25th percentile and the 75th percentile, respectively. Whiskers represent the maximum and minimum values.

<https://doi.org/10.1371/journal.pone.0183470.g002>

gene was *SvCYP*, regardless of the method used for cDNA priming, which agreed with its highest variation in C_t values (Fig 2), except the random primed HF_BLR dataset, wherein *SvCOG* was the least stable.

The stability values calculated by geNorm (S1 Fig) were in general agreement with RefFinder, although the random HF_BLR dataset was ordered slightly differently by these programs. Ranking orders generated by RefFinder and NormFinder (S1 Table) showed more differences, but they agreed in the placement of at least two candidates in the first three positions.

Validation of reference gene candidates using RNA-seq data

To confirm the results obtained by statistical analyses of RT-qPCR estimations of gene expression values, we utilized RNA-seq data from flower bud and leaf transcriptomes obtained from 5 F and 3 H plants. We calculated read coverage of the respective amplicon regions and

Table 3. RefFinder output. Recommended comprehensive ranking of candidate reference gene primer pairs in analyzed datasets from the most stable (1) to the least stable (9,10) genes.

rank	oligo_HF_BLR	stability	oligo_H_BLRP	stability	random_HF_BLR	stability	random_H_BLRP	stability
1	Sv_GAPDH 1	1.19	Sv_GAPDH 1	1.00	Sv_ACT	1.19	Sv_ACT	1.57
2	Sv_ACT	1.41	Sv_ACT	1.86	Sv_GAPDH 1	2.6	Sv_GAPDH 1	2.11
3	Sv_COG	3.41	Sv_ETF 1	3.13	Sv_ETF 1	2.21	Sv_ETF 1	2.45
4	Sv_GAPDH 2	3.94	Sv_GAPDH 2	4.12	Sv_GAPDH 2	3.72	Sv_GAPDH 2	3.72
5	Sv_ETF 1	4.68	Sv_ETF 2	5.00	Sv_TUB A	5.23	Sv_18S rRNA	4.30
6	Sv_TUB A	5.96	Sv_COG	5.5	Sv_ETF 2	6.24	Sv_COG	5.23
7	Sv_ETF 2	6.74	Sv_TUB B	7.24	Sv_18S rRNA	7.33	Sv_ETF 2	6.24
8	Sv_TUB B	8.00	Sv_TUB A	7.74	Sv_TUB B	7.45	Sv_TUB B	8.24
9	Sv_CYP	9.00	Sv_CYP	9.00	Sv_CYP	8.97	Sv_TUB A	8.74
					Sv_COG	9.46	Sv_CYP	10.00

*RefFinder stability values represent geometric means of ranking values

<https://doi.org/10.1371/journal.pone.0183470.t003>

ordered candidate genes according to CV (Table 4). This analysis designated *SvACT* as the most stably expressed gene of our survey in concordance with RT-qPCR-based assays. This candidate reference gene was followed by *SvGAPDH* (primer pair *Sv_GAPDH 1*). As with the RT-qPCR results, the least stable candidate appeared to be *SvCYP*. The ranking generated from RNA-seq data agreed with the RefFinder output obtained for oligo_HF_BL dataset, which was constructed using the same cDNA priming method as used for RNA-seq. In contrast, RefFinder output based on random-primed cDNA differed from RNA-seq ranking, particularly on the stability of *Sv_COG*.

Quantification of *SvMSH1* expression

To demonstrate the utility of the selected reference genes for the normalization of gene expression across various tissues of *S. vulgaris*, we estimated *MSH1* transcript levels in pollen, flower buds, leaves and roots (Fig 3).

The *SvACT* and *SvGAPDH* (primer pair *Sv_GAPDH 1*) reference genes provided generally consistent results with both oligo(dT)- and random-primed cDNA. *SvMSH1* expression was highest in pollen and lowest in roots, where higher standard errors were reported in random-primed cDNA than in oligo(dT)-primed cDNA. Although roots were thoroughly washed

Table 4. Supporting RNA-seq analysis. Rank of analysed genes (primer pairs) according to their coefficients of variation (CV) for RPKM normalized transcript coverage. Maximum fold change (*mfc*) and sample standard deviation (*s*²) are also presented. Oligo(dT)-cDNA libraries were obtained from buds and leaves from H and F plants. RefFinder output for HF_BL is shown for the comparison.

rank	RNA-seq oligo_HF_BL	RPKM	mfc	s ²	CV	RefFinder oligo_HF_BL	RefFinder random_HF_BL
1	Sv_ACT	310	0.55	44	0.14	Sv_ACT	Sv_GAPDH 1
2	Sv_GAPDH 1	453	1.0	122	0.27	Sv_GAPDH 1	Sv_ACT
3	Sv_COG	4.5	2.5	1.5	0.33	Sv_COG	Sv_GAPDH 2
4	Sv_TUB B	12	3.3	4.6	0.38	Sv_GAPDH 2	Sv_ETF 1
5	Sv_GAPDH 2	61	2.7	25	0.41	Sv_ETF 1	Sv_TUB A
6	Sv_TUB A	41	4.1	25	0.61	Sv_TUB A	Sv_ETF 2
7	Sv_ETF 1	275	6.9	178	0.64	Sv_ETF 2	Sv_CYP
8	Sv_ETF 2	311	5.0	205	0.66	Sv_TUB B	Sv_TUB B
9	Sv_CYP	36	400	41	1.13	Sv_CYP	Sv_COG

<https://doi.org/10.1371/journal.pone.0183470.t004>




Fig 3. Relative expression of the *MSH 1* gene. Vegetative organs and pollen of *S. vulgaris* were normalized with SvGAPDH 1 or Sv_ACT primer pairs. **A:** Oligo(dT)-primed cDNAs. **B:** Random hexamer-primed cDNAs. Results from three independent measurements are presented with mean values and standard errors.

<https://doi.org/10.1371/journal.pone.0183470.g003>

before RNA extraction, trace amounts of soil might have interfered with random-primed cDNA synthesis in some samples, which could cause higher variation among the specimens.

There were no statistically significant differences between F and H plants.

Discussion

The choice of appropriate reference genes for the normalization of transcript levels is an essential prerequisite for obtaining accurate RT-qPCR results. Candidate reference genes may be evaluated experimentally, using various statistical software approaches [60, 61, 62, 63]. The increasing availability of comprehensive transcriptomic data has also made *in silico* analyses of gene expression variation possible, as was recently demonstrated in *Silene latifolia* [4]. We also used read coverages calculated from leaf and flower bud transcriptomes of *S. vulgaris* to independently support the expression stability of candidate reference genes. Ultimately, we primarily relied on the experimental approach, because we aimed to find reference genes expressed invariantly across tissues and organs of *S. vulgaris*, including pollen, for which RNA-seq data were not available.

The SvACT and SvGAPDH genes were the least variable candidates across tissues and organs of *S. vulgaris*, regardless of whether pollen was included as determined by RefFinder (Table 3). They also exhibited the lowest CV based on transcriptomic data from leaves and flower buds (Table 4). These two genes were stably expressed with cDNAs generated by both oligo(dT)- and random-priming.

However, the method of cDNA construction appears to have affected the more variable genes—particularly SvCOG. This gene was the least stable candidate in random-primed cDNA from leaves, flower buds and roots, because of low SvCOG transcript levels. As the majority of random-primed cDNA is reverse transcribed from rRNA, the portion of mRNA-derived cDNA is much lower than in oligo(dT)-primed cDNA. Very low concentrations of SvCOG cDNA in the random primed samples made its RT-qPCR measurements less accurate and reproducible.

The RefFinder tool [59] which compiles four algorithms: geNorm [60], NormFinder [61], BestKeeper [62] and the comparative delta-Ct method [63] did not evaluate candidate genes congruently with standalone geNorm and NormFinder software. This discrepancy among various analytical tools has been described previously [23, 65, 66, 67, 68]. However, despite modest inconsistencies, Sv_ACT and Sv_GAPDH 1 primer pairs were consistently ranked among the first three most stable reference genes, no matter which analytical tool was applied.

Both *GAPDH* [5, 23, 48, 49, 50, 65] and *ACT* [38, 39, 40] are often used as reference genes for gene expression quantitation in angiosperms, including the genus *Silene* [4]. The *ACT* gene was reported to be affected by stress conditions [69], but our sample sets did not include stressed individuals. As the stability of reference genes has to be evaluated under specific experimental parameters [56], the selection of reference genes appropriate under stress in *S. vulgaris* would require further validation.

18S rRNA is sometimes utilized for expression normalization in angiosperms [70, 71], but there are also numerous reports about its expression instability [13, 14]. 18S rRNA exhibited variable transcript levels in *S. vulgaris* despite very high expression and cannot be recommended as a reference gene.

We used the *SvACT* and *SvGAPDH* genes to normalize *SvMSH1* transcript levels in oligo(dT)-primed and random primed cDNA. The two methods of cDNA preparation provided generally consistent results, particularly when *SvGAPDH* has been used as a reference. The *MSH1* gene not only influences organelle genome stability, but also impacts DNA methylation, flowering time, branching pattern and abiotic stress tolerance in *Arabidopsis* [72]. It will be beneficial for the study of cyto-nuclear interactions in *S. vulgaris* to estimate both nucleus-encoded *MSH1* transcripts and organellar transcripts in the same cDNA specimens.

In conclusion, we recommend two reference genes, *SvACT* and *SvGAPDH*, to be applied in transcript level estimations across various organs of F and H individuals of *S. vulgaris*. They were selected from a broader set of candidates by an experimental approach and are suitable for random primed as well as oligo(dT)-primed cDNA.

Supporting information

S1 Fig. The outputs from geNorm depicting stability values. A. oligo(dT)-primed cDNA, B. random hexamer-primed cDNA. (PDF)

S1 Table. NormFinder outputs. The expression stability values for oligo(dT)- and random-primed cDNA. (DOCX)

Acknowledgments

We thank Kateřina Haškovcová for excellent technical assistance.

Author Contributions

Conceptualization: Pavla Koloušková, Helena Štorchová.

Data curation: Pavla Koloušková, James D. Stone, Helena Štorchová.

Formal analysis: Pavla Koloušková, James D. Stone, Helena Štorchová.

Funding acquisition: Helena Štorchová.

Methodology: Pavla Koloušková, James D. Stone.

Project administration: Helena Štorchová.

Validation: Helena Štorchová.

Writing – original draft: Pavla Koloušková, James D. Stone, Helena Štorchová.

Writing – review & editing: James D. Stone, Helena Štorchová.

References

1. Wang H, Wang J, Jiang J, Chen S, Guan Z, Liao Y, et al. Reference genes for normalizing transcription in diploid and tetraploid *Arabidopsis*. *Sci Rep* 2014; 4:6781. <https://doi.org/10.1038/srep06781> PMID: 25345678
2. Expósito-Rodríguez M, Borges AA, Borges-Pérez A, Pérez JA. Selection of internal control genes for quantitative real-time RT-PCR studies during tomato development process. *BMC Plant Biol* 2008; 8:131. <https://doi.org/10.1186/1471-2229-8-131> PMID: 19102748
3. Zhu X, Li X, Chen WX, Chen J, Lu W, Chen L, et al. Evaluation of new reference genes in papaya for accurate transcript normalization under different experimental conditions. *PloS One* 2012; 7:e444405. <https://doi.org/10.1371/journal.pone.0044405> PMID: 22952972
4. Zemp N, Minder A, Widmer A. Identification of internal reference genes for gene expression normalization between the two sexes in dioecious white campion. *PloS One* 2014; 9:e92893. <https://doi.org/10.1371/journal.pone.0092893> PMID: 24675788
5. Yang Z, Chen Y, Hu B, Tan Z, Huang B. Identification and validation of reference genes for quantification of target gene expression with quantitative real-time PCR for tall fescue under four abiotic stresses. *PloS One* 2015; 10:e0119569. <https://doi.org/10.1371/journal.pone.0119569> PMID: 25786207
6. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 2009; 55(4): 611–622. <https://doi.org/10.1373/clinchem.2008.112797> PMID: 19246619
7. Boava LP, Laia ML, Jacob TR, Dabbas KM, Gonçalves JF, Ferro JA, et al. Selection of endogenous genes for gene expression studies in *Eucalyptus* under biotic (*Puccinia psidii*) and abiotic (acibenzolar-S-methyl) stresses using RT-qPCR. *BMC Res Notes* 2010; 3:43. <https://doi.org/10.1186/1756-0500-3-43> PMID: 20181283
8. Carvalho K, de Campos MK, Pereira LF, Vieira LG. Reference gene selection for real-time quantitative polymerase chain reaction normalization in "Swingle" citrumelo under drought stress. *Anal Biochem* 2010; 402(2):197–199. <https://doi.org/10.1016/j.ab.2010.03.038> PMID: 20363209
9. Fernández M, Villarreal C, Balbontín C, Valenzuela S. Validation of reference genes for real-time qRT-PCR normalization during cold acclimation in *Eucalyptus globulus*. *Trees-Struct Funct* 2010; 24(6):1109–1116.
10. Garg R, Sahoo A, Tyagi AK, Jain M. Validation of internal control genes for quantitative gene expression studies in chickpea (*Cicer arietinum* L.). *Biochem Biophys Res Commun* 2010; 396(2):283–288. <https://doi.org/10.1016/j.bbrc.2010.04.079> PMID: 20399753
11. Lee JM, Roche JR, Donaghy DJ, Thrush A, Sathish P. Validation of reference genes for quantitative RTP-CR studies of gene expression in perennial ryegrass (*Lolium perenne* L.). *BMC Mol Biol* 2010; 11:8. <https://doi.org/10.1186/1471-2199-11-8> PMID: 20089196
12. Libault M, Thibivilliers S, Bilgin DD, Radwan O, Benitez M, Clough SJ, et al. Identification of four soybean reference genes for gene expression normalization. *Plant Genome* 2008; 1(1):44–54.
13. Xiao XL, Wu XM, Ma JB, Li PB, Li TT, Yao YA. Systematic assessment of reference genes for RT-qPCR across plant species under salt stress and drought stress. *Acta Physiol Plant* 2015; 37(9). <https://doi.org/10.1007/s11738-015-1922-8>
14. Chen X, Mao YJ, Huang SW, Ni J, Lu WL, Hou JY, et al. Selection of Suitable Reference Genes for Quantitative Real-time PCR in *Sapium sebiferum*. *Front Plant Sci* 2017; 8:637. <https://doi.org/10.3389/fpls.2017.00637> PMID: 28523004
15. Wang ML, Li QH, Xin HH, Chen X, Zhu XJ, Li XH. Reliable reference genes for normalization of gene expression data in tea plants (*Camellia sinensis*) exposed to metal stresses. *PloS One* 2017; 4:e0175863. <https://doi.org/10.1371/journal.pone.0175863> PMID: 28453515
16. Long XY, Wang JR, Ouellet T, Rocheleau H, Wei YU, Pu ZE, et al. Genome-wide identification and evaluation of novel internal control genes for Q-PCR based transcript normalization in wheat. *Plant Mol Biol* 2010; 74(3):307–311. <https://doi.org/10.1007/s11103-010-9666-8> PMID: 20658259
17. Jain M. Genome-wide identification of novel internal control genes for normalization of gene expression during various stages of development in rice. *Plant Sci* 2009; 176(5):702–706.

18. Ovesná J, Kučera L, Vaculová K, Štryplová K, Svobodová I, Milella L. Validation of the β -amy1 transcription profiling assay and selection of reference genes suited for a RT-qPCR assay in developing barley caryopsis. *Plos One* 2012; 7(7): e41886. <https://doi.org/10.1371/journal.pone.0041886> PMID: 22860024
19. Podevin N, Krauss A, Henry I., Swennen R, Remy S. Selection and validation of reference genes for quantitative RT-PCR expression studies of the non-model crop *Musa*. *Mol Breed* 2012; 30(3):1237–1252. <https://doi.org/10.1007/s11032-012-9711-1> PMID: 23024595
20. Yeap WC, Loo JM, Wong YC, Kulaveerasingam H. Evaluation of suitable reference genes for qRT-PCR gene expression normalization in reproductive, vegetative tissues and during fruit development in oil palm. *Plant Cell Tiss Organ Cult* 2014; 116(1): 55–66.
21. Freitas NC, Barreto HG, Fernandes-Brum CN, Moreira RO, Chalfun A, Paiva LV. Validation of reference genes for qPCR analysis of *Coffea arabica* L. somatic embryogenesis-related tissues. *Plant Cell Tiss Organ Cult* 2017; 128(3):663–678.
22. Yang Y, Hou S, Cui G, Chen S, Wei J, Huang L. Characterization of reference genes for quantitative real-time PCR analysis in various tissues of *Salvia miltiorrhiza*. *Mol Biol Rep* 2010; 37(1):507–513. <https://doi.org/10.1007/s11033-009-9703-3> PMID: 19680786
23. Karuppaiya P, Yan XX, Liao W, Wu J, Chen F, Tang L. Identification and validation of superior reference gene for gene expression normalization via RT-qPCR in staminate and pistillate flowers of *Jatropha curcas*—A biodiesel plant. *PloS One* 2017; 12:e0172460. <https://doi.org/10.1371/journal.pone.0172460> PMID: 28234941
24. Li WG, Zhang LH, Zhang YD, Wang GD, Song DY, Zhang YW. Selection and validation of appropriate reference genes for quantitative real-time PCR normalization in staminate and perfect flowers of andromonoecious *Taihangia rupestris*. *Front Plant Sci* 2017; 8:729. <https://doi.org/10.3389/fpls.2017.00729> PMID: 28579993
25. Bernasconi G, Antonovics J, Biere A, Charlesworth D, Delph LF, Filatov DA, et al. *Silene* as a model system in ecology and evolution. *Heredity* 2009; 103(1): 5–14. <https://doi.org/10.1038/hdy.2009.34> PMID: 19367316
26. Hanson MR, Bentolila S. Interactions of mitochondrial and nuclear genes that affect male gametophyte development. *Plant Cell* 2004; 16(S): S154–S169.
27. Budar F, Touzet P, De Paepe R. The nucleo-mitochondrial conflict in cytoplasmic male sterilities revisited. *Genetica* 2003; 117(1):3–26. PMID: 12656568
28. Sloan DB, Müller K, McCauley DE, Taylor DR, Štorchová H. Intraspecific variation in mitochondrial genome sequence, structure, and gene content in *Silene vulgaris*, an angiosperm with pervasive cytoplasmic male sterility. *New Phytol* 2012; 196(4):1228–1239. <https://doi.org/10.1111/j.1469-8137.2012.04340.x> PMID: 23009072
29. Sloan DB, Keller SR, Berardi AE, Sanderson BJ, Karpovich JF, Taylor DR. De novo transcriptome assembly and polymorphism detection in the flowering plant *Silene vulgaris* (Caryophyllaceae). *Mol Ecol Resour* 2012; 12(2):333–343. <https://doi.org/10.1111/j.1755-0998.2011.03079.x> PMID: 21999839
30. Blavet N, Charif D, Oger-Desfeux C, Marais GA, Widmer A. Comparative high-throughput transcriptome sequencing and development of SIESTa, the *Silene* EST annotation database. *BMC Genomics* 2011; 12:376. <https://doi.org/10.1186/1471-2164-12-376> PMID: 21791039
31. Gagliardi D, Leaver CJ. Polyadenylation accelerates the degradation of the mitochondrial mRNA associated with cytoplasmic male sterility in sunflower. *EMBO J* 1999; 18(13):3757–3766. <https://doi.org/10.1093/emboj/18.13.3757> PMID: 10393190
32. Lupold DS, Caoile AGFS, Stern DB. Polyadenylation occurs at multiple sites in maize mitochondrial *cox2* mRNA and is independent of editing status. *Plant Cell* 1999; 11(8):1565–1578.
33. Abdelnoor RV, Yule R, Elo A, Christensen AC, Meyer-Gauen G, Mackenzie SA. Substoichiometric shifting in the plant mitochondrial genome is influenced by a gene homologous to MutS. *Proc Natl Acad Sci U S A* 2003; 100(10):5968–5973. <https://doi.org/10.1073/pnas.1037651100> PMID: 12730382
34. Xu YZ, Arrieta-Montiel MP, Viridi KS, de Paula WB, Widhalm JR, Basset GJ, et al. MutS HOMOLOG1 is a nucleoid protein that alters mitochondrial and plastid properties and plant response to high light. *Plant Cell* 2011; 23(9):3428–3441. <https://doi.org/10.1105/tpc.111.089136> PMID: 21934144
35. González-Verdejo CI, Die JV, Nadal S, Jiménez-Marín A, Moreno MT, Román B. Selection of housekeeping genes for normalization by real-time RT-PCR: Analysis of Or-MYB1 gene expression in *Orobancha ramosa* development. *Anal Biochem* 2008; 379(2):176–181. <https://doi.org/10.1016/j.ab.2008.05.003> PMID: 18503743
36. Jain M, Nijhawan A, Tyagi AK, Khurana JP. Validation of housekeeping genes as internal control for studying gene expression in rice by quantitative real-time PCR. *Biochem Biophys Res Commun* 2006; 345(2):646–651. <https://doi.org/10.1016/j.bbrc.2006.04.140> PMID: 16690022

37. Kim BR, Nam HY, Kim SU, Kim SI, Chang YJ. Normalization of reverse transcription quantitative-PCR with housekeeping genes in rice. *Biotechnol Lett* 2003; 25(21):1869–1872. PMID: [14677714](#)
38. Maroufi A, Van Bockstaele E, De Loose M. Validation of reference genes for gene expression analysis in chicory (*Cichorium intybus*) using quantitative real-time PCR. *BMC Mol Biol* 2010; 11:15. <https://doi.org/10.1186/1471-2199-11-15> PMID: [20156357](#)
39. Cháb D, Kolář J, Olson MS, Štorchová H. Two *FLOWERING LOCUS T (FT)* homologs in *Chenopodium rubrum* differ in expression patterns. *Planta* 2008; 228(6):929–940. <https://doi.org/10.1007/s00425-008-0792-3> PMID: [18654796](#)
40. Li QF, Sun SSM, Yuan DY, Yu HX, Gu MH, Liu QQ. Validation of candidate reference genes for the accurate normalization of real-time quantitative RT-PCR data in rice during seed development. *Plant Mol Biol Rep* 2010; 28(1):49–57.
41. Hoenemann C, Hohe A. Selection of reference genes for normalization of quantitative real-time PCR in cell cultures of *Cyclamen persicum*. *Electron J Biotechnol* 2011; 14(1):8. <https://doi.org/10.2225/vol14-issue1-fulltext-8>
42. Jian B, Liu B, Bi YR, Hou WS, Wu CX, Han TF. Validation of internal control for gene expression study in soybean by quantitative real-time PCR. *BMC Mol Biol* 2008; 9:59. <https://doi.org/10.1186/1471-2199-9-59> PMID: [18573215](#)
43. Lin YL, Lai Z.X. Reference gene selection for qPCR analysis during somatic embryogenesis in longan tree. *Plant Sci* 2010; 178(4):359–365.
44. Mallona I, Lischewski S, Weiss J, Hause B, Egea-Cortines M. Validation of reference genes for quantitative real-time PCR during leaf and flower development in *Petunia hybrida*. *BMC Plant Biol* 2010; 10:4. <https://doi.org/10.1186/1471-2229-10-4> PMID: [20056000](#)
45. Nicot N, Hausman JF, Hoffmann L, Le Evers D. Housekeeping gene selection for real-time RT-PCR normalization in potato during biotic and abiotic stress. *J Exp Bot* 2005; 56(421):2907–2914. <https://doi.org/10.1093/jxb/eri285> PMID: [16188960](#)
46. Silveira ED, Alves-Ferreira M, Guimarães LA, da Silva FR, Carneiro VTD. Selection of reference genes for quantitative real-time PCR expression studies in the apomictic and sexual grass *Brachiaria brizantha*. *BMC Plant Biol* 2009; 9:84. <https://doi.org/10.1186/1471-2229-9-84> PMID: [19573233](#)
47. Wan H, Zhao Z, Qian C, Sui Y, Malik AA, Chen J. Selection of appropriate reference genes for gene expression studies by quantitative real-time polymerase chain reaction in cucumber. *Anal Biochem* 2010; 399(2):257–261. <https://doi.org/10.1016/j.ab.2009.12.008> PMID: [20005862](#)
48. Barsalobres-Cavallari CF, Severino FE, Maluf MP, Maia IG. Identification of suitable internal control genes for expression studies in *Coffea arabica* under different experimental conditions. *BMC Mol Biol* 2009; 10:1. <https://doi.org/10.1186/1471-2199-10-1> PMID: [19126214](#)
49. Huis R, Hawkins S, Neutelings G. Selection of reference genes for quantitative gene expression normalization in flax (*Linum usitatissimum* L.). *BMC Plant Biol* 2010; 10:71. <https://doi.org/10.1186/1471-2229-10-71> PMID: [20403198](#)
50. Reid KE, Olsson N, Schlosser J, Peng F, Lund ST. An optimized grapevine RNA isolation procedure and statistical determination of reference genes for real-time RT-PCR during berry development. *BMC Plant Biol* 2006; 6:27. <https://doi.org/10.1186/1471-2229-6-27> PMID: [17105665](#)
51. Hong SY, Seo PJ, Yang MS, Xiang F, Park CM. Exploring valid reference genes for gene expression studies in *Brachypodium distachyon* by real-time PCR. *BMC Plant Biol* 2008; 8:112. <https://doi.org/10.1186/1471-2229-8-112> PMID: [18992143](#)
52. Hu RB, Fan CM, Li HY, Zhang QZ, Fu YF. Evaluation of putative reference genes for gene expression normalization in soybean by quantitative real-time RT-PCR. *BMC Mol Biol* 2009; 10:93. <https://doi.org/10.1186/1471-2199-10-93> PMID: [19785741](#)
53. de Almeida MR, Ruedell CM, Ricachenevsky FK, Sperotto RA, Pasquali G, Fett-Neto AG. Reference gene selection for quantitative reverse transcription-polymerase chain reaction normalization during in vitro adventitious rooting in *Eucalyptus globulus* Labill. *BMC Mol Biol* 2010; 11:73. <https://doi.org/10.1186/1471-2199-11-73> PMID: [20854682](#)
54. Artico S, Nardeli SM, Brilhante O, Grossi De-Sa MF, Alves-Ferreira M. Identification and evaluation of new reference genes in *Gossypium hirsutum* for accurate normalization of real-time quantitative RT-PCR data. *BMC Plant Biol* 2010; 10:49. <https://doi.org/10.1186/1471-2229-10-49> PMID: [20302670](#)
55. Dombrowski JE, Martin RC. Evaluation of reference genes for quantitative RT-PCR in *Lolium temulentum* under abiotic stress. *Plant Sci* 2009; 176(3):390–396.
56. Gimeno J, Eattock N, Van Deynze A, Blumwald E. Selection and validation of reference genes for gene expression analysis in switchgrass (*Panicum virgatum*) using quantitative real-time RT-PCR. *PLoS One* 2014; 9(3): e91474. <https://doi.org/10.1371/journal.pone.0091474> PMID: [24621568](#)

57. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 2012; 9(4):357–359. <https://doi.org/10.1038/nmeth.1923> PMID: 22388286
58. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 2010; 26(6):841–842. <https://doi.org/10.1093/bioinformatics/btq033> PMID: 20110278
59. Xie F, Xiao P, Chen D, Xu L, Zhang B. miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Mol. Biol* 2012; 80(1):75–84.
60. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 2002; 3(7):1–34.
61. Andersen CL, Jensen JL, Ørntoft TF. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res* 2004; 64(15):5245–5250. <https://doi.org/10.1158/0008-5472.CAN-04-0496> PMID: 15289330
62. Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper—Excel-based tool using pair-wise correlations. *Biotechnol Lett* 2004; 26(6):509–515. PMID: 15127793
63. Silver N, Best S, Jiang J, Thein SL. Selection of housekeeping genes for gene expression studies in human reticulocytes using real-time PCR. *BMC Mol Biol* 2006; 7:33. <https://doi.org/10.1186/1471-2199-7-33> PMID: 17026756
64. Havlová M, Dobrev PI, Motyka V, Štorchová H, Libus J, Dobrá J, et al. The role of cytokinins in responses to water deficit in tobacco plants over-expressing *trans*-zeatin O-glucosyltransferase gene under *35S* or *SAG12* promoters. *Plant Cell Environ* 2008; 31(3): 341–353. <https://doi.org/10.1111/j.1365-3040.2007.01766.x> PMID: 18088334
65. de Andrade LM, Brito MD, Peixoto RF, Marchiori PER, Nobile PM, Martins APB, et al. Reference genes for normalization of qPCR assays in sugarcane plants under water deficit. *Plant Methods* 2017; 13:28. <https://doi.org/10.1186/s13007-017-0178-2> PMID: 28428808
66. Wu J, Zhang JH, Pan Y, Huang HH, Lou XZ, Tong ZK. Identification and evaluation of reference genes for normalization in quantitative real-time PCR analysis in the premodel tree *Betula luminifera*. *J For Res* 2017; 28(2):273–282.
67. Sun HP, Li F, Ruan QM, Zhong XH. Identification and validation of reference genes for quantitative real-time PCR studies in *Hedera helix* L. *Plant Physiol Biochem* 2016; 108:286–294. <https://doi.org/10.1016/j.plaphy.2016.07.022> PMID: 27474936
68. Niu XP, Chen MX, Huang XY, Chen HH, Tao AF, Xu JT, et al. Reference gene selection for qRT-PCR normalization analysis in kenaf (*Hibiscus cannabinus* L.) under abiotic stress and hormonal stimuli. *Front Plant Sci* 2017; 8:771. <https://doi.org/10.3389/fpls.2017.00771> PMID: 28553304
69. Wallström SV, Aidemark M, Escobar MA, Rasmusson AG. An alternatively spliced domain of the *NDC1* NAD(P)H dehydrogenase gene strongly influences the expression of the *ACTIN2* reference gene in *Arabidopsis thaliana*. *Plant Sci* 2012; 183:190–196. <https://doi.org/10.1016/j.plantsci.2011.08.011> PMID: 22195593
70. Wang XM, Fu YY, Ban LP, Wang Z, Feng GY, Li J, et al. Selection of reliable reference genes for quantitative real-time RT-PCR in alfalfa. *Genes Genet Syst* 2015; 90(3):175–180. <https://doi.org/10.1266/ggs.90.175> PMID: 26510572
71. Huang YX, Tan HX, Yu J, Chen Y, Guo ZY, Wang GQ, et al. Stable Internal Reference Genes for Normalizing Real-Time Quantitative PCR in *Baphicacanthus cusia* under Hormonal Stimuli and UV Irradiation, and in Different Plant Organs. *Front Plant Sci* 2017; 8:668. <https://doi.org/10.3389/fpls.2017.00668> PMID: 28515733
72. Virdi KS, Laurie JD, Xu YZ, Yu J, Shao MR, Sanchez R, et al. Arabidopsis *MSH1* mutation alters the epigenome and produces heritable changes in plant growth. *Nat Commun* 2015; 6:6386. <https://doi.org/10.1038/ncomms7386> PMID: 25722057