

ORIGINAL RESEARCH

Pooled analysis of drug-related interstitial lung disease and/or pneumonitis in nine trastuzumab deruxtecan monotherapy studies[☆]

C. A. Powell^{1*}, S. Modi², H. Iwata³, S. Takahashi⁴, E. F. Smit⁵, S. Siena^{6,7}, D.-Y. Chang⁸, E. Macpherson⁹, A. Qin¹⁰, J. Singh¹⁰, C. Taitt¹⁰, N. Shire⁹ & D. Ross Camidge¹¹

¹Catherine and Henry J. Gaisman Division of Pulmonary Critical Care and Sleep Medicine, Icahn School of Medicine at Mount Sinai, New York; ²Memorial Sloan Kettering Cancer Center, New York, USA; ³Aichi Cancer Center Hospital, Nagoya, Aichi; ⁴Medical Oncology, The Cancer Institute Hospital of JFCR, Koto, Tokyo, Japan; ⁵Department of Thoracic Oncology, Netherlands Cancer Institute, Amsterdam, the Netherlands; ⁶Department of Oncology and Hemato-Oncology, Università degli Studi di Milano, Milan; ⁷Niguarda Cancer Center, Grande Ospedale Metropolitano Niguarda, Milan, Italy; ⁸National Taiwan University Hospital, Taipei City, Taiwan; ⁹AstraZeneca Pharmaceuticals, Gaithersburg; ¹⁰Daiichi Sankyo Inc., Basking Ridge; ¹¹University of Colorado Cancer Center, Aurora, USA



Available online 11 August 2022

Introduction: This pooled analysis of nine phase I and II trastuzumab deruxtecan (T-DXd) monotherapy studies described drug-related interstitial lung disease (ILD)/pneumonitis in patients treated with T-DXd.

Methods: Patients who received T-DXd across nine studies were included. Investigator-assessed ILD/pneumonitis events were retrospectively reviewed by an independent adjudication committee; events adjudicated as drug-related ILD/pneumonitis are summarized.

Results: The analysis included 1150 patients (breast cancer, 44.3%; gastric cancer, 25.6%; lung cancer, 17.7%; colorectal cancer, 9.3%; other cancer, 3.0%). Median treatment duration was 5.8 (range, 0.7-56.3) months, with a median of 4 (range, 1-27) prior lines of therapy. The overall incidence of adjudicated drug-related ILD/pneumonitis was 15.4% (grade 5, 2.2%). Most patients with ILD/pneumonitis experienced low-grade events (grade 1 or 2, 77.4%); 87.0% had their first event within 12 months [median, 5.4 (range, <0.1-46.8) months] of their first dose of T-DXd. Based on data review, adjudicated ILD/pneumonitis onset occurred earlier than identified by investigators for 53.2% of events [median difference in onset date, 43 (range, 1-499) days]. Stepwise Cox regression identified several baseline factors potentially associated with increased risk of adjudicated drug-related ILD/pneumonitis: age <65 years, enrollment in Japan, T-DXd dose >6.4 mg/kg, oxygen saturation <95%, moderate/severe renal impairment, presence of lung comorbidities, and time since initial diagnosis >4 years.

Conclusions: In this pooled analysis of heavily treated patients, the incidence of ILD/pneumonitis was 15.4%, with most being low grade and occurring in the first 12 months of treatment. The benefit–risk of T-DXd treatment is positive; however, some patients may be at increased risk of developing ILD/pneumonitis, and further investigation is needed to confirm ILD/pneumonitis risk factors. Close monitoring and proactive management of ILD/pneumonitis are warranted for all.

Key words: trastuzumab deruxtecan, adverse event, interstitial lung disease, pneumonitis, HER2

INTRODUCTION

Trastuzumab deruxtecan (T-DXd) is an antibody–drug conjugate (ADC) composed of an anti-human epidermal growth factor receptor 2 (HER2) humanized monoclonal antibody, a

tetrapeptide-based cleavable linker, and a topoisomerase I inhibitor payload, which is present at a high drug-to-antibody ratio (~8 : 1).^{1,2} The linker is stable in plasma but selectively cleaved by cathepsins, which may be up-regulated in cancer cells.²⁻⁵ The released payload is membrane permeable, allowing for an antitumor bystander effect, but has a short half-life, potentially minimizing systemic exposure.^{2,4}

T-DXd is being investigated in a series of trials (DESTINY) in several indications. Data from the phase II DESTINY-Breast01 (NCT03248492; DS8201-A-U201) and DESTINY-Gastric01 (NCT03329690; DS8201-A-J202) studies led to approvals of T-DXd in multiple countries for HER2-positive unresectable or metastatic breast cancer that progressed

*Correspondence to: Dr Charles A. Powell, Catherine and Henry J. Gaisman Division of Pulmonary Critical Care and Sleep Medicine, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Place, Box 1232, New York, NY 10029-5674, USA. Tel: +1-212-241-4280

E-mail: charles.powell@mssm.edu (C. A. Powell).

[☆]Note: Previous presentations: An earlier data cut-off was presented previously (Powell CA, et al. Presented online at: AACR Annual Meeting 2021; April 10-15, 2021. Abstract CT167).

2059-7029/© 2022 The Authors. Published by Elsevier Ltd on behalf of European Society for Medical Oncology. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

on two or more prior therapies (accelerated approval in the USA) and HER2-positive locally advanced or metastatic gastric cancer.⁶⁻¹² Recently, T-DXd has been approved in the USA as a second-line treatment of patients with HER2-positive unresectable or metastatic breast cancer based on data from the randomized phase III DESTINY-Breast03 (NCT03529110; DS8201-A-U302) study of T-DXd versus trastuzumab emtansine (T-DM1) in patients with HER2-positive metastatic breast cancer.^{7,13} Clinical investigation is ongoing in several other populations.¹⁴⁻¹⁸

Interstitial lung disease (ILD)/pneumonitis is a known risk with a variety of cancer therapies, including multiple ADCs.¹⁹⁻²³ T-DXd has demonstrated a generally manageable safety profile across the DESTINY clinical program, with hematologic and gastrointestinal adverse events (AEs) being the most common; ILD/pneumonitis has been identified as an AE of special interest.^{1,6-12,14,18,24-31} The mechanism of T-DXd-related ILD/pneumonitis has not been fully delineated. A recent study in cynomolgus monkeys suggested that alveolar macrophage uptake and redistribution of T-DXd could be involved.³²

Study-specific guidelines for monitoring, evaluating, and managing ILD/pneumonitis in T-DXd clinical trials, including dose modification and supportive care recommendations, were updated in 2019 (Table 1) with more specific details regarding dosages and duration of steroid treatment, including for grade 1 events, than previous guidelines.^{11,14,33} The approved labels for T-DXd also include warnings regarding ILD/pneumonitis and guidelines for its management.⁶⁻¹⁰

To characterize T-DXd-related ILD/pneumonitis, we conducted a pooled analysis of nine phase I and II T-DXd monotherapy studies across multiple tumor types: DS8201-A-J101 (NCT02564900), DS8201-A-J102 (NCT03366428), DS8201-A-A103 (NCT03368196), DS8201-A-A104 (NCT03383692), DESTINY-Breast01, DESTINY-Gastric01, DESTINY-CRC01 (NCT03384940; DS8201-A-J203), DESTINY-Lung01 (NCT03505710; DS8201-A-U204), and DESTINY-Gastric02 (NCT04014075; DS8201-A-U205). This pooled analysis includes data from both before and after the updated guidelines were published and provides a snapshot of the current diagnosis and management of ILD/pneumonitis in T-DXd trials.

MATERIALS AND METHODS

Study populations

Data for these analyses were derived from four phase I and five phase II T-DXd monotherapy studies, which are detailed in Figure 1 and Supplementary Table S1, available at <https://doi.org/10.1016/j.esmooop.2022.100554>. T-DXd doses varied across studies; these analyses included data from patients who received T-DXd 5.4, 6.4, 7.4, or 8.0 mg/kg every 3 weeks (q3w).

Adjudication committee

An independent adjudication committee (AC) was established in November 2017 after the first fatal ILD/pneumonitis event suspected to be T-DXd related was reported by

an investigator during a clinical trial.³⁴ The AC retrospectively reviews all potential ILD/pneumonitis cases [i.e. any AE reported under a defined set of preferred terms (Supplementary material, available at <https://doi.org/10.1016/j.esmooop.2022.100554>)] using imaging [including all available chest computed tomography (CT) images related to the potential ILD/pneumonitis case, such as those acquired at baseline before administration of study drug and subsequent chest CT imaging studies] and clinical data (from baseline to the time of the potential ILD/pneumonitis case) to independently assess whether the reported event was ILD/pneumonitis and, if so, whether it was related to T-DXd. All ILD/pneumonitis events adjudicated as T-DXd related were analyzed for this report.

The AC assessed the grade and date of onset of each event. Grading of ILD/pneumonitis events was based on criteria for grading of pneumonitis in the National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE) version 5.0 (Supplementary material, available at <https://doi.org/10.1016/j.esmooop.2022.100554>).³⁵ In most cases where steroids were administered for asymptomatic events, the ILD/pneumonitis event was adjudicated as grade 2 based on steroid use per CTCAE grading criteria. For cases that were adjudicated after the issuance of current guidelines, all asymptomatic events were considered to be grade 1, regardless of steroid use, consistent with those recommendations (as detailed in the Supplementary material, available at <https://doi.org/10.1016/j.esmooop.2022.100554>). In addition, patients who died and had an ongoing ILD/pneumonitis event assessed by the AC as related to the study drug at the time of death were assessed to determine whether death was due to ILD/pneumonitis.

Statistical analysis

Time of exposure to T-DXd and adjudicated drug-related ILD/pneumonitis by tumor type and grade were summarized. Patients with multiple adjudicated drug-related ILD/pneumonitis events were included in the summary statistics based on the event with the worst grade.

A Kaplan–Meier analysis of the time to the first adjudicated drug-related ILD/pneumonitis event, defined as the time from the first T-DXd dose until adjudicated ILD/pneumonitis onset, was carried out. Treatment discontinuations due to reasons other than ILD/pneumonitis were included as competing events for those who did not have an ILD/pneumonitis event.

Investigator-assessed versus AC-assessed onset dates were compared. For patients who had multiple ILD/pneumonitis events as reported by the investigator that were identified by the AC as a continuous event, the earliest onset reported by the investigator was used for the comparison of onset dates.

The analysis of corticosteroid use in patients with adjudicated drug-related ILD/pneumonitis considered any corticosteroids, regardless of the type, dose, number of doses, or reason for administration, initiated within 90 days of the adjudicated ILD/pneumonitis onset date.

Table 1. Recommended guidelines for management of T-DXd–induced interstitial lung disease (updated in 2019) ¹⁴			
	Grade 1	Grade 2	Grade 3 or 4
Work-up	<ul style="list-style-type: none"> • If a patient develops radiographic changes potentially consistent with ILD/pneumonitis or develops an acute onset of new or worsening pulmonary or other related signs/symptoms such as dyspnea, cough, or fever, rule out ILD/pneumonitis • If the AE is confirmed to have an etiology other than ILD/pneumonitis, follow routine clinical practice • Evaluations should include: <ul style="list-style-type: none"> ◦ High-resolution CT ◦ Pulmonologist consultation (infectious disease consultation as clinically indicated) ◦ Blood culture and CBC. Other blood tests could be considered as needed ◦ Consider bronchoscopy and bronchoalveolar lavage if clinically indicated and feasible ◦ Pulmonary function tests and pulse oximetry (SpO₂) ◦ Arterial blood gases if clinically indicated ◦ One blood sample collection for PK analysis as soon as ILD/pneumonitis is suspected, if feasible ◦ Other tests could be considered, as needed • If the AE is confirmed to be ILD/pneumonitis, follow the ILD/pneumonitis management guidance as outlined below • All events of ILD/pneumonitis regardless of severity or seriousness will be followed until resolution, including after drug discontinuation 		
Dose modification	<ul style="list-style-type: none"> • The administration of T-DXd must be interrupted • T-DXd can be restarted only if the event is fully resolved to grade 0: <ul style="list-style-type: none"> ◦ If resolved in ≤28 days from day of onset, maintain dose ◦ If resolved in >28 days from day of onset, reduce dose 1 level • However, if the grade 1 ILD/pneumonitis occurs beyond day 22 and has not resolved within 49 days from the last infusion, the drug should be discontinued 	Permanently discontinue patient from T-DXd treatment	Permanently discontinue patient from T-DXd treatment
Toxicity management	<ul style="list-style-type: none"> • Monitor and closely follow up in 2-7 days for onset of clinical symptoms and pulse oximetry • Consider follow-up imaging in 1-2 weeks (or as clinically indicated) • Consider starting systemic steroids (e.g. ≥0.5 mg/kg/day of prednisone or equivalent) until improvement, followed by gradual taper over ≥4 weeks • If diagnostic observations worsen despite initiation of corticosteroids, follow grade 2 guidelines. (If the patient is asymptomatic, the toxicity should still be considered as grade 1 even if glucocorticoids were given) 	<ul style="list-style-type: none"> • Promptly start systemic glucocorticoids (e.g. ≥1 mg/kg/day of prednisone or equivalent) for ≥14 days or until complete resolution of clinical and chest CT findings, then followed by gradual taper over ≥4 weeks • Monitor symptoms closely • Re-image as clinically indicated • If clinical or diagnostic observations worsen or do not improve in 5 days: <ul style="list-style-type: none"> ◦ Consider increasing dose of steroids (e.g. 2 mg/kg/day of prednisone or equivalent), switching administration to i.v. (e.g. methylprednisolone) ◦ Reconsider additional work-up for alternative etiologies as described above ◦ Escalate care as clinically indicated 	<ul style="list-style-type: none"> • Hospitalization required • Promptly initiate empirical high-dose methylprednisolone i.v. treatment (e.g. 500-1000 mg/day for 3 days), followed by ≥1 mg/kg/day of prednisone (or equivalent) for ≥14 days or until complete resolution of clinical and chest CT findings, then followed by gradual taper over ≥4 weeks • Re-image as clinically indicated • If still no improvement within 3-5 days: <ul style="list-style-type: none"> ◦ Reconsider additional work-up for alternative etiologies as described above ◦ Consider other immunosuppressants and/or treat per local practice

AE, adverse event; CBC, complete blood count; CT, computed tomography; ILD, interstitial lung disease; i.v., intravenous; PK, pharmacokinetics; q3w, every 3 weeks; T-DXd, trastuzumab deruxtecan.

From *New England Journal of Medicine*, Li BT, Smit ET, Goto Y, et al. Trastuzumab deruxtecan in *HER2*-mutant non–small-cell lung cancer. Volume 386, Issue 3, Pages 241-251. Copyright © 2021 Massachusetts Medical Society. Reprinted with permission from Massachusetts Medical Society.

A multivariate stepwise Cox regression model, stratified by tumor type (breast versus non-breast cancer), was used to explore the association between baseline factors and the time to adjudicated drug-related ILD/pneumonitis, with a stepwise variable selection entry criterion of $P < 0.05$ and remain criterion of $P < 0.10$. Baseline factors included in the model were age group, sex, country, Eastern Cooperative Oncology Group performance status, weight, presence of lung cancer or lung metastases/lymphangitis carcinomatosis, prior chest/lung radiotherapy, presence of lung comorbidity (defined as asthma, chronic obstructive pulmonary disease, prior ILD/pneumonitis, pulmonary fibrosis, pulmonary emphysema, and radiation pneumonitis), renal function, white blood cell count, albumin category, lines of therapy in locally advanced/metastatic setting, time since disease diagnosis category, time from end date of last

anticancer therapy to first infusion of T-DXd category, dose category, and oxygen saturation (SpO₂) category. The categorical cut-offs for potential risk factors were based on medical judgment and practical reasons (e.g. the cut-off of 95% for SpO₂ was selected based on an exploratory analysis of a smaller dataset). This analysis was exploratory and hypothesis generating in nature (see [Supplementary material](https://doi.org/10.1016/j.esmooop.2022.100554), available at <https://doi.org/10.1016/j.esmooop.2022.100554> for more information).

A *post hoc* exploratory analysis was conducted using the Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula, as recommended by the National Kidney Foundation's Kidney Disease Outcomes Quality Initiative,³⁶ to estimate glomerular filtration rate as the method for determining renal function at baseline, instead of the Cockcroft–Gault formula (see [Supplementary material](https://doi.org/10.1016/j.esmooop.2022.100554),




Figure 1. Studies and patients included in the pooled analysis. The colored bar on each arrow indicates the time of patient enrollment and the gray is follow-up.^{1,11,12,14,26,30,31,33,34,43}

All studies noted here are active but no longer recruiting, except for DESTINY-Gastric01 and DESTINY-CRC01, which were completed in 2020.^{12,30} Note that most patients were enrolled before the implementation of toxicity management guidelines.

ILD, interstitial lung disease; NSCLC, non-small-cell lung cancer; Q, quarter.

^aOnly patients who received trastuzumab deruxtecan 5.4, 6.4, 7.4, or 8.0 mg/kg every 3 weeks are included.

available at <https://doi.org/10.1016/j.esmooop.2022.100554> for more information).³⁷

Ethics

All included clinical trials were carried out in accordance with the Declaration of Helsinki and the International Conference on Harmonisation Guidelines for Good Clinical Practice. All patients provided written informed consent before enrollment.

Data sharing statement

Anonymized individual participant data (IPD) and applicable supporting clinical study documents may be available upon request at <https://vivli.org/>. In cases where clinical study data and supporting documents are provided pursuant to our company policies and procedures, Daiichi Sankyo Companies will continue to protect the privacy of company and our clinical study subjects. Details on data sharing criteria and the procedure for requesting access can be found at this web address: <https://vivli.org/ourmember/daiichi-sankyo/>.

RESULTS

Patients and baseline characteristics

Data as of 21 December 2020 were pooled from 1150 patients who received one or more dose of ≥ 5.4 mg/kg T-DXd monotherapy. Median age was 60.0 years (range, 20-96

years); additional baseline characteristics are summarized in Table 2. The most common tumor type was breast cancer [44.3% (510/1150)], followed by gastric cancer [25.6% (294/1150)] (Table 2). Patients had a median of 4 prior treatment regimens (range, 1-27). Most patients [93.9% (1080/1150)] had baseline SpO₂ $\geq 95\%$.

In all patients, the median treatment duration was 5.8 months (range, 0.7-56.3 months) (Table 2 and Supplementary Table S2, available at <https://doi.org/10.1016/j.esmooop.2022.100554>). Nearly half of the patients [49.3% (567/1150)] were treated for >6 months; 24.1% (277/1150) and 7.7% (89/1150) were treated for >12 and >24 months, respectively. Most patients treated for >12 months [69.0% (191/277)] had breast cancer.

ILD/pneumonitis events and adjudication

A total of 276 potential ILD/pneumonitis events were sent for adjudication; 84.8% (234/276) were adjudicated as ILD/pneumonitis (including 224 drug-related and 10 non-drug-related events), 11.6% (32/276) were adjudicated as not being ILD/pneumonitis, and 3.6% (10/276) were pending adjudication at the data cut-off date.

Overall, the AC identified drug-related ILD/pneumonitis in 177 of 1150 patients (15.4%) who had a total of 224 ILD/pneumonitis events. Most of these patients with ILD/pneumonitis [137/177 (77.4%); 11.9% (137/1150) of patients overall] had low-grade (worst grade of 1 or 2) ILD/pneumonitis; 15 patients (1.3%) had a grade 3 or 4 event,

Table 2. Baseline characteristics and T-DXd treatment	
	N = 1150
Age, median (range), years	60.0 (20-96)
≥65 years, n (%)	396 (34.4)
Female, n (%)	755 (65.7)
Country, n (%)	
Japan	506 (44.0)
Non-Japan	644 (56.0)
ECOG PS, n (%)	
0	583 (50.7)
1/2	565 (49.1)/2 (0.2)
Tumor type, n (%) ^a	
Breast cancer	510 (44.3)
Gastric cancer	294 (25.6)
Lung cancer	203 (17.7)
Colorectal cancer	107 (9.3)
Other cancer	34 (3.0)
HER2 expression, n (%) ^b	
Breast cancer	
HER2 overexpressing	398 (34.6)
HER2 low expressing	110 (9.6)
Missing	2 (0.2)
Gastric cancer	
HER2 overexpressing	199 (17.3)
HER2 low expressing	44 (3.8)
Missing	51 (4.4)
Lung cancer	
IHC 3+	38 (3.3)
IHC 2+	89 (7.7)
IHC 1+	13 (1.1)
Missing	63 (5.5)
Colorectal cancer	
IHC 3+	43 (3.7)
IHC 2+	30 (2.6)
IHC 1+	15 (1.3)
Missing	19 (1.7)
Any HER2 mutation, n (%)	
Breast cancer	5 (0.4)
Gastric cancer	0
Lung cancer	75 (6.5)
Colorectal cancer	6 (0.5)
Age by tumor type, median (range), years ^a	
Breast cancer (n = 510)	56 (28-96)
Gastric cancer (n = 294)	65 (20-82)
Lung cancer (n = 203)	62 (23-88)
Colorectal cancer (n = 107)	59 (27-79)
Other cancer (n = 34)	58 (31-76)
Lung cancer or lung metastasis/lymphangitis carcinomatosis at baseline, n (%)	738 (64.2)
Lung comorbidities, n (%) ^c	81 (7.0)
Prior chest/lung radiotherapy, n (%)	190 (16.5)
Time since disease diagnosis, median (range), years ^d	2.98 (<0.1-30.9)
0 to ≤4 years, n (%)	624 (54.3)
>4 years, n (%)	403 (35.0)
No. of prior regimens, median (range) ^d	4.0 (1-27)
Lines of therapy in locally advanced/metastatic setting, n (%) ^d	
1-2	405 (35.2)
3-10	674 (58.6)
>10	40 (3.5)
Time from end date of last anticancer therapy to first infusion of T-DXd, median (range), months ^d	1.41 (0.3-54.5)
<Median (1.41 months), n (%)	518 (45.0)
≥Median (1.41 months), n (%)	531 (46.2)
T-DXd dose, n (%)	
5.4 mg/kg q3w	315 (27.4)
6.4 mg/kg q3w	808 (70.3)
>6.4 mg/kg q3w	27 (2.3)

Continued

Table 2. Continued	
	N = 1150
Duration of treatment, median (range), months	5.81 (0.7-56.3)
0-6 months, n (%)	583 (50.7)
>6-12 months, n (%)	290 (25.2)
>12-24 months, n (%)	188 (16.3)
>24 months, n (%)	89 (7.7)
No. of treatment cycles, median (range)	8.0 (1-76)
WBC count, median (range), ×10 ⁹ /l ^d	5.88 (0.9-77.5)
Weight, median (range), kg	59.70 (27.3-265.8)
Renal function, n (%) ^e	
Normal (serum creatinine clearance ≥90 ml/min)	470 (40.9)
Mild impairment (serum creatinine clearance ≥60 to <90 ml/min)	458 (39.8)
Moderate impairment (serum creatinine clearance ≥30 to <60 ml/min)	193 (16.8)
Severe impairment (serum creatinine clearance <30 ml/min)	3 (0.3)
Missing	26 (2.3)
Baseline albumin, median (range), g/l	39.0 (22-55)
Normal (≥3.5 g/dl), n (%)	937 (81.5)
Mild (≥3.0 to <3.5 g/dl), n (%)	151 (13.1)
Moderate (≥2.5 to <3.0 g/dl), n (%)	46 (4.0)
Severe (<2.5 g/dl), n (%)	3 (0.3)
Missing, n (%)	13 (1.1)
SpO ₂ , n (%)	
≥95%	1080 (93.9)
<95%	57 (5.0)
Missing	13 (1.1)

ECOG PS, Eastern Cooperative Oncology Group performance status; HER2, human epidermal growth factor receptor 2; IHC, immunohistochemistry; q3w, every 3 weeks; SpO₂, oxygen saturation; T-DXd, trastuzumab deruxtecan; WBC, white blood cell.

^aTumor type was missing for two patients.

^bHER2 expression is evaluated using either central or local results according to the protocol defined enrollment criteria. Overexpressing was defined as IHC 3+ or *in situ* hybridization (ISH) positive for breast cancer, IHC 3+ or IHC 2+/ISH-positive for gastric cancer. Low expressing was defined as IHC 2+/ISH-negative or IHC 1+/not ISH-positive for breast cancer, or IHC 2+/ISH-negative or IHC 1+ for gastric cancer.

^cIncludes asthma, chronic obstructive pulmonary disease, prior interstitial lung disease/pneumonitis, pulmonary fibrosis, pulmonary emphysema, and radiation pneumonitis.

^dDue to differences in data collection among the studies, some data were not collected for all patients; thus, the number of patients and percentages may not add up to 100% of the population.

^eRenal function calculated based on creatinine clearance using the Cockcroft–Gault formula.

and 25 (2.2%) had a grade 5 event (Table 3). Overall, 87.0% (154/177) had a first adjudicated drug-related ILD/pneumonitis event within 12 months of starting T-DXd; 13.0% (23/177) had a first adjudicated drug-related ILD/pneumonitis event >12 months after starting T-DXd (Figure 2A).

The median time to adjudicated drug-related ILD/pneumonitis onset was 5.4 months (range, <0.1-46.8 months). The median time to grade 5 onset was 3.2 months (range, <0.1-20.8 months). One patient with preexisting ILD/pneumonitis was adjudicated as having grade 5 ILD/pneumonitis. The ILD/pneumonitis onset date was assessed based on radiographic changes visualized by the AC and clinical features, when applicable. The AC identified ILD/pneumonitis onset earlier than investigators for 53.2% of events [99/186; median difference, 43 days (range, 1-499 days)] (Supplementary Table S3, available at <https://doi.org/10.1016/j.esmoop.2022.100554>). Among patients with

Table 3. Adjudicated drug-related ILD/pneumonitis by tumor type and grade^a

n (%)	Grade 1	Grade 2	Grade 3	Grade 4	Grade 5	Total
All patients (N = 1150)	48 (4.2)	89 (7.7)	14 (1.2)	1 (0.1)	25 (2.2)	177 (15.4)
Breast cancer (n = 510)	32 (6.3)	51 (10.0)	7 (1.4)	0	15 (2.9)	105 (20.6)
HER2-positive breast cancer treated with T-DXd 5.4 mg/kg q3w (n = 245) ^b	9 (3.7)	22 (9.0)	2 (0.8)	0	7 (2.9)	40 (16.3)
Gastric cancer (n = 294)	5 (1.7)	15 (5.1)	3 (1.0)	1 (0.3)	1 (0.3)	25 (8.5)
Lung cancer (n = 203) ^c	7 (3.4)	16 (7.9)	2 (1.0)	0	6 (3.0)	31 (15.3)
Colorectal cancer (n = 107)	0	5 (4.7)	1 (0.9)	0	3 (2.8)	9 (8.4)
Other cancer (n = 34)	4 (11.8)	2 (5.9)	1 (2.9)	0	0	7 (20.6)

HER2, human epidermal growth factor receptor 2; ILD, interstitial lung disease; q3w, every 3 weeks; T-DXd, trastuzumab deruxtecan.

^aPatients with multiple ILD/pneumonitis events are listed only once in this table, based on the event with the highest grade.

^bThe HER2-positive breast cancer population (n = 245) is a subset of the entire breast cancer population (n = 510).

^cAll patients with lung cancer received 6.4 mg/kg q3w of T-DXd.

earlier adjudicated onset according to the AC rather than the investigator, the distribution of investigator-assessed outcomes (based on the worst-grade event) appeared generally similar to the outcomes in patients for whom the investigator-reported and AC-assessed onset dates were the same (Supplementary Table S4, available at <https://doi.org/10.1016/j.esmoop.2022.100554>).

Among the 177 patients with AC-assessed drug-related ILD/pneumonitis, the first event was reported as grade 1 by the investigator in 42.9% of patients (76/177). Of these 76 patients with grade 1 ILD/pneumonitis, 47 were subsequently rechallenged with T-DXd as recommended (Table 1), and 3 of these 47 had a subsequent adjudicated (worst grade of 1) drug-related ILD/pneumonitis event (with onset ranging from 84 to 177 days after onset of the first adjudicated event).

Cox multivariate regression analysis

Stepwise Cox regression evaluated whether there were associations between potential factors of interest and hazard of any-grade adjudicated drug-related ILD/pneumonitis. Seven baseline factors of interest were identified based on significance of association with the incidence of ILD/pneumonitis: age, enrollment in Japan, T-DXd dose, SpO₂, moderate/severe renal impairment (based on the Cockcroft–Gault formula³⁷), presence of lung comorbidities (not including lung cancer), and time since initial diagnosis (Figure 2B).

In the *post hoc* sensitivity analysis using the CKD-EPI formula³⁶ rather than the Cockcroft–Gault formula, baseline renal function (as a categorical factor) was not found to be associated with the risk of adjudicated drug-related ILD/pneumonitis (Supplementary Figure S1, available at <https://doi.org/10.1016/j.esmoop.2022.100554>).

Additional analyses of clinical importance

For 80.2% of grade 2-4 adjudicated drug-related ILD/pneumonitis events (85/106) and 84.6% of grade 5 events (22/26; 1 patient who died was listed with 2 events), one or more dose of corticosteroids was administered within 90 days of the adjudicated ILD/pneumonitis onset date. Due to the limitation of data collection, it is unknown how many of these patients received the optimal steroid treatment per

the published ILD/pneumonitis management guidelines and study protocols.

The majority of patients (80.3%; 924/1150) did not receive prior immune checkpoint inhibitors (ICIs) (Supplementary Table S5, available at <https://doi.org/10.1016/j.esmoop.2022.100554>). The rate of any-grade adjudicated drug-related ILD/pneumonitis was 15.9% (147/924) in patients without prior ICI use [grade ≥3 in 3.6% (33/924)] and 13.3% (30/225) in patients with prior ICI use [grade ≥3 in 3.1% (7/225)]. Gastric cancer [n = 72/294 (24.5%)] and lung cancer [n = 135/203 (66.5%)] cohorts had the largest numbers of patients with prior ICI use, compared with <10 patients in each of the other cohorts. Adjudicated drug-related ILD/pneumonitis rates according to prior ICI use for each tumor type are summarized in Supplementary Table S5, available at <https://doi.org/10.1016/j.esmoop.2022.100554>.

DISCUSSION

T-DXd has shown antitumor activity in HER2-positive metastatic breast, gastric, and lung cancer; HER2-low metastatic breast cancer; HER2-mutant metastatic lung cancer; and other tumor types.^{12,14,18,27,28,38} This pooled analysis represents the most comprehensive evaluation of ILD/pneumonitis from the T-DXd clinical program, comprising 1150 heavily pretreated patients from nine phase I and II clinical trials. Overall, 177 patients (15.4%) experienced adjudicated drug-related ILD/pneumonitis, with most (77.4%) experiencing grade 1 or 2 events. In most patients (87.0%), the event occurred within 12 months of the first dose of T-DXd; median treatment duration was 5.8 months (range, 0.7-56.3 months). The longer T-DXd exposure in patients with breast cancer versus other tumor types may be due to these patients having longer progression-free survival and thus a longer treatment duration. It is notable that the median time to the first ILD/pneumonitis event was shorter for grade 5 events compared with the overall median time, suggesting that further investigation of this issue may be warranted. This represents the first data on T-DXd rechallenge following resolution of grade 1 ILD/pneumonitis, with only 3 of 47 rechallenged patients having recurrence of ILD/pneumonitis. Future research is warranted to better understand rechallenge with T-DXd.




Figure 2. Analysis of adjudicated drug-related ILD/pneumonitis events. (A) Kaplan–Meier analysis of time to first adjudicated drug-related ILD/pneumonitis event. Among 177 patients who had ILD/pneumonitis, 154 (87.0%) had a first ILD/pneumonitis event within 12 months of starting treatment. The median time to adjudicated ILD/pneumonitis onset among those with ILD/pneumonitis was 5.4 months (range, <0.1–46.8 months). The median treatment duration in all patients in the pool was 5.8 months (range, 0.7–56.3 months), and 24.1% of all patients remained on treatment for >12 months. Treatment discontinuations due to reasons other than ILD/pneumonitis were included as a competing event. (B) Multivariate stepwise Cox regression analysis, final model.

The AC frequently identified ILD/pneumonitis onset as occurring earlier than did investigators, suggesting a need for additional ongoing vigilance. It is important to note, however, that the AC retrospectively reviewed patient records, with access to all clinical data and imaging scans during their assessment, whereas the investigator may not have had access to the same information during their assessment. While treating patients receiving T-DXd, health care providers should review prior CT scans for an assessment of ILD/pneumonitis along with their review of all medical records.

The rates of ILD/pneumonitis recently reported from the phase III randomized DESTINY-Breast03 trial of T-DXd as a second-line treatment (>70% of patients had received two or fewer prior lines of therapy in the metastatic setting) in patients with HER2-positive metastatic breast cancer were lower than those reported here.¹³ In DESTINY-Breast03, 10.5% of patients (27/257) treated with T-DXd experienced adjudicated drug-related ILD/pneumonitis, which was predominantly grade 1 or 2 [92.6% of patients (25/27)], with no grade 4 or 5 events.¹³ Increased knowledge of ILD/pneumonitis and implementation of ILD/pneumonitis monitoring, diagnosis, and management guidelines may explain why there were no grade 4 or 5 events since ILD/pneumonitis events may have been identified and treated early before they progressed.²³ DESTINY-Breast03 investigated T-DXd as an earlier line of therapy than the trials included in this pooled analysis [median prior lines of therapy, 4 (range, 1-27)], which implies that perhaps there is a lower rate of ILD/pneumonitis in patients who are less heavily pretreated, although line of therapy was not identified as a potential factor of interest in the multivariate analysis reported here. The present analysis reflects a heavily pretreated population overall, including data from multiple tumor types, which may also partially explain differences in the rates of ILD/pneumonitis observed between the pooled analysis and DESTINY-Breast03. Several ongoing trials will continue to define the ILD/pneumonitis landscape in T-DXd.

A multivariate analysis was conducted to analyze factors potentially associated with ILD/pneumonitis. The analysis identified seven potential factors of interest with an increased hazard of drug-related ILD/pneumonitis; however, the clinical relevance of some of these factors remains unclear. Doses of T-DXd >6.4 mg/kg q3w had a significant association with increased ILD/pneumonitis hazard; for this reason, subsequent trials of T-DXd have treated patients at a dose of ≤6.4 mg/kg q3w.^{18,39,40}

Most patients (70.3%) included in this pooled analysis received T-DXd 6.4 mg/kg q3w, which is the recommended dose approved for patients with certain types of gastric cancer and the dose being investigated in lung and colorectal cancer.^{6,7,14,31} Importantly, we did not find a significant difference in risk of ILD/pneumonitis between the 5.4 mg/kg q3w and the 6.4 mg/kg q3w doses in the multivariate Cox regression analysis. Thus, these data indicating a limited risk imposed by the 6.4 mg/kg q3w dose compared with the 5.4 mg/kg q3w dose of T-DXd provide valuable knowledge to clinicians regarding T-DXd-related ILD/pneumonitis when treating patients with multiple tumor types.

Baseline SpO₂ <95% also posed a significant risk, but this was likely driven by a few patients with particularly low baseline SpO₂. It is important to note that the 95% SpO₂ threshold was chosen for practical (e.g. for sufficient patient numbers in each subset) rather than biological or clinical reasons, which precludes comprehensive assessment of the impact of SpO₂.

Moderate or severe baseline renal impairment was also identified as a factor of interest in the preplanned multivariate analysis that used the Cockcroft–Gault formula; however, a subsequent *post hoc* analysis using the CKD-EPI formula showed a lack of association between severity of renal impairment (categorical) and ILD/pneumonitis risk. This analysis was completed because the CKD-EPI formula is Food and Drug Administration recommended for patients with renal impairment.⁴¹ The identification of renal function continues to be a clinical factor of interest for T-DXd-related ILD/pneumonitis and has been identified as a risk factor for ILD/pneumonitis related to other cancer therapies²³; it is being explored in ongoing trials, including the randomized DESTINY-Breast03 trial.

The multivariate analysis suggested that younger patients (<65 years old) had a higher hazard of ILD/pneumonitis (after adjusting for other covariates selected by the model). This is not intuitive but may be due to unmeasured confounding factors among other clinical characteristics.

Patients treated with T-DXd in Japan had a higher hazard of ILD/pneumonitis compared with those treated outside of Japan. It is unknown whether this is due to biological factors or differences in monitoring and management practices in Japan; however, it is consistent with prior studies suggesting that other drug-induced lung injuries are more common in Japan than elsewhere.^{20,23,33,42}

The presence of lung comorbidities was identified as a factor of interest in the multivariate model; however, lung

Factors included in the final model were age group, sex, country, Eastern Cooperative Oncology Group performance status, baseline weight, presence of lung cancer or lung metastasis/lymphangitis carcinomatosis at baseline, prior chest/lung radiotherapy, presence of lung comorbidity, baseline renal function, baseline white blood cell count, baseline albumin category, number of prior lines of therapy in the locally advanced/metastatic setting, time since disease diagnosis category, time since the end date of the last anticancer therapy to the first infusion of trastuzumab deruxtecan category, dose category, and baseline SpO₂ category. Of these, seven factors were identified as factors of interest.

HER2, human epidermal growth factor receptor 2; ILD, interstitial lung disease; q3w, every 3 weeks; Ref, reference; SpO₂, oxygen saturation.

^aHazard ratios are presented relative to the reference categories indicated.

^bIncludes asthma, chronic obstructive pulmonary disease, prior interstitial lung disease/pneumonitis, pulmonary fibrosis, pulmonary emphysema, and radiation pneumonitis.

^cDue to differences in data collection among the studies, some data were not collected for all patients; thus, the number of patients may not add up to the total population.

^dDetermined by Cockcroft–Gault formula.³⁷

cancer and lung metastases/lymphangitic carcinomatosis at baseline and prior chest/lung radiotherapy were not associated with ILD/pneumonitis in this analysis. While drug-related ILD/pneumonitis could be hypothesized to be related to a direct depot effect from binding to tumor deposits in the lung, the presence of ILD/pneumonitis in patients without lung involvement argues against this.

A *post hoc* analysis according to prior ICI use showed numerically similar rates of adjudicated drug-related ILD/pneumonitis in patients with or without prior ICI use.

Limitations

In this analysis, the exact background rate of ILD/pneumonitis could not be determined because the AC did not review all patients; rather, a broad set of AE terms, including but not limited to ‘pneumonitis’ and ‘ILD,’ was used to trigger adjudication. Additionally, data suggest that steroid management was not optimal, with a delay in the detection of ILD/pneumonitis and dosing of steroids that was inconsistent with the recommendations in the ILD/pneumonitis treatment guidelines (e.g. many patients with grade ≥ 2 ILD/pneumonitis did not receive corticosteroids). Because steroid use (e.g. duration and route of administration) was inconsistently recorded in the clinical trial databases, we were not able to definitively distinguish cases in which steroids were administered in accordance with the current guidelines from those in which patients may have received an incomplete course of steroids or whose steroid treatment was unrelated to ILD/pneumonitis (e.g. topical steroids). The new guidelines for managing T-DXd-related ILD/pneumonitis that were implemented in December 2019¹¹ were provided to health care providers and patients, along with specific education. However, many of the adjudicated ILD/pneumonitis events in these studies occurred before these guidelines were implemented.

The thresholds for the factors assessed by Cox multivariate regression were preselected based on medical judgment for practical reasons, which may impede the assessment of their impact on the risk of ILD/pneumonitis.

These analyses were also limited by the heterogeneity of the population (e.g. multiple tumor types, doses given) and duration of treatment. Due to this data heterogeneity, it is not possible to risk stratify patients. Nevertheless, the potential clinical factors of interest identified in these analyses are informative for clinicians on what to be aware of while treating patients with T-DXd. Future analyses of more homogeneous populations from randomized clinical trials will help confirm these findings and allow for further understanding of any relevant risk factors (e.g. specific to different tumor types).

Conclusions

The overall incidence of adjudicated drug-related ILD/pneumonitis in T-DXd-treated patients was 15.4% across all doses and tumor types in this heavily pretreated population. Most cases were low grade, with a median time to onset of 5.4 months (range, <0.1-46.8 months). The median treatment duration was 5.8 months (range, 0.7-56.3 months).

Risk-over-time assessments suggest that although ILD/pneumonitis development may require some cumulative exposure, a significant risk plateau was achieved after ~ 12 months, and those who will develop this complication usually do so within 12 months of the first dose of T-DXd. This is based primarily on longer treatment duration in patients with breast cancer and suggests some individuals are inherently at lower risk of these events despite prolonged exposure. Longer T-DXd treatment duration and follow-up in patients with other tumor types are needed to confirm the observed trend. The monitoring, diagnosis, and management of ILD/pneumonitis is an area of continuing improvement; to this end, new toxicity guidelines have been implemented, and education has been provided to health care providers and patients. Patient awareness and ongoing education are critical elements to aid in early detection. In the DESTINY-Breast03 trial, lower rates of ILD/pneumonitis have been observed, with no patients experiencing a grade 4 or 5 event; these outcomes are likely due to a confluence of factors, including the less heavily pretreated study population and updated guidance for ILD/pneumonitis monitoring and management.¹³ Rechallenge with T-DXd after complete resolution of grade 1 events is possible and warrants further exploration; however, rechallenge is not recommended for all patients (e.g. patients with grade ≥ 2 ILD/pneumonitis). Potential clinical factors of interest for ILD/pneumonitis may include low SpO₂, lung comorbidities, and renal impairment. Specific risk factors should be confirmed in ongoing and future trials. Phase III randomized controlled trials across multiple tumor types and in earlier lines of therapy are ongoing, and this analysis further supports the benefit–risk profile of T-DXd in advanced cancer.

ACKNOWLEDGEMENTS

Medical writing support was provided by JoAnna Anderson, PhD, CMPP, of ArticulateScience, LLC, and funded by AstraZeneca in accordance with Good Publication Practice (GPP3) guidelines (<http://www.ismpp.org/gpp3>).

FUNDING

This work was supported by AstraZeneca Pharmaceuticals (no grant number) and Daiichi Sankyo Company (no grant number). Financial support for medical writing was provided by AstraZeneca. The manuscript was reviewed for medical accuracy by AstraZeneca and Daiichi Sankyo; however, the authors retained full control of the content and made the final decisions for all aspects of this article. In March 2019, AstraZeneca entered into a global development and commercialization collaboration agreement with Daiichi Sankyo for trastuzumab deruxtecan (T-DXd; DS-8201).

DISCLOSURE

All authors received nonfinancial support (assistance with manuscript preparation) from ArticulateScience, LLC, funded by AstraZeneca. Additional disclosures are as follows: CAP reports personal fees from Daiichi Sankyo, AstraZeneca, and Voluntas, outside of the submitted work. SM reports consulting or advisory roles for Daiichi Sankyo,

MacroGenics, and AstraZeneca; speakers bureau involvement with Genentech, Daiichi Sankyo, AstraZeneca, and Seagen; travel fees from Genentech and Daiichi Sankyo; honoraria from Novartis; and research funding from Roche/Genentech, Novartis, Seagen, Synta, and Daiichi Sankyo. HI reports grants from Daiichi Sankyo, Novartis, AstraZeneca, Pfizer, Eli Lilly Japan, Chugai, Eisai, Kyowa Kirin, Merck Sharp & Dohme (MSD), GSK, Nippon Kayaku, and Bayer; and personal fees from Daiichi Sankyo, Novartis, AstraZeneca, Pfizer, Eli Lilly Japan, Chugai, Eisai, and Kyowa Kirin. ST reports grants and personal fees from Daiichi Sankyo, Eisai, Novartis, Taiho, MSD, Chugai, Bayer, and AstraZeneca. EFS reports consulting or advisory roles for Lilly, AstraZeneca, Boehringer Ingelheim, Roche/Genentech, Bristol Myers Squibb, Merck KGaA, MSD Oncology, Takeda, and Bayer; and research funding from Boehringer Ingelheim, Bayer, Roche/Genentech, AstraZeneca, and Bristol Myers Squibb. SS reports personal fees from Amgen, Roche/Genentech, Bayer, Bristol Myers Squibb, Daiichi Sankyo, Merck KGaA, Seagen, Checkmab, and AstraZeneca. DYC reports personal fees from Novartis, Pfizer, Eli Lilly, Roche, AstraZeneca, Sanofi, Amgen, and Daiichi Sankyo; and travel fees from Pfizer, Daiichi Sankyo, and Eisai. EM and NS are employees of AstraZeneca and may own stock or stock options in that company. AQ, JS, and CT are employees of Daiichi Sankyo and may own stock or stock options in that company. DRC reports personal fees from Daiichi Sankyo.

REFERENCES

- Doi T, Shitara K, Naito Y, et al. Safety, pharmacokinetics, and anti-tumour activity of trastuzumab deruxtecan (DS-8201), a HER2-targeting antibody-drug conjugate, in patients with advanced breast and gastric or gastro-oesophageal tumours: a phase 1 dose-escalation study. *Lancet Oncol*. 2017;18(11):1512-1522.
- Ogitani Y, Aida T, Hagihara K, et al. DS-8201a, a novel HER2-targeting ADC with a novel DNA topoisomerase I inhibitor, demonstrates a promising antitumor efficacy with differentiation from T-DM1. *Clin Cancer Res*. 2016;22(20):5097-5108.
- Aggarwal N, Sloane BF. Cathepsin B: multiple roles in cancer. *Proteomics Clin Appl*. 2014;8(5-6):427-437.
- Nakada T, Sugihara K, Jikoh T, Abe Y, Agatsuma T. The latest research and development into the antibody-drug conjugate, [fam-] trastuzumab deruxtecan (DS-8201a), for HER2 cancer therapy. *Chem Pharm Bull (Tokyo)*. 2019;67(3):173-185.
- Ruan J, Zheng H, Fu W, Zhao P, Su N, Luo R. Increased expression of cathepsin L: a novel independent prognostic marker of worse outcome in hepatocellular carcinoma patients. *PLoS One*. 2014;9(11):e112136.
- Enhertu (trastuzumab deruxtecan). *Summary of product characteristics*. Daiichi Sankyo Co, Ltd; 2020. Available at <https://www.pmda.go.jp/files/000238707.pdf>. Accessed February 17, 2022.
- Enhertu (fam-trastuzumab deruxtecan-nxki). *Prescribing information*. Daiichi Sankyo, Inc; 2022. Available at <https://daiichisankyo.us/prescribing-information-portlet/getPIContent?productName=Enhertu&inline=true>. Accessed May 18, 2022.
- Enhertu (trastuzumab deruxtecan). *Summary of product characteristics*. Daiichi Sankyo Europe GmbH; 2021. Available at https://www.ema.europa.eu/en/documents/product-information/enhertu-epar-product-information_en.pdf. Accessed February 17, 2022.
- Enhertu (trastuzumab deruxtecan). *Summary of product characteristics*. Daiichi Sankyo UK Ltd; 2021. Available at <https://www.medicines.org.uk/emc/product/12135/smpc/print>. Accessed February 17, 2022.
- Enhertu (trastuzumab deruxtecan). *Product monograph*. AstraZeneca Canada Inc; 2021. Available at <https://daiichisankyo.us/prescribing-information-portlet/getPIContent?productName=Enhertu&inline=true>. Accessed February 17, 2022.
- Modi S, Saura C, Yamashita T, et al. Trastuzumab deruxtecan in previously treated HER2-positive breast cancer. *N Engl J Med*. 2020;382(7):610-621.
- Shitara K, Bang YJ, Iwasa S, et al. Trastuzumab deruxtecan in previously treated HER2-positive gastric cancer. *N Engl J Med*. 2020;382(25):2419-2430.
- Cortes J, Kim SB, Chung WP, et al. Trastuzumab deruxtecan versus trastuzumab emtansine for breast cancer. *N Engl J Med*. 2022;386(12):1143-1154.
- Li BT, Smit EF, Goto Y, et al. Trastuzumab deruxtecan in HER2-mutant non-small-cell lung cancer. *N Engl J Med*. 2022;386(3):241-251.
- Fuchs CS, Shahidi J, Mathew L, Qin A, Cutsem EV. A phase II trial of [fam-] trastuzumab deruxtecan (T-DXd, DS-8201a) in subjects with HER2-positive, unresectable, or metastatic gastric or gastroesophageal junction (GEJ) adenocarcinoma. *J Clin Oncol*. 2020;38(suppl 4): Abstract TPS460.
- Janjigian YY, Viglianti N, Liu F, Mendoza-Naranjo A, Croydon L. A phase Ib/II, multicenter, open-label, dose-escalation, and dose-expansion study evaluating trastuzumab deruxtecan (T-DXd, DS-8201) monotherapy and combinations in patients with HER2-overexpressing gastric cancer (DESTINY-Gastric03). *J Clin Oncol*. 2021;39(suppl 3): Abstract TPS261.
- ClinicalTrials.gov. Trastuzumab deruxtecan in participants with HER2-overexpressing advanced or metastatic colorectal cancer (DESTINY-CRC02). Available at, <https://clinicaltrials.gov/ct2/show/NCT04744831>. Accessed May 13, 2021.
- Modi S, Jacot W, Yamashita T, et al. Trastuzumab deruxtecan in previously treated HER2-low advanced breast cancer. *N Engl J Med*. 2022;387(1):9-20.
- Camus P, Kudoh S, Ebina M. Interstitial lung disease associated with drug therapy. *Br J Cancer*. 2004;91. Suppl 2(suppl 2):S18-S23.
- Skeoch S, Weatherley N, Swift AJ, et al. Drug-induced interstitial lung disease: a systematic review. *J Clin Med*. 2018;7(10):356.
- Jänne PA, Baik CS, Su W-C, et al. Efficacy and safety of patritumab deruxtecan (HER3-DXd) in EGFR inhibitor-resistant, EGFR-mutated non-small cell lung cancer. *Cancer Discov*. 2022;12(1):74-89.
- Spira A, Lisberg A, Sands J, et al. OA03. 03 Datoptomab Deruxtecan (Dato-DXd; DS-1062), a TROP2 ADC, in patients with advanced NSCLC: updated results of TROPION-PanTumor01 phase 1 study. *J Thorac Oncol*. 2021;16(3):S106-S107.
- Conte P, Ascierto PA, Patelli G, et al. Drug-induced interstitial lung disease during cancer therapies: expert opinion on diagnosis and treatment. *ESMO Open*. 2022;7(2):100404.
- Modi S, Park H, Murthy RK, et al. Antitumor activity and safety of trastuzumab deruxtecan in patients with HER2-low-expressing advanced breast cancer: results from a phase Ib study. *J Clin Oncol*. 2020;38(17):1887-1896.
- Shitara K, Iwata H, Takahashi S, et al. Trastuzumab deruxtecan (DS-8201a) in patients with advanced HER2-positive gastric cancer: a dose-expansion, phase 1 study. *Lancet Oncol*. 2019;20(6):827-836.
- Tamura K, Tsurutani J, Takahashi S, et al. Trastuzumab deruxtecan (DS-8201a) in patients with advanced HER2-positive breast cancer previously treated with trastuzumab emtansine: a dose-expansion, phase 1 study. *Lancet Oncol*. 2019;20(6):816-826.
- Tsurutani J, Iwata H, Krop I, et al. Targeting HER2 with trastuzumab deruxtecan: a dose-expansion, phase 1 study in multiple advanced solid tumors. *Cancer Discov*. 2020;10(5):688-701.
- Modi S, Saura C, Yamashita T, et al. Updated results from DESTINY-Breast01, a phase 2 trial of trastuzumab deruxtecan (T-DXd) in HER2 positive metastatic breast cancer. Presented at San Antonio Breast Cancer Symposium. December 8-11, 2020; San Antonio, TX, virtual. Abstract PD3-06.
- Powell CA, Modi S, Iwata H, et al. Pooled analysis of drug-related interstitial lung disease in 8 single-arm trastuzumab deruxtecan studies. Presented at AACR Annual Meeting 2021; April 10-15, 2021, virtual. Abstract CT167.
- Siena S, Bartolomeo MD, Raghav KPS, et al. A phase II, multicenter, open-label study of trastuzumab deruxtecan (T-DXd; DS-8201) in patients (pts) with HER2-expressing metastatic colorectal cancer (mCRC):

- DESTINY-CRC01. Presented at American Society of Clinical Oncology Annual Meeting 2020. May 29-31, 2020; virtual. Abstract 4000.
31. Siena S, Di Bartolomeo M, Raghav K, et al. Trastuzumab deruxtecan (DS-8201) in patients with HER2-expressing metastatic colorectal cancer (DESTINY-CRC01): a multicentre, open-label, phase 2 trial. *Lancet Oncol*. 2021;22(6):779-789.
 32. Kumagai K, Aida T, Tsuchiya Y, Kishino Y, Kai K, Mori K. Interstitial pneumonitis related to trastuzumab deruxtecan, a human epidermal growth factor receptor 2-targeting Ab-drug conjugate, in monkeys. *Cancer Sci*. 2020;111(12):4636-4645.
 33. Kubo K, Azuma A, Kanazawa M, et al. Consensus statement for the diagnosis and treatment of drug-induced lung injuries. *Respir Investig*. 2013;51(4):260-277.
 34. Center for Drug Evaluation and Research. Multi-discipline review: Enhertu. Available at https://www.accessdata.fda.gov/drugsatfda_docs/nda/2019/761139Orig1s000MultidisciplineR.pdf. Accessed April 29, 2021.
 35. National Cancer Institute. Common Terminology Criteria for Adverse Events (CTCAE) version 5.0. Available at https://ctep.cancer.gov/protocol-development/electronic_applications/docs/ctcae_v5_quick_reference_5x7.pdf. Accessed July 27, 2021.
 36. Vassalotti JA, Centor R, Turner BJ, Greer RC, Choi M, Sequist TD. Practical approach to detection and management of chronic kidney disease for the primary care clinician. *Am J Med*. 2016;129(2):153-162.e7.
 37. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976;16(1):31-41.
 38. Nakagawa K, Nagasaka M, Felip E, et al. OA04.05 Trastuzumab deruxtecan in HER2-overexpressing metastatic non-small cell lung cancer: interim results of DESTINY-Lung01. *J Thorac Oncol*. 2021;16(3):S109-S110.
 39. Takahashi S, Karayama M, Takahashi M, et al. Pharmacokinetics, safety, and efficacy of trastuzumab deruxtecan with concomitant ritonavir or itraconazole in patients with HER2-expressing advanced solid tumors. *Clin Cancer Res*. 2021;37:5771-5780.
 40. Bardia A, Barrios C, Dent R, et al. Abstract OT-03-09: Trastuzumab deruxtecan (T-DXd; DS-8201) vs investigator's choice of chemotherapy in patients with hormone receptor-positive (HR+), HER2 low metastatic breast cancer whose disease has progressed on endocrine therapy in the metastatic setting: a randomized, global phase 3 trial (DESTINY-Breast06). *Cancer Res*. 2021;81(suppl 4):OT-03-09.
 41. US Food and Drug Administration (2020). Guidance for industry: pharmacokinetics in patients with impaired renal function-study design, data analysis, and impact on dosing. Available at <https://www.fda.gov/media/78573/download>. Accessed November 14, 2021.
 42. Iwasa E, Fujiyoshi Y, Kubota Y, et al. Interstitial lung disease as an adverse drug reaction in Japan: exploration of regulatory actions as a basis for high reporting. *Drug Saf*. 2020;43(11):1121-1131.
 43. Kotani D, Shitara K. Trastuzumab deruxtecan for the treatment of patients with HER2-positive gastric cancer. *Ther Adv in Med Oncol*. 2021;13:1758835920986518.