



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Molecular docking of potential SARS-CoV-2 papain-like protease inhibitors



Daoqun Li ^{a, b, c}, Junwen Luan ^{a, b, c}, Leiliang Zhang ^{a, b, c, *}

^a Department of Laboratory Medicine, The First Affiliated Hospital of Shandong First Medical University, Jinan, China

^b Institute of Basic Medicine, Shandong First Medical University and Shandong Academy of Medical Sciences, Jinan, China

^c Science and Technology Innovation Center, Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan, China

ARTICLE INFO

Article history:

Received 2 November 2020

Accepted 17 November 2020

Available online 28 November 2020

Keywords:

SARS-CoV-2

COVID-19

Papain-like protease

Molecular docking

ABSTRACT

SARS-CoV-2 papain-like protease is considered as an important potential target for anti-SARS-CoV-2 drug discovery due to its crucial roles in viral spread and innate immunity. Here, we have utilized an *in silico* molecular docking approach to identify the possible inhibitors of the SARS-CoV-2 papain-like protease, by screening 21 antiviral, antifungal and anticancer compounds. Among them, Neobavaisoflavone has the highest binding energy for SARS-CoV-2 papain-like protease. These molecules could bind near the SARS-CoV-2 papain-like protease crucial catalytic triad, ubiquitination and ISGylation residues: Trp106, Asn109, Cys111, Met208, Lys232, Pro247, Tyr268, Gln269, His272, Asp286 and Thr301. Because blocking the papain-like protease is an important strategy in fighting against viruses, these compounds might be promising candidates for therapeutic intervention against COVID-19.

© 2020 Elsevier Inc. All rights reserved.

1. Introduction

The coronavirus disease 2019 (COVID-19) pandemic caused by the new severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) poses an unprecedented challenge [1,2]. Globally, as of October 21, 2020, more than 40.6 million cases have been reported to the World Health Organization, and 1,121,843 deaths have occurred in more than 235 countries. The scale of the pandemic and the rapidly spread make COVID-19 one of the most serious diseases facing mankind so far.

At present, there is no vaccine or approved antiviral treatment for SARS-CoV-2. In the extensive research on the development of new antiviral drugs for COVID-19, main protease of SARS-CoV-2, which plays an important role in blocking viral infection, has been identified as a potential target [3]. Another attractive antiviral target is papain-like protease (PL^{PRO}) non-structural protein 3 (nsP3), an essential component of the replicase-transcriptase complex [4]. Its topological structure is divided into ubiquitin-like (UBL), thumb, palm, and fingers [5]. It process the viral polyprotein by recognizing the tetrapeptide Leu-X-Gly-Gly motif, and

functions as de-ubiquitinating and de-ISGylating enzyme to suppresses inflammation and antiviral signaling [6,7]. Therefore, inhibition of PL^{PRO} activity can prevent virus replication and destroy its role in host immune response evasion, making it a promising target for antiviral drugs. Recently, an inhibitor for PL^{PRO} GRL-0617 has been proved to be an effective therapeutic agent for SARS-CoV PL^{PRO} [8,9].

In current study, the inhibitory effects towards PL^{PRO} for 21 compounds, including azvudine, aminoethyl, bavachinin, corylifol A, chromen, darunavir, disulfiram, ebselen, efavirenz, famciclovir, GRL-0617, isobavachalcone, mercaptopurine, neobavaisoflavone, oseltamivir, papyriflavonol A, psoralidin, ribavirin, sofosbuvir, tioguanine and 4'-O-methylbavachalcone were investigated by molecular docking. These compounds are antiviral, antifungal or anticancer drugs. They have been proved to be highly effective against HIV, HCV, influenza virus, coronavirus, antifungal and anticancer agents [10–21]. The focus of our study is on the repurposing and identification of compounds, aiming to accelerate the identification of potential drugs for COVID-19 treatment.

Abbreviations: COVID-19, Corona Virus Disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; PL^{PRO}, papain-like protease; nsP3, non-structural protein 3; UBL, ubiquitin-like; DS, Discovery Studio; -Se-S-, selenylsulfide.

* Corresponding author. Department of Laboratory Medicine, The First Affiliated Hospital of Shandong First Medical University, Jinan, China.

E-mail address: armzhang@hotmail.com (L. Zhang).

2. Materials and methods

2.1. Optimization of the 3D coordinates of the ligands

The 3D coordinates of the 21 drug ligands, azvudine (CSID: 24717759), aminoethyl (CSID: 29361441), bavachinin (CSID: 8512670), corylifol A (CSID: 22913562), chromen (CSID: 57578897), darunavir (CSID: 184733), disulfiram (CSID: 3005), ebselen (CSID: 3082), efavirenz (CSID: 57715), famciclovir (CSID: 3207), GRL-0617 (Chemical Book ID: CB74667969), isobavachalcone (CSID: 4444667), mercaptopurine (CSID: 580869), neobavaisoflavone (CSID: 4478223), oseltamivir (CSID: 58540), papyriflavonol A (CSID: 8518529), psoralidin (CSID: 4445118), ribavirin (CSID: 34439), sofosbuvir (CSID: 26286922), tioguanine (CSID: 2005804), and 4'-O-methylbavachalcone (CSID: 24845874) were obtained from ChemSpider (<http://www.chemspider.com/>) and Chemical Book (<https://www.chemicalbook.com/>) as a .mol file. The code within the parenthesis signifies the ChemSpider ID and ChemicalBook Number of the respective molecules. The secondary structure shown is the predicted by DSSP for SARS-CoV-2 (PDB: 7JIW), *Rhinolophus affinis* (GenBank: KF569996.1), SARS (PDB: 4OW0), MERS (PDB: 4RF1) and MHV (PDB: 5WFI) PL^{pro}. Similarity and alignment calculations were performed using MEGA-X and ClustalW.

2.2. Drug-like properties of the ligands

The drug likeliness of a molecule is indicated by the Lipinski's rule of five parameters (molecular weight ranges from 152.18 Da to 547.66 Da, no more than 5 hydrogen bond donors, no. of hydrogen bond acceptors should be less than 10 and logP should not be greater than 5). The Lipinski's rule of five parameters was obtained from the SWISSADME server (www.swissadme.ch/index.php). The chemical structures, chemical formula and the Lipinski's rule parameters of the ligands are listed in Table 1.

2.3. Structure and preparation of PL^{pro} for docking studies

SARS-CoV-2 papain-like protease (PDB: 6W9C, chains A, B, C) was used as the template for molecular docking. The 6W9C is solved by cryo-electron microscopy with 2.70 Å resolution, which is a trimer of three homologous amino acid chains (A, B, and C). Water molecules and ligands were removed from the PBD using Pymol.

2.4. Molecular docking analyses and visualization

Blind dockings were conducted based on docking ligands into a protein target using Discovery Studio (BIOVIA, 2016). In the key steps of the dockings, the protein remained rigid, while the ligand was fully flexible. Energy minimization of the protein was performed using the DS with CHARMM force field to relax the structure and remove the steric overlap. Then, polar hydrogen atoms were added. The centroid of the site model was used to define the active site box required by LibDock. The SARS-CoV-2 PL^{pro}-binding site in the X-ray crystal structure of ligand complex was determined by co-crystallization (X: -46.0103, Y: 14.2997, and Z: 29.9464), with a radius of 13.8 Å. The ligand conformation generation method was selected as "best", the number of binding site hotspots was set to 100, and other parameters were set by default. After each compound was docked, the best conformations and LibDock score were obtained. Considering their interaction energy and binding free energy, the compounds were screened. The binding modes (3D or 2D ligand-receptor interaction simulation map) of the selected compounds were analyzed.
















3. Results and discussion

Docking and scoring software is used widely to enhance the drug design and predict the interaction between drugs and macromolecules in the pharmaceutical products. The docking method involves searching for binding sites on the whole macromolecular surface. Therefore, the papain-like protease of COVID-19, SARS-CoV-2 PL^{pro} was used for blind docking analysis of some known drugs and bioactive substances. Fig. 1A shows the overall workflow of Discovery Studio (DS) LibDock algorithm. Then SARS-CoV-2 PL^{pro} structure was submitted to the SARS-CoV-2 PL^{pro} structure hotspots methods to predict the protein-ligand binding sites.

In addition to the recommend drugs for COVID-19, we have also conducted molecular docking studies with SARS-CoV-2 PL^{pro} for various natural compounds, antiviral drugs, antifungal drugs and anticancer drug (Table 1). Among SARS-CoV-2 PL^{pro}, SARS PL^{pro}, MERS PL^{pro}, *Rhinolophus affinis* PL^{pro}, and MHV PL^{pro}, classic catalytic triad, two different ubiquitin/ubiquitin-like conjugates and ISG15 modified protein binding sites are quite conservative (Fig. 1B). And some of them may be related to protein stability, kinetics, ligand binding and catalytic performance [9,22–24]. Amino acid sequence alignment showed that SARS-CoV-2 and SARS PL^{pro} (82.86%) as well as *Rhinolophus affinis* PL^{pro} (83.49%) were highly similar, and there were significant differences between SARS-CoV-2 and MERS (29.84%) and MHV PL^{pro} (29.52%) (Fig. 1C). Therefore, SARS-CoV-2 PL^{pro} is more closely related to SARS and *Rhinolophus affinis* than MERS and MHV. Examples include the active conservative catalytic trinity of Cys111, His272 and Asp286. The catalytic Trinity accepts Cys111 in the form of -SH as nucleophilic agent, His272 as general acid and alkali, and Asp286 promotes the deprotonation of Cys111 [25,26].

Aminoethyl, corylifol A, chromen, darunavir, ebselen, GRL-0617, isobavachalcone, papyriflavonol A, sofosbuvir and 4'-O-methylbavachalcone shows the formed interactions among the drug ligands near the catalytic triad. Aminoethyl and darunavir form conventional hydrogen bonds with Asp286 (Fig. 2A and B). In addition, aminoethyl constructs ten van der waals with Leu162, Gly163, Val165, Asn267, Tyr268, His272, Gly287, Ala288, Leu289 and Thr301, two π -alkyl interactions with Lys105 and Pro248, two π -sigma with Asp164 and Tyr264, and two unfavorable donor-donor interactions of Trp106 and Tyr273. Darunavir constructs two carbon hydrogen bond with Asn267 and Tyr268, eighteen van der waals with Ala107, Asp108, Asn109, Lys157, Gly163, Asp164, Arg166, Glu167, Met208, Ala246, Pro247, Tyr264, Gly266, Gln269, Tyr273, Gly287, Leu289 and Thr301, four π -alkyl interactions with Lys105, Trp106, Leu162 and Pro248. However, the donor-donor interaction between Darunavir and Ala288 is unfavorable. Darunavir may block the binding of PL^{pro} with target protein and exert its enzymatic activity. Chromen and corylifol A has almost the same site binding pattern, forming two conventional hydrogen bonds with Lys105 and Ala288 (Fig. 2C and D), a carbon hydrogen bond with Tyr268, ten van der waals with Lys92, Lys94, Leu162, Gly163, Asp164, Arg166, Glu167, Tyr264, Asn267 and Gly287, three π -alkyl interactions with Trp93, Ala107 and Leu289, a π -sigma with Trp106, and a unfavorable acceptor-acceptor sites with Asp286. Ebselen is an organic selenium compound that mimics the activity of glutathione. This compound forms a covalent bond between the selenium atom and thiols in the cysteine residue by forming a selenylsulfide (-Se-S-) linkage to inhibit HIV-1 capsid dimerization [27]. In the paper, we found that ebselen constructed a conventional hydrogen bonds with Lys105 (Fig. 2E), ten van der waals with Trp106, Ala107, Asp164, Glu167, Asn267, Tyr268, Tyr273, Asp286, Ala288 and Leu289. Although ebselen could not form -Se-S- with Cys111, it interacts with the PL^{pro} active site. It GRL-0617 interacts with the residues in the active site through hydrogen bonding with

Table 1
Candidate drugs properties screened and dockscore through LibDock in the study.

No.	Compounds	Structure	Treatment	Mass	Dockscore
1	Azvadine		HIV	286.22	83.52
2	Aminoethyl		HIV	312.41	65.25
3	Bavachinin		SARS	338.40	82.65
4	Corylifol A		SARS	390.47	89.15
5	Chromen		H1N1, H3N2	390.47	88.00
6	Darunavir		HIV	547.66	99.38
7	Disulfiram		HIV, MERS, SARS	296.54	36.89
8	Ebselen		HIV	274.18	69.10
9	Efavirenz		HIV	315.68	69.73
10	Famciclovir		VZV	321.33	75.77
11	GRL-0617		SARS	304.39	67.70
12	Isobavachalcone		SARS	324.37	52.81
13	Mercaptopurine		SARS	152.18	60.27
14	Neobavaisoflavone		SARS	322.35	110.53
15	Oseltamivir		AIV, H1N1, IBV	312.41	94.09




Fig. 3. The minimum docked poses of the compounds along with their corresponding ubiquitination and ISGylation interaction plots within the active site of SARS-CoV-2 PL^{PRO}.

in the Fig. 3K. It may affect the host ubiquitination and ISGylation events through the bond interaction with the N-terminal UBL domain.

The results of blind molecular docking of 21 different classes of compounds showed that they might be potential anti-SARS-CoV-2 drugs because they bind to the active site of the papain-like protease of SARS-CoV-2 PL^{pro}. Inhibition of this enzyme/protease is the key to blocking virus replication and reproduction. The inhibitor binding sites in SARS-CoV-2 protease are composed of key residues Cys111, His272 and Asn286 as evident from the recent study on the GRL-0617 and ebselen inhibitors for SARS-CoV-2 PL^{pro} [4,29]. Through the blind docking study of all 21 molecules with SARS-CoV-2 protease, we found that these molecules are usually surrounded by the above mentioned residues, ubiquitination and ISGylation sites, which clearly indicates that those molecules can prevent the replication of SARS-CoV-2 virus and evade the host immune response. The changes areas of the important interacting residues in the ligands and protease active sites were listed in Supplementary Table 1. Although blind docking studies have shown that all molecules can be used as potential drugs for the treatment of COVID-19, based on the estimated binding energy (ΔG) value, we can infer that among all the compounds studied, neobavaisoflavone (with the highest negative minimum ΔG value (–459 kcal/mol) may be the best potential inhibitor of SARS-CoV-2, followed by oseltamivir (–121.55 kcal/mol) > sofosbuvir (–119.44 kcal/mol) > famciclovir (–85.61 kcal/mol) > isobavachalcone (–84.75 kcal/mol) > tioguanine (–78.64 kcal/mol) > chromen (–67.92 kcal/mol) > efavirenz (–66.98 kcal/mol) > papyriflavonol A (–51.99 kcal/mol) > ebselen (–50.99 kcal/mol) > corylifol A (–46.78 kcal/mol) > mercaptopurine (–46.61 kcal/mol) > 4'-O-methylbavachalcone (–42.64 kcal/mol) > azvudine (–33.1 kcal/mol) > psoralidin (–29.89 kcal/mol) > ribavirin (–26.49 kcal/mol) > bavachinin (–25.59 kcal/mol) > disulfiram (–24.84 kcal/mol) > GRL-0617 (–24.62 kcal/mol) > aminoethyl (–20.87 kcal/mol) > darunavir (–8.74 kcal/mol) (Fig. 4).

In this study, the free energy of binding energy of compounds and PL^{pro} was ranged from –459 kcal/mol to –8.74 kcal/mol,




Fig. 4. The binding energy of the interaction between SARS-CoV-2 PL^{pro} and 21 compounds.

indicating that those compounds could interact with the SARS-CoV-2 PL^{pro}. Our finding also suggests the potential binding modes with known therapeutic drugs for COVID-19. Since this study was performed using computational methods, special attention should be paid to these results and further experiments *in vivo* and *in vitro* are needed.

Funding

This work was supported by grants from National Key Plan for Research and Development of China [2016YFD0500300], National Natural Science Foundation of China [82072270 and 81871663], National Major S & T Project for the Prevention and Treatment of Major Infectious Diseases in China [2017ZX10004206-007], the Innovation Project of Shandong Academy of Medical Sciences, and Academic Promotion Programme of Shandong First Medical University [2019LJ001].

Declaration of competing interest

We declare no competing interests.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.bbrc.2020.11.083>.

Transparency document

Transparency document related to this article can be found online at <https://doi.org/10.1016/j.bbrc.2020.11.083>

References

- [1] N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu, P. Niu, F. Zhan, X. Ma, D. Wang, W. Xu, G. Wu, G.F. Gao, W. Tan, I. China Novel Coronavirus, T. Research, A novel coronavirus from patients with pneumonia in China, 2019, *N. Engl. J. Med.* 382 (2020) 727–733.
- [2] Y. Lu, L. Zhang, Social media WeChat infers the development trend of COVID-19, *J. Infect.* 81 (2020) e82–e83.
- [3] M.M. Ghahremanpour, J. Tirado-Rives, M. Deshmukh, J.A. Ippolito, C.H. Zhang, I.C. de Vaca, M.E. Liosi, K.S. Anderson, W.L. Jorgensen, Identification of 14 known drugs as inhibitors of the main protease of SARS-CoV-2, *bioRxiv* (2020), <https://doi.org/10.1101/2020.08.28.271957>.
- [4] X. Gao, B. Qin, P. Chen, K. Zhu, P. Hou, J.A. Wojdyla, M. Wang, S. Cui, Crystal structure of SARS-CoV-2 papain-like protease, *Acta Pharm. Sin. B* (2020), <https://doi.org/10.1016/j.apbsb.2020.08.014>.
- [5] Y.K. Bosken, T. Cholko, Y.C. Lou, K.P. Wu, C.A. Chang, Insights into dynamics of inhibitor and ubiquitin-like protein binding in SARS-CoV-2 papain-like protease, *Front. Mol. Biosci.* 7 (2020) 174.
- [6] T. Klemm, G. Ebert, D.J. Calleja, C.C. Allison, L.W. Richardson, J.P. Bernardini, B.G. Lu, N.W. Kuchel, C. Grohmann, Y. Shibata, Z.Y. Gan, J.P. Cooney, M. Doerflinger, A.E. Au, T.R. Blackmore, G.J. van der Heden van Noort, P.P. Geurink, H. Ovaa, J. Newman, A. Riboldi-Tunnicliffe, P.E. Czabotar, J.P. Mitchell, R. Feltham, B.C. Lechtenberg, K.N. Lowes, G. Dewson, M. Pellegrini, G. Lessene, D. Komander, Mechanism and Inhibition of the Papain-like Protease, PL^{pro}, of SARS-CoV-2, *EMBO J.* 2020, e106275.
- [7] Y.M. Baez-Santos, S.E. St John, A.D. Mesecar, The SARS-coronavirus papain-like protease: structure, function and inhibition by designed antiviral compounds, *Antivir. Res.* 115 (2015) 21–38.
- [8] D. Shin, R. Mukherjee, D. Grew, D. Bojkova, K. Baek, A. Bhattacharya, L. Schulz, M. Widera, A.R. Mehdipour, G. Tascher, P.P. Geurink, A. Wilhelm, G.J. van der Heden van Noort, H. Ovaa, S. Muller, K.P. Knobeloch, K. Rajalingam, B.A. Schulman, J. Cinatl, G. Hummer, S. Ciesek, I. Dikic, Papain-like protease regulates SARS-CoV-2 viral spread and innate immunity, *Nature* 587 (2020) 657–662.
- [9] K. Ratia, S. Pegan, J. Takayama, K. Sleeman, M. Coughlin, S. Baliji, R. Chaudhuri, W. Fu, B.S. Prabhakar, M.E. Johnson, S.C. Baker, A.K. Ghosh, A.D. Mesecar, A noncovalent class of papain-like protease/deubiquitinase inhibitors blocks SARS virus replication, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 16119–16124.
- [10] N. Xu, J. Yang, B. Zheng, Y. Zhang, Y. Cao, C. Huan, S. Wang, J. Chang, W. Zhang, The pyrimidine analog FNC potently inhibits the replication of multiple enteroviruses, *J. Virol.* 94 (2020).
- [11] M.H. Lin, D.C. Moses, C.H. Hsieh, S.C. Cheng, Y.H. Chen, C.Y. Sun, C.Y. Chou,

- Disulfiram can inhibit MERS and SARS coronavirus papain-like proteases via different modes, *Antivir. Res.* 150 (2018) 155–163.
- [12] J.Y. Park, H.J. Yuk, H.W. Ryu, S.H. Lim, K.S. Kim, K.H. Park, Y.B. Ryu, W.S. Lee, Evaluation of polyphenols from *Broussonetia papyrifera* as coronavirus protease inhibitors, *J. Enzym. Inhib. Med. Chem.* 32 (2017) 504–515.
- [13] C. Seniya, A. Yadav, G.J. Khan, N.K. Sah, In-silico studies show potent inhibition of HIV-1 reverse transcriptase activity by a herbal drug, *IEEE ACM Trans. Comput. Biol. Bioinf* 12 (2015) 1355–1364.
- [14] D.W. Kim, K.H. Seo, M.J. Curtis-Long, K.Y. Oh, J.W. Oh, J.K. Cho, K.H. Lee, K.H. Park, Phenolic phytochemical displaying SARS-CoV papain-like protease inhibition from the seeds of *Psoralea corylifolia*, *J. Enzym. Inhib. Med. Chem.* 29 (2014) 59–63.
- [15] Z. Jin, X. Du, Y. Xu, Y. Deng, M. Liu, Y. Zhao, B. Zhang, X. Li, L. Zhang, C. Peng, Y. Duan, J. Yu, L. Wang, K. Yang, F. Liu, R. Jiang, X. Yang, T. You, X. Liu, X. Yang, F. Bai, H. Liu, X. Liu, L.W. Guddat, W. Xu, G. Xiao, C. Qin, Z. Shi, H. Jiang, Z. Rao, H. Yang, Structure of M(pro) from SARS-CoV-2 and discovery of its inhibitors, *Nature* 582 (2020) 289–293.
- [16] S. Yuan, H. Chu, K. Zhang, J. Ye, K. Singh, R.Y. Kao, B.K. Chow, J. Zhou, B.J. Zheng, A novel small-molecule compound disrupts influenza A virus PB2 cap-binding and inhibits viral replication, *J. Antimicrob. Chemother.* 71 (2016) 2489–2497.
- [17] P.L. Williams, C. Yildirim, E.G. Chadwick, R.B. Van Dyke, R. Smith, K.F. Correia, A. DiPerna, G.R. Seage, R. Hazra 3rd, C.S. Crowell, A.R.T.T.s.o.t.P.H.I.V.A.C., Surveillance Monitoring for, Association of maternal antiretroviral use with microcephaly in children who are HIV-exposed but uninfected (SMARTT): a prospective cohort study, *Lancet HIV* 7 (2020) e49–e58.
- [18] J.R. Semaan, M. Parmar, Fanciclovir, StatPearls, Treasure Island (FL), 2020.
- [19] C.Y. Chou, C.H. Chien, Y.S. Han, M.T. Prebanda, H.P. Hsieh, B. Turk, G.G. Chang, X. Chen, Thiopurine analogues inhibit papain-like protease of severe acute respiratory syndrome coronavirus, *Biochem. Pharmacol.* 75 (2008) 1601–1609.
- [20] A.N. Geisman, V.T. Valuev-Elliston, A.A. Ozerov, A.L. Khandzhinskaya, A.O. Chizhov, S.N. Kochetkov, C. Pannecouque, L. Naesens, K.L. Seley-Radtke, M.S. Novikov, 1,6-Bis[(benzyloxy)methyl]uracil derivatives—Novel antivirals with activity against HIV-1 and influenza H1N1 virus, *Bioorg. Med. Chem.* 24 (2016) 2476–2485.
- [21] A. Cazares-Cortazar, L.A. Uribe-Noguez, J.A. Mata-Marin, J. Gaytan-Martinez, M. de la Luz Martinez-Rodriguez, P.E. Villavicencio-Ferrel, A. Chaparro-Sanchez, S. Mauss, A. Ocana-Mondragon, A decrease in hepatitis C virus RNA to undetectable levels in chronic hepatitis C patients after PegIFNalpha + RVB or sofosbuvir + NS5A inhibitor treatment is associated with decreased insulin resistance and persistent oxidative stress, *Arch. Virol.* 165 (2020) 2759–2766.
- [22] M. Kandeel, Y. Kitade, M. Fayed, K.N. Venugopala, A. Ibrahim, The emerging SARS-CoV-2 papain-like protease: its relationship with recent coronavirus epidemics, *J. Med. Virol.* (2020), <https://doi.org/10.1002/jmv.26497>.
- [23] B.T. Freitas, I.A. Durie, J. Murray, J.E. Longo, H.C. Miller, D. Crich, R.J. Hogan, R.A. Tripp, S.D. Pegan, Characterization and noncovalent inhibition of the deubiquitinase and deISGylase activity of SARS-CoV-2 papain-like protease, *ACS Infect. Dis.* 6 (2020) 2099–2109.
- [24] K. Ratia, A. Kilianski, Y.M. Baez-Santos, S.C. Baker, A. Mesecar, Structural Basis for the Ubiquitin-Linkage Specificity and deISGylating activity of SARS-CoV papain-like protease, *PLoS Pathog.* 10 (2014), e1004113.
- [25] D. Niemeyer, K. Mosbauer, E.M. Klein, A. Sieberg, R.C. Mettelman, A.M. Mielech, R. Dijkman, S.C. Baker, C. Drosten, M.A. Muller, The papain-like protease determines a virulence trait that varies among members of the SARS-coronavirus species, *PLoS Pathog.* 14 (2018), e1007296.
- [26] H. Su, H. Su, F. Zhou, Z. Huang, X. Ma, K. Natarajan, M. Zhang, Y. Huang, Molecular insights into small molecule drug discovery for SARS-CoV-2, *Angew Chem. Int. Ed. Engl.* (2020), <https://doi.org/10.1002/anie.202008835>.
- [27] S. Thenin-Houssier, I.M. de Vera, L. Pedro-Rosa, A. Brady, A. Richard, B. Konnick, S. Opp, C. Buffone, J. Fuhrmann, S. Kota, B. Billack, M. Pietka-Ottlik, T. Tellinghuisen, H. Choe, T. Spicer, L. Scampavia, F. Diaz-Griffero, D.J. Kojetin, S.T. Valente, Ebselen, a small-molecule capsid inhibitor of HIV-1 replication, *Antimicrob. Agents Chemother.* 60 (2016) 2195–2208.
- [28] X. Chen, C.Y. Chou, G.G. Chang, Thiopurine analogue inhibitors of severe acute respiratory syndrome-coronavirus papain-like protease, a deubiquitinating and deISGylating enzyme, *Antivir. Chem. Chemother.* 19 (2009) 151–156.
- [29] W.R. Mikolaj Zmudzinski, Kamila Olech, Jaroslaw Granda, Mirosław Giurg, Małgorzata Burda-Grabowska, Linlin Zhang, Xinyuanyuan Sun, Zongyang Lv, Digant Nayak, Małgorzata Kesik-Brodacka, Shaun K. Olsen, Rolf Hilgenfeld, Marcin Drag, Ebselen Derivatives Are Very Potent Dual Inhibitors of SARS-CoV-2 Proteases - PLpro and Mpro in in Vitro Studies, *bioRxiv : the Preprint Server for Biology*, 2020.