

Clinical Images**It's Not Behçet's!**

Yen Chen Liu, BS, Amisha Desai, MD, Bryan Lee, MD, and Malathi Srinivasan, MD

School of Medicine, University of California, Davis, Sacramento, CA, USA.

KEY WORDS: Erythema multiforme; Behçet's syndrome; Stevens-Johnson disease.

J Gen Intern Med 26(5):559–60

DOI: 10.1007/s11606-010-1546-z

© The Author(s) 2010. This article is published with open access at Springerlink.com

A healthy 28 year-old man with isolated oral candidiasis at 17 and 26 years, on no medications, and having a recent URI 1 week before, presented with acute-onset progressive, painful oral mucositis and odynophagia (Fig. 1). His temperature was 101.6°F. The white blood cell count was 19.5, with 4.4% lymphocytes, 226CD4. Oral nystatin and IV caspofungin were started. Subsequently, the patient developed injected sclera with bilateral subconjunctival hemorrhages (Fig. 2) and marginated, erythematous, targetoid genital papules with vesicular centers (Fig. 3). His oral, ocular and genital lesions corresponded to the classic distribution of Behçet's syndrome,



Figure 2. Subconjunctival hemorrhages and ocular ulcerations.



Figure 1. Oral ulcerations.

initially placing this at the top of the differential diagnosis, followed by erythema multiforme (EM) and herpes simplex. However, generalized oral mucositis, conjunctivitis and, most importantly, targetoid lesions are not common in Behcet's syndrome. Behcet's syndrome frequently presents with aphthous ulcers sparing the outer lips, genital ulcers and uveitis.



Figure 3. Scrotal and penile shaft ulcerations.

Received April 27, 2010

Revised July 8, 2010

Accepted October 6, 2010

Published online November 30, 2010

Subsequent biopsy of the targetoid lesions demonstrated EM, with vacuolizing keratinocytes and tagging lymphocytes. Skin lesions and pain resolved after 1 week of high-dose methylprednisolone without complications. Repeat complete blood count (CBC) showed resolution of the lymphocytopenia. EM is a self-limited hypersensitivity reaction to external triggers, most notably HSV, *Mycoplasma pneumoniae* and medications^{1,2}. In this patient, HIV, HSV, *M.pneumonia* and *Legionella* studies were negative. Traditionally, EM has been placed within the hypersensitivity reaction spectrum, which includes Stevens-Johnson syndrome and toxic epidermal necrolysis³. However, there is suggestive evidence that EM may be a distinct entity. Treatment targets underlying infection and removing offending agents. Currently, systemic steroids are frequently used, but remain controversial.

Acknowledgement: All members who contributed to this submission are already indicated in the list of authors.

Funders: None (no internal or external funding of any kind was given).

Prior presentations: None. This submission has never been presented before.

Conflict of Interest: None disclosed.

Open Access: This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Corresponding Author: Yen Chen Liu, BS; School of Medicine, University of California, Davis, 4150 V Street, Suite 2400, Sacramento, CA 95817, USA (e-mail: yenliu@ucdavis.edu).

REFERENCES

1. Ayangco L, Rogers RS III. Oral manifestations of erythema multiforme. *Dermatol Clin.* 2003;21:195–205.
2. Volcheck GW. Clinical evaluation and management of drug hypersensitivity. *Immunol Allergy Clin North Am.* 2004;24:357–71.
3. Williams PM, Conklin RJ. Erythema multiforme: a review and contrast from Stevens-Johnson syndrome/toxic epidermal necrolysis. *Dent Clin North Am.* 2005;49:67–76.