

## CASE REPORT

# The challenge of triple intestinal parasite infections in immigrants—A call for comprehensive differential diagnosis

Emmanuel Edwar Siddig<sup>1</sup>  | Ayman Ahmed<sup>2</sup>

<sup>1</sup>Faculty of Medical Laboratory Sciences, University of Khartoum, Khartoum, Sudan

<sup>2</sup>Institute of Endemic Diseases, University of Khartoum, Khartoum, Sudan

**Correspondence**

Emmanuel Edwar Siddig, Faculty of Medical Laboratory Sciences, University of Khartoum, Khartoum, Sudan.  
Email: [emanwelleds389@gmail.com](mailto:emanwelleds389@gmail.com)

**Key Clinical Message**

Screening for parasitic infections among displaced populations including migrants, refugees, and internally displaced persons is crucial to prevent long-term complications associated with untreated infections. Additionally, prompt treatment with appropriate medications like praziquantel and albendazole can lead to significant improvement and successful patient outcomes. This case highlights the necessity of heightened awareness, early detection, and effective management of parasitic infections to enhance public health efforts and minimize the spread of these diseases. Implementing multisectoral One Health will improve human, animal, and environmental health globally using limited resources.

**Abstract**

Parasitic infections including *Schistosoma* spp., hookworm, and *Strongyloides* spp. are commonly co-existing diseases in underdeveloped countries including South Sudan and Sudan. These infections are often presented with no specific symptoms, therefore they remain undetected and eventually leading to long-term complications. Latent parasitic infections among adults particularly displaced population like migrants and refugees have recently gained global attention. In this case report, we present the case of a young male immigrant from South Sudan with multi-parasitic infections. The patient was admitted to a hospital complaining of gastrointestinal symptoms of hematemesis and melena, which were accompanied by signs of liver and spleen enlargement, coupled with low blood cell counts. While liver function tests showed no abnormalities, stool examinations confirmed the presence of *S. mansoni*, hook worm eggs, and Rhabditiform first-stage larvae of *S. stercoralis*. The patient was treated with medications such as praziquantel and albendazole, which resulted in full recovery and discharge from the hospital. This case highlights the importance of considering multiple infections among patients with gastrointestinal symptoms, especially in endemic areas. Cost-effective prevention and control strategy require the implementation

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Clinical Case Reports* published by John Wiley & Sons Ltd.

of a multisectoral One Health strategy for the surveillance, prevention, and control of parasitic infections in human, animals, and the environment.

#### KEYWORDS

global health, hookworm, massive drugs administration, migrant populations, one health, *Schistosoma* spp., *Strongyloides* spp.

## 1 | INTRODUCTION

Intestinal parasites are causing a significant global health concern, with over 3.5 billion people affected worldwide.<sup>1-3</sup> The infestation rates vary depending on the country of residence and individual demographics related to populations' exposure. Common intestinal parasites include soil-transmitted helminths such as *Ascaris lumbricoides*, *Trichuris trichiura*, hookworm, and protozoa like *Giardia intestinalis* and *Entamoeba* spp.<sup>4,5</sup> Least developed countries in particular, face higher prevalence and burden rates of these infections due to factors related to poverty and underdevelopment including limited access to clean water, poor hygiene and sanitation practices, malnutrition, and hot and humid climates.<sup>6,7</sup> This combination of conditions and practices create an ideal environment for the transmission, spread, and local establishment of intestinal parasites.<sup>8-10</sup> Sudan and South Sudan are predominantly endemic with soil-transmitted helminths and other parasitic diseases.<sup>10-23</sup>

Additionally, infection with intestinal parasites poses a serious health, socio-cultural, and economic burdens in endemic countries, while in industrialized nations, mainly immigrants, seasonal workers, and nomadic populations are the primary groups at risk.<sup>24,25</sup> Transmission of parasites can occur through direct person-to-person contact or contact with contaminated sources like food, water, or soil.<sup>26</sup> In high burden areas, co-infections with multiple parasites are increasingly occurring due to the poor living conditions, malnutrition, and weakened immuno-system.<sup>27-29</sup> The prevalence of dual or triple infections poses a significant public health challenge, as it can lead to higher morbidity and complicate treatment protocols. Studies have shown that dual infections, such as schistosomiasis with soil-transmitted helminths, increase the risk of severe anemia and nutritional deficiencies, especially in children and pregnant women.<sup>30</sup> It lead to high morbidity, mortality, disability, and enhance the susceptibility to other infections.<sup>27-29</sup> Children, teens, and adult, are at higher risk of adverse health outcomes with multiple parasitic infections compared to single infestations.<sup>31</sup> Here, we report case to highlight how poor socioeconomic conditions and disadvantaged living situations of immigrants can profoundly affect health outcomes even in developed countries.

## 2 | CASE HISTORY


A male freelancer, aged 31, from South Sudan, was admitted to Alban Jadeed Teaching Hospital in Khartoum, Capital city of Sudan in February 2023 after a job trip to Sudan, due to hematemesis and melena. Upon reviewing his medical history, it was found that he had no prior history of jaundice, hepatitis, or alcohol abuse. However, the patient had previously experienced diarrheal syndrome years ago, but underlining causative left uninvestigated.

## 3 | METHODS

Clinical examination showed that the patient had ascites and hepatosplenomegaly. Baseline investigations revealed a low white blood cell count of 2200 cells/ $\mu$ L, platelets of 50,000/ $\mu$ L, and hematocrit of 25%. Viral screening for hepatitis B, C, and HIV all yielded negative results. Stool examination indicated the presence of *S. mansoni* eggs with the characteristic broad lateral spine, as well as Rhabditiform first-stage larvae of *S. stercoralis* and Hookworm egg (Figure 1). Abdominal ultrasound showed hepatosplenomegaly, with the left lobe of the liver was the most affected site and central sonolucencies distinct from the true cirrhotic pattern, a portal vein diameter of 18 mm, and a moderate amount of peritoneal fluid. Esophagogastroduodenoscopy and colonoscopy were subsequently performed in response to the previous findings.

## 4 | CONCLUSION AND RESULT

Esophagogastroduodenoscopy revealed 2nd and 3rd grades esophageal varices with active hemorrhage, resulting in the patient's referral to Ibn Sinna Hospital for the placement of a Sengstaken Blakemore tube for 12 h. The patient subsequently underwent endoscopic sclerotherapy with 5% ethanolamine oleate twice, which helped improve his condition. A colonoscopy revealed a pale and atrophic colonic mucosa with no ulcers or polyps present. In addition to endoscopic sclerotherapy for the esophageal varices, the patient was treated with



**FIGURE 1** Showing (A) The *Schistosoma mansoni* eggs with the characteristic broad lateral spine, (B) Hookworm egg, and (C) Rhabditiform first-stage larvae of *Strongyloides stercoralis* (Iodine preparation).

praziquantel (20 mg/kg) orally, three times a day for 1 day, for *S. mansoni* infection, and albendazole (400 mg) twice a day for 3 days for *S. stercoralis* infection. The patient's condition improved and he was discharged from the hospital. During the follow up visits, repeated stool examinations for ova and rhabditiform larvae showed negative results, and over the following months, no relapse of hemorrhage occurred. This indicating full recovery of the patient.

## 5 | DISCUSSION

This case report highlights the importance of a thorough investigation of patients presenting with gastrointestinal symptoms, such as hematemesis and melena. Particularly for those live or had recent visit to areas hyper endemic with intestinal parasites. This underscores the need for raising awareness among the local communities and visitors about the locally endemic diseases through improving reporting and timely sharing of health data publicly. This will further guide healthcare providers in improving their differential diagnosis and advice for those who plan to travel to endemic area about precautions and prophylaxis they should take.

In this case report, the young male was admitted to the hospital with signs of liver and spleen enlargement, coupled with low blood cell counts. While the liver function tests showed no abnormality, the stool exam confirmed triple infections with *S. mansoni*, and hook worm eggs and Rhabditiform first-stage larvae of *S. stercoralis*. Interestingly, the hepatosplenic schistosomiasis observed in this case represents the severe complicated type of *S. mansoni* infection, with the development of hemorrhaging from esophageal varices.<sup>32</sup> This is the most commonly encountered and critical complication.

Based on the initial differential diagnosis of this patient's symptoms, we could have concluded various syndromes, which include hepatitis, liver cirrhosis, or even malignancy.<sup>33,34</sup> However, the ultrasound showed a distinct central sonolucencies and the absence of a true cirrhotic pattern. The patient's viral screening and liver function tests were unremarkable. The etiology of the patient's symptoms and unique shift in the clinical manifestations could be attributed to the interplay between the triple infections and the body immuno-response.

Additionally, the presence of multiple infections in a single patient, particularly in the context of parasitic diseases, necessitates an understanding of the underlying risk factors that contribute to such co-infections. In this specific case, several factors played a crucial role in the patient's susceptibility to triple infections with *S. mansoni*, hookworm, and *S. stercoralis*. Environmental factors, such as poor sanitation and inadequate access to clean water, create breeding grounds for parasites. Socioeconomic status significantly influences risk, as limited resources can lead to higher rates of exposure to contaminated sources. The patient's status as an immigrant from South Sudan highlights the impact of migration; displacement often results in crowded living conditions, further heightening the risk. Cultural practices related to hygiene and sanitation can also contribute to infection risk, while a compromised immune state due to chronic infections or malnutrition creates an environment where multiple infections can co-exist. Lastly, a lack of awareness and education about parasitic transmission and prevention increases vulnerability.

Although the patient is an immigrant from South Sudan, however, prior to the separation in 2010, South Sudan and Sudan were a single country, and both countries are heavily endemic with intestinal parasites, therefore, it could not be exclusively confirmed that his acquired these infections at his home country.<sup>35</sup> Specially

that, schistosomiasis and strongyloidiasis both can persist in patients for a long while undetected.

The final diagnosis of chronic hepatosplenic schistosomiasis was established based on the following criteria: (1) Identification of *S. mansoni* egg in the stool sample; (2) presence of portal hypertension and esophageal varices with normal liver function test and no evidence of hepatic cirrhosis; (3) Travel and residency history of the patient in hyper endemic countries.

The early detection of these triple infections was challenged by the fact that infection with *S. stercoralis* can happen without symptoms, especially in areas of high endemicity.<sup>36</sup> This is exactly the case with our patient as indicated by that; the larva was detected incidentally in his stool sample. The presence of this additional parasitic infection was indicated and confirmed through stool examination and there is no dermatologic evidence of Strongyloides (such as *Larva currens*), as well as the lack of pulmonary or gastrointestinal manifestations like steatorrhea, malabsorption or protein losing enteropathy. This indicates that the triple infections has altogether altered the clinical presentation of symptoms to look non-like any of these infections separately.

In Sudan, both of these infections are endemic, and schistosomiasis affects around 50% of the population in certain regions.<sup>37</sup> In order to improve public health measures relating to parasitic infections like *S. mansoni*, hook worm and *S. stercoralis*, several key steps are essential. The first and most important measure is to raise awareness among communities at risk and healthcare providers regarding the local risk, prevalence, and mode of transmission, personal protection and prevention measures from these parasitic infections.<sup>38</sup> Public health education campaigns should be implemented complemented with massive and social media sessions to engage the community and emphasize the importance of proper sanitation, hygiene, and safe practices for food, water, and defecation.<sup>38</sup> These interventions should be particularly intensified among populations living in crowded settings and suboptimal conditions such as camps for refugees and internally displaced persons (IDPs) to reduce the risk of infection. Additionally, this should be supported with systematic screening and treatment programs or massive drugs administration (MDA) program for treatment and prevention among people at high-risk either due to previous or current exposure to sources of potential contamination.<sup>39</sup> Specially that, the currently ongoing war in the country has distributed the originally fragile health system including diagnostic services, surveillance system, and healthcare and medication services throughout the country.<sup>40–43</sup> While on the hand, it has created a suitable environment for the emergence of invasive pathogens and vectors, intensified the transmission and dynamic of

infections, and increased the vulnerability of the forcibly displaced persons that represents over 85% of the country population.<sup>40–42</sup> In the current situation it would be more strategic to invest in a multisectoral collaborative framework that develop and implement a One Health strategy. The One Health strategy implements cost-effective strategic planning and intervention for improving human, animal, and the environment health through capitalizing on prevent and control interventions and reduce the exposure of vulnerable population to risk factors rather than curative medicine.<sup>44,45</sup> This multisectoral framework should bring health, agricultural, animal resources, metrological and climate, education, humanitarian, and development sectors altogether to maximize the impacts through prioritization, integration, for a better use of resources.<sup>46</sup>

In conclusion, this case documents the complications of having multiple infections on the same time and how this alter the clinical presentation of diseases. It also highlights the importance of considering co-infection and/or multiple infections especially for parasitic infections that have similar manifestation. Therefore, it might be a good strategy for healthcare providers to keep going with the differential diagnosis after making the first detection particularly among patients presenting with gastrointestinal symptoms in endemic areas. Although this might seem to cost more resources, however, considering that the golden standard tool for the diagnosis of intestinal parasites, implementing comprehensive screening for additional infection will not cost anything more than a few extra-minutes. Considering the zoonotic nature of parasitic infections, implementing One Health strategy will substantially reduce the resources needed for the surveillance, prevention, and control of these infections among human, animal, and the environment. This will improve the health and socioeconomic of poor communities and animals.

#### AUTHOR CONTRIBUTIONS

**Emmanuel Edwar Siddig:** Conceptualization; data curation; formal analysis; investigation; methodology; supervision; validation; visualization; writing – original draft; writing – review and editing. **Ayman Ahmed:** Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; supervision; validation; visualization; writing – original draft; writing – review and editing.

#### FUNDING INFORMATION

The authors declare no funding for this study.

#### CONFLICT OF INTEREST STATEMENT

The author reports no conflicts of interest in this work.




## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

## ORCID

Emmanuel Edwar Siddig  <https://orcid.org/0000-0001-6314-7374>

## REFERENCES

- Hajare ST, Gobena RK, Chauhan NM, Erniso F. Prevalence of intestinal parasite infections and their associated factors among food handlers working in selected catering establishments from Bule hora, Ethiopia. *Biomed Res Int*. 2021;19(2021):6669742.
- Sharifi-Sarasiabi K, Heydari-Hengami M, Shokri A, HosseiniTeshnizi S. Prevalence of intestinal parasitic infection in food handlers of Iran: a systematic review and meta-analysis. *Vet Med Sci*. 2021;7(6):2450-2462.
- Mama M, Alemu G. Prevalence and factors associated with intestinal parasitic infections among food handlers of Southern Ethiopia: Cross Sectional Study. *BMC Public Health*. 2016;1(16):105.
- Haque R. Human intestinal parasites. *J Health Popul Nutr*. 2007;25(4):387-391.
- Aiemjoy K, Gebresillasie S, Stoller NE, et al. Epidemiology of soil-transmitted helminth and intestinal protozoan infections in preschool-aged children in the Amhara region of Ethiopia. *Am J Trop Med Hyg*. 2017;96(4):866-872.
- Prüss-Ustün A, Wolf J, Bartram J, et al. Burden of disease from inadequate water, sanitation and hygiene for selected adverse health outcomes: an updated analysis with a focus on low- and middle-income countries. *Int J Hyg Environ Health*. 2019;222(5):765-777.
- Shears P. Poverty and infection in the developing world: healthcare-related infections and infection control in the tropics. *J Hosp Infect*. 2007;67(3):217-224.
- Feleke BE, Beyene MB, Feleke TE, Jember TH, Abera B. Intestinal parasitic infection among household contacts of primary cases, a comparative cross-sectional study. *PLoS One*. 2019;14(10):e0221190.
- Liang S, Seto EY, Remais JV, et al. Environmental effects on parasitic disease transmission exemplified by schistosomiasis in western China. *Proc Natl Acad Sci USA*. 2007;104(17):7110-7115.
- Abou-Zeid AH, Abkar TA, Mohamed RO. Schistosomiasis and soil-transmitted helminths among an adult population in a war affected area, southern Kordofan state, Sudan. *Parasit Vectors*. 2012;3(5):133.
- Mohamed MA, Siddig EE, Elaagip AH, Edris AM, Nasr AA. Parasitic contamination of fresh vegetables sold at central markets in Khartoum state, Sudan. *Ann Clin Microbiol Antimicrob*. 2016;11(15):17.
- Altahir O, AbdElbagi H, Abubakr M, Siddig EE, Ahmed A, Mohamed NS. Blood meal profile and positivity rate with malaria parasites among different malaria vectors in Sudan. *Malar J*. 2022;21(1):124.
- Siddig EE, Mohamed NS, Ahmed A. Severe coinfection of dengue and malaria: a case report. *Clin Case Reports*. 2024;12(6):e9079.
- Siddig EE, Misbah El-Sadig S, Eltigani HF, Musa AM, Mohamed NS, Ahmed A. Delayed cerebellar ataxia induced by plasmodium falciparum malaria: a rare complication. *Clin Case Reports*. 2023;11(10):e8053.
- Siddig EE, Ahmed A. When parasites stray from the path: a curious case of ectopic cutaneous Schistosoma haematobium. *QJM*. 2023;116(9):794-795.
- Ahmed A, Hemaida MA, Hagelnur AA, Eltigani HF, Siddig EE. Sudden emergence and spread of cutaneous larva migrans in Sudan: a case series calls for urgent actions. *IDCases*. 2023;5(32):e01789.
- Ahmed A, Abubakr M, Ali Y, Siddig EE, Mohamed NS. Vector control strategy for Anopheles stephensi in Africa. *Lancet Microbe*. 2022;3(6):e403.
- Mohamed NS, AbdElbagi H, Elsadig AR, et al. Assessment of genetic diversity of plasmodium falciparum circumsporozoite protein in Sudan: the RTS,S leading malaria vaccine candidate. *Malar J*. 2021;20(1):436.
- Siddig EE, Hay R. Laboratory-based diagnosis of scabies: a review of the current status. *Trans R Soc Trop Med Hyg*. 2022;116(1):4-9.
- Mohamed NS, Ali Y, Muneer MS, Siddig EE, Sibley CH, Ahmed A. Malaria epidemic in humanitarian crisis settings the case of South Kordofan state. *Sudan J Infect Dev Ctries*. 2021;15(1):168-171.
- Mohamed NS, Abdelbagi H, Osman HA, et al. A snapshot of plasmodium falciparum malaria drug resistance markers in Sudan: a pilot study. *BMC Res Notes*. 2020;13(1):512.
- Ali Y, Siddig EE, Mohamed N, Ahmed A. Rift Valley fever and malaria co-infection: a case report. *Clin Case Reports*. 2023;11(9):e7926. doi:10.1002/ccr3.7926
- Magambo JK, Zeyhle E, Wachira TM. Prevalence of intestinal parasites among children in southern Sudan. *East Afr Med J*. 1998;75(5):288-290.
- Herrick JA, Nordstrom M, Maloney P, et al. Parasitic infections represent a significant health threat among recent immigrants in Chicago. *Parasitol Res*. 2020;119(3):1139-1148.
- Steverding D. The spreading of parasites by human migratory activities. *Virulence*. 2020;11(1):1177-1191.
- Rogawa A, Ogata S, Mougi A. Parasite transmission between trophic levels stabilizes predator-prey interaction. *Sci Rep*. 2018;8(1):12246.
- Viney ME, Graham AL. Patterns and processes in parasite coinfection. *Adv Parasitol*. 2013;82:321-369.
- Shin JH, Won EJ, Jung JS, et al. A co-infection case report of taenia saginata in a patient with subclinical clonorchiasis confirmed by the combination of diagnostic tools. *BMC Infect Dis*. 2019;19(1):115.
- Intra J, Sarto C, Manuli E, Vannini PM, Brambilla P. Multiple parasitic infestation in a nine-month-old patient: a case report. *Helminthologia*. 2019;56(1):53-56.
- Mengist HM, Zewdie O, Belew A. Intestinal helminthic infection and anemia among pregnant women attending ante-natal

- care (ANC) in east Wollega, Oromia, Ethiopia. *BMC Res Notes*. 2017;10(1):440. doi:[10.1186/s13104-017-2770-y](https://doi.org/10.1186/s13104-017-2770-y)
31. Fauziah N, Aviani JK, Agrianfanny YN, Fatimah SN. Intestinal parasitic infection and nutritional status in children under five years old: a systematic review. *Trop Med Infect Dis*. 2022;7(11):371.
  32. Gunda DW, Kilonzo SB, Manyiri PM, Peck RN, Mazigo HD. Morbidity and mortality due to *Schistosoma mansoni* related periportal fibrosis: could early diagnosis of varices improve the outcome following available treatment modalities in sub Saharan Africa? A scoping review. *Trop Med Infect Dis*. 2020;5(1):20.
  33. Ferreira AO, Sousa HT, Brito J, Rosa L. Upper gastrointestinal bleeding in cirrhosis: varix or no varix? *BMJ Case Rep*. 2013;29:bcr2013008815.
  34. Dulger AC, Kemik O, Begenik H, Sumer A, Bulut D, Bulut G. Synchronous esophageal squamous cell carcinoma and esophageal variceal bleeding due to idiopathic portal hypertension: a case report. *Gastroenterology Res*. 2011;4(2):84-87.
  35. Marnell F, Guillet A, Holland C. A survey of the intestinal helminths of refugees in juba, Sudan. *Ann Trop Med Parasitol*. 1992;86(4):387-393.
  36. Yeh MY, Aggarwal S, Carrig M, et al. *Strongyloides stercoralis* infection in humans: a narrative review of the most neglected parasitic disease. *Cureus*. 2023;15(10):e46908.
  37. Arndts K, Elfaki TEM, Jelden N, et al. *Schistosoma mansoni* PCR+ –infected individuals in The Sudan present elevated systemic levels of chemokines when compared to uninfected and egg+ cohorts. *Clin Exp Immunol*. 2019;196(3):364-373.
  38. Garn JV, Wilkers JL, Meehan AA, et al. Interventions to improve water, sanitation, and hygiene for preventing soil-transmitted helminth infection. *Cochrane Database Syst Rev*. 2022;6(6):CD012199.
  39. Webster JP, Molyneux DH, Hotez PJ, Fenwick A. The contribution of mass drug administration to global health: past, present and future. *Philos Trans R Soc Lond Ser B Biol Sci*. 2014;369(1645):20130434.
  40. Siddig EE, Eltigani HF, Ahmed A. Healing the unseen wounds: Sudan's humanitarian crisis traumatizing a nation. *Asian J Psychiatr*. 2023;89:103764. doi:[10.1016/j.ajp.2023.103764](https://doi.org/10.1016/j.ajp.2023.103764)
  41. Siddig EE, Eltigani HF, Ali ET, Bongomin F, Ahmed A. Sustaining hope amid struggle: the plight of cancer patients in Sudan's ongoing war. *J Cancer Policy*. 2023;38:100444.
  42. Siddig EE, Eltigani HF, Ahmed A. Urgent call to protect children and their health in Sudan. *BMJ*. 2023;9(382):1799.
  43. Siddig EE, Aradaib IE, Ahmed A. A study of case management challenge for black grain eumycetoma during the ongoing war in Sudan. *Clin Case Reports*. 2024;12(9):e9438. doi:[10.1002/ccr3.9438](https://doi.org/10.1002/ccr3.9438)
  44. Pitt SJ, Gunn A. The one health concept. *Br J Biomed Sci*. 2024;15(81):12366.
  45. Ahmed A, Ali Y, Ibrahim NA, et al. One health response for Rift Valley fever outbreak in Sudan. *Preprints*. 2024;10:2024081358. doi:[10.20944/preprints202408.1358.v1](https://doi.org/10.20944/preprints202408.1358.v1)
  46. Remera E, Rwagasore E, Muvunyi CM, Ahmed A. Emergence of the first molecularly confirmed outbreak of Rift Valley fever among humans in Rwanda, calls for institutionalizing the one health strategy. *IJID One Health*. 2024;1(4):100035.

**How to cite this article:** Siddig EE, Ahmed A. The challenge of triple intestinal parasite infections in immigrants—A call for comprehensive differential diagnosis. *Clin Case Rep*. 2024;12:e9549. doi:[10.1002/ccr3.9549](https://doi.org/10.1002/ccr3.9549)