



Transport of Folded Proteins by the Tat System

Kelly M. Frain¹ · Colin Robinson¹ · Jan Maarten van Dijl²

Published online: 10 August 2019

© The Author(s) 2019

Abstract

The twin-arginine protein translocation (Tat) system has been characterized in bacteria, archaea and the chloroplast thylakoidal membrane. This system is distinct from other protein transport systems with respect to two key features. Firstly, it accepts cargo proteins with an N-terminal signal peptide that carries the canonical twin-arginine motif, which is essential for transport. Second, the Tat system only accepts and translocates fully folded cargo proteins across the respective membrane. Here, we review the core essential features of folded protein transport via the bacterial Tat system, using the three-component TatABC system of *Escherichia coli* and the two-component TatAC systems of *Bacillus subtilis* as the main examples. In particular, we address features of twin-arginine signal peptides, the essential Tat components and how they assemble into different complexes, mechanistic features and energetics of Tat-dependent protein translocation, cytoplasmic chaperoning of Tat cargo proteins, and the remarkable proofreading capabilities of the Tat system. In doing so, we present the current state of our understanding of Tat-dependent protein translocation across biological membranes, which may serve as a lead for future investigations.

Keywords TatA · TatB · TatC · Twin-arginine · *Bacillus subtilis* · *Escherichia coli*

1 Introduction

To function correctly and efficiently, every cell needs to be highly organised, tightly regulated and compartmentalised. Proteins are essential macromolecules synthesised by ribosomes in the cytoplasm that often require localisation to a particular subcellular compartment before they can carry out their respective functions. Their proper formation, targeting and activity are imperative to the survival of the cell. This requirement for correct localisation particularly applies to proteins that take part in the acquisition of nutrients, energy

transduction, cell-to-cell communication and cellular locomotion. On average, 20–30% of proteins synthesised in the bacterial cytoplasm are destined for extra-cytoplasmic locations [1]. They therefore have to pass a cell membrane composed of a tightly sealed lipid bilayer intent on keeping the cell structurally sound and impenetrable. Therefore, specialised transport systems have evolved within the cell membrane to allow proteins to cross this barrier. Each system made up of critical components is as specialised as the protein cargo it will transport. However common features tie protein transport systems together, which guarantee cell regulation and safety. These include a gated pore, an energy requirement to drive cargo proteins through the membrane, and the use of signal peptides that direct the cargo protein to the correct translocase and the correct location.

Two major transport systems exist for protein translocation across the bacterial cytoplasmic membrane, namely the general secretory (Sec) pathway and the twin-arginine translocation (Tat) pathway (Fig. 1). The Sec pathway facilitates export of the majority of bacterial proteins, whereas the Tat pathway is quite restricted. For instance, it transports ~30 proteins in *Escherichia coli* and only four in *Bacillus subtilis* [2]. Further, each protein is fully folded in the

✉ Jan Maarten van Dijl
j.m.van.dijl01@umcg.nl

Kelly M. Frain
kmf@mbg.au.dk

Colin Robinson
C.Robinson-504@kent.ac.uk

¹ The School of Biosciences, University of Kent,
Canterbury CT2 7NZ, UK

² Department of Medical Microbiology, University Medical
Center Groningen, University of Groningen (UMCG),
Hanzeplein 1, P.O. Box 30001, 9700 RB Groningen,
The Netherlands




Fig. 1 The Sec- and Tat-dependent protein transport pathways. The Sec pathway is the dominant pathway for protein export from the bacterial cytoplasm. It accepts and translocates cargo proteins across the plasma membrane in a loosely folded or unfolded state, here exemplified with the precursor of the outer membrane protein A of *E. coli* (OmpA). Targeting and folding control of the cargo protein is supported by cytoplasmic targeting factors, such as SecB. The Sec machinery itself is composed of the SecYEG channel and the translocation ATPase SecA, which converts chemical energy in the form of ATP into a driving force that pushes the cargo protein through the membrane. Additionally, translocation may be powered by the trans-

membrane proton gradient. At the *trans*-side of the membrane, the translocated protein folds into its active and protease-resistant final conformation. In contrast to the Sec pathway, the Tat pathway transports fully folded cofactor-containing proteins across the membrane, here exemplified with the precursor of the Tat cargo TorA. Cofactor insertion and folding may be aided by Redox Enzyme Maturation Proteins (REMPS), such as TorD in the case of TorA. The Tat translocase may consist of the three components TatA, TatB and TatC (*E. coli*), or of TatA and TatC components only (*B. subtilis*). Protein transport via Tat is powered by the transmembrane proton-motive force

cytoplasm prior to export via Tat, whereas Sec can only export unfolded proteins.

2 Protein Targeting Via the Twin-Arginine Signal Peptide

To ensure proteins are appropriately directed into the Sec or Tat pathways and to initiate the translocation process, specific signal peptides are present on the N-terminus of each protein. On the *trans* side of the membrane the signal peptide is cleaved by a signal peptidase to liberate just the mature protein [3–7]. The amino acid sequences of signal peptides differ substantially, but they are all composed of a positively charged N-terminal N-domain, a hydrophobic H-domain and a C-terminal C-domain with an Ala-x-Ala signal peptidase cleavage site [3, 8] (Fig. 2). Further, the N-regions of Tat signal peptides contain the canonical twin-arginine motif S-R-R-x-F-L-K (where x is a polar amino acid) [9]. The importance of additional conserved amino

acids in the Tat-motif depends on the cargo protein and varies in different bacteria [10]. However, RR-residues are close to invariant and key to efficient protein export. In particular, the charge-neutral substitution of RR to KK blocks Tat export completely [11]. Yet, a single Arg to Lys mutation only slows down the rate of translocation in most bacteria [12]. In chloroplast thylakoids where the Tat pathway also exists, an RR to KR substitution is tolerated, while a RR to RK substitution precludes transport [12–14]. A single substitution of Arg to Glu has been reported as tolerated too [15]. Of note, the TtrB subunit of the tetrathionate reductase in *Enterobacteriaceae* is the only known native Tat cargo to have a KR-motif [16]. Aside from the RR-motif, other residues within the larger twin-arginine signal peptide are also important. In particular, the Phe residue is present in 80% of Tat-motifs, and substitutions showed a highly hydrophobic residue is essential at this position [11].

Tat signal peptides comprise about 30 residues in most organisms. Hence they are longer than Sec signal peptides, which comprise about 17 to 24 residues [17]. Tat signal




Fig. 2 Sec- and Tat-specific signal peptides. N-terminal signal peptides direct proteins to the Sec and Tat translocases in the membrane. They have a conserved tripartite structure, consisting of a positively charged N-region (indicated by ‘white residues’ in one-letter code), a hydrophobic H-region (red) and a C-region (green) that contains the Ala-X-Ala recognition site for signal peptidase. Cleavage by signal peptidase, C-terminally from the Ala-X-Ala sequence, liberates the mature protein (pink) from the membrane. Twin-arginine signal pep-

ptides, as exemplified by the TorA signal peptide, contain the canonical twin-arginine motif at the interface of the N- and C-regions. Their H-region is longer and less hydrophobic than that of Sec-type signal peptides, and N-terminally of the C-region there are often positively charged residues that serve in Sec-avoidance. Notably, Sec-type signal peptides, here exemplified by the OmpA signal peptide, are usually much shorter than twin-arginine signal peptides

ptides are also overall less hydrophobic than Sec signal peptides, which serves to avoid protein targeting to the Sec pathway [18]. Additionally, the C-domain of Tat signal peptides may include basic residues N-terminally of the A-x-A motif, which contribute to Sec avoidance (Fig. 2) [19, 20].

3 The Twin-Arginine Translocation Pathway

In the early 1990’s, an alternative translocase was discovered in the thylakoid membrane of chloroplasts, which worked in parallel to the Sec pathway [21]. Initially this pathway was named the Δ pH-dependent pathway due to its unusual sole requirement of a transmembrane proton gradient for translocation [22]. Three membrane proteins were soon identified in thylakoids as essential for translocation of fully folded proteins via the Δ pH-dependent pathway [23], namely Tha4 [24], Hcf106 [25] and cpTatC [26]. Subsequently, homologous proteins were identified in some bacteria, archaea and even mitochondria [27, 28]. In *E. coli*, the homologues of Tha4, Hcf106 and cpTatC were also shown to be required for export of proteins with twin-arginine signal peptides and, therefore, they were respectively named TatA, TatB and TatC [9, 29–31].

Combined studies on the thylakoidal and bacterial Tat pathways showed that their function is to transport a subset of complex fully folded proteins that require cofactor insertion or immediate oligomerisation [8, 32]. Today, Tat-translocated proteins have been shown to participate in many processes including energy metabolism, cell division, cell envelope biogenesis, quorum sensing, motility, symbiosis and pathogenesis [33–36]. Tat can even export complex heterologous proteins that are Sec-incompatible, like the tightly folded dihydrofolate reductase with bound methotrexate [37], the green fluorescent protein (GFP) [38], and

several bio-pharmaceutically relevant human proteins [39]. Another intriguing attribute of the Tat pathway is that it can detect unfolded or mutated proteins, and reject them for export [40, 41].

Based on the number of Tat components involved in protein translocation, essentially two types of ‘translocases’ can be distinguished. The prototype Tat translocase that is active in thylakoids and *E. coli*, consists of the afore-mentioned TatABC components. Further, the minimal Tat translocases, as typified in *Bacillus* species consist of TatA and TatC components only. The types of translocases will be discussed in the following paragraphs.

4 The *E. coli* Tat Translocase

The *E. coli* *tatABCD* operon encodes the core components of this bacterium’s Tat system (Table 1). All four genes are constitutively expressed, but the expression level of *tatA* exceeds that of *tatB* 25-fold that of *tatC* 50-fold [42]. This difference is mirrored in the final component make-up of the Tat translocase in the plasma membrane. The *tatE* gene is constitutively expressed from another chromosomal locus. The *tatB* and *tatE* genes are thought to originate from gene duplications of *tatA* [28, 43]. Although Δ *tatABCDE* strains are viable, the mutants show various defects including impaired septation, decreased motility and an increased sensitivity to detergent [44].

TatA (9.6 kDa) is the most abundant component of the Tat complex, most likely responsible for forming the translocase channel [45]. *E. coli* has a core TatA protein, but it also involves the TatA-like proteins, TatB and TatE [28]. Of note, TatE can substitute TatA [43]. TatA, TatB and TatE are similar in structure with a short N-terminal domain that is exposed to the periplasm [46], a single transmembrane

Table 1 Comparison of *E. coli* and *B. subtilis* Tat proteins and Tat complexes including their estimated molecular masses (kDa)

<i>E. coli</i>				<i>B. subtilis</i>			
Protein/complex	Gene product molecular mass (kDa)	Complex molecular mass (kDa)	Ref.	Protein/complex	Gene product molecular mass (kDa)	Complex molecular mass (kDa)	Ref.
TatA	9.6	100–500	[114]	TatAd	7.4	160/270	[77]
				TatAy	6	200	[159]
TatE	7		[43]	TatAc	6.7	100	[78]
TatB	18.4	< 100	[111]				
TatC	28.9	220	[111]	TatCd	28	66–100	[78]
				TatCy	28.9	66	[78]
TatBC		430	[111]	TatAdCd		230/350	[77]
TatABC		580	[104]	TatAyCy		200	[159]
TatABC + substrate		600	[104]	TatAcCd		230	[78]
				TatAcCy		200	[78]

helix, an amphipathic helix in the cytoplasm [47], and an unstructured cytoplasmic C-domain [48] (Fig. 3). Surprisingly, not many mutations in TatA block export, but there are a few instances. In particular, Gly33 in the “hinge region” is critical for TatA function [49], and the transmembrane

helix and various residues in the amphipathic helix are also important [50, 51].

TatE (7 kDa) is a much smaller than TatA [9]. Given the smaller size and ~100-fold lower abundance than TatA, it was initially believed TatE has no real function in the Tat




Fig. 3 Membrane topology and structures of the TatA, TatB and TatC proteins. The Tat translocase of *E. coli* consists of three components, namely TatA, TatB and TatC. TatB and TatC form a receptor complex for cargo proteins, whereas TatA is the main facilitator for protein translocation across the membrane. TatB is missing from the two-component Tat translocases as encountered in *B. subtilis*. The upper half of the Figure shows a traditional representation of the membrane

topology of TatA/E, TatB and TatC based on molecular biological analyses. The lower half of the Figure shows ribbon presentations of the structures of TatA, TatB and TatC as adopted from the RCSB Protein Data Bank (<http://www.rcsb.org/structure/>). These structures have the following PDB accession codes: TatA—2LZR (solution NMR structure [48]); TatB—2MI2 (solution NMR structure [54]); and TatC—4HTS (crystal structure [63])

translocon [42]. More recently however, it was shown TatE could substitute TatA [43], and that it is recruited to the Tat translocase [52]. Importantly, TatE was shown to interact with the Tat signal peptide and to even partially prevent premature cleavage of the TorA signal peptide [53].

The role of TatB (18.4 kDa) is to bind the Tat signal peptide and, thereafter, the mature protein. Despite only sharing 20% sequence identity to TatA and being nearly double TatA's size, TatB is predicted to have a very similar structure and topology (Fig. 3) [50]. Specifically, TatB has a slightly longer amphipathic helix and a longer unstructured C-terminal region [54, 55]. Mutations in TatB's hinge region and amphipathic helix cause translocation defects [56]. Of note, particular amino acid substitutions in TatA's N-terminus allow replacement of TatB by TatA [57] [58], supporting the notion that TatB originated from TatA and subsequently specialized [5, 59].

TatC is the largest (28.9 kDa) and best-conserved protein in the Tat complex that aids cargo binding [60, 61]. The structure of TatC is very different to other Tat components as it has 6 transmembrane helices and an N-in C-in topology (Fig. 3) [62]. The crystallisation of TatC from *Aquifex aeolicus*, which shares 40% sequence identity to *E. coli* TatC, revealed the relative positions of the transmembrane domains [63]. Together, they take the shape of a baseball glove or cupped hand with very restricted structural flexibility [64]. Notably, a conserved Glu residue (Glu170 in *E. coli*) is positioned close to the signal peptide binding pocket in the plane of the membrane and potentially perturbs the bilayer structure [12, 64, 65]. Additional residues needed for TatC function reside in the cytoplasmic N-region and the first cytoplasmic loop [61, 66].

5 The *B. subtilis* Tat Translocase

The Tat translocase can minimally function with just TatA and TatC [5, 67–69]. Interestingly, the Gram-positive bacterium *B. subtilis* has two minimal Tat translocases encoded by the *tatAdCd* and *tatAyCy* operons, which work in parallel and with different cargo specificities (Table 1) [5]. *TatAdCd* has only one known cargo protein, PhoD, which is co-expressed with the translocase under phosphate-deprived conditions [68, 70]. *TatAyCy* is constitutively expressed, along with its cargo proteins EfeB (YwnN), QcrA and YkuE [5, 71–74]. The third TatA gene of *B. subtilis*, *tatAc*, is constitutively expressed from another locus, and was shown to serve a non-essential function in protein translocation via the *TatAyCy* [5, 75].

B. subtilis *TatAd* and *TatAy* are bifunctional, meaning that they act at the same time as *E. coli* TatA and TatB. Interestingly, *B. subtilis* *TatAd* can replace TatA and TatB in *E. coli* [76], whereas *TatAc* expressed in *E. coli* can

functionally replace TatA and TatE and form active translocases with *TatCd* and *TatCy* [77, 78]. This suggests that, despite species-specific features, the translocation mechanism employed by Tat is conserved across species [76, 79].

Structural studies on *B. subtilis* *TatAd* (7.4 kDa) have confirmed its 'L-shape' arrangement in the membrane [80–82]. By itself, *TatAd* oligomerizes to complexes of ~270 kDa and, together with *TatCd* (28 kDa), *TatAd* forms complexes of ~230 kDa in which *TatAd* is stabilized by *TatCd* [83–85]. Although the structure of *TatAd* resembles that of *E. coli* *TatA*, the effects of particular amino acid substitutions differ for the two proteins [47, 86]. Notably, mutagenesis of the *TatAd* N-terminus blocks protein translocation in *E. coli* *tatB* mutant cells, indicating that the N-terminal residues of *TatAd* are needed for *TatB* substitution [83].

Like *TatAd*, *TatAy* (6 kDa) has a structure similar to that of *E. coli* *TatA* [83, 86]. In particular, the conserved Pro2 residue in the N-terminus of *TatAy* and its hinge region are required for protein export [75, 86]. Complexes of *TatAy* alone and *TatAyCy* have a molecular mass of ~200 kDa [87]. Intriguingly, a P2A mutation leads to the formation of large fibrils composed of *TatAy* and *TatCy*, suggesting that Pro2 serves a role in the termination of complex assembly [88].

TatCd and *TatCy* (28/28.9 kDa) resemble *E. coli* *TatC*, having six transmembrane helices [87, 85]. Further, the N-terminus, the first cytoplasmic loop and the C-terminal tail of *TatCd* and *TatCy* are important for protein export, but the relevance of different conserved residues depends on the cargo [89, 90].

TatAc (6.7 kDa) of *B. subtilis* shares significant sequence similarity with *E. coli* *TatE*, and it can actually form active Tat complexes with *TatA* and *TatB*, or with *TatCd* and *TatCy* when expressed in *E. coli* (Table 1) [75–78, 87]. Nevertheless, *TatAc* cannot replace *TatAd* or *TatAy* for protein translocation in *B. subtilis*, where it was shown to assist protein translocation by *TatAyCy* [75].

6 TatA and TatA/BC Complexes

While the Tat system can handle cargo proteins of up to 150 kDa [91], the Tat components are much smaller. This implies that they need to assemble into larger complexes that can facilitate membrane passage of larger cargo proteins [92]. Indeed, two types of Tat complexes were identified, namely *TatA(B)C* and *TatA*. In *E. coli* and thylakoids, membrane-embedded *TatBC* complexes are believed to bind cargo proteins, whereas recruitment of *TatA* complexes is required to facilitate their membrane passage [93–95]. In *B. subtilis*, the cargo receptor function of *TatBC* complexes is fulfilled by *TatAdCd* or *TatAyCy* complexes. Notably, the *TatA* complexes by themselves, especially those of *B. subtilis* (Table 1), are too small and homogeneous to allow

passage of most cargo proteins [68, 77, 78]. The TatA-TatA/BC assemblies are thought to disassemble upon completed cargo translocation [96].

As shown by cross-linking studies, within TatBC complexes, TatC is first to interact with the N-region of a twin-arginine signal peptide [94, 97, 98]. Subsequently, deep insertion of the signal peptide into TatC will follow, leading to interaction of the H-domain with the transmembrane segment of TatB [54, 94, 99]. In turn, this leads to exposure of the signal peptidase cleave site in the C-region to signal peptidase on the *trans*-side of the membrane [63, 100–102]. Intriguingly, several lines of evidence, suggest that more than one cargo protein can be bound by assemblies of seven TatBC copies [60, 103–105]. Within these TatBC assemblies, TatC monomers have two TatB contact sites [61, 99, 102, 106, 107]. Further, the transmembrane segment of TatB appears to be positioned close to the site where translocase oligomerization is initiated by TatA, which suggests that TatB could serve as a regulatory surrogate of TatA [108–110]. The latter would be in line with the fact that TatB is absent from minimal TatAC translocases as encountered in *B. subtilis*. Furthermore, cross-linking analyses show that cargo docking via the signal peptide leads to conformational changes that rearrange TatC's binding site for TatA and TatB [52]. Binding of a signal peptide changes the arrangement of TatC from head-to-tail to tail-to-tail [106].

TatBC complexes contain small amounts of TatA that may serve as points of TatA nucleation for forming the active translocase [111, 112]. Most TatA molecules are, however, present in TatA complexes. The TatA complexes of *E. coli* are very heterogeneous, ranging from 100 to 500 kDa with intermediate size intervals of ~34 kDa [48, 54, 113–115]. In contrast, TatAc, TatAd and TatAy complexes in *Bacillus* are much smaller with molecular masses of ~100, ~270 and ~200 kDa, respectively [76, 77].

7 Tat Translocation Mechanism

Despite almost three decades of research, the Tat translocation mechanism is still incompletely understood. As outlined above, cargo translocation is initiated at TatA/BC complexes and then facilitated by TatA [113]. This may involve either pore formation [116] or membrane weakening [43, 117].

Based on low-resolution EM images, it was proposed that TatA complexes have a pore of 8.5–13 nm that might accommodate cargo proteins of varying size [116, 118]. This pore would be closed by a lid at the cytoplasmic side membrane, resembling a 'trap door', which could swing open with the help of a conserved Gly residue in the hinge region of TatA to allow cargo passage [118, 119]. In this scenario, after cargo docking onto TatBC, TatA would be recruited to form an oligomeric ring conforming to the size of the cargo

protein [120, 121]. Although this model appears attractive, the trap door concept has not been confirmed in other studies [46, 48, 122]. Moreover, complexes of the TatA paralogue TatE (50–110 kDa) appear too small for pore formation [43].

More recently, it was proposed that TatA complexes might serve to weaken the membrane [48, 106, 117, 123]. This would relate to the relatively short transmembrane domain of TatA that can locally restrict the membrane thickness. This membrane weakening would only occur upon cargo binding and interaction of the mature part of the cargo protein with the amphipathic helix of TatA [94, 99, 123–126]. In the absence of cargo, the membrane weakening would not take place as immersion of the amphipathic helix of TatA in the membrane would preserve the membrane integrity, as was shown for the thylakoidal TatA [122].

As mentioned above, protein translocation via Tat is exclusively driven by the proton-motive force, which consists of the ΔpH and the electric potential $\Delta\psi$ across the membrane [127]. Early studies into the energetics of Tat were performed in vitro with the plant thylakoid system. In the presence of light and a ΔpH , but in the absence of nucleotides, the photosystem II oxygen-evolving Tat cargo protein tOE23 was still exported [128]. In addition, a phage shock protein PspA, involved in maintaining the proton-motive force, was found to increase Tat translocation in bacteria [129]. However, in vivo studies in the green alga *Chlamydomonas reinhardtii* showed that the system can still transport proteins without a thylakoidal ΔpH , which can be explained by the fact that the Tat pathway can use both the ΔpH and $\Delta\psi$ [130, 131]. As a consequence of this equivalency, an antiporter mechanism was suggested where a coupling of H^+ flow and protein transport has been suggested [132]. Of note, the counterflow of protons necessary for Tat protein export was estimated to amount about 7.9×10^4 protons per molecule [133]. This is equivalent to 10,000 ATP molecules, 3% of the energy produced by a chloroplast, so it is a considerable cost to the cell.

With regards to individual steps of the translocation mechanism, in vitro studies have shown that the proton-motive force is not required for protein targeting or protein binding to TatBC, but for the more advanced binding stages and oligomerisation of TatA [94, 134]. For thylakoids it was proposed that the ΔpH could potentially protonate TatA (Tha4) at the Glu10 residue, making it energetically feasible to move up in the membrane to its docking site in TatC (Gln234) [112]. However, in an earlier study this Glu10 residue was replaced with Gln, as well as with Ala or Asp, and all of these changes severely reduced the ability of TatA to facilitate protein transport [135]. While this shows the importance of the Glu10 residue and a negative charge at this position for translocation activity, it is not certain whether this implies a role of Glu10 in sensing the thylakoidal luminal pH through protonation, or whether Glu10 forms a salt bridge with a basic

residue somewhere else [135]. It is also still unclear how the assembly of TatA in *E. coli* is facilitated by the proton-motive force, as it has been shown through in vitro studies that the transport driving force is largely provided by the $\Delta\psi$ [136]. In fact, these studies indicate that two distinct $\Delta\psi$ -dependent steps drive protein transport: a first step would involve a $\Delta\psi$ of relatively high magnitude that may be short-lived, and a second step of longer duration would require a $\Delta\psi$ of relatively low magnitude. When the $\Delta\psi$ was increased, so was the transport speed [94, 136]. This raises the question, how exactly the $\Delta\psi$ drives protein transport via Tat in *E. coli* and why this is apparently different in thylakoids, where the ΔpH represents the driving force for protein transport. A conceivable scenario is that movement of certain charged regions within the membrane-embedded *E. coli* Tat proteins could be induced by a $\Delta\psi$, whereas this process would be induced by the ΔpH in the chloroplast thylakoidal membrane. Altogether, it is presently still unclear whether a potential across the membrane drives charge movements or whether proton transport by Tat takes place.

8 Chaperoning of Tat Cargo Proteins

One of the major hallmarks of the Tat pathway is its ability to selectively transport fully folded cofactor-containing proteins. To this end, the system involves different mechanisms. Translocation of particular cargo proteins requires the aid of dedicated chaperones, known as redox enzyme maturation proteins (REMPs) [137, 138]. An example of a Tat cargo protein involving a REMF for export is the oxidoreductase trimethylamine-*N*-oxide (TMAO) reductase (TorA; Fig. 1). This enzyme is encoded by the *torCAD* operon, where *torA* encodes the TMAO reductase, *torC* its haem-binding quinol oxidase and *torD* its REMF. In particular, TorD recognizes and binds the h- and c-regions of the TorA signal peptide most likely as a dimer [139–141]. Following signal peptide binding, TorD guides TorA export via Tat in a GTP-dependent manner. In this scenario, the affinity of TorD for GTP increases upon signal peptide binding, and GTP presumably controls cycles of signal peptide binding and release of TorD, thereby preventing premature protein degradation, coordinating cofactor assembly and foreseeing other maturation steps, such as membrane targeting and interaction [139, 140]. This coordination occurs until the pre-protein interacts with the Tat machinery.

9 Proofreading of the Folding State of Tat Cargo

The proofreading exhibited by the *E. coli* and *B. subtilis* Tat pathways is highly stringent to ensure misfolded proteins are not exported. Thus, the Tat complex rejects and may

sometimes even degrade cargo proteins within the cytosol, although such degradation may also occur independently of the Tat system [142–144]. To note, the thylakoidal Tat system seems to have a less stringent ‘proofreading’ system as unfolded proteins are also imported [37].

To explore mechanisms of Tat proofreading, particular attention has been attributed to cofactor insertion. The native *E. coli* Tat cargo proteins NrfC and NapG were mutated to prevent their central cofactor FeS binding. Indeed this alteration blocked export [143]. The *B. subtilis* Rieske iron-sulphur cluster protein QcrA was also mutated to either stop cofactor binding or disulphide bond formation [145]. Here, a proofreading hierarchy was uncovered: mutant’s defective in disulphide bonding were quickly degraded, whereas those defective in cofactor binding accumulated in the cytoplasm and membrane. Two heme-binding proteins have also been investigated for proofreading. First, cytochrome C was shown to require heme insertion for export [146]. Subsequently, proofreading was investigated with the synthetic BT6 maquette protein, which binds two hemes and is Tat-dependently secreted in *E. coli* when provided with a TorA signal peptide [147]. His residues in BT6 were replaced with Ala to prevent heme binding. This showed that export was completely blocked if heme binding was completely prevented. Binding of one heme by BT6 allowed some export, whereas good export was observed when two hemes were bound [147]. Altogether, these findings suggest that Tat can somehow sense a protein’s conformational stability.

The requirement for conformational stability was further studied in vivo and in vitro with non-native Tat cargo proteins, such as *E. coli* PhoA and scFv or Fab antibody fragments. Export of these proteins only occurred in oxidizing conditions allowing disulphide bond formation prior to their interaction with the Tat machinery [40]. Nevertheless, some proteins provided with a twin-arginine signal peptide, like human growth hormone (hGH), scFv’s and interferon $\alpha 2b$, were exported to the periplasm without their disulphide bonds formed [148]. For hGH it was shown that this protein can form a near native state in absence of its two disulphide bonds. This is reminiscent of observations on the CueO protein of *E. coli*, which can still be exported via Tat without its bound copper cofactor. This probably relates to the fact that CueO without bound copper is structurally close to identical to CueO with bound copper [149].

Several studies in both bacteria and plants have used varying lengths of FG repeats from the yeast nuclear pore protein Nsp1p to probe the structural constraints for Tat-dependent export. These repeats intrinsically lack structure and are hydrophilic. Fused to a Tat signal peptide, export studies demonstrated that with increasing protein length, the translocation efficiency decreased: segments of 100–120 amino acids were tolerated, but a short hydrophobic stretch stopped export [150, 151]. Unstructured linkers were also

placed between the signal peptide and the N-terminus of a mature Tat cargo protein and, surprisingly, an unstructured linker length of 110 amino acids was exported [152]. These findings imply that, despite the generally strict folding requirement for Tat cargo proteins, short unstructured polypeptide regions can be tolerated in particular protein contexts.

A recent study used scFv mutants [153], which were structurally defined, to identify what the *E. coli* Tat machinery recognizes as ‘unfolded’ and rejects for export [154]. Tat tolerated significant changes in hydrophobicity and charge, but did not export the scFv with an unstructured tail or without cytoplasmic disulphide bond formation via the so-called CyDisCo system. CyDisCo comprises the yeast mitochondrial thiol oxidase Erv1p plus the human protein disulfide isomerase PDI that, together, confer the ability to catalyse cytoplasmic disulphide bond formation.

Since it is still unclear what exactly the Tat complex rejects as misfolded, a key question is how the Tat complex rejects certain proteins. Tat proteins, misfolded or not, both interact with the Tat translocase. For example the PhoA protein provided with a twin-arginine signal peptide has been co-purified with TatBC [41]. This gave rise to the idea Tat does not innately have an inbuilt ‘proofreading’ mechanism, but rather an efficient degradation system that clears the Tat translocase. Indeed, the *B. subtilis* protease WprA was shown to interact directly with the Tat machinery and to be essential for protein secretion via TatAyCy [155, 156].

Lastly, in vitro site-specific photo cross-linking experiments revealed that unfolded TorA-PhoA associated with the Tat translocase, and that the interaction with the TatBC receptor site was perturbed as if the cargo was not correctly inserted into the binding socket [157]. This invoked the TatBC complex in proofreading of the cargo protein. This view is consistent with the identification of so-called quality control suppression (QCS) mutations within *E. coli* TatABC, which gave rise to less stringent proofreading [158]. The majority of these QCS mutations were confined to the unstructured or loop regions of TatABC, showing that proofreading at some level is undertaken by the Tat translocase.

10 Conclusion

In recent years, the core components of the Tat protein translocation systems have been identified, biochemically characterized and structurally defined. Yet, the precise mechanism by which Tat translocates proteins across the bacterial cytoplasmic membrane is still elusive due to the fact that high-resolution structural data of protein-translocating Tat complexes is currently missing. It can be anticipated that with the advent of novel high-resolution techniques for structural analyses of large protein complexes many of the

so far unanswered fundamental questions in the Tat field can be tackled and answered.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Holland IB (2004) Translocation of bacterial proteins—an overview. *Biochim Biophys Acta Mol Cell Res* 1694:5–16
- Palmer T, Sargent F, Berks BC (2010) The Tat protein export pathway. *EcoSal Plus*. <https://doi.org/10.1128/ecosalplus.4.3.2>
- von Heijne G (1990) The signal peptide. *J Membr Biol* 115:195–201
- Lüke I, Handford JI, Palmer T, Sargent F (2009) Proteolytic processing of *Escherichia coli* twin-arginine signal peptides by LepB. *Arch Microbiol* 191:919–925
- Jongbloed JDH, Grieger U, Antelmann H et al (2004) Two minimal Tat translocases in *Bacillus*. *Mol Microbiol* 54:1319–1325
- Dalbey RE, Wang P, van Dijl JM (2012) Membrane proteases in the bacterial protein secretion and quality control pathway. *Microbiol Mol Biol Rev* 76:311–330
- Sakaguchi M, Tomiyoshi R, Kuroiwa T et al (1992) Functions of signal and signal-anchor sequences are determined by the balance between the hydrophobic segment and the N-terminal charge. *Proc Natl Acad Sci* 89:16–19
- Berks BC (1996) A common export pathway for proteins binding complex redox cofactors? *Mol Microbiol* 22:393–404
- Sargent F, Bogsch EG, Stanley NR et al (1998) Overlapping functions of components of a bacterial Sec-independent protein export pathway. *EMBO J* 17:3640–3650
- Berks BC, Palmer T, Sargent F (2003) The Tat protein translocation pathway and its role in microbial physiology. *Adv Microb Physiol* 47:187–254
- Stanley NR, Palmer T, Berks BC (2000) The twin arginine consensus motif of Tat signal peptides is involved in Sec-independent protein targeting in *Escherichia coli*. *J Biol Chem* 275:11591–11596
- Buchanan G, Sargent F, Berks BC, Palmer T (2002) A genetic screen for suppressors of *Escherichia coli* Tat signal peptide mutations establishes a critical role for the second arginine within the twin-arginine motif. *Arch Microbiol* 177:107–112
- Chaddock AM, Mant A, Karnachov I et al (1995) A new type of signal peptide: central role of a twin-arginine motif in transfer signals for the delta pH-dependent thylakoidal protein translocase. *EMBO J* 14:2715–2722
- Halbig D, Hou B, Freudl R et al (1999) Bacterial proteins carrying twin-R signal peptides are specifically targeted by the ΔpH-dependent transport machinery of the thylakoid membrane system. *FEBS Lett* 447:95–98
- DeLisa MP, Samuelson P, Palmer T, Georgiou G (2002) Genetic analysis of the twin arginine translocator secretion pathway in bacteria. *J Biol Chem* 277:29825–29831
- Hinsley AP, Stanley NR, Palmer T, Berks BC (2001) A naturally occurring bacterial Tat signal peptide lacking one of the “invariant” arginine residues of the consensus targeting motif. *FEBS Lett* 497:45–49

17. Kipping M, Lilie H, Lindenstrauß U et al (2003) Structural studies on a twin-arginine signal sequence. *FEBS Lett* 550:18–22
18. Cristóbal S, De Gier JW, Nielsen H, Von Heijne G (1999) Competition between Sec- and TAT-dependent protein translocation in *Escherichia coli*. *EMBO J* 18:2982–2990
19. Bogsch E, Brink S, Robinson C (1997) Pathway specificity for a Δ pH-dependent precursor thylakoid lumen protein is governed by a “Sec-avoidance” motif in the transfer peptide and a “Sec-incompatible” mature protein. *EMBO J* 16:3851–3859
20. Tullman-ercek D, Delisa MP, Kawarasaki Y et al (2009) NIH Public Access 282:8309–8316
21. Mould RM, Robinson C (1991) A proton gradient is required for the transport of two luminal oxygen-evolving proteins across the thylakoid membrane. *J Biol Chem* 266:12189–12193
22. Klösgen RB, Brock IW, Herrmann RG, Robinson C (1992) Proton gradient-driven import of the 16 kDa oxygen-evolving complex protein as the full precursor protein by isolated thylakoids. *Plant Mol Biol* 18:1031–1034
23. Clark SA, Theg SM (1997) A folded protein can be transported across the chloroplast envelope and thylakoid membranes. *Mol Biol Cell* 8:923–934
24. Mori H, Summer EJ, Ma X, Cline K (1999) Component specificity for the thylakoidal Sec and Δ pH-dependent protein transport pathways. *J Cell Biol* 146:45–55
25. Settles AM, Yonetani A, Baron A et al (1997) Sec-independent protein translocation by the maize Hcf106 protein. *Science* 278:1467–1470
26. Cline K, Mori H (2001) Thylakoid Δ pH-dependent precursor proteins bind to a cpTatC-Hcf106 complex before Tha4-dependent transport. *J Cell Biol* 154:719–729
27. Wu LF, Ize B, Chanal A, Quentin Y, Fichant G (2000) Bacterial twin-arginine signal peptide-dependent protein translocation pathway: evolution and mechanism. *J Mol Microbiol Biotechnol* 2:179–189
28. Yen MR, Tseng YH, Nguyen EH et al (2002) Sequence and phylogenetic analyses of the twin-arginine targeting (Tat) protein export system. *Arch Microbiol* 177:441–450
29. Bogsch EG, Sargent F, Stanley NR et al (1998) An essential component of a novel bacterial protein export system with homologues in plastids and mitochondria. *J Biol Chem* 273:18003–18006
30. Sargent F, Stanley NR, Berks BC, Palmer T (1999) Sec-independent protein translocation in *Escherichia coli*. A distinct and pivotal role for the TatB protein. *J Biol Chem* 274:36073–36082
31. Weiner JH, Bilous PT, Shaw GM et al (1998) A novel and ubiquitous system for membrane targeting and secretion of cofactor-containing proteins. *Cell* 93:93–101
32. Rodrigue A, Chanal A, Beck B, Muller M, Wu LF (1999) Cotranslocation of a periplasmic enzyme complex by a hitchhiker mechanism through the bacterial tat pathway. *J Biol Chem* 274:13223–13228
33. Palmer T, Sargent F, Berks BC (2005) Export of complex cofactor-containing proteins by the bacterial Tat pathway. *Trends Microbiol* 13:175–180
34. Bernhardt TG, De Boer PAJ (2003) The *Escherichia coli* amidase AmiC is a periplasmic septal ring component exported via the twin-arginine transport pathway. *Mol Microbiol* 48:1171–1182
35. Ize B, Stanley NR, Buchanan G, Palmer T (2003) Role of the *Escherichia coli* Tat pathway in outer membrane integrity. *Mol Microbiol* 48:1183–1193
36. Ding Z, Christie PJ (2003) *Agrobacterium tumefaciens* twin-arginine-dependent translocation is important for virulence, flagellation, and chemotaxis but not type IV secretion. *J Bacteriol* 185:760–771
37. Hynds PJ, Robinson D, Robinson C (1998) The Sec-independent twin-arginine translocation system can transport both tightly folded and misfolded proteins across the thylakoid membrane. *J Biol Chem* 273:34868–34874
38. Santini CL, Bernadac A, Zhang M et al (2001) Translocation of jellyfish green fluorescent protein via the Tat system of *Escherichia coli* and change of its periplasmic localization in response to osmotic up-shock. *J Biol Chem* 276:8159–8164
39. Alanen HI, Walker KL, Lourdes Velez Suberbie M et al (2014) Efficient export of human growth hormone, interferon α 2b and antibody fragments to the periplasm by the *Escherichia coli* Tat pathway in the absence of prior disulfide bond formation. *Biochim Biophys Acta Mol Cell Res* 1853:756–763
40. Delisa MP, Tullman D, Georgiou G (2003) Folding quality control in the export of proteins by the bacterial twin-arginine translocation pathway. *PNAS* 100:6115–6120
41. Richter S, Brüser T (2005) Targeting of unfolded PhoA to the TAT translocon of *Escherichia coli*. *J Biol Chem* 280:42723–42730
42. Jack RL, Sargent F, Berks BC et al (2001) Constitutive expression of *Escherichia coli* tat genes indicates an important role for the twin-arginine translocase during aerobic and anaerobic growth. *J Bacteriol* 183:1801–1804
43. Baglieri J, Beck D, Vasisht N et al (2012) Structure of TatA paralog, TatE, suggests a structurally homogeneous form of Tat protein translocase that transports folded proteins of differing diameter. *J Biol Chem* 287:7335–7344
44. Stanley NR, Findlay K, Berks BC, Palmer T (2001) *Escherichia coli* strains blocked in Tat-dependent protein export exhibit pleiotropic defects in the cell envelope. *J Bacteriol* 183:139–144
45. Sargent F, Gohlke U, De Leeuw E et al (2001) Purified components of the *Escherichia coli* Tat protein transport system form a double-layered ring structure. *Eur J Biochem* 268:3361–3367
46. Koch S, Fritsch MJ, Buchanan G, Palmer T (2012) *Escherichia coli* TatA and TatB proteins have N-out, C-in topology in intact cells. *J Biol Chem* 287:14420–14431
47. Hu Y, Zhao E, Li H et al (2010) Solution NMR structure of the TatA component of the twin-arginine protein transport system from gram-positive bacterium *Bacillus subtilis*. *J Am Chem Soc* 132:15942–15944
48. Rodriguez F, Rouse SL, Tait CE et al (2013) Structural model for the protein-translocating element of the twin-arginine transport system. *Proc Natl Acad Sci USA* 110:E1092–E1101
49. Barrett CML, Ray N, Thomas JD et al (2003) Quantitative export of a reporter protein, GFP, by the twin-arginine translocation pathway in *Escherichia coli*. *Biochem Biophys Res Commun* 304:279–284
50. Hicks MG, De Leeuw E, Porcelli I et al (2003) The *Escherichia coli* twin-arginine translocase: conserved residues of TatA and TatB family components involved in protein transport. *FEBS Lett* 539:61–67
51. Greene NP, Porcelli I, Buchanan G et al (2007) Cysteine scanning mutagenesis and disulfide mapping studies of the TatA component of the bacterial twin arginine translocase. *J Biol Chem* 282:23937–23945
52. Eimer E, Fröbel J, Blümmel AS, Müller M (2015) TatE as a regular constituent of bacterial twin-arginine protein translocases. *J Biol Chem* 290:29281–29289
53. Eimer E, Kao WC, Fröbel J et al (2018) Unanticipated functional diversity among the TatA-type components of the Tat protein translocase. *Sci Rep* 8:1–12
54. Zhang Y, Wang L, Hu Y, Jin C (2014) Solution structure of the TatB component of the twin-arginine translocation system. *Biochim Biophys Acta Biomembr* 1838:1881–1888

55. Lee PA, Buchanan G, Stanley NR et al (2002) Truncation analysis of TatA and TatB defines the minimal functional units required for protein translocation. *J Bacteriol* 184:5871–5879
56. Barrett CML, Mathers JE, Robinson C (2003) Identification of key regions within the *Escherichia coli* TatAB subunits. *FEBS Lett* 537:42–46
57. Blaudeck N, Kreutzenbeck P, Müller M et al (2005) Isolation and characterization of bifunctional *Escherichia coli* TatA mutant proteins that allow efficient Tat-dependent protein translocation in the absence of TatB. *J Biol Chem* 280:3426–3432
58. Barrett CML, Freudl R, Robinson C (2007) Twin arginine translocation (Tat)-dependent export in the apparent absence of TatABC or TAtA complexes using modified *Escherichia coli* TAtA subunits that substitute for TatB. *J Biol Chem* 282:36206–36213
59. Jongbloed JDH, Van Der Ploeg R, Van Dijl JM (2006) Bifunctional TAtA subunits in minimal Tat protein translocases [2]. *Trends Microbiol* 14:2–4
60. Tarry MJ, Schäfer E, Chen S et al (2009) Structural analysis of substrate binding by the TatBC component of the twin-arginine protein transport system. *Proc Natl Acad Sci USA* 106:13284–13289
61. Kneuper H, Maldonado B, Jäger F et al (2012) Molecular dissection of TatC defines critical regions essential for protein transport and a TatB-TatC contact site. *Mol Microbiol* 85:945–961
62. Behrendt J, Standar K, Lindenstrauß U, Brüser T (2004) Topological studies on the twin-arginine translocase component TatC. *FEMS Microbiol Lett* 234:303–308
63. Ramasamy S, Abrol R, Suloway CJM, Clemons WM (2013) The glove-like structure of the conserved membrane protein TatC provides insight into signal sequence recognition in twin-arginine translocation. *Structure* 21:777–788
64. Rollauer SE, Tarry MJ, Graham JE, Jääskeläinen M, Jäger F, Johnson S, Krehenbrink M, Liu SM, Lukey MJ, Marcoux J, McDowell MA, Rodriguez F, Roversi P, Stansfeld PJ, Robinson CV, Sansom MS, Palmer T, Högbom M, Berks BC, Lea SM (2012) Structure of the TatC core of the twin-arginine protein transport system. *Nature* 492(7428):210–214. <https://doi.org/10.1038/nature11683>
65. Blümmel AS, Drepper F, Knapp B et al (2017) Structural features of the TatC membrane protein that determine docking and insertion of a twin-arginine signal peptide. *J Biol Chem* 292:21320–21329
66. Holzapfel E, Eisner G, Alami M et al (2007) The entire N-terminal half of TatC is involved in twin-arginine precursor binding. *Biochemistry* 46:2892–2898
67. Goosens VJ, Monteferrante CG, Van Dijl JM (2014) The Tat system of Gram-positive bacteria. *Biochim Biophys Acta Mol Cell Res* 1843:1698–1706
68. Beck D, Vasisht N, Baglieri J et al (2009) Subcellular localization of TatAd of *Bacillus subtilis* depends on the presence of TatCd or TatCy. *J Biol Chem* 191:4410–4418
69. Schaerlaekens K, Schierová M, Lammertyn E et al (2001) Twin-arginine translocation pathway in *Streptomyces lividans*. *J Bacteriol* 183:6727–6732
70. Jongbloed JDH, Martin U, Antelmann H et al (2000) TatC is a specificity determinant for protein secretion via the twin-arginine translocation pathway. *J Biol Chem* 275:41350–41357
71. van der Ploeg R, Mäder U, Homuth G et al (2011) Environmental salinity determines the specificity and need for tat-dependent secretion of the YwbN protein in *Bacillus subtilis*. *PLoS ONE* 6:18140
72. Monteferrante CG, Miethke M, Van Der Ploeg R et al (2012) Specific targeting of the metallophosphoesterase YkuE to the *Bacillus* cell wall requires the twin-arginine translocation system. *J Biol Chem* 287:29789–29800
73. Miethke M, Monteferrante CG, Marahiel MA, van Dijl JM (2013) The *Bacillus subtilis* EfeUOB transporter is essential for high-affinity acquisition of ferrous and ferric iron. *Biochim Biophys Acta Mol Cell Res* 1833:2267–2278
74. Goosens VJ, Otto A, Glasner C et al (2013) Novel twin-arginine translocation pathway-dependent phenotypes of *Bacillus subtilis* unveiled by quantitative proteomics. *J Proteome Res* 12:796–807
75. Goosens VJ, De-San-Eustaquio-Campillo A, Carballido-López R, van Dijl JM (2015) A Tat ménage à trois—the role of *Bacillus subtilis* TAtA in twin-arginine protein translocation. *Biochim Biophys Acta Mol Cell Res* 1853:2745–2753
76. Barnett JP, Eijlander RT, Kuipers OP, Robinson C (2008) A minimal tat system from a gram-positive organism: a bifunctional TAtA subunit participates in discrete TatAC and TAtA complexes. *J Biol Chem* 283:2534–2542
77. Monteferrante CG, Baglieri J, Robinson C, van Dijl JM (2012) TAtA, the third TAtA subunit of *Bacillus subtilis*, can form active twin-arginine translocases with the TatCd and TatCy subunits. *Appl Environ Microbiol* 78:4999–5001
78. Beck D, Vasisht N, Baglieri J et al (2013) Ultrastructural characterisation of *Bacillus subtilis* TAtA complexes suggests they are too small to form homooligomeric translocation pores. *Biochim Biophys Acta Mol Cell Res* 1833:1811–1819
79. Porcelli I, De Leeuw E, Wallis R et al (2002) Characterization and membrane assembly of the TAtA component of the *Escherichia coli* twin-arginine protein transport system. *Biochemistry* 41:13690–13697
80. Müller SD, De Angelis AA, Walther TH et al (2007) Structural characterization of the pore forming protein TatAd of the twin-arginine translocase in membranes by solid-state 15 N-NMR. *Biochim Biophys Acta Biomembr* 1768:3071–3079
81. Walther TH, Grage SL, Roth N, Ulrich AS (2010) Membrane alignment of the pore-forming component TatAd of the twin-arginine translocase from *Bacillus subtilis* resolved by solid-state NMR spectroscopy. *J Am Chem Soc* 132:15945–15956
82. Lange C, Müller SD, Walther TH et al (2007) Structure analysis of the protein translocating channel TAtA in membranes using a multi-construct approach. *Biochim Biophys Acta Biomembr* 1768:2627–2634
83. Barnett JP, Lawrence J, Mendel S, Robinson C (2011) Expression of the bifunctional *Bacillus subtilis* TatAd protein in *Escherichia coli* reveals distinct TAtA/B-family and TatB-specific domains. *Arch Microbiol* 193:583–594
84. Schreiber S, Stengel R, Westermann M et al (2006) Affinity of TatCd for TAtA delucidates its receptor function in the *Bacillus subtilis* twin arginine translocation (Tat) translocase system. *J Biol Chem* 281:19977–19984
85. Nolandt OV, Walther TH, Roth S et al (2009) Structure analysis of the membrane protein TatCd from the Tat system of *B. subtilis* by circular dichroism. *Biochim Biophys Acta Biomembr* 1788:2238–2244
86. Van Der Ploeg R, Barnett JP, Vasisht N et al (2011) Salt sensitivity of minimal twin arginine translocases. *J Biol Chem* 286:43759–43770
87. Goosens VJ, van Dijl JM (2017) Twin-arginine protein translocation. *Curr Top Microbiol Immunol* 404:69–94
88. Patel R, Vasilev C, Beck D et al (2014) A mutation leading to super-assembly of twin-arginine translocase (Tat) protein complexes. *Biochim Biophys Acta Mol Cell Res* 1843:1978–1986
89. Eijlander RT, Jongbloed JDH, Kuipers OP (2009) Relaxed specificity of the *Bacillus subtilis* TAtAcd translocase in Tat-dependent protein secretion. *J Bacteriol* 91:196–202
90. Eijlander RT, Kolbusz MA, Berendsen EM, Kuipers OP (2009) Effects of altered TatC proteins on protein secretion efficiency

- via the twin-arginine translocation pathway of *Bacillus subtilis*. *Microbiology* 155:1776–1785
91. Berks BC, Sargent F, Palmer T (2000) The Tat protein export pathway. *Mol Microbiol* 35:260–274
 92. Rose P, Fröbel J, Graumann PL, Müller M (2013) Substrate-dependent assembly of the Tat translocase as observed in live *Escherichia coli* cells. *PLoS ONE* 8:1–17
 93. Berks BC, Lea SM, Stansfeld PJ (2014) Structural biology of tat protein transport. *Curr Opin Struct Biol* 27:32–37
 94. Alami M, Lüke I, Deitermann S et al (2003) Differential interactions between a twin-arginine signal peptide and its translocase in *Escherichia coli*. *Mol Cell* 12:937–946
 95. Celedon JM, Cline K (2012) Stoichiometry for binding and transport by the twin Arginine translocation system. *J Cell Biol* 197:523–534
 96. Alcock F, Baker MAB, Greene NP et al (2013) Live cell imaging shows reversible assembly of the TatA component of the twin-arginine protein transport system. *Proc Natl Acad Sci* 110:E3650–E3659
 97. Gérard F, Cline K (2006) Efficient twin arginine translocation (Tat) pathway transport of a precursor protein covalently anchored to its initial cpTatC binding site. *J Biol Chem* 281:6130–6135
 98. Kreutzenbeck P, Kröger C, Lausberg F et al (2007) *Escherichia coli* twin arginine (Tat) mutant translocases possessing relaxed signal peptide recognition specificities. *J Biol Chem* 282:7903–7911
 99. Maurer C, Panahandeh S, Jungkamp A-C et al (2010) TatB functions as an oligomeric binding site for folded Tat precursor proteins. *Mol Biol Cell* 21:4151–4161
 100. Zoufaly S, Fröbel J, Rose P et al (2012) Mapping precursor-binding site on TatC subunit of twin arginine-specific protein translocase by site-specific photo cross-linking. *J Biol Chem* 287:13430–13441
 101. Fröbel J, Rose P, Lausberg F et al (2012) Transmembrane insertion of twin-arginine signal peptides is driven by TatC and regulated by TatB. *Nat Commun* 3:1311
 102. Blümmel AS, Haag LA, Eimer E et al (2015) Initial assembly steps of a translocase for folded proteins. *Nat Commun* 6:7234
 103. Bolhuis A, Mathers JE, Thomas JD et al (2001) TatB and TatC form a functional and structural unit of the twin-arginine translocase from *Escherichia coli*. *J Biol Chem* 276:20213–20219
 104. Behrendt J, Brüser T (2014) The TatBC complex of the tat protein translocase in *Escherichia coli* and its transition to the substrate-bound TatABC complex. *Biochemistry* 53:2344–2354
 105. Ma X, Cline K (2010) Multiple precursor proteins bind individual Tat receptor complexes and are collectively transported. *EMBO J* 29:1477–1488
 106. Rollauer SE, Tarry MJ, Graham JE et al (2012) Structure of the TatC core of the twin-arginine protein transport system. *Nature* 492:210–214
 107. Lee PA, Orriss GL, Buchanan G et al (2006) Cysteine-scanning mutagenesis and disulfide mapping studies of the conserved domain of the twin-arginine translocase TatB component. *J Biol Chem* 281:34072–34085
 108. Cline K (2015) Mechanistic aspects of folded protein transport by the twin arginine translocase (Tat). *J Biol Chem* 290:16530–16538
 109. Alcock F, Stansfeld PJ, Basit H et al (2016) Assembling the tat protein translocase. *Elife* 5:e20718
 110. Habersetzer J, Moore K, Cherry J et al (2017) Substrate-triggered position switching of TatA and TatB during Tat transport in *Escherichia coli*. *Open Biol* 7:8
 111. Orriss GL, Tarry MJ, Ize B et al (2007) TatBC, TatB, and TatC form structurally autonomous units within the twin arginine protein transport system of *Escherichia coli*. *FEBS Lett* 581:4091–4097
 112. Aldridge C, Ma X, Gerard F, Cline K (2014) Substrate-gated docking of pore subunit Tha4 in the TatC cavity initiates Tat translocase assembly. *J Cell Biol* 205:51–65
 113. Hauer RS, Schlesier R, Heilmann K et al (2013) Enough is enough: TatA demand during Tat-dependent protein transport. *Biochim Biophys Acta Mol Cell Res* 1833:957–965
 114. Oates J, Barrett CML, Barnett JP et al (2005) The *Escherichia coli* twin-arginine translocation apparatus incorporates a distinct form of TatABC complex, spectrum of modular TatA complexes and minor TatAB complex. *J Mol Biol* 346:295–305
 115. White GF, Schermann SM, Bradley J et al (2010) Subunit organization in the TatA complex of the twin arginine protein translocase: a site-directed EPR spin labeling study. *J Biol Chem* 285:2294–2301
 116. Gohlke U, Pullan L, McDevitt CA et al (2005) The TatA component of the twin-arginine protein transport system forms channel complexes of variable diameter. *Proc Natl Acad Sci USA* 102:10482–10486
 117. Brüser T, Sanders C (2003) An alternative model of the twin arginine translocation system. *Microbiol Res* 158:7–17
 118. Gouffi K, Gérard F, Santini CL, Wu LF (2004) Dual topology of the *Escherichia coli* TatA protein. *J Biol Chem* 279:11608–11615
 119. Walther TH, Gottselig C, Grage SL et al (2013) Folding and self-assembly of the TatA translocation pore based on a charge zipper mechanism. *Cell* 152:316–326
 120. Dabney-Smith C, Mori H, Cline K (2006) Oligomers of Tha4 organize at the thylakoid Tat translocase during protein transport. *J Biol Chem* 281:5476–5483
 121. Chan CS, Zlomslis MR, Tieleman DP, Turner RJ (2007) The TatA subunit of *Escherichia coli* twin-arginine translocase has an N-in topology. *Biochemistry* 46:7396–7404
 122. Aldridge C, Storm A, Cline K, Dabney-Smith C (2012) The chloroplast twin arginine transport (Tat) component, Tha4, undergoes conformational changes leading to Tat protein transport. *J Biol Chem* 287:34752–34763
 123. Taubert J, Hou B, Risselada HJ et al (2015) TatBC-Independent TatA/Tat substrate interactions contribute to transport efficiency. *PLoS ONE* 10:1–24
 124. Pal D, Fite K, Dabney-Smith C (2013) Direct interaction between a precursor mature domain and transport component Tha4 during twin arginine transport of chloroplasts. *Plant Physiol* 161:990–1001
 125. Taubert J, Brüser T (2014) Twin-arginine translocation-arresting protein regions contact TatA and TatB. *Biol Chem* 395:827–836
 126. Hou B, Heidrich ES, Mehner-Breitfeld D, Brüser T (2018) The TatA component of the twin-arginine translocation system locally weakens the cytoplasmic membrane of *Escherichia coli* upon protein substrate binding. *J Biol Chem* 293:7592–7605
 127. Mould RM, Shackleton JB, Robinson C (1991) Transport of proteins into chloroplasts. Requirements for the efficient import of two luminal oxygen-evolving complex proteins into isolated thylakoids. *J Biol Chem* 266:17286–17289
 128. Mori H, Cline K (2002) A twin arginine signal peptide and the pH gradient trigger reversible assembly of the thylakoid Δ pH/Tat translocase. *J Cell Biol* 157:205–210
 129. DeLisa MP, Lee P, Palmer T, Georgiou G (2004) Phage shock protein PspA of *Escherichia coli* relieves saturation of protein export via the Tat pathway. *J Bacteriol* 186:366–373
 130. Finazzi G, Chasen C, Wollman FA, De Vitry C (2003) Thylakoid targeting of Tat passenger proteins shows no Δ pH dependence in vivo. *EMBO J* 22:807–815
 131. Braun NA, Davis AW, Theg SM (2007) The chloroplast tat pathway utilizes the transmembrane electric potential as an energy source. *Biophys J* 93:1993–1998

132. Musser SM, Theg SM (2000) Proton transfer limits protein translocation rate by the thylakoid $\Delta\text{pH}/\text{Tat}$ machinery. *Biochemistry* 39:8228–8233
133. Alder NN, Theg SM (2003) Energetics of protein transport across biological membranes: a study of the thylakoid ΔpH -dependent/cpTat pathway. *Cell* 112:231–242
134. Gérard F, Cline K (2007) The thylakoid proton gradient promotes an advanced stage of signal peptide binding deep within the Tat pathway receptor complex. *J Biol Chem* 282:5263–5272
135. Dabney-Smith C, Mori H, Cline K (2003) Requirement of a Tha4-conserved transmembrane glutamate in thylakoid Tat translocase assembly revealed by biochemical complementation. *J Biol Chem* 278:43027–43033
136. Bageshwar UK, Musser SM (2007) Two electrical potential-dependent steps are required for transport by the *Escherichia coli* Tat machinery. *J Cell Biol* 179:87–99
137. Turner RJ, Papish AL, Sargent F (2004) Sequence analysis of bacterial redox enzyme maturation proteins (REMPs). *Can J Microbiol* 50:225–238
138. Chan CS, Chang L, Rommens KL, Turner RJ (2009) Differential interactions between tat-specific redox enzyme peptides and their chaperones. *J Bacteriol* 191:2091–2101
139. Jack RL, Buchanan G, Dubini A, Hatzixanthis K, Palmer T, Sargent F (2004) Coordinating assembly and export of complex bacterial proteins. *EMBO J* 23:3962–3972
140. Hatzixanthis K, Clarke TA, Oubrie A, Richardson DJ, Turner RJ, Sargent F (2005) Signal peptide-chaperone interactions on the twin-arginine protein transport pathway. *Proc Natl Acad Sci USA* 102:8460–8465
141. Tranier S, Mortier-Barrière I, Ilbert M et al (2002) Characterization and multiple molecular forms of TorD from *Shewanella massilia*, the putative chaperone of the molybdoenzyme TorA. *Protein Sci* 11:2148–2157
142. Halbig D, Wiegert T, Blaudeck N et al (1999) The efficient export of NADP-containing glucose-fructose oxidoreductase to the periplasm of *Zymomonas mobilis* depends both on an intact twin-arginine motif in the signal peptide and on the generation of a structural export signal induced by cofactor bind. *Eur J Biochem* 263:543–551
143. Matos CFRO, Robinson C, Di Cola A (2008) The Tat system proofreads FeS protein substrates and directly initiates the disposal of rejected molecules. *EMBO J* 27:2055–2063
144. Lindenstrauß U, Matos CF, Graubner W, Robinson C, Brüser T (2010) Malfolded recombinant Tat substrates are Tat-independently degraded in *Escherichia coli*. *FEBS Lett* 584:3644–3648
145. Goosens VJ, Monteferrante CG, Van Dijl JM (2014) Co-factor insertion and disulfide bond requirements for twin-Arginine translocase-dependent export of the *Bacillus subtilis* Rieske protein QcrA. *J Biol Chem* 289:13124–13131
146. Sanders C, Wethkamp N, Lill H (2001) Transport of cytochrome c derivatives by the bacterial Tat protein translocation system. *Mol Microbiol* 41:241–246
147. Sutherland GA, Grayson KJ, Adams NBP et al (2018) Probing the quality control mechanism of the *Escherichia coli* twin-arginine translocase with folding variants of a *de novo*—designed heme protein. *J Biol Chem* 293:6672–6681
148. Alanen HI, Walker KL, Lourdes Velez Suberbie M et al (2015) Efficient export of human growth hormone, interferon $\alpha 2\text{b}$ and antibody fragments to the periplasm by the *Escherichia coli* Tat pathway in the absence of prior disulfide bond formation. *Biochim Biophys Acta Mol Cell Res* 1853:756–763
149. Stolle P, Hou B, Brüser T (2016) The Tat substrate CueO is transported in an incomplete folding state. *J Biol Chem* 291:13520–13528
150. Cline K, McCaffery M (2007) Evidence for a dynamic and transient pathway through the TAT protein transport machinery. *EMBO J* 26:3039–3049
151. Richter S, Lindenstrauß U, Lücke C et al (2007) Functional tat transport of unstructured, small, hydrophilic proteins. *J Biol Chem* 282:33257–33264
152. Lindenstrauß U, Brüser T (2009) Tat transport of linker-containing proteins in *Escherichia coli*. *FEMS Microbiol Lett* 295:135–140
153. Austerberry JI, Dajani R, Panova S et al (2017) The effect of charge mutations on the stability and aggregation of a human single chain Fv fragment. *Eur J Pharm Biopharm* 115:18–30
154. Jones AS, Austerberry JI, Dajani R et al (2016) Proofreading of substrate structure by the Twin-Arginine Translocase is highly dependent on substrate conformational flexibility but surprisingly tolerant of surface charge and hydrophobicity changes. *Biochim Biophys Acta Mol Cell Res* 1863:3116–3124
155. Monteferrante CG, Mackichan C, Marchadier E et al (2013) Mapping the twin-arginine protein translocation network of *Bacillus subtilis*. *Proteomics* 13:800–811
156. Krishnappa L, Monteferrante CG, van Dijl JM (2012) Degradation of the twin-arginine translocation substrate YwBn by extracytoplasmic proteases of *Bacillus subtilis*. *Appl Environ Microbiol* 78:7801–7804
157. Panahandeh S, Maurer C, Moser M et al (2008) Following the path of a twin-arginine precursor along the TatABC translocase of *Escherichia coli*. *J Biol Chem* 283:33267–33275
158. Rocco MA, Waraho-Zhmayev D, DeLisa MP (2012) Twin-arginine translocase mutations that suppress folding quality control and permit export of misfolded substrate proteins. *Proc Natl Acad Sci* 109:13392–13397
159. Barnett JP, Van Der Ploeg R, Eijlander RT, Nenninger A, Mendel S, Rozeboom R, Kuipers OP, Van Dijl JM, Robinson C (2009) The twin-arginine translocation (Tat) systems from *Bacillus subtilis* display a conserved mode of complex organization and similar substrate recognition requirements. *FEBS J* 276:232–243

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.