

ORIGINAL RESEARCH

Continuous Education of Basic Life Support (BLS) through Social Media; a Quasi-Experimental Study

Seyyed Mahdi Zia Ziabari¹, Vahid Monsef Kasmaei¹, Lida Khoshgozaran^{1*}, Maryam Shakiba²

1. Department of Emergency Medicine, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

2. Guilan Road Trauma Research Center, School of Health, Guilan University of Medical Sciences, Rasht, Iran.

Received: September 2018; Accepted: October 2018; Published online: 20 January 2019

Abstract: **Introduction:** It is imperative, if not fundamental, for medical students to receive continuing education in Basic life support (BLS). This present study aimed to investigate the effectiveness of continuing education using social media in this regard. **Methods:** In this quasi-experimental study, a questionnaire containing 20 items about BLS awareness was completed by the interns who were trained monthly in the emergency medicine training course. Then half of the interns received distant education through Telegram software up to three months, and the other did not receive any further education. Again, both groups completed the questionnaire at the end of the third month. **Results:** 100 medical interns with the mean age of 25.05 ± 1.26 years were allocated to two groups of 50 subjects (65.0% female). Two groups were similar regarding the mean age ($p = 0.304$), gender ($p = 0.529$), mean awareness score at the beginning of the study ($p = 0.916$), and average pre-internship exam score ($p = 0.080$). The mean awareness score of medical interns increased significantly in intervention (11.44 ± 2.37 to 14.88 ± 1.97 , $p < 0.0001$) and control (11.38 ± 3.22 to 12.54 ± 3.04 , $p < 0.0001$) groups at second examination. But mean difference of awareness score, before and after the education, was significantly higher in intervention group (3.44 ± 1.48 versus 1.16 ± 1.51 ; $p < 0.0001$). **Conclusion:** The increase in the BLS awareness score was significantly higher in medical interns who were continuously educated using Telegram social media software.

Keywords: Education; cardiopulmonary resuscitation; mobile application; computer-assisted instruction

© Copyright (2019) Shahid Beheshti University of Medical Sciences

Cite this article as: Zia Ziabari SM, Monsef Kasmaei V, Khoshgozaran L, Shakiba M. Continuous Education of Basic Life Support (BLS) through Social Media; a Quasi-Experimental Study. Arch Acad Emerg Med. 2019; 7(1): e4.

1. Introduction

Despite many advances in medicine, sudden cardiac arrest remains a leading cause of death in most countries worldwide (1). Subsequent outcomes of cardiac arrest depend on basic measures such as chest compression, defibrillator, and more advanced actions (2). The importance of cardiopulmonary resuscitation (CPR) has been well documented, so that a good CPR can reduce mortality up to 50% (3). In general, the ability to diagnose and treat a respiratory or cardiac arrest is a basic medical skill that all health care providers including medical students should have these skills. They should learn to carry out CPR and pre-hospital activities and increase their skills during educational courses (4, 5). The

measures for basic life support (BLS) for adolescents outside the hospital include rapid detection of cardiac arrest, quick activation of the emergency medical services (EMS), pulse check, CPR, and rapid defibrillation with an automated external defibrillator (AED) (6). Since the quality of CPR depends on the learner memorizing and applying the scientific and practical skills, continuing education should be provided in this regard (7). The use of e-learning in medical education has had good results and has been effective (8, 9). In this way, electronic tools and software are used as a means to improve learning, communication and interactions, and increasingly to train medical professionals (10, 11). The advancement in e-learning technology and the use of computers and online training has made it possible to save costs and transfer data without any time limit. Also, the person-to-person training capabilities meet the learner's needs well (12, 13). Although the BLS course is held every two years, researches have shown that quality of BLS clinical skills begin to decrease after 3 months (14-16). Studies showed that the

*Corresponding Author: Lida Khoshgozaran, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran. E-mail: Lida.khoshgozaran@gmail.com Tel: 00989111414384






Figure 1: Mean awareness score in intervention and control groups before and after the education ($p < 0.0001$).

students had more tendency and self-confidence to perform CPR after multiple educational sessions (17, 18). Researches have also shown that watching an educational video clip on cell phone to remind and review BLS content increases the skill and self-confidence to carry out CPR and the use of AED in rescuers (18). Considering the above mentioned, we decide to investigate the effectiveness of continuous education of BLS through social media (Telegram software) on medical interns.

2. Methods

2.1. Study design and setting

This quasi-experimental study, was conducted on medical interns who were trained monthly in the emergency medicine training course of Poursina Hospital, Rasht, Iran, from August 2016 to July 2017. All medical interns entered the study with informed consent, and reviewing the research project was carried out by the Ethics Committee of the Research Unit of Guilan University of Medical Sciences (IR.GUMS.REC.1396.69). the study was registered in Iranian Registry of clinical trials with number: IRCT20110818007369N5.

2.2. Participants

All medical interns who were passed the one month emergency medicine educational program in mentioned hospital during the study period were enrolled. Dissatisfaction with continuing to participate in the study, lack of access to smartphone, not using of Telegram software were among the exclusion criteria. There was no any sex or age limitation. The medical interns were given an ethical commitment that their personal information would remain confidential and they also gave a moral commitment to not share the contents of the classroom and the software until the end of the study with other interns.

2.3. Data gathering

To assess the students' awareness regarding the BLS at the beginning and end of the study, a questionnaire containing 20 items was used (19). The Persian version of this questionnaire was translated into English, and it was again translated into Persian by a native. To determine the quantitative validity, content validity ratio (CVR) and content validity index (CVI) were used. The score for cut-off point in Lawshe table for 10 experts was 0.62. In the questionnaire, the CVR scores of all questions were 0.8 to 1, which is confirmed in terms of CVR index. Also, in terms of the CVI, the lowest CVI was 0.7 and its maximum was 1 in the three dimensions of relevance, simplicity and clarity. Therefore, questions with a CVI of between 0.7 and 0.8 were subject to partial review, and the remaining questions due to the high CVI remained unchanged. To determine the internal consistency of the questions, the Richardson coefficient 20 was used, with the obtained alpha value equal to 0.996 (more than 0.7), indicating the internal consistency of the questions in the calculation of students' awareness to the BLS algorithm.

2.4. Intervention

Mentioned questionnaire was first distributed among medical interns, and they were asked to answer the questions based on their information and to enter data including their name, age, gender, and average pre-internship exam score in the questionnaire. Based on the answers to the questions, a score of 1-20 was attributed to the students (correct answer to each question had 1 score, the wrong answer was zero, and negative score did not belong to the wrong answers). The scores obtained from the questionnaire were gained with algebraic scores from each question (zero or one), and the students' awareness was measured based on these scores. After completing the questionnaire, all of the interns received an in-person tutorial using PowerPoint software by the teachers of Rasht Poursina hospital, and an emergency medicine resident, based on the 2015 AHA educational guideline on BLS for adults and children (1, 6, 20-22). After being trained at the hospital, half of the interns who were selected sequentially and non-randomly, received distant continuing education for up to three months through social media (Telegram software, which is comprehensive and accessible in Iran). This training was conducted in a person-to-person manner by sending abstracts of taught materials in the classroom based on the 2015 AHA guidelines, which was presented in short texts, photographs and educational videos related to how to perform BLS. The other half of the participants did not receive any further training after the in-person education offered at the hospital. Then, at the end of the third month after holding the first in-person class, both groups received the questionnaire through the Telegram software, and the aware-

Table 1: Baseline characteristics of case (continuous education with social media) and control (regular education) groups

Variable	Value	P values
Gender (male/female ratio)		
Case	31/19 (1.63)	0.529
Control	34/16(2.12)	
Age (year)		
Case	24.92 ± 1.01	0.304
Control	25.18 ± 1.46	
Awareness score		
Case	11.44 ± 2.37	0.916
Control	11.38 ± 3.22	
Pre-internship exam score		
Case	15.97 ± 0.81	0.080
Control	16.24 ± 0.71	

Data are presented as mean ± standard deviation.

ness on BLS was measured by refilling the same questionnaire.

2.5. Statistical Analysis

We calculated sample size using altman monogram with α : 0.01, β : 0.2 and d^2 : 0.9. The allocation ratio was 1:1, and finally 50 students entered in each group. Data were analysed using mean and standard deviation. Normality of data was evaluated using skewness and kurtosis indices. Paired t-test and independent t-test were used to compare means between the groups. All analyses were performed in SPSS version 19 and a significant level was considered as below 0.05.

3. Results

119 medical interns were studied. 19 out of 119 students were excluded (3 were not using the Telegram software, 4 were not satisfied to participate to the study, 12 did not fill the questionnaire after 3 months). Finally 100 medical interns with the mean age of 25.05 ± 1.26 (23 - 31) years were allocated to two groups of 50 subjects (65.0% female). Table 1 compare the baseline characteristics of two groups. Two groups were similar regarding the mean age ($p = 0.304$), gender ($p = 0.529$), mean awareness score at the beginning of the study ($p = 0.916$), and average pre-internship exam score ($p = 0.080$). The mean awareness score of medical interns increased significantly in intervention (11.44 ± 2.37 to 14.88 ± 1.97 , $p < 0.0001$) and control (11.38 ± 3.22 to 12.54 ± 3.04 , $p < 0.0001$) groups at second examination (figure 1). But mean difference of awareness score, before and after the education, was significantly higher in intervention group (3.44 ± 1.48 versus 1.16 ± 1.51 ; $p < 0.0001$).

4. Discussion

The findings of this study showed that the increase in the BLS awareness score was significantly higher in medical interns who were continuously educated using Telegram social media software. An interventional study by Khorasani et al. showed that the mean score of awareness and practice before and after the primary emergency care education workshops indicated a significant difference in practice before and after taking part in workshop in dentistry interns (23). Ji Yun Ahn et al. showed that watching an educational video clip in cell phone in order to remind and review the BLS materials increased skill and self-confidence to perform CPR and using AED in rescuers after three months (18). In our study, the impact of this continuous education was seen positively, although there was no need to use a memory card and a reminder message to view clips and educational pictures, and all interns could have enough time to read, understand, and answer person-to-person questions using Telegram software within three months. In a randomized controlled trial, Eric L. Einspruch et al. in the United States, compared memorizing CPR skills in conventional education to teaching through 22-minute self-learning videos. In both groups, the decrease amount in skills after two months was generally the same. In the second assessment two months later, the subjects of the self-learning group were better than the conventional group in all skills, except for the emergency call, which were the same for both groups. In the assessment with manikin, the self-learning group was better than the conventional training group in the ventilation volume, but both groups performed equally in proper hand positioning in chest compression and massage depth (7). Contrary to this study showing a decrease in skills after two months, the awareness of interns' after three-month continuing education did not reduce, but increased significantly compared to the control group in our study. In general, the ability to diagnose and treat a respiratory or cardiac arrest is a basic medical skill that all medical students should have the skills. They should learn CPR and pre-hospital activities and increase their skills during educational courses (4, 5). Given that these skills and awareness about how to conduct appropriate BLS are forgotten over time, and according to AHA 2015, studies have demonstrated that deterioration of BLS skills in as little as 3 months after initial training (14-16), therefore, it should be emphasized that in order to maintain and improve clinical skills, they need to be adequately memorized with repeated training (7). With the help of information and communication technology capabilities, students are able to provide information at any time and place quickly. The learner can have enough time to read, understand, and answer the questions in a person to person manner, in spite of his/her work, family, disability and geographical location; this encourages and



motivates him/her (24). Considering the widespread use of modern Telegram software in society especially in the student population, this study examined the effect of person to person and continuing education of CPR operation using Telegram among medical students. Since all students were spending their internship in different parts, the software provided continuous communication with each other.

5. Limitation

The weaknesses of this study were the lack of randomization and the impossibility of blinding due to the nature of the study, and the lack of facilities for examining the practical skills of students at the same time. Since it is not possible to examine skills in addition to knowledge, it is suggested that future studies consider these two issues together.

6. Conclusion

The results of this study indicate the positive effect of continuing education of BLS through Telegram software on the awareness of medical students. This finding can demonstrate the positive role of using modern communication networks and the possibility of using them in the student community.

7. Appendix

7.1. Acknowledgements

This research was supported by Guilan Road Trauma Research Center. We thank our colleagues from Rasht Poursina hospital who provided insight and expertise that greatly assisted the research.

7.2. Authors' contribution

Seyyed Mahdi Zia Ziabari devised the study, the main conceptual ideas and proof outline. Lida Khoshgozaran worked out almost all of the technical details. Maryam Shakiba performed analysis and interpretation of the data and Vahid Monsef Kasmaei took the lead in charge of overall direction and planning. All authors discussed the results and commented on the manuscript.

7.3. Funding/Support

None.

7.4. Conflict of interest

We know of no conflicts of interest associated with this publication, and there has been no significant financial support for this work that could have influenced its outcome. As Corresponding Author, I confirm that the manuscript has been read and approved for submission by all the named authors.

References

1. Travers AH RT, Bobrow BJ, et al. . CPR overview: American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2010;122(18):676-84.
2. Abella SB AP, Eigel B, et al. Reducing barriers for implementation of bystander-initiated cardiopulmonary resuscitation. *Circulation*. 2008;117(5):704-9.
3. Huikuri HV, Castellanos A, Myerburg RJ. Sudden death due to cardiac arrhythmias. *New England Journal of Medicine*. 2001;345(20):1473-82.
4. Meena Kumari K MBa, Suhas Alur S, Pavan Madhukar Bhat, Siddharth Bansal. Clinical awareness of do's and don'ts of cardiopulmonary resuscitation (CPR) among university medical students-A Questionnaire Study. *Journal of clinical and diagnostic research: JCDR*. 2014;8(7):MC08.
5. Einspruch EL LB, Aufderheide TP, Nichol G, Becker L. Retention of CPR skills learned in a traditional AHA Heart-saver course versus 30-min video self-training: a controlled randomized study. *Resuscitation*. 2007;74(3):476-86.
6. Kleinman ME BE, Goldberger ZD, Swor RA, Terry M, Bobrow BJ, Gazmuri RJ, Travers AH, Rea T. Part 5: Adult basic life support and cardiopulmonary resuscitation quality: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(18 suppl 2):S414-S35.
7. Bhanji F DA, Wolff MS, Flores GE, Halamek LP, Berman JM, Sinz EH, Cheng A. Part 14: education: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(18 suppl 2):S561-S73.
8. Elizabeth JC JJ, Jennifer RB, Matthew P, Rosalind LS, Paula RW and Munir P. A pilot randomised controlled trial to assess the utility of an e-learning package that trains users in adverse drug reaction causality. *International Journal of Pharmacy Practice*. 2015;23(6):447-55.
9. Gordon M, Chandratilake M, Baker P. Improved junior paediatric prescribing skills after a short e-learning intervention: a randomised controlled trial. *Archives of disease in childhood*. 2011;96(12):1191-4.
10. Ziva C RR, Petra AW and Josip C. Factors influencing the implementation, adoption, use, sustainability and scalability of eLearning for family medicine specialty training: a systematic review protocol. *Systematic reviews*. 2016;5(1):179.
11. Sangra A VD, Cabrera N. Building an inclusive definition of e-learning: An approach to the conceptual framework. *The International Review of Research in Open and Dis-*



- tributed Learning. 2012;13(2):145-59.
12. Triola MM, Huwendiek S, Levinson AJ, Cook DA. New directions in e-learning research in health professions education: report of two symposia. *Medical teacher*. 2012;34(1):e15-e20.
 13. Campbell J ea. A universal truth: no health without a workforce. *Global Health Workforce Alliance World Health Organization*. 2013.
 14. Bhanji F MM, Sinz E, Rodgers DL, McNeil MA, Hoadley TA, Meeks RA, Hamilton ME, Meaney PA, Hunt EA, Nadkarni VM, Hazinski ME. Part 16: education, implementation, and teams: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010;122(Suppl 3):S920-S933.
 15. Soar J MM, Bhanji F, Billi JE, Dennett J, Finn J, Ma MH, Perkins GD, Rodgers DL, Hazinski ME, Jacobs I, Morley PT. Part 12: education, implementation, and teams: 2010 international consensus on cardiopulmonary resuscitation and emergency cardiovascular care science with treatment recommendations. *Resuscitation*. 2010;81(1):e288-e332.
 16. Mancini ME SJ, Bhanji F, Billi JE, Dennett J, Finn J, Ma MH, Perkins GD, Rodgers DL, Hazinski ME, Jacobs I, Morley PT. Education, Implementation, and Teams Chapter Collaborators. Part 12: education, implementation, and teams: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation*. 2010;122(suppl 2):S539-S81.
 17. Oermann MH K-ES, Odom-Maryon T. Effects of monthly practice on nursing students' CPR psychomotor skill performance. *Resuscitation*. 2011;82(4):447-53.
 18. Ahn JY CG, Shon YD, Park SM, Kang KH. Effect of a reminder video using a mobile phone on the retention of CPR and AED skills in lay responders. *Resuscitation*. 2011;82(12):1543-7.
 19. Chandrasekaran S KS, Bhat SA, Saravanakumar, Shabbir PM, Chandrasekaran V. Awareness of basic life support among medical, dental, nursing students and doctors. *Indian journal of Anaesthesia*. 2010;54(2):121-6.
 20. Atkins DL BS, Duff JP, Gonzales JC, Hunt EA, Joyner BL, Meaney PA, Niles DE, Samson RA, Schexnayder SM. Part 11: pediatric basic life support and cardiopulmonary resuscitation quality: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(18 suppl 2):S519-S25.
 21. O'Connor RE AAA, Brady WJ, Ghaemmaghami CA, Menon V, Welsford M, Shuster M. Part 9: acute coronary syndromes: 2015 American Heart Association Guidelines Update for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2015;132(suppl 2):S483-S500.
 22. Markenson D FJ, Chameides L, Cassan P, Chung K-L, Epstein J, Gonzales L, Herrington RA, Pellegrino JL, Ratcliff N, Singer A. Part 17: first aid: 2010 American Heart Association and American Red Cross Guidelines for First Aid. *Circulation*. 2010;122(18 suppl 3):S934-S46.
 23. Khorasani M TM, Hamadzadeh H, Bakhshi M. Effect of Emergency Primary Care Training Workshops: A Survey on 45 Iranian Dental School Interns. *Journal of international oral health: JIOH*. 2015;7(Suppl 1):18-21.
 24. Zarif Sanaee N. Assessing the criteria for the quality and effectiveness of e-Learning in higher education. *Interdisciplinary Journal of Virtual Learning in Medical Sciences (IJVLMS)*. 2011;1(3):24-32.

