

Journal of Surgical Case Reports, 2018;5, 1

doi: 10.1093/jscr/rjy110 Corrigendum

CORRIGENDUM

Primary choriocarcinoma of the liver: a rare, but important differential diagnosis of liver lesions

Alexander Kohler^{1,*†}, Thilo Welsch^{1,†}, Anne-Kathrin Sturm^{2,†}, Gustavo B. Baretton^{2,†}, Christoph Reissfelder^{1,†}, Jürgen Weitz^{1,†}, and Carina Riediger^{1,†}

¹Department of Visceral, Thoracic and Vascular Surgery, University Hospital Carl Gustav Carus, TU Dresden, 01307 Dresden, Germany and ²Institute for Pathology, University Hospital Carl Gustav Carus, TU Dresden, 01307 Dresden, Germany

*Correspondence address. Department of Visceral, Thoracic and Vascular Surgery, University Hospital Carl Gustav Carus, TU Dresden, Fetscherstr. 74, 01307 Dresden, Germany. Tel: +49-351-4581-9401, Mobile +49-17683249848; Fax: +49-351-458-5375; E-mail: alexander.kohler@uniklinikum-dresden.de

Journal of Surgical Case Reports, 2018;4, 1–4. doi: 10.1093/jscr/rjy068

This article has been corrected to address the following error:

In the first published version, Figure 4 showed an extended right hemihepatectomy. The patient was treated with an extended left hemihepatectomy. Therefore Figure 4 has been replaced to reflect this and has been reproduced below.

The authors apologize for this error.




Figure 1: Liver diagram showing lesion (circle) and resection line (Segments I-VI were resected).

[†]All mentioned authors have contributed to this paper, with surgical, clinical and theoretical contributions.

Published by Oxford University Press and JSCR Publishing Ltd. All rights reserved. © The Author(s) 2018.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/ licenses/by-nc/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com