






# BMJ Open Interventions for physician prescribers of opioids for chronic non-cancer pain: protocol for an overview of systematic reviews

Erica Wennberg <sup>1,2</sup>, Sarah B Windle,<sup>1,3</sup> Kristian B Filion <sup>1,3,4</sup>, Genevieve Gore,<sup>5</sup> Brett D Thombs <sup>1,3,4,6,7,8,9</sup>, Irina Kudrina,<sup>10,11</sup> Elena Paraskevopoulos,<sup>12,13</sup> Marc O Martel,<sup>10,14</sup> Jonathan Kimmelman <sup>9</sup>, Sonia Johnson,<sup>15</sup> Andrew Taylor,<sup>15</sup> Mark J Eisenberg <sup>1,3,4,16</sup>

**To cite:** Wennberg E, Windle SB, Filion KB, *et al*. Interventions for physician prescribers of opioids for chronic non-cancer pain: protocol for an overview of systematic reviews. *BMJ Open* 2022;**12**:e060964. doi:10.1136/bmjopen-2022-060964

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-060964>).

Received 09 January 2022  
Accepted 02 March 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

## Correspondence to

Dr Mark J Eisenberg;  
[mark.eisenberg@ladydavis.ca](mailto:mark.eisenberg@ladydavis.ca)

## ABSTRACT

**Introduction** Interventions targeting behaviours of physician prescribers of opioids for chronic non-cancer pain have been introduced to combat the opioid crisis. Systematic reviews have evaluated effects of specific interventions (eg, prescriber education, prescription drug monitoring programmes) on patient and population health outcomes and prescriber behaviour. Integration of findings across intervention types is needed to better understand the effects of prescriber-targeted interventions.

**Methods and analysis** We will conduct an overview of systematic reviews. Eligible systematic reviews will include primary studies that evaluated any intervention targeting the behaviours of physician prescribers of opioids for chronic non-cancer pain in an outpatient or mixed setting, compared with no intervention, usual practice or another active or control intervention. Eligible outcomes will pertain to the intervention effect on patient and population health or opioid prescribing behaviour. We will search MEDLINE, Embase and PsycInfo via Ovid; the Cochrane Database of Systematic Reviews and Epistemonikos from inception. We will also hand search reference lists for additional publications. Screening and data extraction will be conducted independently by two reviewers, with disagreements resolved by consensus or consultation with a third reviewer. The risk of bias of included systematic reviews will be assessed in duplicate by two reviewers using the Risk of Bias in Systematic Reviews tool. Results will be synthesised narratively by intervention type and grouped by outcome. To assist with result interpretation, outcomes will be labelled as intended or unintended according to intervention objectives, and as positive, negative, evidence of no effect or inconclusive evidence according to effect on the population (for patient and population health outcomes) or intervention objectives (for prescriber outcomes).

**Ethics and dissemination** As the proposed study will use published data, ethics approval is not required. Dissemination of results will be achieved through publication of a manuscript in a peer-reviewed journal and conference presentations.

**PROSPERO registration number** CRD42020156815.

## Strengths and limitations of this study

- The overview of systematic review methodology will enable examination of the diverse body of evidence contained across systematic reviews of interventions targeting physician prescribers of opioids for chronic non-cancer pain.
- Design of the protocol was guided by Chapter V of the *Cochrane Handbook for Systematic Reviews of Interventions*, along with elements from additional guidance documents for overviews of systematic reviews.
- Limitations of this study relate to those of the overview of systematic review methodology; namely, restriction of the interventions and outcomes synthesised to those captured in available systematic reviews and risk of systematic reviews' conclusions being affected by publication bias.

## INTRODUCTION

To combat the ongoing opioid crisis in North America, countries and jurisdictions have introduced interventions targeting the behaviours of physician prescribers of opioids for chronic non-cancer pain (CNCP) (pain lasting over 3 months not associated with a cancer diagnosis<sup>1</sup>). A wide range of interventions fall under this category, including prescriber education, prescription drug monitoring programmes (PDMPs), pain clinic legislation (eg, laws requiring that physician pain clinic owners be board certified in pain management) and clinical guidelines.<sup>2</sup> As these interventions have the potential to alter the way in which opioids are prescribed, it is highly important to consider not only the effects of these interventions on prescriber behaviour but also on patient and population health. Numerous systematic reviews have evaluated the effects of interventions

targeting physician opioid prescribers for CNCP on opioid prescriber behaviours and outcomes among patients with CNCP and the general population.<sup>3–6</sup> These systematic reviews vary not only in their populations and outcomes of interest but also in the specific interventions evaluated (eg, PDMPs). While the variability in these reviews' areas of focus means a wealth of information is spread across them, it makes it difficult to consider their findings holistically. A systematic synthesis of this heterogeneous systematic review evidence has yet to be performed and would be of great value in better understanding the effect of prescriber-targeted interventions on both patient and population health and prescriber behaviour. Therefore, we will perform an overview of systematic reviews of the effect of interventions targeting the behaviours of physician opioid prescribers for CNCP in adults on patient and population health and prescriber behaviour.

## OBJECTIVE

Our objective is to synthesise the systematic review evidence on the effect of interventions targeting the behaviours of physician opioid prescribers for CNCP in adults on patient and population health and prescriber behaviour.

## METHODS AND ANALYSIS

This overview of systematic reviews will be guided by Chapter V of the *Cochrane Handbook for Systematic Reviews of Interventions*,<sup>7</sup> along with elements from additional guidance documents described in a recent review.<sup>8</sup> The overview of systematic review methodology was chosen to examine evidence across systematic reviews of interventions targeting physician prescribers of opioids for CNCP, as these systematic reviews address different outcomes.<sup>7</sup> Our overview will be reported according to the Preferred Reporting Items for Overviews of systematic reviews including harms pilot checklist.<sup>9</sup> It has been registered on PROSPERO. Important protocol amendments will be documented in PROSPERO.

### Eligibility criteria

#### Population

This overview will be restricted to systematic reviews of studies conducted in healthcare professionals who prescribe opioids, with a focus on physician opioid prescribers (table 1). For the purposes of this overview, 'physician opioid prescribers' will be defined as medical doctors who prescribe opioids. Eligible systematic reviews will include primary studies evaluating interventions targeted exclusively at physician opioid prescribers or targeted at multiple healthcare professional populations including physician opioid prescribers. Reviews of interventions targeted at multiple healthcare professional populations must include studies in which these interventions are delivered specifically or in part to physician opioid prescribers. Reviews limited to studies

of interventions delivered exclusively to non-physician healthcare professionals (eg, dentists, nurse practitioners, physician assistants, pharmacists) will be ineligible, as will reviews limited to studies of interventions delivered exclusively or in part to patients (eg, structured pain management programmes). Reviews that include some studies in eligible populations and some studies in ineligible populations will be included provided they report at least one outcome specific to an eligible population.

#### Intervention

We will include systematic reviews of any type of intervention(s) aimed at impacting opioid prescribing behaviour, with a focus on those aimed at impacting opioid prescribing behaviour for adult CNCP in an outpatient setting. Examples of eligible interventions include PDMPs, prescriber education (eg, online courses, workshops and telementoring programmes such as Project ECHO (Extension for Community Healthcare Outcomes)<sup>10</sup>), pain clinic legislation, clinical guidelines (eg, the 2017 Canadian Guideline for Opioids for Chronic Non-Cancer Pain<sup>11</sup>) evaluated as interventions and interventions relating to naloxone coprescription with opioids (eg, naloxone education for prescribers and naloxone coprescription requirements). Eligible systematic reviews will include primary studies of interventions targeted exclusively at impacting opioid prescribing behaviour for adult CNCP in an outpatient or mixed outpatient/inpatient setting or targeted at impacting prescribing behaviour for multiple opioid prescription indications including adult CNCP in an outpatient/mixed setting (eg, adult CNCP in addition to other pain indications or opioid use disorder). For interventions targeting multiple prescription indications, eligible reviews must include primary studies specific to opioid prescribing in the context of adult CNCP or studies in a mixed prescription indication context that includes adult CNCP. For interventions targeting a mixed prescription setting, eligible reviews will include primary studies in an exclusively outpatient setting or in a mixed setting. Reviews limited to studies of interventions exclusively targeting paediatric and non-CNCP prescription indications (eg, acute pain, postsurgical pain, opioid use disorder) or palliative pain management will be excluded, as will reviews limited to studies exclusively targeting prescribing in an inpatient setting. Interventions exclusively targeting opioid prescription for cancer pain will be excluded as opioid prescription guidelines and use patterns differ between chronic non-cancer and cancer pains. Interventions targeting opioid prescription within opioid treatment programmes will not be eligible. Reviews which include some studies of eligible interventions and some studies of ineligible interventions will be eligible provided they report at least one outcome specific to an eligible intervention or group of interventions. We will not restrict by intervention components or method of delivery.

**Table 1** Eligibility criteria for overview of systematic reviews of the effect of interventions targeting behaviours of physician prescribers of opioids for adult chronic non-cancer pain on prescriber behaviour and patient and population health

PICO element	Inclusion	Exclusion
Population	This overview will be restricted to systematic reviews of studies conducted in healthcare professionals who prescribe opioids, with a focus on physician opioid prescribers (medical doctors who prescribe opioids). Eligible systematic reviews will include primary studies evaluating interventions targeted exclusively at physician opioid prescribers or targeted at multiple healthcare professional populations including physician opioid prescribers. Reviews of interventions targeted at multiple healthcare professional populations must include studies in which these interventions are delivered specifically or in part to physician opioid prescribers. Reviews that include some studies in eligible populations and some studies in ineligible populations will be included provided they report at least one outcome specific to an eligible population.	Reviews limited to studies of interventions delivered exclusively to non-physician healthcare professionals (dentists, nurse practitioners, physician assistants, pharmacists, etc.) Reviews limited to studies of interventions delivered exclusively or in part to patients (eg, structured pain management programmes).
Intervention	We will include systematic reviews of any type of intervention(s) aimed at impacting opioid prescribing behaviour, with a focus on those aimed at impacting opioid prescribing behaviour for adult CNCP in an outpatient setting. Examples of eligible interventions include PDMPs, prescriber education (eg, online courses, workshops, and tele-mentoring programmes such as Project ECHO), pain clinic legislation, clinical guidelines (eg, the 2017 Canadian Guideline for Opioids for Chronic Non-Cancer Pain) evaluated as interventions and interventions relating to naloxone coprescription with opioids (eg, naloxone education for prescribers and naloxone coprescription requirements). Eligible systematic reviews will include primary studies of interventions targeted exclusively at impacting opioid prescribing behaviour for adult CNCP in an outpatient/mixed setting or targeted at impacting prescribing behaviour for multiple opioid prescription indications including adult CNCP in an outpatient/mixed setting (eg, adult CNCP in addition to other pain indications or opioid use disorder). For interventions targeting multiple prescription indications, eligible reviews must include primary studies specific to opioid prescribing in the context of adult CNCP or studies in a mixed prescription indication context that includes adult CNCP. For interventions targeting a mixed prescription setting, eligible reviews will include primary studies in an exclusively outpatient setting or in a mixed outpatient/inpatient setting. Reviews that include some studies of eligible interventions and some studies of ineligible interventions will be included provided they report at least one outcome specific to an eligible intervention or group of interventions.	Reviews limited to studies of interventions not aimed at impacting opioid prescribing behaviour. Reviews limited to studies exclusively targeting non-adult CNCP prescription indications (eg, acute pain, post-surgical pain, cancer pain, paediatric CNCP, opioid use disorder) or palliative pain management. Reviews limited to studies exclusively targeting prescribing in an inpatient setting. Reviews that do not report any outcomes specific to an eligible intervention or group of interventions.
Comparators	Eligible systematic reviews may include one or both of the following types of primary studies: a. Comparative studies that evaluated intervention effect against no intervention, usual care procedures or other active (eg, prescriber education vs clinical guideline implementation) or control (eg, attention control) interventions b. Non-comparative studies (eg, time series without comparator).	

Continued

Table 1 Continued

PICO element	Inclusion	Exclusion
Outcomes	<p>Eligible systematic reviews will report at least one outcome pertaining to intervention effect on patient and population health or opioid prescribing behaviour.</p> <p>Eligible patient and population health outcomes will include:</p> <ol style="list-style-type: none"> <li>Changes in patient-reported health and pain outcomes (eg, changes in patient-reported physical functioning, quality of life and pain outcomes, including both measures of pain intensity/severity and pain interference with functioning).</li> <li>Changes in pharmaceutical or non-pharmaceutical opioid (eg, heroin)-related morbidity and mortality (eg, changes in prevalence or incidence of fatal and non-fatal opioid overdose, opioid-related hospitalisations and opioid-related emergency department visits, overall or by specific drug; changes in incidence of opioid abuse treatment initiation or inpatient admissions for opioid abuse treatment).</li> <li>Changes in prevalence or incidence of self-reported non-medical prescription opioid use or non-pharmaceutical opioid use.</li> </ol> <p>Eligible opioid prescribing behaviour outcomes will include:</p> <ol style="list-style-type: none"> <li>Changes in opioid prescribing practices (eg, changes in incidence or prevalence of opioid prescriptions, overall, by specific drug or by release type (eg, short-acting vs long-acting/extended release); changes in average duration or dosage of individual opioid prescriptions; changes in coprescription of naloxone with opioids (eg, changes in incidence or number of naloxone prescriptions); changes in number of overlapping opioid and benzodiazepine prescriptions (eg, changes in number of patients with benzodiazepine and opioid prescriptions overlapping by at least one common day)).</li> <li>Changes in rates of prescribing of and referrals to alternative pain management therapies (eg, changes in number of non-opioid analgesic prescriptions, changes in number of referrals to physical therapy).</li> <li>Changes in intervention adherence, where these constitute a measure of intervention effect and a change in prescribing behaviour (eg, changes in prescriber adherence to CNCP opioid prescribing guideline recommendations following an educational intervention designed to improve prescriber adherence to said recommendations).</li> </ol>	<p>Systematic reviews that exclusively report outcomes not related to intervention effect on patient and population health or opioid prescribing behaviour, for example,</p> <ul style="list-style-type: none"> <li>Feasibility</li> <li>Acceptability (including healthcare professional and public perceptions of and attitudes towards interventions)</li> <li>Cost-effectiveness</li> <li>Intervention adherence (where this does not constitute a measure of intervention effect)</li> </ul>
Study design	<p>Systematic reviews with or without meta-analysis. Reviews must meet the following criteria to be considered systematic:</p> <ol style="list-style-type: none"> <li>Methods are described, including a systematic search with inclusion/exclusion criteria.</li> <li>Formal risk of bias assessment of included studies was performed (eg, using the Cochrane Risk of Bias tool), with individual results reported for each study and item/domain of the tool.</li> </ol> <p>We will include systematic reviews with or without meta-analysis. Data may be derived from any primary study type (eg, experimental or observational) conducted in humans.</p>	<p>Any review or study that does not meet the criteria of a systematic review, including:</p> <ul style="list-style-type: none"> <li>Overviews of systematic reviews</li> <li>Non-systematic reviews</li> <li>Primary studies</li> <li>Commentaries</li> </ul>
Forms of publication	<p>Language: English*</p> <p>Systematic review abstracts and conference proceedings will be eligible provided they meet the aforementioned systematic review criteria and include sufficient detail to enable extraction of risk of bias assessments per study and tool domain/item.</p> <p>*English-language abstracts of non-English language publications will not be eligible for inclusion, as records will be assessed for eligibility on the basis of the most complete version of the publication.</p>	<p>Non-English language publications</p>

CNCP, chronic non-cancer pain; ECHO, Extension for Community Healthcare Outcomes; PDMP, prescription drug monitoring programmes.

### Comparators

Eligible systematic reviews may include one or both of the following types of primary studies: (a) comparative studies that compared the intervention of interest against no intervention, usual care procedures or other active (eg, prescriber education vs clinical guideline implementation) or control (eg, attention control) interventions; or (b) non-comparative studies (eg, time series without comparator).

### Outcomes

Eligible systematic reviews will report outcomes pertaining to intervention effect on patient and population health or opioid prescribing behaviour. Systematic reviews of intervention feasibility, acceptability (including health-care professional and public perceptions of and attitudes towards interventions) and cost-effectiveness will be excluded.

Eligible patient and population health outcomes will include:

1. Changes in patient-reported health and pain outcomes (eg, changes in patient-reported physical functioning, quality of life and pain outcomes, including both measures of pain intensity/severity and pain interference with functioning). These outcomes have been identified as core outcome domains among patients with chronic pain.<sup>12</sup>
2. Changes in pharmaceutical or non-pharmaceutical opioid (eg, heroin)-related morbidity and mortality (eg, changes in prevalence or incidence of fatal and non-fatal opioid overdose, opioid-related hospitalisations and opioid-related emergency department visits, overall or by specific drug; changes in incidence of opioid abuse treatment initiation or inpatient admissions for opioid abuse treatment).
3. Changes in prevalence or incidence of self-reported non-medical prescription opioid use or non-pharmaceutical opioid use.

Eligible opioid prescribing behaviour outcomes will include:

1. Changes in opioid prescribing practices (eg, changes in incidence or prevalence of opioid prescriptions, overall, by specific drug, or by release type (eg, short-acting vs long-acting/extended release); changes in average duration or dosage of individual opioid prescriptions; changes in coprescription of naloxone with opioids (eg, changes in incidence or number of naloxone prescriptions); changes in number of overlapping opioid and benzodiazepine prescriptions (eg, changes in number of patients with benzodiazepine and opioid prescriptions overlapping by at least 1 common day)).
2. Changes in rates of prescribing of and referrals to alternative pain management therapies (eg, changes in number of non-opioid analgesic prescriptions, changes in number of referrals to physical therapy).
3. Changes in intervention adherence, where these constitute a measure of intervention effect and a change in prescribing behaviour (eg, changes in prescriber

adherence to CNCP opioid prescribing guideline recommendations following an educational intervention designed to improve prescriber adherence to said recommendations).

### Design

Inclusion will be restricted to systematic reviews with or without meta-analysis. The following criteria will be used to define eligibility as a systematic review: (1) methods are described, including a systematic search with inclusion/exclusion criteria and (2) formal risk of bias assessment of included studies was performed (eg, using the Cochrane Risk of Bias tool), with individual results reported for each study and each item/domain of the tool. We will include systematic reviews with and without meta-analysis. Data may be derived from any primary study type (eg, randomised controlled trials or non-randomised studies of interventions) conducted in humans.

### Forms of publication

Studies will be restricted to English-language publications. Systematic review abstracts and conference proceedings will be included provided they meet the aforementioned systematic review criteria and contain sufficient detail to enable extraction of risk of bias assessments by study and tool domain/item. English-language abstracts of non-English language publications will not be eligible for inclusion, as records will be assessed for eligibility on the basis of the most complete version of the publication.

### Data sources

We will search the following databases from inception: MEDLINE, Embase and PsycInfo via Ovid; the Cochrane Database of Systematic Reviews and Epistemonikos. Reference lists of included publications will be hand searched for eligible publications not identified in the search. We will not conduct an additional search for primary studies. If eligible systematic reviews are available only in protocol form, we will contact the authors to inquire whether a prepublication version of the manuscript is available.

### Search strategy

The search was designed and will be executed by an experienced health sciences librarian (GG). Prior to execution, it will be peer reviewed using Peer Review of Electronic Search Strategies.<sup>13</sup> The search is tailored to each database and includes a combination of subject headings and terms related to opioids and prescribers, as applicable. We will apply a librarian-modified version of the PubMed systematic review filter, which includes additional search terms from the Canadian Agency for Drugs and Technologies in Health systematic review filter. Preliminary search strategies for all five databases are presented in [tables 2–6](#).

### Study selection

Search results from each database will be downloaded into EndNote and subsequently imported into DistillerSR (Evidence Partners, Ottawa, Canada). Duplicates

**Table 2** Search strategy (MEDLINE via Ovid)

Search number	Description
1	exp analgesics, opioid/ or exp opioid-related disorders/ or (narcotic* or opiate* or opioid* or acetylmethadol or alfentanil or anileridine or Belladonna or Benzomorphan* or bezitramide or buprenorphine or butorphanol or Codeine or Dextromethorphan or Dextromoramide or Dextropropoxyphene or dezocine or Diamorphine or dihydrocodeine or Diphenylpropylamine or Ethylmorphine or Fentanyl* or Heroin or Hydrocodon* or Hydromorphon* or ketobemidone or levacetylmethadol or Meperidine or Meptazinol or methadone or Morphan* or Morphine* or nalbuphine or nicomorphine or normethadone or Opium or Oripavine or Oxycodone or Oxymorphone or Papaveretum or Pentazocine or pethidin* or Phenazocine or Phenoperidine or phentanyl or Phenylpiperidine or Piritramide or remifentanil or Sufentanil or sulfentanil or sulfentanyl or tapentadol or Tilidine or Tramadol*).mp. or (analgesic*).ti.
2	practice patterns, physicians'/ or exp prescriptions/ or exp prescription drug monitoring programs/ or (doctor* or physician* or surgeon* or dispens* or prescribe* or prescribing or deprescrib* or overprescri* or prescription* or script? or stewardship* or refill* or taper*).mp.
3	1 and 2
4	systematic review/ or meta analysis/ or "systematic review as topic"/ or exp "meta-analysis as topic"/ or technology assessment, biomedical/
5	(meta analy* or metaanaly* or technology assessment* or hta or htas or ((evidence or mixed method* or rapid or systematic) adj3 (overview or review or metareview or metasynthesis))).ti. or (cochrane database of systematic reviews or technology assessment*).jw.
6	4 or 5
7	3 and 6

will be identified and removed in DistillerSR. Screening will proceed through a three-stage process in DistillerSR. Two reviewers will first independently screen the titles of identified citations for eligibility. Citations considered potentially eligible by either reviewer in the title stage will move on to abstract screening. Two reviewers will then independently screen the abstracts of potentially eligible citations. Citations considered potentially eligible by one or both reviewers in the abstract stage will be retrieved in full text, and the full text will then be reviewed for eligibility independently by two reviewers. Disagreements after full-text review will be resolved by consensus or consultation with a third reviewer, as necessary. The publications remaining after full-text review will be included in the overview of reviews. Publications excluded during the full-text review will be presented in the final manuscript in a table that includes the rationale for exclusion.

Overlap in primary studies is expected among eligible reviews addressing the same research question. We will address overlap between eligible reviews in a series of steps, beginning with creation of citation matrices to identify systematic reviews with complete overlap.<sup>14</sup> Separate citation matrices will be created for each intervention

**Table 3** Search strategy (Embase via Ovid)

Search number	Description
1	exp narcotic analgesic agent/
2	controlled substance/
3	(narcotic* or opiate* or opioid* or acetylmethadol or alfentanil or anileridine or Belladonna or Benzomorphan* or bezitramide or buprenorphine or butorphanol or Codeine or Dextromethorphan or Dextromoramide or Dextropropoxyphene or dezocine or Diamorphine or dihydrocodeine or Diphenylpropylamine or Ethylmorphine or Fentanyl* or Heroin or Hydrocodon* or Hydromorphon* or ketobemidone or levacetylmethadol or Meperidine or Meptazinol or methadone or Morphan* or Morphine* or nalbuphine or nicomorphine or normethadone or Opium or Oripavine or Oxycodone or Oxymorphone or Papaveretum or Pentazocine or pethidin* or Phenazocine or Phenoperidine or phentanyl or Phenylpiperidine or Piritramide or remifentanil or Sufentanil or sulfentanil or sulfentanyl or tapentadol or Tilidine or Tramadol*).mp. or analgesic*.ti.
4	1 or 2 or 3
5	prescription/ or prescription drug monitoring program/ or (doctor* or physician* or surgeon* or dispens* or prescribe* or prescribing or deprescrib* or overprescri* or prescription* or script? or stewardship* or refill* or taper*).mp.
6	4 and 5
7	systematic review/ or exp meta analysis/ or "systematic review (topic)"/ or "meta analysis (topic)"/ or biomedical technology assessment/
8	(meta analy* or metaanaly* or technology assessment* or hta or htas or ((evidence or mixed method* or rapid or systematic) adj3 (review or metareview or metasynthesis))).ti.
9	(cochrane database of systematic review or technology assessment*).jw.
10	7 or 8 or 9
11	6 and 10

type (eg, PDMPs) to avoid underestimation of the degree of overlap, as some systematic reviews may include more than one intervention type. Complete overlap will be defined as two reviews that include all the same citations, or one review that includes all the citations of another. Each member of a pair of reviews with complete overlap will be assessed for exclusion based on meeting one of the following conditions: (a) reports on no unique outcome area(s), contains no unique citations and is at higher risk of bias compared with the other review or (b) reports on no unique outcome area(s), contains no unique citations, is at similar or higher risk of bias and is less recent compared with the other review (eg, a systematic review that has been updated).<sup>15 16</sup> These decisions will be made by two reviewers and will be tracked in a table that presents the characteristics of excluded reviews. In all other cases, reviews with complete overlap will be included.

**Table 4** Search strategy (PsycINFO via Ovid)

Search number	Description
1	exp narcotic drugs/ or (narcotic* or opiate* or opioid* or acetylmethadol or alfentanil or anileridine or Belladonna or Benzomorphan* or bezitramide or buprenorphine or butorphanol or Codeine or Dextromethorphan or Dextromoramide or Dextropropoxyphene or dezocine or Diamorphine or dihydrocodeine or Diphenylpropylamine or Ethylmorphine or Fentanyl* or Heroin or Hydrocodon* or Hydromorphon* or ketobemidone or levacetylmethadol or Meperidine or Meptazinol or methadone or Morphan* or Morphine* or nalbuphine or nicomorphine or normethadone or Opium or Oripavine or Oxycodone or Oxymorphone or Papaveretum or Pentazocine or pethidin* or Phenazocine or Phenoperidine or phentanyl or Phenylpiperidine or Piritramide or remifentanil or Sufentanil or sulfentanil or sulfentanyl or tapentadol or Tilidine or Tramadol*).mp.
2	exp "prescribing (drugs)"/ or prescription drugs/ or (doctor* or physician* or surgeon* or dispens* or overprescri* or prescribe* or prescribing or deprescrib* or prescription* or script? or stewardship* or refill* or taper*).mp.
3	1 and 2
4	meta analysis/
5	(systematic review or meta analysis or metasyntesis).md.
6	(meta analy* or metaanaly* or technology assessment* or hta or htas or ((evidence or mixed method* or rapid or systematic) adj3 (review or metareview or metasyntesis)).ti.
7	or/4-6
8	3 and 7

**Table 5** Search Strategy (Cochrane Database of Systematic Reviews)

Search number	Description
1 Title Abstract Keyword	narcotic* or opiate* or opioid* or acetylmethadol or alfentanil or anileridine or Belladonna or Benzomorphan* or bezitramide or buprenorphine or butorphanol or Codeine or Dextromethorphan or Dextromoramide or Dextropropoxyphene or dezocine or Diamorphine or dihydrocodeine or Diphenylpropylamine or Ethylmorphine or Fentanyl* or Heroin or Hydrocodon* or Hydromorphon* or ketobemidone or levacetylmethadol or Meperidine or Meptazinol or methadone or Morphan* or Morphine* or nalbuphine or nicomorphine or normethadone or Opium or Oripavine or Oxycodone or Oxymorphone or Papaveretum or Pentazocine or pethidin* or Phenazocine or Phenoperidine or phentanyl or Phenylpiperidine or Piritramide or remifentanil or Sufentanil or sulfentanil or sulfentanyl or tapentadol or Tilidine or Tramadol*
2 Title Abstract Keyword	doctor* or physician* or surgeon* or dispens* or prescribe* or prescribing or deprescrib* or overprescri* or prescription* or script* or stewardship* or refill* or taper*
3	1 and 2
Search limits	Cochrane Reviews Cochrane Protocols

**Table 6** Search strategy (Epistemonikos)

Search number	Description
1 (Title/ Abstract)	narcotic* OR opiate* OR opioid* OR acetylmethadol OR alfentanil OR anileridine OR Belladonna OR Benzomorphan* OR bezitramide OR buprenorphine OR butorphanol OR Codeine OR Dextromethorphan OR Dextromoramide OR Dextropropoxyphene OR dezocine OR Diamorphine OR dihydrocodeine OR Diphenylpropylamine OR Ethylmorphine OR Fentanyl* OR Heroin OR Hydrocodon* OR Hydromorphon* OR ketobemidone OR levacetylmethadol OR Meperidine OR Meptazinol OR methadone OR Morphan* OR Morphine* OR nalbuphine OR nicomorphine OR normethadone OR Opium OR Oripavine OR Oxycodone OR Oxymorphone OR Papaveretum OR Pentazocine OR pethidin* OR Phenazocine OR Phenoperidine OR phentanyl OR Phenylpiperidine OR Piritramide OR remifentanil OR Sufentanil OR sulfentanil OR sulfentanyl OR tapentadol OR Tilidine OR Tramadol*
2 (Title/ Abstract)	doctor* OR physician* OR surgeon* OR dispens* OR prescribe* OR prescribing OR deprescrib* OR overprescri* OR prescription* OR script* OR stewardship* OR refill* OR taper*
3	1 and 2
Filters	Publication type: Systematic Review Systematic Review Question: Interventions

**Data extraction**

Data will be extracted independently by two reviewers using pilot-tested forms in DistillerSR. The pilot-testing process will be carried out by two reviewers with a small sample of studies to identify necessary adjustments to the extraction forms and to assess the feasibility of conducting independent extraction. When large amounts of non-numerical data are independently extracted into DistillerSR, it can result in high numbers of conflicts from slight wording differences, resulting in reduced efficiency of the conflict resolution process. If the pilot testing process reveals that independent extraction will be inadvisable for this reason, extraction will instead proceed via initial extraction by a first reviewer and subsequent validation by a second reviewer using the DistillerSR Quality Control function. Otherwise, extraction will proceed independently and disagreements between the two reviewers will be detected in DistillerSR. In either case, disagreements will be resolved by consensus or a third reviewer as necessary.

We will extract the following data on systematic review characteristics: first author, publication year, search period, number of databases searched and names, objectives, inclusion criteria (population, intervention, comparators, outcomes, study design), exclusion criteria, number of included primary studies, total number of participants, risk of bias tool used and source of funding. The number of included primary studies and total number of participants will be extracted by intervention and by outcome. For reviews which report on both eligible and non-eligible interventions or report both

eligible and non-eligible outcomes, we will only extract the number of included primary studies and total number of participants relevant to the eligible intervention(s)/outcome(s). We will also extract the following data on the characteristics of systematic reviews' included primary studies: first author, publication year and risk of bias (as assessed by the systematic review). Primary study characteristics will only be extracted for those studies relevant to our review. Finally, we will extract outcomes pertaining to intervention effect on prescriber behaviour and patient and population health. Outcome data will be extracted as they are presented in the systematic review, including effect estimates, 95% CIs, descriptive statistics (eg, count data, means) and measures of heterogeneity. Both study-level and meta-analytic results will be extracted. We will additionally extract the number of primary studies the results are drawn from and evidence-grade assessments (as available). We will also extract outcome data stratified by sex; gender; ethnicity; Indigenous identity and efficacy, effectiveness and efficiency study design (as available). Where data are missing or confirmation is needed, review authors will be contacted.

#### **Risk of bias assessment of included systematic reviews**

Two reviewers will independently assess the risk of bias of included systematic reviews using the Risk of Bias in Systematic Reviews (ROBIS) tool.<sup>17</sup> ROBIS assesses concerns about bias in the review process in four domains: study eligibility criteria, identification and selection of studies, data collection and study appraisal and synthesis and findings. Each domain includes 5–6 signalling questions to aid in the assessment, leading to a final rating of high, low or unclear concern in each domain. Questions are answered as yes, probably yes, probably no, no or no information. Answers of yes or probably yes to all signalling questions will result in a judgement of low concern for that domain. Answers of yes, probably yes and no information will result in a judgement of unclear concern. Any answer of no or probably no will result in a judgement of high concern. Final assessments in each domain will be used in the assessment of risk of bias in the review, which is determined based on three signalling questions: (1) Did the interpretation of findings address all of the concerns identified in domains 1 to 4; (2) Was the relevance of identified studies to the review's research question appropriately considered and (3) Did the reviewers avoid emphasising results on the basis of their statistical significance. These signalling questions will be answered and interpreted in the same manner as for the individual domains, leading to a judgement of low, high or unclear risk of bias in the review. We will not exclude any systematic reviews on the basis of risk of bias results.

#### **Risk of bias of primary studies contained in included systematic reviews**

We will extract risk of bias assessments performed by included systematic reviews and present them in tabular form. These tables will be grouped by primary study and

will include the systematic review of origin, the tool used and the assessment results. Domain-specific and overall ratings will be extracted. Some primary studies may have more than one risk of bias assessment available due to inclusion in more than one systematic review. For these studies, we will extract and present all available risk of bias assessments.

#### **Data synthesis**

We will use a qualitative, analytical approach to synthesise the evidence. We will create five types of summary tables; one to present characteristics of included systematic reviews, one to present primary study risk of bias assessments performed by included systematic reviews (outlined in the above section), one to present characteristics of interventions investigated by included systematic reviews, one to present ROBIS risk of bias assessments for each systematic review and one to present their results. The table presenting characteristics of included systematic reviews will include first author, publication year, search period, number of databases searched and names, objectives, focus (population, intervention, comparators, outcomes, study design), number of relevant included primary studies and total number of participants (separated by intervention or outcome as applicable), risk of bias tool used and source of funding. The table presenting characteristics of investigated interventions will include interventions' target population(s), target prescription indication(s), target prescription setting(s), major components, objectives and country or jurisdiction of origin. The table presenting ROBIS risk of bias assessments for each systematic review will include scores in each domain (low/high/unclear) and the risk of bias in the review (low/high/unclear). The tables presenting results of included systematic reviews will be grouped by outcome and will include relevant outcome data from each systematic review, the number of included systematic reviews assessing the outcome, the number of primary studies and study participants represented and evidence grade assessments from each systematic review (as available). Separate tables will be created for each intervention type (eg, PDMPs, clinical guidelines) and country of origin as needed (eg, Canadian vs American clinical guidelines), as opioid prescription guidelines and legislation vary by country. When patient and population health outcomes are available for an intervention, these will be made the priority of our synthesis and conclusions to reflect their higher importance compared with prescriber behaviour outcomes in determining best practices.

To assist in the interpretation of our results, we will label outcomes relating to intervention effect as (a) intended or unintended and (b) positive, negative, evidence of no effect or inconclusive evidence. Labelling will be conducted in duplicate by two reviewers, with disagreements resolved via consensus or consultation with a third reviewer as necessary. Labelling outcomes as intended and unintended will enable separation of the intended effects of investigated interventions on a given



population from their potential unintended effects.<sup>18</sup> The categorisation of an outcome as intended or unintended will be determined according to the objectives of the intervention in question, as defined by included publications and summarised in our table of intervention characteristics. Outcomes that align with the objectives of an intervention (ie, planned effects) will be categorised as intended outcomes, and outcomes which do not align with the objectives of an intervention (ie, unplanned effects) will be categorised as unintended outcomes. Labelling outcomes as positive, negative, evidence of no effect and inconclusive evidence will enable identification of the effects of each investigated intervention, including potential benefits and harms in the case of patient and population health outcomes. For outcomes related to an intervention's objectives, categorisation as positive or negative will be determined according to their alignment with intervention objectives. A decrease in overall opioid prescribing rates following the implementation of an intervention designed to reduce opioid prescribing, for example, would be categorised as a positive effect, while an increase in these rates would be categorised as a negative effect. For outcomes unrelated to an intervention's objectives, categorisation as positive or negative will be determined according to the effect they represent on the associated population. For example, an increase in rates of opioid overdose in the general population following the implementation of an intervention would be categorised as a negative effect, while a decrease in these rates would be categorised as a positive effect. Outcomes for which an effect is not demonstrated will be categorised as evidence of no effect if this conclusion is supported by precise estimates that rule out clinically important differences, and inconclusive evidence if insufficient evidence is available to judge whether an effect is present.

### Addressing overlap between included systematic reviews

To address overlap between included systematic reviews, citation matrices that were created for each intervention type in the screening stage will be updated to reflect final inclusion/exclusion decisions. They will then be used to calculate corrected covered area (CCA) scores by intervention type using the following formula<sup>19</sup>:

$$CCA = \frac{N-r}{(r \times c) - r},$$

where N is the total number of primary studies across all reviews (including duplicates), r is the number of unique primary studies across all reviews and c is the number of reviews. The CCA score ranges from 0% to 100%, with a higher CCA score reflecting a higher degree of overlap. Citation matrices will also be created, and CCA scores calculated, within intervention types by outcome category (eg, patient-reported health and pain outcomes).<sup>14</sup> CCA scores for each intervention type overall and by outcome category will be reported in our results tables and taken into account in our synthesis. When CCA scores are high (>15)<sup>19</sup> and findings between reviews are discrepant, reasons for discrepancy will be explored (eg, differences

in methodology, exclusions of studies from meta-analyses) and the findings of reviews that are of lower risk of bias and are more comprehensive will be focused on in our synthesis. When CCA scores are high between reviews and findings are concordant, the probable role of overlap will be noted in our synthesis to reduce the risk of biasing our results.

### Patient and public involvement

This protocol was developed in collaboration with two employees of Health Canada (SJ and AT). They will be involved throughout the systematic review and in dissemination of our findings.

### ETHICS AND DISSEMINATION

As the planned project is an overview of systematic reviews of published data, there are no ethical or safety concerns. Dissemination plans include publication of our results in a peer-reviewed journal and presentation at conferences. We will additionally curate our results for dissemination to non-scientific audiences.

### Author affiliations

- <sup>1</sup>Lady Davis Institute for Medical Research, Montreal, Québec, Canada
- <sup>2</sup>Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada
- <sup>3</sup>Department of Epidemiology, Biostatistics and Occupational Health, McGill University, Montreal, Québec, Canada
- <sup>4</sup>Department of Medicine, McGill University, Montreal, Québec, Canada
- <sup>5</sup>Schulich Library of Science and Engineering, McGill University, Montreal, Québec, Canada
- <sup>6</sup>Department of Psychiatry, McGill University, Montreal, Québec, Canada
- <sup>7</sup>Department of Psychology, McGill University, Montreal, Québec, Canada
- <sup>8</sup>Department of Counselling and Educational Psychology, McGill University, Montreal, Québec, Canada
- <sup>9</sup>Biomedical Ethics Unit, McGill University, Montreal, Québec, Canada
- <sup>10</sup>Department of Anesthesia, McGill University, Montreal, Québec, Canada
- <sup>11</sup>Department of Family Medicine, McGill University, Montreal, Québec, Canada
- <sup>12</sup>Department of Family Medicine, Royal Ottawa Mental Health Centre, Ottawa, Ontario, Canada
- <sup>13</sup>Department of Family Medicine, Queensway Carleton Hospital, Ottawa, Ontario, Canada
- <sup>14</sup>Faculty of Dentistry, McGill University, Montreal, Québec, Canada
- <sup>15</sup>Health Canada, Ottawa, Ontario, Canada
- <sup>16</sup>Division of Cardiology, Sir Mortimer B Davis Jewish General Hospital, Montreal, Québec, Canada

**Contributors** EW and SBW contributed to protocol design and drafted the protocol manuscript. MJE, KBF, GG, BDT, IK, EP, MOM, JK, SJ, and AT contributed to protocol design and revisions, drawing upon their respective areas of expertise. All authors approved the final version of the manuscript.

**Funding** This work was supported by the Canadian Institutes of Health Research (number EOC-162067). SBW is supported by a Canada Graduate Scholarship in Honour of Nelson Mandela from the Canadian Institutes of Health Research and a Doctoral Training Award from the Fonds de recherche Québec—Santé. KBF is supported by a Senior support award from the Fonds de recherche du Québec—Santé and a William Dawson Scholar award from McGill University. IK is supported by LE-250 (from the Quebec Ministry of Health and Social Services - Fonds de recherche du Québec—Santé) award.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not applicable.



**Provenance and peer review** Not commissioned; externally peer reviewed.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

#### ORCID iDs

Erica Wennberg <http://orcid.org/0000-0002-8069-6778>

Kristian B Filion <http://orcid.org/0000-0001-6055-0088>

Brett D Thombs <http://orcid.org/0000-0002-5644-8432>

Jonathan Kimmelman <http://orcid.org/0000-0003-1614-6779>

Mark J Eisenberg <http://orcid.org/0000-0002-1296-0661>

#### REFERENCES

- 1 Task Force on Taxonomy of the International Association for the Study of Pain. *Classification of chronic pain, second edition (revised)*. Seattle (WA): international association for the study of pain, 1994.
- 2 Haegerich TM, Jones CM, Cote P-O, et al. Evidence for state, community and systems-level prevention strategies to address the opioid crisis. *Drug Alcohol Depend* 2019;204:107563.
- 3 Mathieson S, Maher CG, Ferreira GE, et al. Deprescribing opioids in chronic non-cancer pain: systematic review of randomised trials. *Drugs* 2020;80:1563–76.
- 4 Fink DS, Schleimer JP, Sarvet A, et al. Association between prescription drug monitoring programs and nonfatal and fatal drug overdoses: a systematic review. *Ann Intern Med* 2018;168:783–90.
- 5 Shoemaker-Hunt SJ, Wyant BE. The effect of opioid stewardship interventions on key outcomes: a systematic review. *J Patient Saf* 2020;16:S36–41.
- 6 Rhodes E, Wilson M, Robinson A, et al. The effectiveness of prescription drug monitoring programs at reducing opioid-related harms and consequences: a systematic review. *BMC Health Serv Res* 2019;19:784.
- 7 Pollock M, Fernandes RM, Becker LA, et al. Chapter V: Overviews of Reviews. In: Higgins JPT, Thomas J, Chandler J, et al, eds. *Cochrane Handbook for systematic reviews of interventions version 6.0*. Cochrane, 2020.
- 8 Pollock M, Fernandes RM, Becker LA, et al. What guidance is available for researchers conducting overviews of reviews of healthcare interventions? A scoping review and qualitative metasummary. *Syst Rev* 2016;5:190.
- 9 Bougioukas KI, Liakos A, Tsapas A, et al. Preferred reporting items for overviews of systematic reviews including harms checklist: a pilot tool to be used for balanced reporting of benefits and harms. *J Clin Epidemiol* 2018;93:9–24.
- 10 Komaromy M, Duhigg D, Metcalf A, et al. Project echo (extension for community healthcare outcomes): a new model for educating primary care providers about treatment of substance use disorders. *Subst Abuse* 2016;37:20–4.
- 11 Busse JW, Craigie S, Juurlink DN, et al. Guideline for opioid therapy and chronic noncancer pain. *CMAJ* 2017;189:E659–66.
- 12 Dworkin RH, Turk DC, Farrar JT, et al. Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain* 2005;113:9–19.
- 13 McGowan J, Sampson M, Salzwedel DM, et al. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016;75:40–6.
- 14 Hennessy EA, Johnson BT. Examining overlap of included studies in meta-reviews: guidance for using the corrected covered area index. *Res Synth Methods* 2020;11:134–45.
- 15 Pollock A, Farmer SE, Brady MC, et al. Interventions for improving upper limb function after stroke. *Cochrane Database Syst Rev* 2014;23.
- 16 Hersi M, Traversy G, Thombs BD, et al. Effectiveness of stop smoking interventions among adults: protocol for an overview of systematic reviews and an updated systematic review. *Syst Rev* 2019;8:28.
- 17 Whiting P, Savović J, Higgins JPT, et al. ROBIS: a new tool to assess risk of bias in systematic reviews was developed. *J Clin Epidemiol* 2016;69:225–34.
- 18 Sherrill S. Identifying and measuring unintended outcomes. *Eval Program Plann* 1984;7:27–34.
- 19 Pieper D, Antoine S-L, Mathes T, et al. Systematic review finds overlapping reviews were not mentioned in every other overview. *J Clin Epidemiol* 2014;67:368–75.