

RESEARCH ARTICLE

Open Access



Estrogen and soy isoflavonoids decrease sensitivity of medulloblastoma and central nervous system primitive neuroectodermal tumor cells to chemotherapeutic cytotoxicity

Scott M. Belcher^{1,2,3,4*}, Caleb C. Burton³, Clifford J. Cookman⁴, Michelle Kirby², Gabriel L. Miranda³, Fatima O. Saeed⁵ and Kathleen E. Wray³

Abstract

Background: Our previous studies demonstrated that growth and migration of medulloblastoma (MB), the most common malignant brain tumor in children, are stimulated by 17 β -estradiol. The growth stimulating effects of estrogens are mediated through ER β and insulin-like growth factor 1 signaling to inhibit caspase 3 activity and reduce tumor cell apoptosis. The objective of this study was to determine whether estrogens decreased sensitivity of MB cells to cytotoxic actions of chemotherapeutic drugs.

Methods: Using in vitro cell viability and clonogenic survival assays, concentration response analysis was used to determine whether the cytoprotective effects of estradiol protected human D283 Med MB cells from the cytotoxic actions of the MB chemotherapeutic drugs cisplatin, vincristine, or lomustine. Additional experiments were done to determine whether the ER antagonist fulvestrant or the selective ER modulator tamoxifen blocked the cytoprotective actions of estradiol. ER-selective agonists and antagonists were used to define receptor specificity, and the impacts of the soy-derived phytoestrogens genistein, daidzein, and s-equol on chemosensitivity were evaluated.

Results: In D283 Med cells the presence of 10 nM estradiol increased the IC₅₀ for cisplatin-induced inhibition of viability 2-fold from ~5 μ M to >10 μ M. In clonogenic survival assays estradiol decreased the chemosensitivity of D283 Med cells exposed to cisplatin, lomustine and vincristine. The ER β selective agonist DPN and low physiological concentrations of the soy-derived phytoestrogens genistein, daidzein, and s-equol also decreased sensitivity of D283 Med cells to cisplatin. The protective effects of estradiol were blocked by the antiestrogens 4-hydroxytamoxifen, fulvestrant (ICI 182,780) and the ER β selective antagonist PPHTP. Whereas estradiol also decreased chemosensitivity of PFSK-1 cells, estradiol increased sensitivity of Daoy cell to cisplatin, suggesting that ER β mediated effects may vary in different MB celltypes.

(Continued on next page)

* Correspondence: smbch2@ncsu.edu

¹Department of Biological Science and Center for Human Health and the Environment, North Carolina State University, Raleigh, NC, USA

²Department of Pharmacology and Cell Biophysics, University of Cincinnati, Cincinnati, OH, USA

Full list of author information is available at the end of the article



(Continued from previous page)

Conclusions: These findings demonstrate that E2 and environmental estrogens decrease sensitivity of MB to cytotoxic chemotherapeutics, and that ER β selective and non-selective inhibition of estrogen receptor activity blocks these cytoprotective actions. These findings support the therapeutic potential of antiestrogen adjuvant therapies for MB, and findings that soy phytoestrogens also decrease sensitivity of MB cells to cytotoxic chemotherapeutics suggest that decreased exposure to environmental estrogens may benefit MB patient responses to chemotherapy.

Keywords: Chemotherapy, Cytoprotection, Concentration-response, Estrogen, In vitro, Isoflavones, Medulloblastoma, Phytoestrogen

Background

Medulloblastoma (MB) arise from neural precursors of the cerebellum and brainstem and are associated with the 4th ventricle. They are the most common central nervous system (CNS) malignancy in childhood with a peak incidence around 5 years of age [1–4]. These primitive neuroectodermal tumors (PNETs) while rare, with an overall incidence rate of 1.5 per million population, are more common in children 1–9 years of age (affecting 9.6 per million children) compared to adults 19 years of age and older, who have an incidence rate of 0.6 per million [5]. Less commonly, PNETs may develop in the cerebral hemisphere, these tumors are referred to CNS-PNETs. While sharing histological similarities, CNS-PNET tumors are genetically distinct from MB and have an overall incidence rate of 0.62 per million [5–8]. Histopathology grading has classically been used to separate MB into subgroups which differ with regard to biomarker profile and prognosis [9]. These subgroups include classic MB, desmoplastic MB, MB with excessive nodularity (MBEN), large cell MB and anaplastic MB [9–11]. Due to cellular and molecular heterogeneity across histological subgroups, and even within a singular tumor, a newer approach to MB grading has emerged that relies on comparative genome, transcriptome, and epigenetic analysis which may allow improved risk stratification and individualized targeted treatments [12]. By consensus four molecular subgroups are now recognized, they include wingless (WNT), sonic hedgehog (SHH), group 3 and group 4 [12–15]. Each subgroup has a characteristic genetic profile and gene expression patterns that appear to drive tumor progression, predict therapeutic responsiveness and prognosis. Further refinement of these molecular analyses has also found that pediatric and adult MB, are both histologically and genetically different diseases with characteristic differences in mutation accumulation, chromosomal deletion and amplification, and distinctive prognosis and survival rates [16–18].

Advancements in multimodal MB therapy utilizing maximal tumor resection, followed by radiation, and chemotherapy have greatly improved the chances of patient survival with 5 year overall survival rates for MB reaching between 60 and 80% depending on specific

tumor grade or molecular subtype; the survival rate for CNS-PNET patients is approximately 50% [18]. Conventional standard of care for MB most often involves combined radiation and polychemotherapy that results in improved outcomes compared to treatments limited to only tumor excision, radiation therapy or single agent chemotherapy [8, 19, 20]. Cytotoxic chemotherapy treatments for standard risk MB include a combination of cisplatin, cyclophosphamide, lomustine, and vincristine [21]. These agents vary in their mechanism of actions with cisplatin causing apoptosis due to DNA cross-linking, cyclophosphamide and lomustine are DNA alkylating agents, and the vinca alkaloid vincristine inhibits cell division by binding tubulin to inhibit microtubule formation [21]. Despite the success of these combined treatments, greater than 70% of MB survivors experience life-long neurological disabilities that include cognitive, motor, and/or vision impairments, as well as psychosocial dysfunction. Additionally, more than half of survivors also have severe endocrine impairments, which further contribute to a greatly diminished quality of life for MB survivors [22–24]. Thus, there is continued need to refine existing therapy and develop new adjuvant therapies that further improve MB and CNS-PNET cure rates, while at the same time, reduce the life-long adverse effects of both the disease and its treatment [25].

Previous study has demonstrated that growth and migration of MB and CNS-PNET cells and tumors are responsive to estrogen (17 β -estradiol; E2) and other estrogenic compounds [26–30]. Results of Western blot and immunohistochemistry analysis using ER α and ER β specific antibodies, along with pharmacological studies using ER selective agonists and antagonists have demonstrated that human MB tumors and cell lines express predominantly ER β and that estrogen's activity is dependent on ER β and independent of ER α [26–28]. In human MB cell lines and in vivo mouse models of MB, the growth stimulating effects of estrogen were found to be largely independent of increased proliferation [26]. Rather, estrogens were found to activate ER β and increase insulin-like growth factor 1 (IGF1) signaling pathways that act to reduce tumor cell apoptosis [26–28].

Results from additional studies have also demonstrated that the non-selective ER antagonist fulvestrant and the ER β selective antagonist PHTPP inhibited MB cell growth in cultured human MB cell lines [27]. In vivo therapeutic doses of fulvestrant also blocked the growth of MB tumor xenografts in nude mice, and slowed tumor growth and progression in genetic mouse models of MB [26–28]. These findings support the potential efficacy of anti-estrogen treatments (or other interventions that decrease ER activity) as beneficial adjuvants for MB management. The role of ER β in MB pathology however, is controversial in part because the role of ER β in cancer progression is in general poorly understood, with both tumor suppressor and tumor promoting effects of ER β having been reported in different ER expressing tumors [31, 32]. Results from different mouse models of MB have also found that estrogen and ER β activity can decrease MB tumor incidence [33], and in vitro studies have suggested that in the Daoy MB cell line, inhibition of ER β activity decreases sensitivity to cisplatin by enhancing Rad51 mediated DNA repair mechanisms [34]. To clarify the role of estrogens in MB we have used cell viability and clonogenic survival assays to determine whether the cytoprotective effects of E2 protected human D283 Med MB cells from the cytotoxic actions of the MB chemotherapeutic drugs cisplatin, vincristine, or lomustine. The effects of E2 on cisplatin chemosensitivity were also determined in the MB cell line Daoy and a CNS-PNET cell line PFSK-1. Additional experiments were done to determine whether the ER nonselective antagonist fulvestrant or the selective ER antagonist tamoxifen blocked the cytoprotective actions of 17 β -estradiol, and whether other ER selective agonists, and low concentrations of the soy-derived phytoestrogens genistein, daidzein, and s-equal were able to impact D283 Med chemosensitivity.

Methods

Steroids and pharmacological agents

Dimethylsulfoxide (DMSO), 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2H-tetrazolium bromide (MTT), 4-hydroxytamoxifen (4-OHT), 4',7-dihydroxyisoflavonoid (daidzein), 4',5,7-trihydroxyisoflavonoid (genistein) and 17 β -estradiol (E2) were from Sigma-Aldrich (St. Louis, MO). 4',7-dihydroxyisoflavan (s-equal) was from Cayman Chemical (Ann Arbor, MI). Fulvestrant (ICI 182,780), 4,4',4''-(4-propyl-[1H]-pyrazole-1,3,5-triyl)trisphenol (PPT), 2,3-bis(4-Hydroxyphenyl)-propionitrile (DPN) and 4-[2-Phenyl-5,7-bis(trifluoromethyl)pyrazolo [1,5-a]pyrimidin-3-yl]phenol (PHTPP) were from Tocris Bioscience (R&D Systems, Inc., Minneapolis, MN). Cisplatin, vincristine, and lomustine were from Selleck Chemical (Houston, TX). Cisplatin was usually prepared as a 1 mg/mL (3.3 mmole/L) stock solution in PBS. In assays involving hydrophobic ligands in DMSO, cisplatin

toxicity results were normalized using a standard curve comparing D283 Med cytotoxicity in the presence or absence of DMSO [35].

Cell Culture Conditions.

All cell lines were acquired from the American Type Culture Collection, cryopreserved and expanded for analysis. The D283Med cell line (HTB-185) was established from a peritoneal implant and ascetic fluid of a 6 year old male with metastatic medulloblastoma and grows in multicell aggregates in suspension with some adherent cells or on poly-L-lysine coated culture dishes with an epithelial morphology [36]. The Daoy cell line (HTB-186) was isolated from a desmoplastic cerebellar medulloblastoma of a 4 year old male and grows adherent with a polygonal morphology [37]. The PFSK-1 cell line, (CRL-2060) was established from a PNET from the cerebral hemisphere of a 22 month old male, and grows adherent with a fibroblast-like morphology [38]. The unique growth and morphological characteristics of each cell line was retained throughout the duration of the study. Details of cell culture methods were described previously [26–28]. Briefly, D283 Med and Daoy cells were grown in a humidified incubator at 37 °C and 5% CO₂ atmosphere in growth media containing minimum essential media (MEM) with Earle's Balanced Salt Solution (EBSS). Growth media for PFSK-1 cells was RPMI 1640. Media was supplemented with 10% fetal bovine serum (FBS), 4 mM L-glutamine, 100 U/ml penicillin and 100 μ g/ml streptomycin (Thermo Fisher Scientific, Waltham, MA). For general growth and expansion, D283 Med cells were maintained in suspension culture at 0.5 – 1 \times 10⁶ cells/ml. Daoy and PFSK-1 cells were maintained between 20 and 80% confluence. Growth media was refreshed every 2-3 days with cells split at a ratio of 1:5.

For Daoy growth analysis cells from subconfluent cultures were harvested by dissociation with 0.2 mM EDTA in PBS, resuspended in phenol red-free media supplemented with 10% charcoal stripped FBS (CSS). Viable cell numbers were determined by direct cell counting of trypan blue-excluding cells with a hemacytometer. Cells were seeded in triplicate into 60 mm culture dishes (22.06 cm²). Optimization experiments with cells plated at an initial density of 1000, 3000, 10,000, or 20,000 cells per dish in media supplemented with 10% FBS (positive control), 10% CSS plus or minus various concentrations of E2 indicated that 3000 cells per well allowed optimal viability analysis at all time points [28]. Cultures were untreated, or treated with DMSO (0.001%), or 10 nM E2 that was serially diluted into fresh DMSO/PBS vehicle to obtain an equal 0.001% final DMSO concentration in all cultures. At 24, 48, 72 and 96 h post-treatment viable cell numbers were determined by direct counting of trypan blue-excluding cells.

Viability Analysis

Viability was assessed by accumulation of formazan by reduction 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) or 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) in the presence of phenazine methosulfate (CellTiter 96[®] AQueous One Solution Cell Proliferation Assay; Promega) as previously described [39]. To avoid any potential MTT/MTS reduction assay bias, effects of E2 on D283 Med cell viability were confirmed with separate experiments using an Alamar Blue (resazurine) fluorescent dye assay at excitation wavelength of 535 nm (20 nm bandwidth) and an emission wavelength of 590 nm (35 nm bandwidth) [40]. Comparable results were observed for all assays. Regardless of specific assay, growing cells were harvested, counted and resuspended at a desired density in 10% CSS supplemented MEM/EBSS with 10 nM E2, or desired final concentration of fulvestrant or the vehicle control (0.01% DMSO) prior to cisplatin exposure. Cultures were incubated at 37 °C in 5% CO₂ overnight (18-24 h) at which time cells were exposed to the desired final concentration of cisplatin and incubated an additional 48 h prior to viability analysis. For each bioassay D283 Med cells were seeded in 96 well plates at 1×10^5 cells/mL ($1 - 2 \times 10^4$ cells per well) based on results of preliminary experiments to optimize each assay.

Clonogenic assay

Clonogenic/colony forming assays were adapted from published protocols [41, 42]. Exposure to ER ligands were started 24 h prior to determining cell numbers and diluting the cells to a concentration of 500 cells per ml and 1000 cell were seeded into 6-well tissue culture plates in 2 mL of 10% CSS supplemented phenol red-free MEM/EBSS. For D283 Med cells poly-L-lysine coated culture plates were used allowing adherent growth. For chemotherapeutic drug exposures, 0.5 mL of a 5× stock prepared in cell culture media was added to each well. After 6 h (cisplatin) or 24 h (vincristine and lomustine) of exposure, media was aspirated, cells were washed 2 times with chemotherapeutic compound-free media, and then cultured in 2.5 mL of growth media at 37 °C in 5% CO₂ until visible colonies containing >50 cells were observed. Preliminary range finding concentration response analysis was performed with each chemotherapeutic agent for each cell line, at the concentrations used between 5 and 40 clones per plate were typically observe for vehicle cultures. Incubation times were typically between 2 and 3 weeks with growth media refreshed every 2-3 days. Colonies were fixed and stained with 1% methylene blue in 50% ethanol (Fisher Scientific, Pittsburg, PA) or 0.1% coomassie brilliant blue in methanol (Bio-Rad, Hercules, CA). Digital images were captured and colonies

were counted using an Alpha Innotech FluorChem FC2 imager (ProteinSimple, Santa Clara, CA) and Adobe Photoshop (Adobe Systems Inc., San Jose, CA).

Caspase activity


All methods were done as previously described with D283 Med cells seeded into 96 well plates at a density of 1×10^6 cells/ml in phenol red-free MEM/EBSS lacking L-glutamine, 10% CSS and supplemented with increasing concentrations of daidzein, genistein or s-equal [27]. Cells exposed to 10 nM E2 or DMSO vehicle served as controls. Cells were lysed 48 h after seeding with 20 mM Tris-HCl (pH 7.5) with 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, and 1% Triton. Cell lysates were assayed for protein concentration using the BioRad Dc protein assay (Bio-Rad, Hercules, CA). Caspase activity (pmol of pNA hour⁻¹ mg protein⁻¹) from 10 µg of lysate was determined by comparing the amount p-nitroaniline (pNA) liberated from Ac-DEVD-pNA (Enzo Life Sciences, Farmingdale, NY) with a standard curve derived from known concentrations of pNA. Normalized caspase activity for each phytoestrogen are reported as a percentage of the maximal inhibitory effect of 10 nM E2.

Data and statistical analysis

All experiments were repeated a minimum of 3 times. Statistical analysis was conducted using one way ANOVA or two-way ANOVA with Holm-Sidak's multiple comparisons test. A minimal level of statistical significance for differences between groups was $p < .05$ and unless otherwise noted is indicated by *. Concentration response curves and IC₅₀ estimates were generated using a normalized variable slope Hill model. Analysis was performed using GraphPad Prism v6 software (GraphPad Software, Inc., La Jolla, CA).

Results

Compared to vehicle treated D283 Med cells, the cytotoxic effects of cisplatin were decreased in the presence of 10 nM E2 (Fig. 1a-b). When analyzed by an MTS reduction assay the presence of E2 increased the observed IC₅₀ of cisplatin from 5.6 µM (95% CI 4.7 - 6.9) to 14.7 µM (95% CI 10.5 - 20.5; Fig. 1a). Two-way ANOVA revealed a significant effect of cisplatin concentrations [$F(5, 36) = 49.65, p < .0001$], a significant effect of 10 nM E2 [$F(1, 36) = 10.07, p < .0031$], and a significant interaction between cisplatin concentration and E2 exposure [$F(5, 36) = 5.873, p = 0.0005$]. Shown in Fig. 1b are results of independent experiments using the resazurin reduction bioassay as an indicator of D283 Med viability where the IC₅₀ for cisplatin cytotoxicity in control cultures lacking E2 was calculated as 4.8 µM (95% CI 4.1 - 5.7). Revealing that the observed effects were not an assay specific effect, the



presence of 10 nM E2 similarly increased the calculated IC_{50} for cisplatin cytotoxicity to 9.4 μ M (95% CI 7.7 - 11.5). The cytoprotective effects of E2 were blocked by the ER antagonist fulvestrant (ICI 182,780). In the presence of both E2 and fulvestrant, the calculated cisplatin IC_{50} was 4.2 μ M (95% CI 3.2 - 5.39) which was indistinguishable from control (Fig. 1b).

Following the initial characterization studies of the effects of E2 on D283 Med cells in viability assays, a clonogenic colony forming assay [41, 42] in which cytotoxic treatments reproducibly caused 99-99.5% loss of viability was used to characterize in more detail the effects of estrogen the cytotoxicity of cisplatin. Based on preliminary concentration response analysis (Fig. 1c), the effect of 10 nM E2 on chemosensitivity of D283 Med cells to increasing cisplatin concentrations (2, 4, or 9 μ M) was characterized (Fig. 1d-e). At each cisplatin concentration E2 was significantly cytoprotective [$F(1, 18) = 311.6, p < .0001; p < .0001$ for each cisplatin concentration]. To

determine whether the observed protective effect of E2 in D283 Med cells was independent of the cytotoxic mechanism of action, additional experiments were performed to test the impact of E2 on lomustine and vincristine cytotoxicity (Fig. 2). Initial range-finding concentration response analysis in the D283 Med clonogenic assay indicated an IC_{50} for lomustine of 12.1 μ M (95% CI 11.7 - 12.6) (Fig. 2a) and 1.5 nM (95% CI 0.74 - 3.1) for vincristine (Fig. 2b). The presence of E2 significantly protected D283 Med cells from the cytotoxic effect of lomustine [$F(1, 30) = 74.64, p < .0001$] at each concentration tested (Fig. 2c; 10 M $p = .0036$, and $p < .0001$ for 20 and 40 μ M). For vincristine 10 nM E2 also significantly [$F(1, 18) = 196.2, p < .0001$] decreased cytotoxicity at each concentration (Fig. 2d; 5 and 10 nM, $p < .0001$ and $p = .0003$ for 20 nM).

Compared to vehicle treated D283 Med cells exposed to 4 μ M cisplatin, E2 ($p < .0001$) and the ER β selective agonist DPN ($p < .0001$) each increased numbers of




Fig. 2 The effects of E2 exposure on lomustine and vincristine cytotoxicity in D283 Med cells. Initial range finding concentration response analysis of the cytotoxic effects of **a** lomustine ($IC_{50} = 12.1 \mu M$) and **b** vincristine ($IC_{50} = 1.5 nM$) on D283 Med in a clonogenic assay of cell survival. **c** Quantification of surviving colony numbers from clonogenic assays of D283 Med cells exposed to 10, 20, or 40 μM lomustine with or without 10 nM E2 ($n = 6$ for each group). **d** Quantification of colony number from clonogenic assays of D283 Med cells exposed to 5, 10, or 20 nM vincristine with or without 10 nM E2 ($n = 8$ for each group). All results are expressed as mean \pm SEM. Significant differences from the control group were determined by two-way ANOVA followed by Holm-Sidak's post-test analysis and is indicated above the error bars; * $p \leq .05$


surviving colonies compared to cisplatin alone control cultures (Fig. 3a-b). The cytotoxic effect of 4 μM cisplatin was not significantly changed by the ER α selective agonist PPT ($p > .9999$). At a final concentration of 10 nM, the soy isoflavonoids genistein ($p = .0239$), daidzein ($p < .0001$), or the bacterial metabolite of daidzein, (s)-equol ($p < .0001$) each significantly protected D283 Med cells from the cytotoxic effect of cisplatin (Fig. 3b). The relative magnitude of the protective effects for each of the compounds is consistent with their selectivity and potency at ER β [43]. Increasing concentrations of each phytoestrogen significantly [$F(2, 256) = 4.85, p < .0086$] and dose-dependently decreased caspase 3 activity in D283 Med cells. The decrease in caspase activity mirrored the cytoprotective effects seen in the clonogenic assay (Fig. 3b). The differences in the suppression of caspase activity compared to control reached a significant difference in the 10 nM (10^{-8} M) groups for s-equol and daidzein and for genistein at 100 nM (Fig. 3c).

The cytoprotective effects of E2 in D283 Med cells exposed to cisplatin ($p = .0001$) were eliminated by the non-selective ER antagonist fulvestrant (10 nM; ICI; E2 vs. E2/fulvestrant $p = < .0001$; Fig. 3d-e), the selective estrogen receptor modulator 4-OH tamoxifen (1 μM ; E2 vs E2/tamoxifen $p = < .0001$) or the ER β selective antagonist PHTPP (5 μM ; E2 vs E2/PHTPP $p = < .0001$; Fig. 3e). In CNS-PNET derived PFSK-1 cells, 10 nM E2 also resulted

in increased survival [$F(1, 34) = 62.30, p < .0001$], with a clear decrease in sensitivity to the cytotoxic effects observed for all three cisplatin concentrations tested (Fig. 4a). Fulvestrant (10 nM) also blocked the cytoprotective effects of 10 nM E2 ($p = .0213$; Fig. 4b). In contrast to both D283 Med and PFSK-1 cells, the cytotoxic effects of cisplatin were increased in Daoy cells by the presence of 10 nM E2 where a significant decrease [$F(1, 18) = 62.75, p < .0001$] in surviving colony formation was observed in the estrogen treated cultures (Fig. 4c). The increased sensitivity of Daoy to cisplatin in the presence of E2 ($p = .0194$) was also eliminated by fulvestrant ($p = .0012$; Fig. 4d) and 10 nM E2 did not stimulate growth of Daoy cells (Fig. 4e).

Discussion

The use of aggressive multimodal treatments has resulted in increased survival for MB patients, most survivors however suffer from life-long adverse effects that greatly diminish their quality of life [22]. The presented findings demonstrate that E2 can increase the resistance to cytotoxic chemotherapeutics commonly used to treat MB, and that blockade of estrogen receptor activity inhibits this effect. These findings suggest that ER antagonists may be a useful adjuvant approach to current cytotoxic chemotherapy used to treat MB. The cytoprotective effects of estrogens, either endogenous or derived from environmental sources such as diet or estrogenic endocrine



disruptors from medical devices [37, 44], if translatable to MB patients, would require more aggressive chemotherapeutic interventions to achieve a cure in patients with increased levels of estrogenic activities. In previous studies, ER β activation in MB and CNS-PNET tumor cells was found to stimulate cytoprotective mechanisms which decreased caspases 3 activity, and loss of ER β activity inhibited MB tumor growth and increased apoptosis in vivo [27]. The results of the current study lend additional evidence that ER β initiated mechanisms promote MB and CNS-PNET survival by demonstrating that the ER β selective agonist DPN protected D283 Med cells from cisplatin induced cytotoxicity, and that inhibition of ER β by PHTPP blocked the protective effect of E2. Along with demonstrating mechanistic

involvement of ER β , the ability of the anti-estrogen chemotherapeutics fulvestrant and tamoxifen to each block the protective actions of estrogen in MB and CNS-PNET support previous findings which demonstrated that antiestrogen chemotherapeutics block the growth of MB tumors in vivo [26, 27] and that tamoxifen can sensitize MB cells to the cytotoxic effects of the topoisomerase inhibitor etoposide [45].

Specific treatments for MB and CNS-PNETS are constantly evolving. Depending on specific risk stratification, the current standard of care often includes maximal surgical resection that allows preservation of neurological function, postoperative radiation therapy, followed by chemotherapy employing a combination of the DNA crosslinking agent cisplatin, a DNA alkylating agents such




Fig. 4 The effects of E2 exposure on cisplatin cytotoxicity in PFSK-1 CNS-PNET cells and Daoy cells. **a** Quantification of surviving colony numbers from clonogenic assays of PFSK-1 cells exposed to 2, 4 or 9 μM cisplatin with or without 10 nM E2 ($n = 8$ for each cisplatin treatment group except 4 μM where $n = 4$). **b** Quantification of colony number from clonogenic assays of PFSK-1 cells cotreated with 4 μM cisplatin and either vehicle, 10 nM E2, 10 nM fulvestrant (ICI), 10 nM E2 and 10 nM fulvestrant (ICI/E2), $n = 4$ for each group. **c** Quantification of surviving colony numbers from clonogenic assays of Daoy cells exposed to 2, 4 or 9 μM cisplatin with or without 10 nM E2. For each group $n = 4$. **d** Quantification of colony number from clonogenic assays of Daoy cells cotreated with 4 μM cisplatin and either vehicle, 10 nM E2, 10 nM fulvestrant (ICI), 10 nM E2 and 10 nM fulvestrant (ICI/E2), $n = 8$ for each group. **e** Analysis of the effects of 10 nM E2 on viability of Daoy cells. At T_0 3000 cells were plated into 60 mm cell culture dishes, in growth media cells containing 10% CSS. At each indicated time point (hours post treatment) cells were harvested and trypan-excluding cells were counted. Vehicle was 0.0001% DMSO and replacement of CSS with 10% FBS served as a positive control. At each time point $n = 10$ for all treatments. Results are expressed as mean \pm SEM. Significant differences from vehicle control are indicated above the treatment group error bars with individual comparisons indicated above brackets: * $p \leq .05$; ** $p \leq 0.01$; *** $p \leq .001$; NS, not significant

as lomustine, and vincristine, a microtubule inhibitor [21, 46]. Each of these drugs works in different ways to stop the growth of tumor cells, either by killing the cells, or by stopping them from dividing. As for other cancers, treatment for MB and CNS-PNET has leveraged the fundamental understanding that cancer patients given multimodal treatments which include some combination of surgical tumor resection and radiation, plus a single or multiple chemotherapy agents, have improved short and long term outcomes. The combine effect of multiple

cytotoxic treatments arise because each targets different processes involved in tumor cell survival [47, 48]. Targeted cancer therapies based on molecular markers such as endocrine therapy for prostate cancer or inhibiting ER activity in ER-positive breast cancer also benefits from a multimodal treatment approach that can include endocrine based therapies, along with chemotherapy involving single or multiple cytotoxic agents [19, 20, 49–53]. Because E2 increases MB tumor survival through a general cytoprotective mechanism by increasing IGF-signaling [27],

we hypothesized that its cytoprotective effects would decrease the cytotoxic effects of chemotherapeutic agents used for MB treatment independent of their mechanism of action. The presented studies, focused primarily on the most commonly used MB chemotherapy drug cisplatin (a DNA crosslinking agent), found that estrogen and soy-derived phytoestrogens were cytoprotective, typically causing about a 2 fold increase in viability. The cytotoxic effects of both the alkylating agent lomustine and the microtubule inhibitor vincristine were also decreased by E2 in D283 Med MB cells demonstrating that the estrogen-induced cytoprotective mechanisms were in fact independent of the mechanism by which these chemotherapeutic agents act to initiate MB cell death.

We and others have found that Daoy cells express ER β , with little or no active ER α , but the pattern of ER β protein isoform expression is distinctive from other MB and CNS-PNET cells in which E2 has cytoprotective activities [26, 54]. We also previously found that estrogen stimulated the migration of Daoy cells by an ER β -dependent mechanism that was identical to that observed in other MB and CNS-PNET cells [26]. It was found here however, that unlike other MB cells, E2 alone did not increase viability of Daoy cells, instead the presence of E2 increased sensitivity of these cells to cisplatin cytotoxicity. It was also observed that inhibition of estrogen activity with 10 nM fulvestrant (a concentration that is 10-fold more than required to fully inhibit estrogen dependent growth of MCF-7 breast cancer cells [55]) blocked estrogen mediated sensitization of Daoy cells to cisplatin. At this fulvestrant concentration the classical nuclear receptor transactional activities of the ERs are inhibited, suggesting that blockage of ER β activity is responsible for the observed chemoresistance to cisplatin. Fulvestrant however, also acts as a full agonist of rapid ER β signaling in cerebellar granule cell precursors [56]. The impact of fulvestrant agonist activity on rapid estrogen signaling in MB remains to be clearly defined.

Urbanska and colleagues, while not investigating the growth promoting actions of estrogen, previously reported that ER β could interact with nuclear IRS1 to inhibit Rad51 mediated DNA repair mechanisms in Daoy cells, findings that suggested estrogen's ability to increase Daoy cell sensitivity to cisplatin might involve an ER β /IRS1 mediated decrease in Rad51 homologous recombination DNA repair mechanisms [54]. Their additional results from experiments using a higher 10 μ M concentration of fulvestrant ($IC_{50} = 0.29$ nM) found it caused resistance of Daoy and D384 cells to the cytotoxic actions of cisplatin, effects that were not significant in the D283 Med cells [34]. The studies presented here, using lower concentrations of fulvestrant, failed to observe increased sensitivity of Daoy cells to cisplatin. In light of the higher concentrations of fulvestrant used for those

previous experiments, it is possible that the observed protective effects of fulvestrant were not specific and involved activities other than inhibition of ER β . Another possible explanation for decreased cytotoxicity could be that DMSO, if used as a vehicle, was inactivating cisplatin. This possibility cannot be ruled out because specific information regarding vehicle for neither cisplatin nor fulvestrant were not specified [34]. The ability of DMSO to greatly decrease the cytotoxic activity of cisplatin and other platinum chemotherapeutic drugs has previously been characterized in detail [35].

For studies using the Daoy cells as a model of MB, it is also valuable to consider the fact that current molecular profiling and cytogenetic data supports the conclusion that the Daoy cell line, while most closely resembling the SHH molecular subgroup, is markedly different from all primary MB tumor subgroups [57, 58]. Daoy cells are distinctive from MB cell lines like D283 Med that retain the hallmarks of MYC amplification and i17q that are associated with poor clinical outcomes [57, 58]. It is especially notable that the hypertetraploid karyotype of the Daoy cells does not resemble MB karyotypes, and the presence of two X chromosomes and a lack of a Y chromosome, is inconsistent with the sex of the male patient from which the original tumor biopsy was isolated [37] and thus may poorly represent an authentic MB cell type. In spite of those caveats, it is also possible that the observed differences in the impact of E2 on Daoy and D283 Med MB cells may actually reflect the well-known heterogeneity found in MB. It is possible that different molecular subtypes of MB and even different populations of cells in a single patient's tumor may differently respond to ER agonists and antagonists. This raises the interesting possibility that the responses to estrogen in MB are heterogeneous and that some populations of cells are differentially responsive to estrogen's effects.

Conclusions

The presented results demonstrate that the cytoprotective effects of E2, which can be cell line dependent, are clearly chemoprotective in some MB and CNS-PNET cell lines. The results of additional experiments also demonstrated that like E2, low and physiological levels of the soy-derived phytoestrogens genistein, daidzein, and equol can decrease caspase activity in D283 Med MB cells resulting in an estrogen-like inhibition of the cytotoxic actions of cisplatin. The finding that soy phytoestrogens also decrease sensitivity to the cytotoxic actions of cisplatin suggest that attention to decreasing exposures to environmental estrogens that include not only phytochemicals but also estrogenic endocrine disruptors may benefit MB patients undergoing cytotoxic chemotherapy.

Abbreviations

CNS-PNET: Central nervous system primitive neuroectodermal tumors; CSS: Charcoal stripped FBS; DMSO: Dimethylsulfoxide; E2: 17 β -estradiol; EBSS: Earle's Balanced Salt Solution; FBS: Fetal bovine serum; MB: Medulloblastoma; MBEN: Medulloblastoma with excessive nodularity; MEM: Minimum essential media; MTS: 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; pNA: p-nitroaniline; PNETs: Primitive neuroectodermal tumors; SHH: Sonic hedgehog; WNT: Wingless

Acknowledgements

As part of the Department of Pharmacology and Cell Biophysics, Summer Undergraduate Research Program, Katie Wray was a Dalton Zannoni Summer Undergraduate Research Fellow of the American Society of Pharmacology and Experimental Therapeutics. We are grateful for the assistance of Drs. Robert Rapaport and JoEl Schultz and Nancy Thyberg in their dedicated support of graduate and undergraduate training.

Funding

These studies were funded in part by RO1 ES015145 grant from NIEHS and the NCSU Center for Health and Human Environment that is funded by NIEHS award P30ES025128.

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Authors' contributions

SB analyzed and interpreted data, performed cell viability experiments and wrote the manuscript; CB performed and analyzed cell viability and apoptosis experiments; CC performed cell viability assays, assisted in data collection and analysis and contributed early drafts of the manuscript; MK performed and analyzed cell growth studies, GM, FS and KW performed and analyzed cell viability studies. All authors read and approved the final manuscript.

Ethics approval and consent to participate

N/A.

Consent for publication

N/A.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Biological Science and Center for Human Health and the Environment, North Carolina State University, Raleigh, NC, USA. ²Department of Pharmacology and Cell Biophysics, University of Cincinnati, Cincinnati, OH, USA. ³Department of Pharmacology and Cell Biophysics, Summer Undergraduate Research Program University of Cincinnati, Cincinnati, OH, USA. ⁴Department of Pharmacology and Cell Biophysics, Molecular, Cellular and Biochemical Pharmacology PhD Graduate Program, University of Cincinnati, Cincinnati, OH, USA. ⁵Department of Pharmacology and Cell Biophysics, Molecular, Cellular and Biochemical Pharmacology Masters in Safety Pharmacology Training Program, University of Cincinnati, Cincinnati, OH, USA.

Received: 21 October 2016 Accepted: 22 June 2017

Published online: 06 September 2017

References

- Ries LAG, Smith MA, Gurney JG, Linet M, Tamra T, Young JL, Brunin GR, Monograph SP: Cancer incidence and survival among children and adolescents: United States SEER program 1975–1995. NIH Pub No 99–4649 1999.
- Gibson P, Tong Y, Robinson G, Thompson MC, Curle DS, Eden C, et al. Subtypes of medulloblastoma have distinct developmental origins. *Nature*. 2010;468:1095–9.
- Wang J, Wechsler-Reya RJ. The role of stem cells and progenitors in the genesis of medulloblastoma. *Exp Neurol*. 2014;260:69–73.
- Yang ZJ, Ellis T, Markant SL, Read TA, Kessler JD, Bourbonlous M, et al. Medulloblastoma can be initiated by deletion of patched in lineage-restricted progenitors or stem cells. *Cancer Cell*. 2008;14:135–45.
- Smoll NR, Drummond KJ. The incidence of medulloblastomas and primitive neuroectodermal tumours in adults and children. *J Clin Neurosci*. 2012;19:1541–4.
- Pomeroy SL, Tamayo P, Gaasenbeek M, Sturla LM, Angelo M, McLaughlin ME, et al. Prediction of central nervous system embryonal tumour outcome based on gene expression. *Nature*. 2002;415:436–42.
- Dubuc AM, Northcott PA, Mack S, Witt H, Pfister S, Taylor MD. The genetics of pediatric brain tumors. *Curr Neurol Neurosci Reports*. 2010;10:215–23.
- Mueller S, Chang S. Pediatric brain tumors: current treatment strategies and future therapeutic approaches. *Neurotherapeutics*. 2009;6:570–86.
- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, Scheithauer BW, Kleihues P: The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 2007, 114:97–109.
- Ellison D. Classifying the medulloblastoma: insights from morphology and molecular genetics. *Neuropathol Appl Neurobiol*. 2002;28:257–82.
- Gulino A, Arcella A, Giangaspero F. Pathological and molecular heterogeneity of medulloblastoma. *Curr Opin Oncol*. 2008;20:668–75.
- Taylor MD, Northcott PA, Korshunov A, Remke M, Cho YJ, Clifford SC, et al. Molecular subgroups of medulloblastoma: the current consensus. *Acta Neuropathol*. 2012;123:465–72.
- DeSouza RM, Jones BR, Lowis SP, Kurian KM. Pediatric medulloblastoma - update on molecular classification driving targeted therapies. *Front Oncol*. 2014;4:176.
- Li KK, Lau KM, Ng HK. Signaling pathway and molecular subgroups of medulloblastoma. *Int J Clin Exp Pathol*. 2013;6:1211–22.
- Samkari A, White JC, Packer RJ. Medulloblastoma: toward biologically based management. *Semin Pediatr Neurol*. 2015;22:6–13.
- Parsons DW, Li M, Zhang X, Jones S, Leary RJ, Lin JC-H, et al. The genetic landscape of the childhood cancer Medulloblastoma. *Science*. 2011;331:435–9.
- Korshunov A, Remke M, Werft W, Benner A, Ryzhova M, Witt H, et al. Adult and pediatric medulloblastomas are genetically distinct and require different algorithms for molecular risk stratification. *J Clin Oncol*. 2010;28:3054–60.
- Smoll NR. Relative survival of childhood and adult medulloblastomas and primitive neuroectodermal tumors (PNETs). *Cancer*. 2012;118:1313–22.
- Aapro MS. Adjuvant therapy of primary breast cancer: a review of key findings from the 7th international conference, St. Gallen, February 2001. *Oncologist*. 2001;6:376–85.
- Davis TE, Carbone PP. Drug treatment of breast cancer. *Drugs*. 1978;16:441–64.
- Martin AM, Raabe E, Eberhart C, Cohen KJ. Management of Pediatric and Adult Patients with Medulloblastoma. *Curr Treat Options in Oncol*. 2014; 15(4):581–94.
- Frange P, Alapetite C, Gaboriaud G, Bours D, Zucker JM, Zerah M, et al. From childhood to adulthood: long-term outcome of medulloblastoma patients. The Institut curie experience (1980–2000). *J Neuro-Oncol*. 2009;95:271–9.
- Mulhern RK, Merchant TE, Gajjar A, Reddick WE, Kun LE. Late neurocognitive sequelae in survivors of brain tumours in childhood. *Lancet Oncol*. 2004;5:399–408.
- Rose SR, Danish RK, Kearney NS, Schreiber RE, Lustig RH, Burghen GA, et al. ACTH deficiency in childhood cancer survivors. *Pediatr Blood Cancer*. 2005; 45:808–13.
- Belcher SM. Blockade of estrogen receptor signaling to improve outlook for medulloblastoma sufferers. *Future Oncol*. 2009;5:751–4.
- Belcher SM, Ma X, Le HH. Blockade of estrogen receptor signaling inhibits growth and migration of medulloblastoma. *Endocrinology*. 2009;150:1112–21.
- Cookman CJ, Belcher SM. Estrogen receptor-beta up-regulates IGF1R expression and activity to inhibit apoptosis and increase growth of Medulloblastoma. *Endocrinology*. 2015;156:2395–408.
- Kirby M, Zsarnovszky A, Belcher SM. Estrogen receptor expression in a human primitive neuroectodermal tumor cell line from the cerebral cortex: estrogen stimulates rapid ERK1/2 activation and receptor-dependent cell migration. *Biochem Biophys Res Commun*. 2004;319:753–8.

29. Levin ER. Integration of the extranuclear and nuclear actions of estrogen. *Mol Endocrinol*. 2005;19:1951–9.
30. Levin ER. Minireview: Extranuclear steroid receptors: roles in modulation of cell functions. *Mol Endocrinol* (Baltimore, Md). 2011;25:377–84.
31. Murphy LC, Leygue E. The role of estrogen receptor-beta in breast cancer. *Semin Reprod Med*. 2012;30:5–13.
32. Thomas C, Gustafsson JA. The different roles of ER subtypes in cancer biology and therapy. *Nat Rev Cancer*. 2011;11:597–608.
33. Mancuso M, Leonardi S, Giardullo P, Pasquali E, Borra F, Stefano ID, et al. The estrogen receptor beta agonist diarylpropionitrile (DPN) inhibits medulloblastoma development via anti-proliferative and pro-apoptotic pathways. *Cancer Lett*. 2011;308:197–202.
34. Wilk A, Waligorska A, Waligorski P, Ochoa A, Reiss K. Inhibition of ERbeta induces resistance to cisplatin by enhancing Rad51-mediated DNA repair in human medulloblastoma cell lines. *PLoS One*. 2012;7:e33867.
35. Hall MD, Telma KA, Chang KE, Lee TD, Madigan JP, Lloyd JR, et al. Say no to DMSO: dimethylsulfoxide inactivates cisplatin, carboplatin, and other platinum complexes. *Cancer Res*. 2014;74:3913–22.
36. Friedman HS, Burger PC, Bigner SH, Trojanowski JQ, Wikstrand CJ, Halperin EC, et al. Establishment and characterization of the human medulloblastoma cell line and transplantable xenograft D283 med. *J Neuropathol Exp Neurol*. 1985;44:592–605.
37. Jacobsen PF, Jenkyn DJ, Papadimitriou JM. Establishment of a human medulloblastoma cell line and its heterotransplantation into nude mice. *J Neuropathol Exp Neurol*. 1985;44:472–85.
38. Fults D, Pedone CA, Morse HG, Rose JW, McKay RD. Establishment and characterization of a human primitive neuroectodermal tumor cell line from the cerebral hemisphere. *J Neuropathol Exp Neurol*. 1992;51:272–80.
39. Wong JK, Kennedy PR, Belcher SM. Simplified serum- and steroid-free culture conditions for high-throughput viability analysis of primary cultures of cerebellar granule neurons. *J Neurosci Methods*. 2001;110:45–55.
40. O'Brien J, Wilson I, Orton T, Pognan F. Investigation of the Alamar blue (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity. *Eur J Biochem*. 2000;267:5421–6.
41. Franken NA, Rodermond HM, Stap J, Haveman J, van Bree C. Clonogenic assay of cells in vitro. *Nat Protoc*. 2006;1:2315–9.
42. Munshi A, Hobbs M, Meyn RE. Clonogenic cell survival assay. *Methods Mol Med*. 2005;110:21–8.
43. Muthyala RS, Ju YH, Sheng S, Williams LD, Doerge DR, Katzenellenbogen BS, et al. Equol, a natural estrogenic metabolite from soy isoflavones: convenient preparation and resolution of R- and S-equols and their differing binding and biological activity through estrogen receptors alpha and beta. *Bioorg Med Chem*. 2004;12:1559–67.
44. Patisaul HB, Jefferson W. The pros and cons of phytoestrogens. *Front Neuroendocrinol*. 2010;31:400–19.
45. Ramachandran C, Khatib Z, Petkarou A, Fort J, Fonseca HB, Melnick SJ, et al. Tamoxifen modulation of etoposide cytotoxicity involves inhibition of protein kinase C activity and insulin-like growth factor II expression in brain tumor cells. *J Neuro-Oncol*. 2004;67:19–28.
46. De Braganca KC, Packer RJ. Treatment Options for Medulloblastoma and CNS primitive Neuroectodermal tumor (PNET). *Curr Treat Options Neurol*. 2013;15:593–606.
47. DeVita VT, Young RC, Canellos GP. Combination versus single agent chemotherapy: a review of the basis for selection of drug treatment of cancer. *Cancer*. 1975;35:98–110.
48. Muggia FM, Cortes-Funes H, Wasserman TH. Radiotherapy and chemotherapy in combined clinical trials: problems and promise. *Int J Radiat Oncol Biol Phys*. 1978;4:161–71.
49. Albain KS, Barlow WE, Ravdin PM, Farrar WB, Burton GV, Ketchel SJ, et al. Adjuvant chemotherapy and timing of tamoxifen in postmenopausal patients with endocrine-responsive, node-positive breast cancer: a phase 3, open-label, randomised controlled trial. *Lancet*. 2009;374:2055–63.
50. Goel S, Sharma R, Hamilton A, Beith J. LHRH agonists for adjuvant therapy of early breast cancer in premenopausal women. *Cochrane Database Syst Rev*. 2009;(4):CD004562. doi:10.1002/14651858.CD004562.pub3.
51. Goldhirsch A, Coates AS, Colleoni M, Castiglione-Gertsch M, Gelber RD. Adjuvant chemoendocrine therapy in postmenopausal breast cancer: cyclophosphamide, methotrexate, and fluorouracil dose and schedule may make a difference. International breast cancer study group. *J Clin Oncol*. 1998;16:1358–62.
52. Lumachi F, Brunello A, Maruzzo M, Basso U, SMM B. Treatment of estrogen receptor-positive breast cancer. *Curr Med Chem*. 2013;20:596–604.
53. Mehta RS, Barlow WE, Albain KS, Vandenberg TA, Dakhil SR, Tirumali NR, et al. Combination anastrozole and fulvestrant in metastatic breast cancer. *N Engl J Med*. 2012;367:435–44.
54. Urbanska K, Pannizzo P, Lassak A, Gualco E, Surmacz E, Croul S, et al. Estrogen receptor beta-mediated nuclear interaction between IRS-1 and Rad51 inhibits homologous recombination directed DNA repair in medulloblastoma. *J Cell Physiol*. 2009;219:392–401.
55. Wakeling AE, Dukes M, Bowler J. A potent specific pure Antiestrogen with clinical potential. *Cancer Res*. 1991;51:3867–73.
56. Wong JK, Le HH, Zsarnovszky A, Belcher SM. Estrogens and ICI182,780 (Faslodex) modulate mitosis and cell death in immature cerebellar neurons via rapid activation of p44/p42 mitogen-activated protein kinase. *J Neurosci*. 2003;23:4984–95.
57. Xu J, Margol A, Asgharzadeh S, Erdreich-Epstein A. Pediatric brain tumor cell lines. *J Cell Biochem*. 2015;116:218–24.
58. Weeraratne SD, Amani V, Teider N, Pierre-Francois J, Winter D, Kye MJ, et al. Pleiotropic effects of miR-183–96–182 converge to regulate cell survival, proliferation and migration in medulloblastoma. *Acta Neuropathol*. 2012; 123:539–52.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

