

# Bees may drive the reproduction of four sympatric cacti in a vanishing coastal mediterranean-type ecosystem

Pablo C. Guerrero<sup>1</sup>, Claudia A. Antinao<sup>1</sup>, Beatriz Vergara-Meriño<sup>1</sup>, Cristian A. Villagra<sup>2</sup> and Gastón O. Carvallo<sup>3</sup>

<sup>1</sup> Departamento de Botánica, Facultad de Ciencias Naturales y Oceanográficas, Universidad de Concepción, Concepción, Chile

<sup>2</sup> Instituto de Entomología, Universidad Metropolitana de Ciencias de la Educación, Santiago, Chile

<sup>3</sup> Instituto de Biología, Facultad de Ciencias, Pontificia Universidad Católica de Valparaíso, Valparaíso, Chile

## ABSTRACT

**Background:** Sympatric congeneric plants might share pollinators, or each species might avoid competition by evolving specialized traits that generate partitions in pollinator assemblages. In both cases, pollen limitation (a decrease in the quality and quantity of compatible reproductive pollen) can occur, driving the plant mating system to autogamy as a mechanism of reproductive assurance. We assessed the relationships between pollinator assemblages and mating systems in a group of sympatric congeneric plants. We attempted to answer the following questions: (i) How similar are pollinator assemblages among sympatric cactus species? (ii) Which mating systems do sympatric cactus species use?

**Methods:** We studied sympatric *Eriosyce* taxa that inhabit a threatened coastal strip in a mediterranean-type ecosystem in central Chile. We performed field observations on four taxa and characterized pollinators during the years 2016 and 2017. We estimated differences in the pollinator assemblages using the Bray–Curtis index. To elucidate the mating systems, we conducted hand-pollination experiments using three treatments: manual cross-pollination, automatic self-pollination, and control (unmanipulated individuals). We tested differences in seed production for statistical significance using Kruskal–Wallis analysis.

**Results:** *Eriosyce subgibbosa* showed a distinctive pollinator assemblage among the sympatric species that we studied (similarity ranged from 0% to 8%); it was visited by small bees and was the only species that was visited by the giant hummingbird *Patagona gigas*. Pollinator assemblages were similar between *E. chilensis* (year 2016 = 4 species; 2017 = 8) and *E. chilensis* var. *albidiflora* (2016 = 7; 2017 = 4); however, those of *E. curvispina* var. *mutabilis* (2016 = 7; 2017 = 6) were less similar to those of the aforementioned species. *E. curvispina* var. *mutabilis* showed the highest interannual variation in its pollinator assemblage (18% similarity). Reproduction in *E. subgibbosa* largely depends on pollinators, although it showed some degree of autogamy. Autonomous pollination was unfeasible in *E. chilensis*, which depended on flower visitors for its reproductive success. Both *E. chilensis* var. *albidiflora* and *E. curvispina* var. *mutabilis* showed some degree of autogamy.

**Discussion:** We observed differences in pollinator assemblages between *E. subgibbosa* and the remaining *Eriosyce* taxa, which depend on hymenopterans for

Submitted 18 March 2019  
Accepted 10 September 2019  
Published 7 October 2019

Corresponding author  
Pablo C. Guerrero,  
pabloguerrero@udec.cl

Academic editor  
Alastair Culham

Additional Information and  
Declarations can be found on  
page 12

DOI 10.7717/peerj.7865

© Copyright  
2019 Guerrero et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

pollen transfer. Pollinator assemblages showed considerable interannual variation, especially those of *E. subgibbosa* (ornithophilous syndrome) and *E. curvispina* var. *mutabilis* (melitophilous syndrome). Autogamous reproduction in these taxa may act as a reproductive assurance mechanism when pollinator availability is unpredictable. Our study contributes to improving our understanding of the reproductive systems of ecological interactions between threatened species in a Chilean mediterranean-type ecosystem.

**Subjects** Biodiversity, Ecology, Plant Science

**Keywords** Cactaceae, Endemism, Insect pollinators, Plant-animal interactions, Pollen limitation, Hummingbird, Central Chile, Los Molles, Pichidanguí, anthropocene

## INTRODUCTION

Pollinator assemblages are a crucial component of plant reproduction (Eckhart, 1992; Johnson, 2010), driving the evolution of plant mating systems (Barrett, 1998; Kalisz, Vogler & Hanley, 2004; Goodwillie, Kalisz & Eckert, 2005). Different factors, such as the taxonomic and functional composition of pollinator assemblages, pollination rates, and pollen flows, influence the ecological dynamics of plant-pollinator interactions, showing significant variation in spatial and temporal dimensions (Johnson & Steiner, 2000; Herlihy & Eckert, 2005; Weber, 2017). Pollinator assemblage properties in sympatric species could occur in a continuum between the two extremes: sharing a wide range of floral visitors (Marques et al., 2007; Schlüter et al., 2009; Ferreira et al., 2018) or specializing in the use of floral visitors, which reduces long-term competition by pollinators (Stone, Willmer & Rowe, 1988). Ultimately, these scenarios can favor pollen limitation either by deleterious impacts of heterospecific pollen transference, such as microallelopathic effects at the stigma level and clogging (Van Der Niet, Johnson & Linder, 2006; Bodbyl Roels & Kelly, 2011; Barrett & Harder, 2017), or by changeable pollination environments (Eckert & Herlihy, 2004; Kalisz, Vogler & Hanley, 2004; Verdú et al., 2006). Extreme scenarios, such as those mentioned above, decrease pollination quality and quantity which can promote rapid evolutionary changes toward selfing and genetic self-compatibility (Kalisz & Vogler, 2003; Moeller & Geber, 2005; Morgan, Wilson & Knight, 2005).


Partitioning pollinator assemblages (specialization or the use of a restricted number of taxa or functional groups; Johnson & Steiner, 2000) can be considered an advantageous strategy when closely related taxa occur in sympatry (Stone, Willmer & Rowe, 1988), which could reduce heterospecific pollen transference (Kudo & Kasagi, 2005). However, adapting specialized pollinator assemblages may increase plant susceptibility to pollen limitation since pollinators show unpredictable richness and density over time (Johnson & Steiner, 2000). In this scenario, the evolution of selfing strategies may be favored in plants (Fenster et al., 2004; Kalisz, Vogler & Hanley, 2004; Kudo & Kasagi, 2005; Knight et al., 2005; Pérez, Arroyo & Armesto, 2009). Experimental evidence has shown a rapid evolution of the pollination system in angiosperms (Kearns, Inouye & Waser, 1998). Indeed, understanding the evolutionary plasticity of pollination systems may contribute to

buffering anthropogenic impacts, such as species loss or land use changes, that affect plant-pollinator interactions (Hadley & Betts, 2009; Young et al., 2016).

Among angiosperms, the species of the Cactaceae family have one of the most impressive evolutionary labile reproductive systems (Mandujano et al., 2010; Schlumpberger & Renner, 2012; Hernández-Hernández et al., 2014; Guerrero et al., 2019a). Cacti have evolved conspicuous flowers that attract a wide range of animal pollinators, including vertebrates (e.g., bats, hummingbirds, passerine birds, and lizards) and insects, such as moths, flies, and bees (Mandujano et al., 2010; Guerrero et al., 2012). Cacti exhibit geographical covariation between pollinator assemblages and floral morphology (Schlumpberger et al., 2009; Walter, 2010), a relationship that may reinforce the dependence on biotic pollinators to achieve reproductive success (Valiente-Banuet et al., 1997; McIntosh, 2002; Walter, 2008) and explain the high levels of self-incompatibility exhibited in this family (Boyle, 1997; Martínez-Peralta, Márquez-Guzmán & Mandujano, 2014). Self-compatibility and autonomous systems for reproductive assurance in cacti have been observed in cases of pollinator inconstancy (Nassar & Ramírez, 2004).

Cacti exhibit rich variation in reproductive strategies, although, astonishingly, the reproductive system has been studied in only 2% of species in this family (Mandujano et al., 2010). In addition, most studies of cactus pollination have mainly focused on a single plant species (Mandujano et al., 2010; Larrea-Alcázar & López, 2011; Egli & Giorgetta, 2015), ignoring the pollination mechanisms of sympatric species (Fleming, Tuttle & Horner, 1996; Egli & Giorgetta, 2017; Ferreira et al., 2018). Therefore, increasing our knowledge of pollination in sympatric cactus species allows us to identify potential sources of pollen limitation, assess reproductive isolation barriers, and determine mechanisms of seed production. These issues are relevant to identifying demographic bottlenecks in taxa, especially those with high extinction risks. As a way of contributing to filling these gaps in our knowledge of reproductive systems in sympatric plants, we present a study that included four cactus taxa belonging to the *Eriogyne* sect. *Neoporteria* that inhabit a highly threatened ecosystem in coastal central Chile. Specifically, our study attempted to answer the following key questions: (i) How similar are pollinator assemblages among sympatric cactus species? (ii) Which mating systems do sympatric cactus species use?

Since flower morphology potentially predicts the group of pollinators that visit plants (Fenster, Martén-Rodríguez & Schemske, 2009), we expect that *Eriogyne* taxa with funnel-shaped flowers (i.e., *Eriogyne chilensis*, *E. chilensis* var. *albidiflora* and *E. curvispina* var. *mutabilis*; Fig. 1) will be bee pollinated sharing pollinators. In contrast, species that have internal tepals curved inwards forming tubular flowers (i.e., *E. subgibbosa*, Fig. 1), a trait previously suggested as an adaptation to an ornithophilous pollination syndrome (Walter, 2008), may be hummingbird pollinated and thus should present substantial differences in its pollinator assemblage compared to the remaining taxa. We also expect some degree of self-compatibility in all taxa, as previously suggested in a previous study (Walter, 2008).



**Figure 1** Local distribution of the *Erioseye* species between Pichidangui Bay and Los Molles Peninsula, Chile. The colors of the dots on the map and within the insets refer to different taxa: (A) *E. chilensis* var. *albidiflora* (white), (B) *E. chilensis* (red), (C) *E. curvispina* var. *mutabilis* (orange) and (D) *E. subgibbosa* (blue). Map data © 2018 Google, Image DigitalGlobe, CNES / Airbus, Data SIO, NOAA, US Navy, NGA, GEBCO. [Full-size !\[\]\(5f471a71b78d7676bc356df190b88ab4\_img.jpg\) DOI: 10.7717/peerj.7865/fig-1](https://doi.org/10.7717/peerj.7865/fig-1)

## MATERIALS AND METHODS

### Study site

We carried out our study in a coastal strip of central Chile between Pichidangui Bay (32°08'S; 71°32'W) and Los Molles Peninsula (32°14'S; 71°30'W) (Fig. 1). This area is the most significant remnant of coastal Mediterranean-type scrubland; it is dominated by sclerophyllous vegetation (Luebert & Pliscoff, 2006; Alaniz, Galleguillos & Perez-Quezada, 2016) that is characterized by its high plant diversity, with 57% consisting of flowering species endemic to central Chile (Lund & Teillier, 2012). Today, 17 plant taxa from this area are in a threatened status (Lund & Teillier, 2012) due to anthropogenic pressures. The study was conducted in a publicly accessible area (no permit required).

### Study species

We studied all species of the genus *Erioseye* that are present in this area, which are sympatrically distributed with partial or full overlaps in their ranges (Fig. 1; Guerrero et al., 2011): *E. chilensis* (Hildm. ex K. Schum.) Katt., *E. chilensis* var. *albidiflora* (F. Ritter) Katt., *E. subgibbosa* (Haw.) Katt., and *E. curvispina* (Bertero ex Colla) Katt. var. *mutabilis* (F. Ritter) Katt. The first three species belong to the *Neoporteria* section, while the fourth species belongs to the *Horridocactus* section (Guerrero et al., 2019b). Two of these taxa

(*E. chilensis* and *E. chilensis* var. *albidiflora*) are narrow endemics that inhabit exclusively studied areas, and many other species in the family have reduced their potential distribution range through the effects of habitat loss in recent decades (Goettsch *et al.*, 2015). This situation has led to *E. chilensis* being classified as critically endangered (Faundez *et al.*, 2013). Despite having wide distributional ranges, *E. subgibbosa* and *E. curvispina* var. *mutabilis* only have an overlapping distribution in the studied site (Fig. 1). *Eriosyce chilensis* is a globose cactus that exclusively inhabits rocky outcrops near the coast. This taxon has two taxonomic varieties. The nominal species *E. chilensis* has flowers with pink tepals and occurs in all studied areas except in the northernmost portion (near Pichidanguí) (Fig. 1). The second variety is *E. chilensis* var. *albidiflora*, a local endemic that is distributed in the northernmost range of the species and has flowers with white tepals. Both taxonomic varieties of *E. chilensis* have flowers with tepals deflected outwards (funnel-shaped flowers; Fig. 1). *Eriosyce subgibbosa* is a globular-elongated cactus that is distributed on the Pacific coastline in a 600 km range between 31°S and 37°S and presents pink to pale pink tepals that form a narrow tube with the internal tepals curved inwards (tubular flowers; Fig. 1). These taxa occur in the northern and southern ranges of the study site and occupy coastal rocky outcrops and cliffs (Walter, 2008). *Eriosyce chilensis*, *E. chilensis* var. *albidiflora* and *E. subgibbosa* are very similar in their vegetative structures but display more substantial differences in flower morphology (Fig. 1). *Eriosyce subgibbosa* has a rather tubular-shaped hypanthium in comparison with the funnel-shaped flowers of *E. chilensis* and *E. chilensis* var. *albidiflora*. Besides, the latter two species differ in the pigmentation of their flowers, as *E. chilensis* can be seen as pink by human vision, while *E. chilensis* var. *albidiflora* is white and yellowish. Details of diagnostic characters for the determination of Chilean *Eriosyce* are available in Kattermann (1994). Finally, *Eriosyce curvispina* var. *mutabilis* is a globular cactus that is narrowly distributed from 32°S to 32°30'S in coastal terraces and valleys with oceanic influence. This species has a hemicryptic habit, living semi-buried (Kattermann, 1994). This species has showy flowers with outward tepals (similar to those of *E. chilensis*) that are pigmented yellow to orange (Fig. 1). The floral morphology of *E. curvispina* var. *mutabilis* suggest that this species is bee-pollinated.

### Visitors to *Eriosyce* flowers

We performed field observations to determine the main pollinators of each *Eriosyce* taxon during its peak period of flower anthesis: August (*E. subgibbosa*), October (*E. chilensis* and *E. chilensis* var. *albidiflora*) and November (*E. curvispina* var. *mutabilis*). Despite differences in peak flowering times among the species, there is some overlap between individuals in their floral phenology from August to December. Observations for each taxon were performed during two consecutive years (2016 and 2017) in 25 m<sup>2</sup> plots that contained clumps of 2–17 adult individuals ( $12.7 \pm 1.5$  individuals per plot, mean  $\pm$  standard error) with  $2.5 \pm 0.5$  flowers per individual. For each species, the plots were at least 20 m away from each other. We conducted observations on three consecutive days between 8:00 and 20:00 h in periods of 30 min per plot. Four observers were assigned to each species except for *E. curvispina* var. *mutabilis*, for which we had two

observers. Observation times coincided with the flower anthesis of each species; once opened, flowers remained in this state. Previous observations after 20:00 h led us to disregard the presence of nocturnal visitors. This methodology allowed us to cover 540 min of observation each day per observer. We performed a total of 6,480 min of observation per species (540 min  $\times$  4 observers  $\times$  3 consecutive days), except for *E. curvispina* var. *mutabilis*, for which we accumulated 3,240 min of observation.

When animals approached the flowers, we recorded the time they entered the flowers and collected the animals when feasible. Since we do not evaluate the effectiveness of animal visitors as pollinators, we considered all animals that entered the flowers and contacted reproductive parts (stigmas and anthers) to be pollinators. The collected specimens were pin-mounted and taken to the laboratory for later identification by Apoidea expert Dr. Luisa Ruz (Pontifical Catholic University of Valparaíso, Chile). Brochure specimens were deposited in the Entomology Collection of the University of Concepción (MZUC-UCCC), Chile.

Based on flower visits, we built a plant-pollinator matrix for each year. We estimated pollinator richness using matrix randomization ( $N = 9,999$  randomizations) based on models that find  $2 \times 2$  submatrices that can be swapped while holding the totals of the columns (pollinators) and rows (plant individuals) constant (“swap count” algorithm, [Gotelli & Entsminger, 2001](#); [Gotelli & Entsminger, 2003](#)). Additionally, we estimated the similarity of pollinator assemblages among *Eriosyce* cacti, both between years within each taxon and among taxa within each year, using the Bray–Curtis dissimilarity index ([Bray & Curtis, 1957](#)), this index ranges between 0 (two samples share all species) and 1 (two sites do not share any species). Since the Bray–Curtis index (hereafter *BC*) is a dissimilarity index, we reported similarity as  $1 - BC$ . All these analyses were performed using the library *vegan* ([Oksanen et al., 2018](#)) in R Software version 3.5.0 ([R Development Core Team, 2018](#)).

### Dependence from pollinators and mating system

We marked and bagged flower buds of each cactus species during its bloom peak: *E. subgibbosa* in August 2017 ( $N = 70$  individuals); *E. chilensis* var. *albidiflora* ( $N = 32$  individuals) and *E. chilensis* ( $N = 55$  individuals) in October 2016; and *E. curvispina* var. *mutabilis* ( $N = 75$  individuals) in November 2017. We used plastic silk fabric (200  $\times$  200 mm) to bag the flowers, which prevented animals from entering the flowers.

We monitored individuals daily, and as anthesis proceeded, we randomly assigned one of the following treatments to each individual: (i) manual cross-pollination, in which the flowers were pollinated with pollen from individuals located at least five m away from the focal individuals; (ii) automatic self-pollination, in which the flowers were kept bagged; and (iii) the control, in which the bags were removed on the day of anthesis and the flowers were unmanipulated (following [Eckert et al., 2010](#)). For manual cross-pollination, the donor individuals were 2-day-old flowers. Preliminary assays and the results of this study attest that this pollen is viable. Only one flower was treated per focal plant because the pollination of more than one flower may lead to resource limitations ([Charnov, 1979](#)), which may cause seed production to vary among individuals with a different number of treated flowers.

**Table 1** Pollinator richness and number of visits for four *Eriosyce* taxa in two consecutive years, central Chile.

Taxon	Year 2016		Year 2017	
	Species richness	Number of visits	Species richness	Number of visits
<i>E. chilensis</i>	4 (3–8)	32	8 (4–10)	74
<i>E. chilensis</i> var. <i>albidiflora</i>	7 (3–9)	32	4 (4–10)	97
<i>E. curvispina</i> var. <i>mutabilis</i>	7 (4–11)	85	6 (4–10)	118
<i>E. subgibbosa</i>	4 (3–9)	50	2 (2–7)	14
Total	11 (4–10)	199	12 (5–11)	303

**Note:**

Values in parentheses in species richness are 95% confidence intervals estimated after matrix randomization ( $N = 9,999$ ).

Before performing pollination experiments, we assessed the receptivity of stigma by using the bubble production method, which involves immersing the stigmas in a 3% hydrogen peroxide solution and observing bubble production for 2–3 min. This bubbling indicates the activity of stigmatic peroxidases and, hence, the receptivity of the stigma (Armbruster *et al.*, 2002). The receptivity of stigmas was assayed in 1- to 2-day-old flowers ( $N = 10$  flowers from different individuals of each species), and we observed receptivity in 100% of the flowers. All taxa of *Eriosyce* included in this study showed stigma receptivity even before anthesis (we assessed the receptivity of some stigmas in floral buds). We collected tagged fruits for counting seeds four weeks after applying the treatments.

We assessed differences in seed number among treatments for each plant species using a Kruskal–Wallis analysis (Sokal & Rohlf, 1995). When we found significant differences, we performed Nemenyi a posteriori tests (Zar, 1996).

## RESULTS

*Eriosyce* flowers were visited by 16 taxa during two consecutive years; 199 and 303 visits were observed in 2016 and 2017, respectively (Table 1; Appendix 1). In general, after blooming, *Eriosyce* flowers remained open for a period of 4–6 days. The *Eriosyce* species did not differ statistically in their pollinator richness (Table 1), and the similarity of pollinator assemblages was relatively low between year for each taxon (<0.5 of similarity; Table 2). Floral pollinators of *Eriosyce* belonged to two main groups: bees of the superfamily Apoidea (Hymenoptera) and the giant hummingbird *Patagona gigas* (Trochilidae: Apodiformes; Appendix 1). All observed bees were native species ( $N = 14$ ) except for *Apis mellifera* Linnaeus, 1758, with three visits accounted for only 0.9% of visits to *E. chilensis* var. *albidiflora* (Appendix 1).

*Eriosyce subgibbosa* showed the most distinctive pollinator assemblage compared to the remaining taxa (Table 2; Appendix 1), which could be attributed to visits from *P. gigas*, which exclusively visited this cactus species. Pollinator assemblage of *E. subgibbosa* was similar to the other *Eriosyce* species in 0% to 8.9% (year 2016) and 0% to 3.1% (year 2017; Table 2). Contribution of *Patagona gigas* to the total visits received by *E. subgibbosa* was 80% (2016) and 85% (2017). Remaining visits to *E. subgibbosa* were carried out by native bees (Appendix 1). The three other *Eriosyce* species, which have funnel-shaped floral

**Table 2** Similarity of pollinator assemblages (1—Bray–Curtis index) among the *Eriosyce* species during 2016 (below the diagonal) and 2017 (above the diagonal). The diagonal depicts intraspecific similarity between the years 2016 and 2017. The observed similarity indexes were contrasted with null models to test whether they were different from the expected by chance, values that deviated from null expectations are marked with asterisks (\* $P < 0.05$ ), the interval of null models are shown in parentheses.


	<i>E. chilensis</i> var. <i>albidiflora</i>	<i>E. chilensis</i>	<i>E. curvispina</i> var. <i>mutabilis</i>	<i>E. subgibbosa</i>
<i>E. chilensis</i> var. <i>albidiflora</i>	0.403 (0.050–0.851)	0.690 (0.070–0.837)	0.242 (0.027–0.857)	0.018 (0–0.851)
<i>E. chilensis</i>	0.844 (0.051–0.847)	0.453 (0.073–0.833)	0.104* (0.154–0.860)	0 (0–0.718)
<i>E. curvispina</i> var. <i>mutabilis</i>	0.291 (0.165–0.868)	0.342 (0.049–0.843)	0.187 (0.136–0.872)	0.030 (0–0.776)
<i>E. subgibbosa</i>	0.032* (0.064–0.833)	0 (0–0.862)	0.089 (0.050–0.852)	0.438 (0–0.811)

morphology, showed more similar floral assemblages within them, and hummingbird visits were not observed (Table 2). The most similar pollinator assemblages were observed between *E. chilensis* and *E. chilensis* var. *albidiflora* (84.4% and 69.1% for 2016 and 2017, respectively). *Eriosyce curvispina* var. *mutabilis* showed a distinctive pollinator assemblage compared with the other funnel-shaped species and exhibited the highest interannual variation in pollinator composition among all *Eriosyce* taxa (Table 2).

Results of the mating treatment in *E. subgibbosa* indicate significant pollen limitation since manually cross-pollinated flowers produced more seeds than unmanipulated flowers (Kruskal–Wallis  $\chi^2 = 6.987$ ;  $P < 0.031$ ; Fig. 2). Although it was not an overdispersed phenomenon, only one *E. subgibbosa* individual generated seeds under the automatic self-pollination treatment; thus, autonomous selfing cannot be excluded as a reproductive system in this species (Fig. 2). *Eriosyce* species with funnel-shaped flowers showed differences among their mating systems. *Eriosyce curvispina* var. *mutabilis* supported autonomous self-pollination and no differences in seed production were observed among compared treatments ( $\chi^2 = 0.637$ ;  $P = 0.727$ ; Fig. 2). *Eriosyce chilensis* showed allogamous pollination, which may be largely dependent on pollinators ( $\chi^2 = 9.91$ ;  $P < 0.001$ ; Fig. 2). In *E. chilensis* var. *albidiflora*, autonomous pollination had a marginal contribution, whereas allogamous pollination was potentially more important to seed production (Fig. 2;  $\chi^2 = 16.83$ ;  $P < 0.001$ ).

## DISCUSSION

We found contrasting pollinator assemblages among sympatric *Eriosyce* species; they exhibited floral morphology associated with the ornithophilous syndrome (*E. subgibbosa*) and the melitophilous syndrome (the remaining taxa). Interestingly, *E. subgibbosa* was the only species that interacted with the giant hummingbird (*Patagona gigas*), although it also received visits from some relatively small native bees (e.g., *Dialictus* species from Halictidae family were their most frequent visitors). As found in other studies in cacti, plants with ornithophilous syndromes do not necessarily restrict other pollinators from using the resources that they offer (Gorostiague & Ortega-Baes, 2016). The similarity of pollinator assemblages in species funnel-shaped flowers was not homogeneous since *E. chilensis* and *E. chilensis* var. *albidiflora* present the higher similarity while *E. curvispina* var. *mutabilis* was the most dissimilar funnel-shaped species.



**Figure 2** Reproductive output of the *Eriosyce* species after the pollination experiment. (A) Number of seeds (mean  $\pm$  standard error), (B) number of seed (standardized) and (C) proportion of fruits produced. For each panel, the bars indicate the results of the three pollination treatments: unmanipulated plants (Control), manually cross-pollinated plants (Cross) and automatically self-pollinated plants (Self). Taxa names were abbreviated as follows: *E. chilensis* var. *albidiflora* (alb), *E. chilensis* (chi), *E. curvispina* var. *mutabilis* (mut) and *E. subgibbosa* (sub). For seed production, we reported the statistical significance of the Kruskal–Wallis test that compared differences among treatments for each species ( $***P < 0.001$ ;  $**P < 0.01$ ;  $*P < 0.05$ ; ns, non-significant differences). Letters above seed number bars are the results of Nemenyi a posteriori tests; different letters indicate significant differences among compared treatments.

Full-size DOI: 10.7717/peerj.7865/fig-2

A substantial interannual variation in pollinator assemblage was observed, with more than three times fewer total visits (hummingbirds and hymenopterans) to *E. subgibbosa* during the 2nd year of our study. The presence of other plants with massive nectar production may reduce visits to *E. subgibbosa*, which produces less nectar compared to species of the genus *Puya* that are abundant in the study site (Walter, 2008; Hornung-Leoni, González-Gómez & Troncoso, 2013). Plants that produce massive amounts of nectar interfere in the pollination process of plants with less nectar production (Fleming et al., 2001; Chittka & Schürkens, 2001). For instance, *Puya chilensis* Mol. and *P. venusta* Phil. ex Baker. co-occur in the study site and exhibited a massive bloom during the year

2017 (P Guerrero, 2017, personal observations), suggesting a pollination interference against *E. subgibbosa*, because this last species have smaller nectar chambers and therefore produce less nectar (Walter, 2008). In addition, changes in pollinator frequency and diversity may promote reproductive constraints in *E. subgibbosa* during certain years. Indeed, we detected pollen limitation mediated by pollinators since manually cross-pollinated flowers produced a higher number of seeds than unmanipulated flowers. Pollen limitation is a widely represented phenomenon in angiosperms (Bennett et al., 2018), and it acts as a significant evolutionary driver since pollen limitation is a selection agent for mating systems (Totland, 2001). The pollination scenario in *E. subgibbosa* indicates a specialized and interannually erratic assemblage, suggesting a pollen limitation that could be hampered by the existence of selfing, which has been observed in other cactus species (Nassar & Ramírez, 2004). Also, together with natural interannual variation in pollinator visit frequencies, anthropogenic disturbances that occur at a site (e.g., changes in land use, desertification) may negatively impact pollinator visit frequencies in locally endemic cactus (Strong & Williamson, 2007; Martínez-Peralta & Mandujano, 2016). Our results support facultative selfing in *E. subgibbosa*, which may act as a reproductive assurance mechanism that may become increasingly important as the number of *Patagona gigas* visits decreases.

In our study, we found that *Eriosyce* species with funnel-shaped flowers were strictly bee-pollinated, which is consistent with their morphology and flowering peaks, which occur during mid-spring when the increasing day temperature drives increasing pollinator activity. Although the diversity of native bees in the xeric areas of Chile is relatively low compared to other mediterranean ecosystems (Michener, 1979), native bees are a crucial component of the pollination process (Montalva & Ruz, 2010; Medel, González-Browne & Fontúrbel, 2018). Previous studies on the same bees species that visited *Eriosyce* showed that they are generalists in the use of floral resources (Montalva & Ruz, 2010; Michener, 2007), such as *Alloscirtetica lanosa* (Apidae), *Diadasia chilensis* (Apidae), *Xenochilicola diminuta* (Colletidae) and *Caenohalictus cyanopygus* (Halictidae). Similarly, we observed a generalist behavior of bees that visited the studied species; this may favor heterospecific pollen transfer among taxa and may open a window of ecological opportunity for hybrid formation. Compared to the giant hummingbird, insects have been less studied, and thus ranges of Chilean bees are far from being understood. This state of the art is related to the fact that most of the distribution ranges of Chilean bees are only reported at latitudinal level (Montalva & Ruz, 2010). In addition, several Apoidea pollinators observed in this study seem to correspond to new native bee species (i.e., *Anthrenoides* sp. 1, *Liphanthus* sp. 1, L Ruz, 2018, personal communications.) whose distributional ranges and ecological associations have not yet been described. In this sense, the distribution, local abundances, and foraging preferences of the observed bees are topics that require more attention. Otherwise, the role of native bee species in plant reproduction in central Chile will remain deficient.

Bee-pollinated *Eriosyce* species, which consistently displayed funnel-shaped flowers, exhibited variation on pollinator dependence for their reproductive success. Although self-incompatibility has been reported in at least 31 genera of cacti (Boyle, 1997), including

some *Copiapoa* and *Eriosyce* species (Ganders, 1976). We observed some degree of autonomous reproduction in *E. chilensis* var. *albidiflora* and *E. curvispina* var. *mutabilis*, this was expected since a previous study highlighted that species of the *Eriosyce* subsection *Neoporteria* have some degree of self-compatibility (Walter, 2008). Our result is coherent with a recent meta-analysis that showed that mixed mating is frequent among angiosperms (Whitehead et al., 2018). Also, our study supports the existence of a mixed reproductive system in cacti (i.e., *E. curvispina* var. *mutabilis*), that may act as reproductive backup under erratic pollinator environments. It is interesting to observe that funnel-shaped flowers that exhibited higher similarity in their pollinator assemblages (*E. chilensis* and *E. chilensis* var. *albidiflora*), both taxa also occupy the same microhabitat-type (i.e., coastal rocky outcrops and cliffs; Guerrero et al., 2011). Microhabitat conditions may be crucial for maintaining pollinator consistency among taxa and across years (Schatz, 2006; Adedoja et al., 2019). *Eriosyce curvispina* var. *mutabilis* that inhabit coastal terraces showed marked differences in their pollinator assemblage and a mixed mating system, supporting both allogamous and autogamous strategies. Coastal terraces are sites with harsh pollination conditions since the open spaces are subject to wind gusts, pollinator predators may be present, and there may be more frequent anthropogenic disturbances (Carvallo et al., 2019). Under this scenario, it is likely that *E. curvispina* var. *mutabilis* evolved a mixed mating system with some degree of autonomous selfing (Eckert et al., 2010). This may allow *E. curvispina* var. *mutabilis* to maintain some level of seed production during pollinator-depauperate seasons.

The observed variation in mating systems among the sympatric *Eriosyce* species that we studied raises scientific questions about how the pollination syndromes evolved, and whether mating system evolution played a role in the evolutionary origin of the narrow endemic *E. chilensis*. This because species belonging to the *Neoporteria* clade are characterized by an ornithophilous syndrome, but our study revealed strict melitophilous reproduction. This result and the phylogenetic topology of the clade (Guerrero et al., 2019b), suggest the reversal of a plesiomorphic state in the *Neoporteria* from bird pollination back to bee pollination. Previous studies in the Cactaceae have revealed that changes in pollination systems can increase diversification rates by promoting speciation (Schlumpberger & Renner, 2012; Hernández-Hernández et al., 2014). Finally, our study highlights the necessity of further studies that could test the existence and potential effects of inbreeding depression, particularly in species with some degree of autogamous reproduction. Inbreeding depression must be considered in the context of a disturbed landscape because it can promote population and genetic bottlenecks, especially in narrowly endemic species such as those included in this study.

## CONCLUSIONS

We found a close relationship between floral morphology and pollinator assemblages since funnel-shaped flowers were mainly visited by small-sized native bee species and the tubular-shaped flower was visited by the giant hummingbird. The higher similarity in pollinator assemblages within bee pollinated taxa may contribute to the hybrid formation, but to test whether this process is effectively occurring in nature, new analyses based on

molecular data must be performed. Also, our study contributes to highlighting the potential role of self-compatibility when pollinators are scarce or unpredictable as a mechanism of reproductive assurance. However, longer observations are required to test this hypothesis, this mechanism could be critical in the long term persistence since the reduction of pollinator abundance may have negative impacts in plant reproduction success. This is an important issue, especially in the cactus family, which is a threatened group at a global scale that includes several taxa with a narrow distribution.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This study was funded by Fondo Nacional de Desarrollo Científico y Tecnológico FONDECYT, Chile (Grants 1160583 to PCG and 11150301 to GOC), and by Fondo de Investigación del Bosque Nativo (FIBN) (Grant CONAF-009/2015). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

Fondo Nacional de Desarrollo Científico y Tecnológico FONDECYT: 1160583 to PCG and 11150301 to GOC.

Fondo de Investigación del Bosque Nativo (FIBN): CONAF-009/2015.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Pablo C. Guerrero conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, authored or reviewed drafts of the paper, approved the final draft.
- Claudia A. Antinao performed the experiments, analyzed the data, approved the final draft.
- Beatriz Vergara-Meriño performed the experiments, approved the final draft.
- Cristian A. Villagra performed the experiments, authored or reviewed drafts of the paper, approved the final draft.
- Gastón O. Carvallo conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.

### Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

No permit required for doing field work on coastal area.

## Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplemental Files](#).

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.7865#supplemental-information>.

## REFERENCES

- Adedoja O, Dormann CF, Kehinde T, Samways MJ. 2019.** Refuges from fire maintain pollinator-plant interaction networks. *Ecology and Evolution* **9(10)**:5777–5786 DOI [10.1002/ece3.5161](https://doi.org/10.1002/ece3.5161).
- Alaniz AJ, Galleguillos M, Perez-Quezada JF. 2016.** Assessment of quality of input data used to classify ecosystems according to the IUCN Red List methodology: the case of the central Chile hotspot. *Biological Conservation* **204**:378–385 DOI [10.1016/j.biocon.2016.10.038](https://doi.org/10.1016/j.biocon.2016.10.038).
- Armbruster WS, Mulder CPH, Baldwin BG, Kalisz S, Wessa B, Nute H. 2002.** Comparative analysis of late floral development and mating-system evolution in tribe Collinsieae (Scrophulariaceae s.l.). *American Journal of Botany* **89(1)**:37–49 DOI [10.3732/ajb.89.1.37](https://doi.org/10.3732/ajb.89.1.37).
- Barrett SCH. 1998.** The evolution of mating strategies in flowering plants. *Trends in Plant Science* **3(9)**:335–341 DOI [10.1016/S1360-1385\(98\)01299-0](https://doi.org/10.1016/S1360-1385(98)01299-0).
- Barrett SCH, Harder LD. 2017.** The ecology of mating and its evolutionary consequences in seed plants. *Annual Review of Ecology, Evolution, and Systematics* **48(1)**:135–157 DOI [10.1146/annurev-ecolsys-110316-023021](https://doi.org/10.1146/annurev-ecolsys-110316-023021).
- Bennett JM, Steets JA, Burns JH, Durka W, Vamosi JC, Arceo-Gómez G, Burd M, Burkle LA, Ellis AG, Freitas L, Li J, Rodger JG, Wolowski M, Xia J, Ashman T-L, Knight TM. 2018.** GloPL, a global data base on pollen limitation of plant reproduction. *Scientific Data* **5(1)**:180249 DOI [10.1038/sdata.2018.249](https://doi.org/10.1038/sdata.2018.249).
- Bodbyl Roels SA, Kelly JK. 2011.** Rapid evolution caused by pollinator loss in *Mimulus guttatus*. *Evolution* **65(9)**:2541–2552 DOI [10.1111/j.1558-5646.2011.01326.x](https://doi.org/10.1111/j.1558-5646.2011.01326.x).
- Boyle TH. 1997.** The genetics of self-incompatibility in the genus *Schlumbergera* (Cactaceae). *Journal of Heredity* **88(3)**:209–214 DOI [10.1093/oxfordjournals.jhered.a023090](https://doi.org/10.1093/oxfordjournals.jhered.a023090).
- Bray JR, Curtis JT. 1957.** An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs* **27(4)**:325–349 DOI [10.2307/1942268](https://doi.org/10.2307/1942268).
- Carvallo GO, Vergara-Meriño B, Díaz A, Villagra CA, Guerrero PC. 2019.** Rocky outcrops conserve genetic diversity and promote regeneration of a threatened relict tree in a critically endangered ecosystem. *Biodiversity and Conservation* **28(11)**:2805–2824 DOI [10.1007/s10531-019-01797-6](https://doi.org/10.1007/s10531-019-01797-6).
- Charnov EL. 1979.** Simultaneous hermaphroditism and sexual selection. *Proceedings of the National Academy of Sciences of the United States of America* **76(5)**:2480–2484 DOI [10.1073/pnas.76.5.2480](https://doi.org/10.1073/pnas.76.5.2480).
- Chittka L, Schürkens S. 2001.** Successful invasion of a floral market. *Nature* **411(6838)**:653 DOI [10.1038/35079676](https://doi.org/10.1038/35079676).
- Eckert CG, Herlihy CR. 2004.** Using a cost-benefit approach to understand the evolution of self-fertilization in plants: the perplexing case of *Aquilegia canadensis* (Ranunculaceae). *Plant Species Biology* **19(3)**:159–173 DOI [10.1111/j.1442-1984.2004.00112.x](https://doi.org/10.1111/j.1442-1984.2004.00112.x).
- Eckert CG, Kalisz S, Geber MA, Sargent R, Elle E, Cheptou P-O, Goodwillie C, Johnston MO, Kelly JK, Moeller DA, Porcher E, Ree RH, Vallejo-Marín M, Winn AA. 2010.** Plant mating

- systems in a changing world. *Trends in Ecology & Evolution* **25**(1):35–43  
DOI 10.1016/j.tree.2009.06.013.
- Eckhart VM. 1992. Spatio-temporal variation in abundance and variation in foraging behavior of the pollinators of gynodioecious *Phacelia linearis* (Hydrophyllaceae). *Oikos* **64**(3):573–586  
DOI 10.2307/3545177.
- Eggl U, Giorgetta M. 2015. Flowering phenology and observations on the pollination biology of South American cacti. 1. *Denmoza rhodacantha*. *Haseltonia* **20**:3–12 DOI 10.2985/026.020.0103.
- Eggl U, Giorgetta M. 2017. Flowering phenology and observations on the pollination biology of South American cacti. 3. Temporally robust sequential Big Bang flowering of two unrelated sympatric globular cacti in Northern Argentina. *Haseltonia* **23**:97–109  
DOI 10.2985/026.023.0113.
- Faundez L, Guerrero PC, Saldivia P, Walter HE. 2013. *Eriosyce chilensis*. The IUCN Red List of Threatened Species 2013. Available at <https://www.iucnredlist.org/species/151924/576839> (accessed 8 January 2019).
- Fenster CB, Armbruster WS, Wilson P, Dudash MR, Thomson JD. 2004. Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution, and Systematics* **35**(1):375–403  
DOI 10.1146/annurev.ecolsys.34.011802.132347.
- Fenster CB, Martín-Rodríguez S, Schemske DW. 2009. Evolution of floral diversity in *Iochroma* (Solanaceae). *Evolution* **63**(10):2758–2762 DOI 10.1111/j.1558-5646.2009.00730.x.
- Ferreira BHS, Gomes AC, Souza CS, Fabri JR, Sigrist MR. 2018. Pollination and reproductive system of synchronopatric species of Cactaceae (Cactoideae) subject to interspecific flow of pollen: an example of ecological adaptation in the Brazilian Chaco. *Plant Biology* **20**(1):101–112  
DOI 10.1111/plb.12641.
- Fleming TH, Sahley CT, Holland JN, Nason JD, Hamrick JL. 2001. Sonoran desert columnar cacti and the evolution of generalized pollination systems. *Ecological Monographs* **71**(4):511–530  
DOI 10.1890/0012-9615(2001)071[0511:SDCCAT]2.0.CO;2.
- Fleming TH, Tuttle MD, Horner MA. 1996. Pollination biology and the relative importance of nocturnal and diurnal pollinators in three species of sonoran desert columnar cacti. *Southwestern Naturalist* **41**:257–269.
- Ganders FR. 1976. Self-incompatibility in the Cactaceae. *Cactus and Succulent Journal* **38**:39–40.
- Goettsch B, Hilton-Taylor C, Cruz-Piñón G, Duffy JP, Frances A, Hernández HM, Inger R, Pollock C, Schiér J, Superina M, Taylor N, Tognelli M, Abba AM, Arias S, Arreola-Nava HJ, Baker MA, Bárcenas RT, Barrios D, Braun P, Butterworth CA, Búrquez A, Caceres F, Chazaro-Basañez M, Corral-Díaz R, Del Valle Perea M, Demaio PH, Barros WAD, Durán R, Yancas LF, Felger RS, Fitz-Maurice B, Fitz-Maurice WA, Gann G, Gómez-Hinostrosa C, Gonzales-Torres LR, Griffith MP, Guerrero PC, Hammel B, Heil KD, Hernández-Oria JG, Hoffmann M, Ishihara MI, Keisling R, Larocca J, León-De La Luz JL, Loaiza CR, Lowry M, Machado MC, Majure LC, Martínez Ávalos JG, Martorell C, Maschinski J, Méndez E, Mittermeier RA, Nassar JM, Negrón-Ortiz V, Oakley LJ, Ortega-Baes P, Pin Ferreira AB, Pinkava DJ, Porter JM, Puente-Martínez R, Smith M, Manuel Sotomayorel MDCJ, Stuart SN, Tapia Muñoz JL, Terrazas T, Terry M, Trevisson M, Valverde T, Van Devender TR, Véliz-Pérez ME, Walter HE, Wyatt SA, Zappi D, Zavala-Hurtado JA, Gaston KJ. 2015. High proportion of cactus species threatened with extinction. *Nature Plants* **1**(10):15142 DOI 10.1038/nplants.2015.142.
- Goodwillie C, Kalisz S, Eckert CG. 2005. The evolutionary enigma of mixed mating systems in plants: occurrence, theoretical explanations, and empirical evidence. *Annual Review of Ecology, Evolution, and Systematics* **36**(1):47–79 DOI 10.1146/annurev.ecolsys.36.091704.175539.

- Gorostiague P, Ortega-Baes P. 2016.** How specialised is bird pollination in the Cactaceae? *Plant Biology* **18**(1):63–72 DOI [10.1111/plb.12297](https://doi.org/10.1111/plb.12297).
- Gotelli NJ, Entsminger GL. 2001.** Swap and fill algorithms in null model analysis: rethinking the knight's tour. *Oecologia* **129**(2):281–291 DOI [10.1007/s004420100717](https://doi.org/10.1007/s004420100717).
- Gotelli NJ, Entsminger GL. 2003.** Swap algorithms in null model analysis. *Ecology* **84**(2):532–535 DOI [10.1890/0012-9658\(2003\)084\[0532:SAINMA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[0532:SAINMA]2.0.CO;2).
- Guerrero PC, Arroyo MTK, Bustamante RO, Duarte M, Hagemann TK, Walter HE. 2011.** Phylogenetics and predictive distribution modeling provide insights into the geographic divergence of *Eriosyce* subgen. *Neoporteria* (Cactaceae). *Plant Systematics and Evolution* **297**(1–2):113–128 DOI [10.1007/s00606-011-0512-5](https://doi.org/10.1007/s00606-011-0512-5).
- Guerrero PC, Carvallo GO, Nassar JM, Rojas-Sandoval J, Sanz V, Medel R. 2012.** Ecology and evolution of negative and positive interactions in Cactaceae: lessons and pending tasks. *Plant Ecology & Diversity* **5**(2):205–215 DOI [10.1080/17550874.2011.630426](https://doi.org/10.1080/17550874.2011.630426).
- Guerrero PC, Majure LC, Cornejo-Romero A, Hernández-Hernández T. 2019a.** Phylogenetic relationships and evolutionary trends in the Cactus family. *Journal of Heredity* **110**(1):4–21 DOI [10.1093/jhered/esy064](https://doi.org/10.1093/jhered/esy064).
- Guerrero PC, Walter HE, Arroyo MTK, Peña CM, Tamburrino I, De Benedictis M, Larridon I. 2019b.** Molecular phylogeny of the large South American genus *Eriosyce* (Notocactaceae, Cactaceae): generic delimitation and proposed changes in infrageneric and species ranks. *Taxon, Early View* DOI [10.1002/tax.12066](https://doi.org/10.1002/tax.12066).
- Hadley AS, Betts MG. 2009.** Tropical deforestation alters hummingbird movement patterns. *Biology Letters* **5**(2):207–210 DOI [10.1098/rsbl.2008.0691](https://doi.org/10.1098/rsbl.2008.0691).
- Herlihy CR, Eckert CG. 2005.** Evolution of self-fertilization at geographical range margins? A comparison of demographic, floral, and mating system variables in central vs. peripheral populations of *Aquilegia canadensis* (Ranunculaceae). *American Journal of Botany* **92**(4):744–751 DOI [10.3732/ajb.92.4.744](https://doi.org/10.3732/ajb.92.4.744).
- Hernández-Hernández T, Brown JW, Schlumpberger BO, Eguiarte LE, Magallón S. 2014.** Beyond aridification: multiple explanations for the elevated diversification of cacti in the New World Succulent Biome. *New Phytologist* **202**(4):1382–1397 DOI [10.1111/nph.12752](https://doi.org/10.1111/nph.12752).
- Hornung-Leoni CT, González-Gómez PL, Troncoso AJ. 2013.** Morphology, nectar characteristics and avian pollinators in five Andean *Puya* species (Bromeliaceae). *Acta Oecologica* **51**:54–61 DOI [10.1016/j.actao.2013.05.010](https://doi.org/10.1016/j.actao.2013.05.010).
- Johnson SD. 2010.** The pollination niche and its role in the diversification and maintenance of the southern African flora. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**(1539):499–516 DOI [10.1098/rstb.2009.0243](https://doi.org/10.1098/rstb.2009.0243).
- Johnson SD, Steiner K. 2000.** Generalization versus specialization in plant pollination systems. *Trends in Ecology & Evolution* **15**(4):140–143 DOI [10.1016/S0169-5347\(99\)01811-X](https://doi.org/10.1016/S0169-5347(99)01811-X).
- Kalish S, Vogler DW. 2003.** Benefits of autonomous selfing under unpredictable pollinator environments. *Ecology* **84**(11):2928–2942 DOI [10.1890/02-0519](https://doi.org/10.1890/02-0519).
- Kalish S, Vogler DW, Hanley KM. 2004.** Context-dependent autonomous self-fertilization yields reproductive assurance and mixed mating. *Nature* **430**(7002):884–887 DOI [10.1038/nature02776](https://doi.org/10.1038/nature02776).
- Kattermann F. 1994.** *Eriosyce* (Cactaceae): the genus revised and amplified. *Succulent Plant Research 1*. Richmond: Privately published by David Hunt.
- Kearns CA, Inouye DW, Waser NM. 1998.** Endangered mutualisms: the conservation of plant-pollinator interactions. *Annual Review of Ecology and Systematics* **29**(1):83–112 DOI [10.1146/annurev.ecolsys.29.1.83](https://doi.org/10.1146/annurev.ecolsys.29.1.83).

- Knight TM, Steet JA, Vamosi JC, Mazer SJ, Burd M, Campbell DR, Dudash MR, Johnston MO, Mitchell RJ, Ashman TL. 2005.** Pollen limitation of plant reproduction: pattern and process. *Annual Review of Ecology, Evolution, and Systematics* 36(1):467–497 DOI 10.1146/annurev.ecolsys.36.102403.115320.
- Kudo G, Kasagi T. 2005.** Microscale variations in the mating system and heterospecific incompatibility mediated by pollination competition in alpine snowbed plants. *Plant Species Biology* 20(2):93–103 DOI 10.1111/j.1442-1984.2005.00129.x.
- Larrea-Alcázar DM, López RP. 2011.** Pollination biology of *Oreocereus celsianus* (Cactaceae), a columnar cactus inhabiting the high subtropical Andes. *Plant Systematics and Evolution* 295(1–4):129–137 DOI 10.1007/s00606-011-0485-4.
- Luebert F, Pliscoff P. 2006.** *Sinopsis bioclimática y vegetacional de Chile*. Santiago de Chile: Editorial Universitaria, 310.
- Lund R, Teillier S. 2012.** Flora vascular de Los Molles, Región de Valparaíso, Chile. *Chloris Chilensis* 2(2). Available at <http://www.chlorischile.cl/Lund-Los%20Molles/flora%20de%20Los%20Molles.htm> (accessed 16 March 2019).
- Mandujano MC, Carrillo-Ángeles I, Martínez-Peralta C, Golubov J. 2010.** Reproductive biology of Cactaceae. In: Ramawat K, ed. *Desert Plants*. Berlín, Heidelberg: Springer.
- Marques I, Roselló-Graell A, Draper D, Iriando JM. 2007.** Pollination patterns limit hybridization between two sympatric species of *Narcissus* (Amaryllidaceae). *American Journal of Botany* 94(8):1352–1359 DOI 10.3732/ajb.94.8.1352.
- Martínez-Peralta C, Mandujano MC. 2016.** Pollen limitation in a rare cactus with synchronous mass flowering. *Haseltonia* 22:2–8 DOI 10.2985/026.022.0102.
- Martínez-Peralta C, Márquez-Guzmán J, Mandujano MC. 2014.** How common is self-incompatibility across species of the herkogamous genus *Ariocarpus*? *American Journal of Botany* 101(3):530–538 DOI 10.3732/ajb.1400022.
- McIntosh ME. 2002.** Plant size, breeding system, and limits to reproductive success in two sister species of *Ferocactus* (Cactaceae). *Plant Ecology* 162(2):273–288 DOI 10.1023/A:1020329718917.
- Medel R, González-Browne C, Fontúrbel FE. 2018.** Pollination in the Chilean Mediterranean-type ecosystem: a review of current advances and pending tasks. *Plant Biology* 20:89–99 DOI 10.1111/plb.12644.
- Michener CD. 1979.** Biogeography of the bees. *Annals of the Missouri Botanical Garden* 66(3):277–347 DOI 10.2307/2398833.
- Michener CD. 2007.** *The bees of the world*. Baltimore: Johns Hopkins, 972.
- Moeller DA, Geber MA. 2005.** Ecological context of the evolution of self-pollination in *Clarkia xantiana*: population size, plant communities, and reproductive assurance. *Evolution* 59(4):786–799 DOI 10.1554/04-656.
- Montalva J, Ruz L. 2010.** Actualización de la lista sistemática de las abejas chilenas (Hymenoptera: Apoidea). *Revista Chilena de Entomología* 35:15–52.
- Morgan MT, Wilson WG, Knight TM. 2005.** Plant population dynamics, pollinator foraging, and the selection of self-fertilization. *American Naturalist* 166(2):169–183 DOI 10.1086/431317.
- Nassar JM, Ramírez N. 2004.** Reproductive biology of the melon cactus, *Melocactus curvispinus* (Cactaceae). *Plant Systematics and Evolution* 248(1–4):31–44 DOI 10.1007/s00606-004-0193-4.
- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O’Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2018.** *vegan*: Community Ecology Package. R package version 2.5-2. Available at <https://CRAN.R-project.org/package=vegan>.

- Pérez MF, Arroyo MTK, Armesto JJ. 2009. Evolution of autonomous selfing accompanies increased specialization in the pollination system of *Schizanthus* (Solanaceae). *American Journal of Botany* **96**(6):1168–1176 DOI [10.3732/ajb.0800306](https://doi.org/10.3732/ajb.0800306).
- R Development Core Team. 2018. *The R project for statistical computing*. Vienna: R Foundation for Statistical Computing. Available at <https://www.R-project.org>.
- Schatz B. 2006. Fine scale distribution of pollinators explains the occurrence of the natural orchid hybrid X *Orchis bergonii*. *Ecoscience* **13**(1):111–118 DOI [10.2980/1195-6860](https://doi.org/10.2980/1195-6860).
- Schlumpberger BO, Cocucci AA, Moré M, Sérsic AN, Raguso RA. 2009. Extreme variation in floral characters and its consequences for pollinator attraction among populations of an Andean cactus. *Annals of Botany* **103**(9):1489–1500 DOI [10.1093/aob/mcp075](https://doi.org/10.1093/aob/mcp075).
- Schlumpberger BO, Renner SS. 2012. Molecular phylogenetics of *Echinopsis* (Cactaceae): polyphyly at all levels and convergent evolution of pollination modes and growth forms. *American Journal of Botany* **99**(8):1335–1349 DOI [10.3732/ajb.1100288](https://doi.org/10.3732/ajb.1100288).
- Schlüter PM, Ruas PM, Kohl G, Ruas CF, Stuessy TF, Paulus HF. 2009. Genetic patterns and pollination in *Ophrys iricolor* and *O. mesaritica* (Orchidaceae): sympatric evolution by pollinator shift. *Botanical Journal of the Linnean Society* **159**(4):583–598 DOI [10.1111/j.1095-8339.2009.00957.x](https://doi.org/10.1111/j.1095-8339.2009.00957.x).
- Sokal R, Rohlf FJ. 1995. *Biometry*. Third Edition. New York: Freeman.
- Stone GN, Willmer P, Rowe JA. 1988. Partitioning of pollinators during flowering in an African community *Acacia* community. *Ecology* **79**(8):2808–2827 DOI [10.2307/176518](https://doi.org/10.2307/176518).
- Strong AW, Williamson PS. 2007. Breeding system of *Astrophytum asterias*: an endangered cactus. *Southwestern Naturalist* **52**(3):341–346 DOI [10.1894/0038-4909\(2007\)52\[341:BSOAAA\]2.0.CO;2](https://doi.org/10.1894/0038-4909(2007)52[341:BSOAAA]2.0.CO;2).
- Totland Ø. 2001. Environment-dependent pollen limitation and selection on floral traits in an alpine species. *Ecology* **82**(8):2233–2244 DOI [10.1890/0012-9658\(2001\)082\[2233:EDPLAS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[2233:EDPLAS]2.0.CO;2).
- Valiente-Banuet A, Rojas-Martínez A, Casas A, Del Coro Arizmendi M, Dávila P. 1997. Pollination biology of two winter-blooming giant columnar cacti in the Tehuacán Valley, central Mexico. *Journal of Arid Environments* **37**(2):331–341 DOI [10.1006/jare.1997.0267](https://doi.org/10.1006/jare.1997.0267).
- Van Der Niet T, Johnson SD, Linder HP. 2006. Macroevolutionary data suggest a role for reinforcement in pollination system shifts. *Evolution* **60**(8):1596–1601 DOI [10.1554/05-705.1](https://doi.org/10.1554/05-705.1).
- Verdú M, González-Martínez SC, Montilla AI, Mateu I, Pannell JR. 2006. Ovule discounting in an outcrossing, cryptically dioecious tree. *Evolution* **60**(10):2056–2063 DOI [10.1111/j.0014-3820.2006.tb01843.x](https://doi.org/10.1111/j.0014-3820.2006.tb01843.x).
- Walter HE. 2008. Floral biology, phylogeography and systematics of *Eriosyce* subgenus *Neoporteria* (Cactaceae). *Bradleya* **26**(26):75–98 DOI [10.25223/brad.n26.2008.a6](https://doi.org/10.25223/brad.n26.2008.a6).
- Walter HE. 2010. Floral biology of *Echinopsis chiloensis* ssp. *chiloensis* (Cactaceae): evidence for a mixed pollination syndrome. *Flora—Morphology, Distribution, Functional Ecology of Plants* **205**(11):757–763 DOI [10.1016/j.flora.2009.12.038](https://doi.org/10.1016/j.flora.2009.12.038).
- Weber JJ. 2017. The messenger matters: Pollinator functional group influences mating system dynamics. *Molecular Ecology* **26**(16):4113–4115 DOI [10.1111/mec.14191](https://doi.org/10.1111/mec.14191).
- Whitehead MR, Lanfear R, Mitchell RJ, Karron JD. 2018. Plant mating systems often vary widely among populations. *Frontiers in Ecology and Evolution* **6**:1–9 DOI [10.3389/fevo.2018.00038](https://doi.org/10.3389/fevo.2018.00038).
- Young HS, McCauley DJ, Galetti M, Dirzo R. 2016. Patterns, causes, and consequences of anthropocene defaunation. *Annual Review of Ecology, Evolution, and Systematics* **47**(1):333–358 DOI [10.1146/annurev-ecolsys-112414-054142](https://doi.org/10.1146/annurev-ecolsys-112414-054142).
- Zar JH. 1996. *Biostatistical analysis*. Upper Saddle River: Prentice Hall.