

# Ridini­lazole: a novel antimicrobial for *Clostridium difficile* infection

Jonathan C. Cho<sup>a</sup>, Matthew P. Crotty<sup>b</sup>, Joe Pardo<sup>c</sup>

The University of Texas at Tyler, Tyler, TX; Methodist Dallas Medical Center, Dallas, TX; North FL/South GA Veterans Health System, Gainesville, FL, USA

## Abstract

*Clostridium difficile* (*C. difficile*) infection remains a global healthcare threat worldwide and the limited options available for its treatment are of particular concern. Ridini­lazole is one potential future agent, as it demonstrates rapid bactericidal activity against *C. difficile*. Current studies show that ridini­lazole has a lower propensity for collateral damage to the gut microbiome and appears to diminish the production of *C. difficile* toxins. Results from phase II studies demonstrate that patients receiving ridini­lazole had a higher sustained clinical response compared with patients receiving vancomycin (66.7% vs. 42.4%;  $P=0.0004$ ). Adverse reactions were similar between ridini­lazole and vancomycin (40% vs. 56%, respectively), with most being gastrointestinal-related. Nausea (20%) and abdominal pain (12%) were the most commonly reported adverse reactions associated with ridini­lazole. Phase II study results are promising and future availability of phase III trial results will help further delineate the role and value of ridini­lazole.

**Keywords** Ridini­lazole, *Clostridium difficile*, infectious diarrhea

*Ann Gastroenterol* 2019; 32 (2): 134-140

## Introduction

*Clostridium difficile* (*C. difficile*) is recognized as an urgent threat to human health and represents an extremely challenging pathogen, given its impact on the healthcare system [1-5]. *C. difficile* infection (CDI) is recognized as the most common healthcare-associated infection and has spread across the globe [1,6-9]. Efforts in antimicrobial stewardship and infection prevention have demonstrated effectiveness in reducing the burden of CDI [10-13]. Unfortunately, CDI remains a looming presence in modern healthcare. One of the most challenging

aspects of managing patients with *C. difficile* is the pathogen's propensity to cause recurrent infections. Recurrence rates of up to 25% have been reported following treatment with metronidazole or vancomycin [14,15]. The vicious cycle of recurrence continues further, with patients who have one recurrence being at increased risk for another [16].

Treatment of acute CDI is currently limited to the antibiotics metronidazole, vancomycin and fidaxomicin, with bezlotoxumab being an option as adjunctive therapy. Metronidazole has largely fallen out of favor because of potential issues of resistance and inferior clinical response [17]. Vancomycin is currently utilized as first-line therapy; however, recurrence remains a substantial issue with this agent [14,17,18]. Fidaxomicin has demonstrated similar efficacy to that of vancomycin with the benefit of decreased recurrence; however, this agent may not be as effective against the fluoroquinolone-resistant BI/NAP1/027 strain (ribotype 027), the predominant ribotype in North America [19-21]. In view of the problematic nature of currently available treatment options, ridini­lazole represents a potential welcome addition to the armamentarium. This paper will review the pharmacology, spectrum of activity, pharmacokinetics, pharmacodynamics, clinical trials and safety of ridini­lazole.

<sup>a</sup>College of Pharmacy, The University of Texas at Tyler, Tyler, TX (Jonathan C. Cho); <sup>b</sup>Department of Pharmacy, Methodist Dallas Medical Center, Dallas, TX (Matthew P. Crotty); <sup>c</sup>Department of Pharmacy, North FL/South GA Veterans Health System, Gainesville, FL (Joe Pardo), USA

Conflict of Interest: JCC serves on the speakers' bureau for Allergan. MPC served as a consultant for Theravance and Nabriva. The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans Affairs or the United States government

Correspondence to: Jonathan C. Cho, Pharm D, MBA, BCPS, Clinical Assistant Professor, College of Pharmacy, The University of Texas at Tyler, 3900 University Blvd., Tyler TX, 75799, USA, e-mail: jcho@uttyler.edu

Received 27 July 2018; accepted 7 November 2018; published online 13 December 2018

DOI: <https://doi.org/10.20524/aog.2018.0336>

## Data sources

To gather relevant published information, a literature search was performed using PubMed, EMBSCO and Google Scholar electronic databases for relevant publications written

in the English language. Search terms included ridini­lazole, SMT19969, and *C. difficile*. Information was also gathered from abstracts from the Interscience Conference on Antimicrobial Agents and Chemotherapy, IDWeek, American Society for Microbiology Microbe, and the European Congress of Clinical Microbiology and Infectious Diseases.

## Chemistry and pharmacology

Ridini­lazole, previously known as SMT19969, is a novel antibacterial currently under development for the treatment of *C. difficile*. Its chemical name is 2,2'-bis(4-pyridyl)3H,3'H 5,5'-bibenzimidazole. The agent has a unique mechanism of action and is thought to interfere with cell division. It demonstrates rapid bactericidal activity against *C. difficile* [22,23]. A cluster of *C. difficile* genes involved in cell division were found to display an altered expression following exposure to ridini­lazole in a transcriptomic analysis. These findings, combined with previous work demonstrating a filamentous phenotype of *C. difficile* upon exposure to ridini­lazole, suggest that the agent works by targeting cell growth [22,24]. Preclinical animal studies demonstrated negligible systemic exposure of the orally administered agent [25]. Following oral administration of 200 mg b.i.d., ridini­lazole is minimally absorbed from the gastrointestinal lumen, resulting in low systemic levels and high fecal concentrations [26]. The specificity of activity targeting *C. difficile* and limiting collateral damage to the gut microbiota, along with its minimal systemic absorption, make ridini­lazole an exciting agent for treating CDI.

## Microbiologic activity

### Activity against *C. difficile*

Ridini­lazole has demonstrated potent activity against a breadth of *C. difficile* strains *in vitro* [27-32]. In a study of 107 *C. difficile* strains of varying resistance phenotypes, ridini­lazole demonstrated minimum inhibitory concentration (MIC) values (MIC range: 0.015-0.5 mg/L; MIC<sub>50</sub>: 0.03 mg/L; MIC<sub>90</sub>: 0.125 mg/L) lower than those of metronidazole (MIC range: <0.125-2 mg/L; MIC<sub>50</sub>: 0.5 mg/L; MIC<sub>90</sub>: 2 mg/L) and vancomycin (MIC range: 0.5-8 mg/L; MIC<sub>50</sub>: 1 mg/L; MIC<sub>90</sub>: 2 mg/L), and comparable to those of fidaxomicin (MIC range: 0.004-0.125 mg/L; MIC<sub>50</sub>: 0.06 mg/L; MIC<sub>90</sub>: 0.125 mg/L) [29]. Among 200 isolates collected from *C. difficile* toxin-positive stool samples at 6 United States sites, ridini­lazole was similarly found to have lower MIC values (MIC range: 0.12-0.5 mg/L; MIC<sub>50</sub>: 0.12 mg/L; MIC<sub>90</sub>: 0.25 mg/L) than metronidazole (MIC range: 0.12-2 mg/L; MIC<sub>50</sub>: 0.25 mg/L; MIC<sub>90</sub>: 1 mg/L) or vancomycin (MIC range: 0.25-4 mg/L; MIC<sub>50</sub>: 1 mg/L; MIC<sub>90</sub>: 2 mg/L) and was comparable to fidaxomicin (MIC range: 0.015-1 mg/L; MIC<sub>50</sub>: 0.03 mg/L; MIC<sub>90</sub>: 0.125 mg/L) [30].

Mutant prevention concentrations, used to describe the antimicrobial drug concentration required to block the

growth of the least susceptible cell present in a high-density bacterial population and give insight into the likelihood for resistance selection compared with achievable therapeutic concentrations, remained well below ( $\leq 0.25$  mg/L) fecal concentrations of ridini­lazole, suggesting a low propensity for resistance selection among *C. difficile* strains clinically [33]. Following 14 serial passages at sub-MIC concentrations ( $0.5 \times \text{MIC}$ ), no increase in ridini­lazole MIC was noted and spontaneous mutations were highly infrequent, occurring at  $< 3.17 \times 10^{-9}$  [34]. Moreover, no spontaneously ridini­lazole-resistant mutants were identified among 2 clinical strains of *C. difficile* (ribotype 012 and ribotype 027) following 15 serial passages using brain heart infusion broth containing ridini­lazole concentrations ranging from 0.004-64 mg/L [35]. In contrast, Bassères *et al* were able to isolate a stable mutant of *C. difficile* ribotype 027 with a filamentous phenotype following 11 serial passages that stabilized at an MIC of 0.48 mg/L, eight times higher than the initial value of the isolate (0.06 mg/L) [36]. Overall, the development of *C. difficile* resistance to ridini­lazole seems likely to be uncommon.

### Activity against gut microbiota

While exhibiting substantial activity against *C. difficile*, ridini­lazole is largely specific for this bacterium and demonstrates limited activity against gut microbiota. When tested *in vitro* against 350 bacterial isolates representing common intestinal flora, ridini­lazole showed limited activity against gram-negative and gram-positive anaerobes other than *Clostridium* spp., summarized in Table 1. Of note is the demonstrated lack of activity of ridini­lazole against *Bacteroides fragilis* (*B. fragilis*) (MIC range: 512 to >512 mg/L; MIC<sub>50</sub>: >512 mg/L) and *Bifidobacteria* spp. (MIC range: 16 to >512 mg/L; MIC<sub>50</sub>: >512 mg/L). In contrast, vancomycin MICs for *B. fragilis* and *Bifidobacteria* spp. ranged from 32-128 mg/L and 0.5-1 mg/L, respectively [27]. In a subsequent evaluation of 162 strains representing 35 species of *Clostridium* spp. (non-*C. difficile*), ridini­lazole showed variable activity MIC range (0.06 to >512 mg/L) depending upon species and strain [37]. In an *in vitro* CDI human gut model, ridini­lazole demonstrated minimal impact on viable cell counts of indigenous gut microbiota other than total *Clostridium* spp. [32].

In fecal samples from healthy adult males who were given ridini­lazole 200 or 500 mg b.i.d. in a Phase I trial, clostridial viable counts were below the lower limits of detection for viable counting and other gut microbiota were largely unchanged [38]. Further phase I evaluation similarly showed a limited impact of ridini­lazole on gut microbiota other than *Clostridium* spp. Counts of *Bacteroides* spp. increased (2 log by day 9), while counts of lactobacilli decreased (1 log by day 4, remaining constant until day 9). Minimal change was observed in counts of *Bifidobacterium* spp. and total anaerobes, while counts of total aerobes and lactose fermenting *Enterobacteriaceae* increased (2 log by day 9) [26]. An evaluation of fecal samples from 82 patients enrolled in a phase II trial similarly demonstrated a limited impact of ridini­lazole on gut microbiota, except for a 1 log<sub>10</sub> reduction on day 10 of therapy for *Clostridium leptum*. In

**Table 1** *In vitro* susceptibility of various gastrointestinal bacteria to ridinilazole and select comparator agents [27,34]

Ribotype	Agent	MIC range (mg/L)	MIC <sub>50</sub> (mg/L)	MIC <sub>90</sub> (mg/L)
<i>Bifidobacterium</i> species	Ridinilazole	16→512	64, >512	128, >512
	Fidaxomicin	≥0.03-0.25	0.125	0.125
	Vancomycin	0.5-1	1	1
	Metronidazole	2→512	32	128
<i>Bacteroides fragilis</i>	Ridinilazole	512→512	>512	>512
	Fidaxomicin	>512	>512	>512
	Vancomycin	32-128	64	64
	Metronidazole	0.5-2	1	2
<i>Clostridium innocuum</i>	Ridinilazole	0.06-1	0.25	1
	Fidaxomicin	128-512	256	256
	Vancomycin	16	16	16
	Metronidazole	0.5-16	1	2
<i>Clostridium ramosum</i>	Ridinilazole	128→512	>512	>512
	Fidaxomicin	>512	>512	>512
	Vancomycin	4	4	4
	Metronidazole	0.5-8	0.5	1
<i>Enterococcus faecalis</i>	Ridinilazole	128→512	>512	>512
	Fidaxomicin	1-8	8	8
	Vancomycin	1-4	1	4
	Metronidazole	>512	>512	>512
<i>Enterococcus faecium</i>	Ridinilazole	64→512	128	128
	Fidaxomicin	0.5-16	8	>128
	Vancomycin	0.5-256	0.5	256
	Metronidazole	256→512	>512	>512
<i>Lactobacillus</i> species	Ridinilazole	0.06→512	16, 128	>512
	Fidaxomicin	0.25→512	8	>512
	Vancomycin	0.5→512	256	>512
	Metronidazole	2→512	>512	>512
<i>Peptostreptococcus anaerobius</i>	Ridinilazole	0.125-128	64	64
	Fidaxomicin	≥0.03	≥0.03	≥0.03
	Vancomycin	0.5	0.5	0.5
	Metronidazole	0.125-1	0.5	1
<i>Prevotella</i> species	Ridinilazole	32→512	>512	>512
	Fidaxomicin	64→512	>512	>512
	Vancomycin	64→512	128	512
	Metronidazole	0.25-1	0.5	1

MIC, minimum inhibitory concentration

contrast, vancomycin exposure led to significantly decreased bacterial counts of *Bacteroides* spp., *Prevotella* spp., *Clostridium coccoides* and *Clostridium leptum* from day 1 to 10 by approximately 3 log<sub>10</sub> copies per gram of stool. This substantial impact led to increased bacterial counts of *Enterobacteriaceae* during vancomycin treatment, but not during ridinilazole treatment [39]. A separate exploratory phase II trial evaluated

the impact of ridinilazole (n=14) and fidaxomicin (n=13) on the host gut microbiome in a randomized, open label study. The results suggested that ridinilazole was more preserving of the gut microbiome compared with fidaxomicin. Specifically, fidaxomicin exposure led to a reduction in 10 of the bacterial families analyzed compared to only 2 bacterial families with reduced populations among samples exposed to ridinilazole.

Moreover, fidaxomicin exposure led to a reduction in the population of bacterial families belonging to the Firmicutes, thought to have a role in protection against CDI [40]. An additional phase II trial compared the composition of fecal microbiota between patients administered ridini­lazole and vancomycin using 16S rDNA at baseline, day 5 of therapy, day 10 (end of therapy), day 25 and at the end of the study. Significant reductions in percent relative abundance were modest and were only found within the Firmicutes phylum among samples from ridini­lazole-exposed patients. Within the Firmicutes, vancomycin affected more taxa than ridini­lazole. Additionally, vancomycin exposure led to substantial decreases in the abundance of Actinobacteria, Bacteroidetes, and Proteobacteria. Interestingly the fecal microbiota returned to baseline more quickly among ridini­lazole-treated patients [41]. The targeted activity of ridini­lazole has demonstrated a lower propensity for collateral damage to the gut microbiome in multiple models and human fecal evaluations and highlights a promising aspect of this new agent.

### Pharmacokinetics

A phase I study (n=56) was conducted to elucidate the pharmacokinetics and safety of ridini­lazole in healthy volunteers after single and multiple oral doses taken with or without food [26]. All doses studied were well-tolerated and any adverse effects potentially due to ridini­lazole were mild. In part one, placebo or single ascending doses of ridini­lazole (2, 20, 100, 400, 1000 or 2000 mg) were administered. Plasma levels following doses up to 2000 mg in the fasted state were generally undetectable. The 1000 mg dose was studied in the fasted and fed state. A single oral dose of 1000 mg in the fed state produced low but detectable plasma concentrations in all 6 subjects in that study arm (range: 0.102-0.296 ng/mL). Maximum plasma concentrations were observed 4 h following administration. In part two, ridini­lazole was administered in the fed state at a dose of 200 mg or 500 mg b.i.d. for 10 days. By Day 10, low plasma ridini­lazole concentrations were observed in most test subjects (range: 0.105-0.305 ng/mL) with  $T_{max}$  again occurring at 4 hours post-dose. In those receiving 200 mg b.i.d., the mean area under the plasma concentration-time curve from time zero up to the last quantifiable concentration was 0.670 ng.h/mL (range: 0.524-1.30), with average maximum plasma concentrations of 0.141 ng/mL (range: 0.108-0.243). These findings are consistent with those of prior animal studies and subsequent phase II human data, collectively indicating that ridini­lazole is mostly contained within the gastrointestinal tract following oral administration [42-44].

Fecal concentrations of ridini­lazole were also measured in the phase I study. Ridini­lazole was primarily excreted in the feces as unchanged parent drug, which accounted for 97% of the total peak area. Mean fecal concentrations increased with increasing doses. For subjects receiving 200 mg b.i.d., mean (range) fecal concentrations on Day 5 and Day 10 were 1466  $\mu$ g/g (847-2390  $\mu$ g/g) and 1364  $\mu$ g/g (783-1980  $\mu$ g/g), respectively [26]. Similar fecal concentrations were observed

in a phase II trial, wherein the mean fecal concentration on treatment day 10 was 1373  $\mu$ g/g [44]. The observed fecal concentrations across clinical and preclinical studies are significantly above the  $MIC_{90}$  of ridini­lazole for *C. difficile* (0.125-0.25 mg/L) [29,30].

### Pharmacodynamics

*C. difficile* isolates exposed to ridini­lazole exhibit elongated morphology. Though exposed cells may continue to replicate DNA initially, septum formation is halted, which ultimately impairs cell division [22]. The pharmacodynamic effects of ridini­lazole have been studied *in vitro* against a multitude of *C. difficile* strains. Ridini­lazole was tested against 82 clinical *C. difficile* isolates collected in the United Kingdom [28]. The  $MIC_{90}$  was 0.125 mg/L, 16- to 32-fold lower than  $MIC_{90}$  values for metronidazole and vancomycin. Killing kinetics were compared against fidaxomicin and vancomycin for three different *C. difficile* strains: BI1 (ribotype 027), 630 (ribotype 012) and 5325 (ribotype 078). For the BI1 strain, vancomycin was bactericidal at 2 $\times$  MIC. Fidaxomicin was bacteriostatic at 1-10 $\times$  MIC (1.5-2.0  $\log_{10}$  reduction at 24 h) and bactericidal at 20 $\times$  MIC. Ridini­lazole was bactericidal at all concentrations tested, with reduction in viable *C. difficile* counts to below the limit of detection by 24 h. Killing by ridini­lazole did not appear dependent on drug concentration. Ridini­lazole was bactericidal against strains 630 and 5325 at  $\geq 5\times$  MIC and  $\geq 10\times$  MIC, respectively. The post-antibiotic effect (PAE) for ridini­lazole against all strains was >20 h at 20 $\times$  MIC, a concentration that should be exceeded in patients taking 200 mg by mouth b.i.d. [26,44]. Fidaxomicin also exhibited a PAE for all strains, while vancomycin was generally bacteriostatic with minimal PAE (0-2 h) [28].

Bassères *et al* examined *in vitro* production of toxins A and B in *C. difficile* strains exposed to sub-MIC (0.125 $\times$ -0.5 $\times$ ) and supra-MIC (4 $\times$  or 40 $\times$  MIC) concentrations of ridini­lazole [22]. Toxin B production was suppressed below the limit of detection with all concentrations of ridini­lazole tested. Toxin A levels were reduced by 75% to >90% at varying ridini­lazole concentrations. Cytotoxin levels were similarly reduced in an *in vitro* gut model of CDI [32]. Additionally, interleukin (IL)-8 release from eukaryotic cell lines exposed to antibiotic-treated *C. difficile* isolates were measured [22]. IL-8 production was reduced by 25-74% following exposure to *C. difficile* that had been treated with ridini­lazole, suggesting that ridini­lazole-mediated toxin reduction may dampen host inflammatory responses. Metronidazole and vancomycin had minimal impact on toxin production, and IL-8 production after exposure to metronidazole or vancomycin was similar to that of controls [22]. In a phase II study, 100 patients with CDI were randomized to 10 days of treatment with either ridini­lazole or vancomycin and fecal concentrations of calprotectin and lactoferrin were measured to determine bowel inflammation at baseline, day 5, and day 10 (end of treatment) [45]. When all disease severities were analyzed together at day 10, concentrations of both toxins were reduced similarly from baseline in both the ridini­lazole and



vancomycin groups. In contrast, ridinilazole demonstrated a greater reduction in lactoferrin (1.93 vs. 0.62 log<sub>10</sub>) and calprotectin (1.70 vs. 0.22 log<sub>10</sub>) from baseline to day 10 when only severe disease was analyzed [45].

The efficacy of ridinilazole versus comparator agents has also been investigated in animal models. The hamster model of clindamycin-induced CDI is a validated method for evaluating the potential of new CDI therapies [46]. In this model, hamsters are rendered susceptible to CDI by oral administration of clindamycin followed by infection with *C. difficile* spores. This protocol produces a severe model of CDI with subject death generally occurring within 3 days in the absence of treatment. In two hamster-model experiments using the epidemic BI/NAP1 (ribotype 025) strain, survival rates with ridinilazole treatment at a total daily dose of 50 mg/kg were 70% and 100% vs. 60% and 10% with vancomycin [42,43]. Fidaxomicin was also effective against the epidemic strain, with 80-90% of animals surviving [42]. In a study arm using the non-epidemic VA11 clinical strain [43], ridinilazole was associated with better survival rates than vancomycin (80-95% vs. 50%). In a study using the non-epidemic ribotype 012 strain [42], all vancomycin-treated animals survived during dosing, but there was a 0% survival rate in the follow-up period. Fidaxomicin conferred some protection, but overall survival rates across different fidaxomicin dosing groups were low, ranging from 0-40%. Ridinilazole conferred significant protection against ribotype 012, with survival rates of 80-100% across dosing groups. Further, following administration of high ridinilazole doses (25 mg/kg b.i.d.), no spores were isolated in animals with ribotype 027 infection from day 7 onwards, although a significant number of animals had spores isolated in the ribotype 012 arm [42].

## Safety and efficacy

The safety and efficacy of ridinilazole were compared to those of vancomycin in a phase II, double-blind, active-controlled study at 33 various sites in the United States and Canada [44]. Patients were included in the study if they met the definition of having CDI, had a positive diagnostic test for *C. difficile* or had received more than 24 h of CDI treatment prior to initiation of the study agent. Included patients were randomized 1:1 to receive either ridinilazole 200 mg orally b.i.d. or vancomycin 125 mg orally q.i.d. for 10 days. The primary endpoint was sustained clinical response, which was defined as clinical cure at test of cure (TOC) (day 12-14) and absence of CDI recurrence 30 days after end of treatment (day 10-11). Secondary endpoints and objectives included time to hospital discharge, time to diarrhea resolution, and tolerability of ridinilazole compared with vancomycin. The primary efficacy analysis was performed on 69 patients in the modified intent-to-treat (mITT) population. Patients receiving ridinilazole had a higher sustained clinical response compared with patients receiving vancomycin in the mITT population (66.7% vs. 42.4%;  $P=0.0004$ ). Ridinilazole was determined to be non-inferior to vancomycin; 36 (77.8%) and

23 (67.7%) patients in the ridinilazole and vancomycin groups, respectively, had a clinical response at TOC. The median time to diarrhea resolution was 4 and 5 days (hazard ratio [HR] 1.19; 90% confidence interval [CI] 0.76-1.87) and median time to hospital discharge was 5 and 7 days (HR 0.99; 90%CI 0.34-2.91) for the ridinilazole and vancomycin groups, respectively. CDI recurrence was documented in 14.3% of patients receiving ridinilazole and 34.8% of patients receiving vancomycin (-16.2%, 90%CI -35.5 to 3).

Ridinilazole displayed adverse reactions similar to those of vancomycin, with most being gastrointestinal-related (40% vs. 56%, respectively) [44]. Three patients had severe adverse reactions that may have been related to the study drug: one patient in the ridinilazole arm had hypokalemia and two patients in the vancomycin arm had serious reactions: one septic shock and moderate hematemesis and the other elevated liver enzymes and diarrhea. Nausea (20%) and abdominal pain (12%) were the most commonly reported adverse reactions associated with ridinilazole. These results were similar to those seen in phase I studies [26].

## Concluding remarks

*C. difficile* infection remains a global healthcare threat with a propensity to cause recurrent disease. Currently, treatment options for *C. difficile* are limited and ridinilazole seems to be a promising potential agent for the treatment of CDI. Ridinilazole has also demonstrated a lower propensity for collateral damage to the gut microbiome and appears to diminish the production of *C. difficile* toxins and subsequent bowel inflammation, which may prove advantageous in managing severe CDI. Current phase II results are promising and future availability of phase III trial results will help further delineate the role and value of ridinilazole.

## References

1. Kyne L, Hamel MB, Polavaram R, Kelly CP. Health care costs and mortality associated with nosocomial diarrhea due to *Clostridium difficile*. *Clin Infect Dis* 2002;**34**:346-353.
2. O'Brien JA, Lahue BJ, Caro JJ, Davidson DM. The emerging infectious challenge of *Clostridium difficile*-associated disease in Massachusetts hospitals: clinical and economic consequences. *Infect Control Hosp Epidemiol* 2007;**28**:1219-1227.
3. Dubberke ER, Olsen MA. Burden of *Clostridium difficile* on the healthcare system. *Clin Infect Dis* 2012;**55**(Suppl 2):S88-S92.
4. Centers for Disease Control and Prevention. Antibiotic resistance threats in the United States 2013. <http://www.cdc.gov/drugresistance/pdf/ar-threats-2013-508.pdf> [Accessed 11 November, 2018].
5. Lessa FC, Mu Y, Bamberg WM, et al. Burden of *Clostridium difficile* infection in the United States. *N Engl J Med* 2015;**372**:825-834.
6. He M, Miyajima F, Roberts P, et al. Emergence and global spread of epidemic healthcare-associated *Clostridium difficile*. *Nat Genet* 2013;**45**:109-113.
7. Jones AM, Kuijper EJ, Wilcox MH. *Clostridium difficile*: a European perspective. *J Infect* 2013;**66**:115-128.

8. Burke KE, Lamont JT. *Clostridium difficile* infection: a worldwide disease. *Gut Liver* 2014;**8**:1-6.
9. Borren NZ, Ghadermarzi S, Hutfless S, Ananthakrishnan AN. The emergence of *Clostridium difficile* infection in Asia: A systematic review and meta-analysis of incidence and impact. *PLoS One* 2017;**12**:e0176797.
10. Climo MW, Israel DS, Wong ES, Williams D, Coudron P, Markowitz SM. Hospital-wide restriction of clindamycin: effect on the incidence of *Clostridium difficile*-associated diarrhea and cost. *Ann Intern Med* 1998;**128**:989-995.
11. Valiquette L, Cossette B, Garant MP, Diab H, Pépin J. Impact of a reduction in the use of high-risk antibiotics on the course of an epidemic of *Clostridium difficile*-associated disease caused by the hypervirulent NAP1/027 strain. *Clin Infect Dis* 2007;**45**(Suppl 2):S112-S121.
12. Dingle KE, Didelot X, Quan TP, et al; Modernising Medical Microbiology Informatics Group. Effects of control interventions on *Clostridium difficile* infection in England: an observational study. *Lancet Infect Dis* 2017;**17**:411-421.
13. Baur D, Gladstone BP, Burkert F, et al. Effect of antibiotic stewardship on the incidence of infection and colonisation with antibiotic-resistant bacteria and *Clostridium difficile* infection: a systematic review and meta-analysis. *Lancet Infect Dis* 2017;**17**:990-1001.
14. Eyre DW, Walker AS, Wyllie D, et al; Infections in Oxfordshire Research Database. Predictors of first recurrence of *Clostridium difficile* infection: implications for initial management. *Clin Infect Dis* 2012;**55**(Suppl 2):S77-S87.
15. Eyre DW, Cule ML, Wilson DJ, et al. Diverse sources of *C. difficile* infection identified on whole-genome sequencing. *N Engl J Med* 2013;**369**:1195-1205.
16. Kelly CP. Can we identify patients at high risk of recurrent *Clostridium difficile* infection? *Clin Microbiol Infect* 2012;**18**(Suppl 6):21-27.
17. McDonald LC, Gerding DN, Johnson S, et al. Clinical Practice Guidelines for *Clostridium difficile* Infection in Adults and Children: 2017 Update by the Infectious Diseases Society of America (IDSA) and Society for Healthcare Epidemiology of America (SHEA). *Clin Infect Dis* 2018;**66**:987-994.
18. Debast SB, Bauer MP, Kuijper EJ. European Society of Clinical Microbiology and Infectious Diseases: update of the treatment guidance document for *Clostridium difficile* infection. *Clin Microbiol Infect* 2014;**20**(Suppl 2):1-26.
19. Louie TJ, Miller MA, Mullane KM, et al; OPT-80-003 Clinical Study Group. Fidaxomicin versus vancomycin for *Clostridium difficile* infection. *N Engl J Med* 2011;**364**:422-431.
20. Crook DW, Walker AS, Kean Y, et al; Study 003/004 Teams. Fidaxomicin versus vancomycin for *Clostridium difficile* infection: meta-analysis of pivotal randomized controlled trials. *Clin Infect Dis* 2012;**55**(Suppl 2):S93-S103.
21. Tickler IA, Goering RV, Whitmore JD, Lynn AN, Persing DH, Tenover FC; Healthcare Associated Infection Consortium. Strain types and antimicrobial resistance patterns of *Clostridium difficile* isolates from the United States, 2011 to 2013. *Antimicrob Agents Chemother* 2014;**58**:4214-4218.
22. Bassères E, Endres BT, Khaleduzzaman M, et al. Impact on toxin production and cell morphology in *Clostridium difficile* by ridiniilazole (SMT19969), a novel treatment for *C. difficile* infection. *J Antimicrob Chemother* 2016;**71**:1245-1251.
23. Vickers RJ, Tillotson G, Goldstein EJ, Citron DM, Garey KW, Wilcox MH. Ridiniilazole: a novel therapy for *Clostridium difficile* infection. *Int J Antimicrob Agents* 2016;**48**:137-143.
24. Bassères E, Endres BT, Vickers R, et al. Transcriptome functional analysis of *Clostridium difficile* exposed to ridiniilazole; insight into potential mechanism of action. In: Abstracts of the Twenty-seventh European Congress of Clinical Microbiology and Infectious Diseases, Vienna, Austria, 2017. Abstract EP0404, Blackwell, Hoboken, NJ, USA.
25. Vickers RJ, Storer R, Tinsley J, et al. SMT19969: Preclinical safety and pharmacokinetics of a novel antibiotic for *Clostridium difficile* infection. In: Abstracts of the Twenty-third European Congress of Clinical Microbiology and Infectious Diseases, Berlin, Germany, 2013, Abstract P1656.
26. Vickers R, Robinson N, Best E, Echols R, Tillotson G, Wilcox M. A randomised phase 1 study to investigate safety, pharmacokinetics and impact on gut microbiota following single and multiple oral doses in healthy male subjects of SMT19969, a novel agent for *Clostridium difficile* infections. *BMC Infect Dis* 2015;**15**:91.
27. Goldstein EJ, Citron DM, Tyrrell KL, Merriam CV. Comparative *in vitro* activities of SMT19969, a new antimicrobial agent, against *Clostridium difficile* and 350 gram-positive and gram-negative aerobic and anaerobic intestinal flora isolates. *Antimicrob Agents Chemother* 2013;**57**:4872-4876.
28. Corbett D, Wise A, Birchall S, et al. *In vitro* susceptibility of *Clostridium difficile* to SMT19969 and comparators, as well as the killing kinetics and post-antibiotic effects of SMT19969 and comparators against *C. difficile*. *J Antimicrob Chemother* 2015;**70**:1751-1756.
29. Freeman J, Vernon J, Vickers R, Wilcox MH. Susceptibility of *Clostridium difficile* isolates of varying antimicrobial resistance phenotypes to smt19969 and 11 comparators. *Antimicrob Agents Chemother* 2016;**60**:689-692.
30. Snyderman DR, McDermott LA, Thorpe CM, et al. *In-vitro* activity of ridiniilazole and comparators against isolates of *Clostridium difficile* obtained from stools of patients as part of US Surveillance in 2014. In: American Society for Microbiology Microbe 2017, New Orleans, LA, USA, Abstract 235.
31. Snyderman DR, McDermott LA, Thorpe CM, et al. Antimicrobial susceptibility and ribotypes of *Clostridium difficile* isolates from a Phase 2 clinical trial of ridiniilazole (SMT19969) and vancomycin. *J Antimicrob Chemother* 2018;**73**:2078-2084.
32. Baines SD, Crowther GS, Freeman J, Todhunter S, Vickers R, Wilcox MH. SMT19969 as a treatment for *Clostridium difficile* infection: an assessment of antimicrobial activity using conventional susceptibility testing and an *in vitro* gut model. *J Antimicrob Chemother* 2015;**70**:182-189.
33. Blondeau JM, Shebelski SD, Vickers R. Mutant prevention concentration of SMT19969 against clinical isolates of *Clostridium difficile*. In: Abstracts of the Fifty-fourth Interscience Conference on Antimicrobial Agents and Chemotherapy, Washington DC, USA, 2014, Abstract F-245.
34. Vickers R, Tinsley J, Wilson F, et al. SMT19969: *In vitro* and *in vivo* evaluation of a novel antibiotic for *Clostridium difficile* infection. In: Abstracts of the Twenty-third European Congress of Clinical Microbiology and Infectious Diseases, Berlin, Germany, 2013, Abstract P1655.
35. Hall D, Hinshaw R, Pillar C, et al. Resistance development studies with ridiniilazole (RDZ) and *Clostridium difficile*. In: American Society for Microbiology Microbe 2016, Boston, MA, USA, Abstract 443.
36. Bassères E, Rashid T, Endres BT, et al. *In vitro* selection of *Clostridium difficile* resistance mutants exposed to ridiniilazole. In: American Society for Microbiology Microbe 2017, New Orleans, LA, USA, Abstract 233.
37. Goldstein EJ, Citron DM, Tyrrell KL. Comparative *in vitro* activities of SMT19969, a new antimicrobial agent, against 162 strains from 35 less frequently recovered intestinal *Clostridium* species: implications for *Clostridium difficile* recurrence. *Antimicrob Agents Chemother* 2014;**58**:1187-1191.
38. Best E, Vickers R, Wilcox MH. Effects of a novel antimicrobial (SMT19969) for treatment of *Clostridium difficile* infection (CDI)

- on gut flora. In: Abstracts of the Fifty-third Interscience Conference on Antimicrobial Agents and Chemotherapy, Denver, Colorado, 2013, Abstract K-167.
39. Chang J, Kane A, Snyderman DR, et al. Ridinilazole preserves major components of the intestinal microbiota during treatment of *Clostridium difficile* infection. In: American Society for Microbiology Microbe 2016, Boston, MA, USA, Abstract LB-116.
  40. Mitra S, Chilton C, Freeman J, et al. Preservation of gut microbiome following ridinilazole versus fidaxomicin treatment of *Clostridium difficile* infection. In: ID Week, San Diego, CA, 2017, Abstract 1842.
  41. Thorpe C, Kane A, Chang J, et al. Ridinilazole treatment for *C. difficile* infection preserves the intestinal microbiota of patients. In: Abstracts of the Twenty-seventh European Congress of Clinical Microbiology and Infectious Diseases, Vienna, Austria, 2017, Abstract P2049.
  42. Sattar A, Thommes P, Payne L, Warn P, Vickers RJ. SMT19969 for *Clostridium difficile* infection (CDI): *in vivo* efficacy compared with fidaxomicin and vancomycin in the hamster model of CDI. *J Antimicrob Chemother* 2015;**70**:1757-1762.
  43. Weiss W, Pulse M, Vickers R. *In vivo* assessment of SMT19969 in a hamster model of *Clostridium difficile* infection. *Antimicrob Agents Chemother* 2014;**58**:5714-5718.
  44. Vickers RJ, Tillotson GS, Nathan R, et al; CoDIFY study group. Efficacy and safety of ridinilazole compared with vancomycin for the treatment of *Clostridium difficile* infection: a phase 2, randomised, double-blind, active-controlled, non-inferiority study. *Lancet Infect Dis* 2017;**17**:735-744.
  45. Vicker RJ, Yan K, Li Y. Ridinilazole for *Clostridium difficile* infection (CDI) - reductions in calprotectin and lactoferrin during therapy. In: American Society for Microbiology Microbe 2016, Boston, MA, USA, Abstract 441.
  46. Douce G, Goulding D. Refinement of the hamster model of *Clostridium difficile* disease. *Methods Mol Biol* 2010;**646**:215-227.