

latrogenic Bidirectional Dissection of the Right Coronary Artery and the Ascending Aorta: The Worst Nightmare for an Interventional Cardiologist

Ziad Dahdouh, MD¹, Vincent Roule, MD¹, Therese Lognone, MD¹, Remi Sabatier, MD¹, Mathieu Bignon, MD¹, Guillaume Malcor, MD¹, Adrien Lemaitre, MD¹, Katrien Blanchart, MD¹, Julien Wain-Hobson, MD¹, Vladimir Saplacan, MD², Fabio Cutone, MD², Dimitrios Buklas, MD², Calin Ivascau, MD², Massimo Massetti, MD², and Gilles Grollier, MD¹

¹Departments of Cardiology and ²Thoracic and Cardiovascular Surgery, CHU de Caen, Caen, France

Korean Circ J 2012;42:504-6 http://dx.doi.org/10.4070/kcj.2012.42.7.504

We would like to correct the sentence as written below. The changes are underlined. The author sincerely regrets these errors

1) In the abstract (last sentence):

"Therefore, an isolated surgical aortic valve replacement was performed." should be replaced by: "Therefore, surgical aortic valve replacement with prosthetic tube graft was performed".

2) In the case paragraph (last sentence):

"An isolated aortic valve replacement was performed but the post-surgical period was complicated by the incidence of uncontrollable septic shock and death occurred at day seven following the procedure."

should be replaced by:

"An aortic valve replacement with prosthetic tube graft was performed but the post-surgical period was complicated by the incidence of uncontrollable septic shock and death occurred at day seven following the procedure."