



OPEN

Characteristics of hydrate-bound gas retrieved at the Kedr mud volcano (southern Lake Baikal)

Akihiro Hachikubo¹✉, Hirotugu Minami¹, Satoshi Yamashita¹, Andrey Khabuev², Alexey Krylov^{2,3,4}, Gennadiy Kalmychkov⁵, Jeffrey Poort⁶, Marc De Batist⁷, Alexandr Chenskiy⁸, Andrey Manakov⁹ & Oleg Khlystov²

We reported the characteristics of hydrate-bound hydrocarbons in lake-bottom sediments at the Kedr mud volcano in Lake Baikal. Twenty hydrate-bearing sediment cores were retrieved, and methane-stable isotopes of hydrate-bound gases ($\delta^{13}\text{C}$ and $\delta^2\text{H}$ of -47.8‰ to -44.0‰ V-PDB and -280.5‰ to -272.8‰ V-SMOW, respectively) indicated their thermogenic origin accompanied with secondary microbial methane. Powder X-ray diffraction patterns of the crystals and molecular composition of the hydrate-bound gases suggested that structure II crystals showed a high concentration of ethane (around 14% of hydrate-bound hydrocarbons), whereas structure I crystals showed a relatively low concentration of ethane (2–5% of hydrate-bound hydrocarbons). These different crystallographic structures comprised complicated layers in the sub-lacustrine sediment, suggesting that the gas hydrates partly dissociate, concentrate ethane and form structure II crystals. We concluded that a high concentration of thermogenic ethane primarily controls the crystallographic structure of gas hydrates and that propane, iso-butane (2-methylpropane) and neopentane (2,2-dimethylpropane) are engaged into crystals in the re-crystallisation process.

Gas hydrates are crystalline clathrate compounds composed of water and gas molecules that are stable at low temperature and high partial pressure of each gas component¹. Natural gas hydrates, which contain methane (C_1) as a major component, exist in sea/lake sediment columns and permafrost layers and are considered to be a possible global source of energy^{2,3}. There are different views on the role of gas hydrates on global warming^{4–6}. They are of concern as a large reservoir of C_1 ; however, the amount of hydrate-bound gas is smaller than that expected previously, which makes them unlikely to cause global warming by dissociation of C_1 ^{5,6}. Moreover, the current understanding of the formation, dissociation and maintenance processes of natural gas hydrates is still incomplete.

Molecular fractionation during formation of gas hydrate crystals occurs according to the size ratio of guest molecules to host cages and the difference in equilibrium pressure of each component of hydrocarbons. Milkov et al.⁷ reported that gas hydrates retrieved at the southern Hydrate Ridge (offshore Oregon) are rich in ethane (C_2) but exclude propane (C_3), because the crystal structure I (sI) cannot encage C_3 . Sassen et al.⁸ studied the gas hydrates retrieved at the Gulf of Mexico and found that they are high in C_2 , C_3 and butane (C_4) but exclude isopentane (*i*- C_5 , 2-methylbutane), because the crystal structure II (sII) cannot encage such large guest molecules. Therefore, molecular composition of natural gas primarily controls the crystallographic structure of gas hydrates.

Natural gas hydrates have been discovered in sub-lacustrine sediments in Lake Baikal, in association with fluid venting at mud volcanoes, pockmarks and cold seeps^{9,10}. Although the Lake Baikal gas hydrates generally belong to sI, higher concentrations of C_2 locally induce the formation of sII as sII gas hydrates form for a particular composition of C_1 and C_2 ^{11,12}. Co-existence of sI and sII hydrates in the same sediment core, retrieved at the

¹Kitami Institute of Technology, 165 Koen-cho, Kitami 090-8507, Japan. ²Limnological Institute, SB RAS, 3 Ulan-Batorskaya St, Irkutsk, Russia 664033. ³Institute of Earth Sciences, St. Petersburg State University, 7-9, Universitetskaya Nab., St. Petersburg, Russia 199034. ⁴VNIIOkeangeologia, Anglyisky prospect 1, St. Petersburg, Russia 190121. ⁵Vinogradov Institute of Geochemistry, SB RAS, 1-a Favorovskiy St, Irkutsk, Russia 664033. ⁶Sorbonne Université, CNRS, Institut des Sciences de la Terre de Paris, ISTeP, 4 place Jussieu, 75005 Paris, France. ⁷Renard Centre of Marine Geology, Ghent University, Krijgslaan 281 s8, 9000 Ghent, Belgium. ⁸Irkutsk National Research Technical University, 83 Lemontov St, Irkutsk, Russia 664074. ⁹Nikolaev Institute of Inorganic Chemistry, SB RAS, 3 Acad. Lavrentiev Ave, Novosibirsk, Russia 630090. ✉email: hachi@mail.kitami-it.ac.jp




Figure 1. Bathymetry map of the Kedr mud volcano (MV) at the southern basin of Lake Baikal. The blue, red, green and white points indicate structure I only, structure II only, both structures I and II and no gas hydrate, respectively. This map was created using Kashmir 3D, Version 9.3.1 (<https://www.kashmir3d.com/>).

Kukuy K-2 mud volcano (MV) in the central Baikal basin (Fig. 1), was reported by Kida et al.^{13,14}. Some formation models for these ‘double structure’ gas hydrates were proposed by Hachikubo et al.¹⁵, Poort et al.¹⁶ and Manakov et al.¹⁷. The hypothesis shared by the latter two papers is that the sII hydrates are formed by re-crystallisation after dissociation of pre-existing sI hydrates. Because C₂ is prone to be engaged in the hydrate phase rather than C₁, C₂ can be concentrated into the hydrate during the re-crystallisation process with C₂-rich sII forming¹⁸.

C₂-rich sII gas hydrates (C₂ concentration ca. 14% of volatile hydrocarbons) have been discovered not only at the Kukuy K-2 MV but also at the Kukuy K-4 MV and the Kukuy K-10 gas hydrate mound, located in the central Baikal basin, and at the PosolBank seep, located in the southern Baikal basin (Fig. 1)^{9,19}. Recently, double-structure gas hydrates were also recovered at the Kedr MV, located in the southern Baikal basin, between 2015 and 2017^{10,20}. This study focused on the characteristics of molecular and stable isotope compositions of hydrate-bound hydrocarbons retrieved at the Kedr MV area to improve our understanding of the formation process of double-structure gas hydrates.

The Kedr MV is located in the southern Baikal basin, at 27 km south of Listvyanka (Fig. 1), and it was separated into two study areas: a complex of mud-volcanic buildings (Kedr-1 area), where mud-volcanic breccia was found²⁰, and a separate hill with gas hydrate in sediment (Kedr-2 area). Multi-beam echosounder data obtained in 2015 revealed that the Kedr MV consists of small mounds and pockmarks, and results of pore water geochemistry suggested the existence of fluid discharge from greater depths²⁰. This part of Lake Baikal is known for the presence of the coal-bearing sediments of the Tankhoy Formation, Oligocene–Miocene age^{21,22}. These deposits would represent an ideal source for upward migrating gas that could lead to the formation of gas hydrates near the lake floor^{20,22}. During expeditions onboard R/V *G. Yu. Vereshchagin* (VER) in 2015–2017, 42 sediment cores, including 20 hydrate-bearing cores, were sampled at the Kedr MV: samples were taken in August–September 2015 (VER15-03), August 2016 (VER16-03) and August 2017 (VER17-03). Sediment cores were retrieved using gravity corer (length 3.5 or 6.1 m). Coring targets were mainly small mounds and pockmarks, along with five peripheral locations (2016St21, 2016St22, 2016St25, 2016St26 and 2016St27) as references.




Figure 2. Gas hydrates in sediment cores recovered from the Kedr-1 and Kedr-2 areas. Upward core direction is toward the left in each. (a) 2015St1GC2 (Kedr-1, structure II), (b) 2015St1GC8 (Kedr-1, structure II), (c) 2015St1GC11 (Kedr-1, structure I), (d) 2015St1GC19 (Kedr-1, structure I) and (e) 2017St13GC3 (Kedr-2, upper layer: structure II, lower layer: structure I).

Results

Gas hydrate crystals. Gas hydrate crystals in the sediment cores displayed massive, granular, plate-like or vein-like inclusions in the sediments (Fig. 2). Several hydrate-bearing sediment cores had solid granules, suggesting the presence of sII crystals¹⁷. PXRD profiles of the gas hydrate samples agreed with the crystal shapes (Fig. 3): the granular shape of 2015St1GC2 and the upper layer of 2017St13GC3 (230 cm below lake floor, cmblf) corresponded to sII, whereas plate-like crystals of the lower layer of 2017St13GC3 (260 cmblf) corresponded to sI. Massive crystals of 2015St1GC8, accompanied with granules, corresponded to sII. All samples included ice Ih in the PXRD profiles, caused by partial dissociation of gas hydrates during the retrieval of the cores and handling onboard.

Molecular and isotopic compositions of hydrate-bound hydrocarbons. Figure 4 shows the relationship between molecular and isotopic compositions of hydrate-bound hydrocarbons for the different sample sites in Lake Baikal. We obtained 84 samples of hydrate-bound gases from the Kedr-1 area and 12 from the Kedr-2 area. Molecular and isotopic compositions of the hydrocarbons in the hydrate-bound gases are listed in the Supplementary Information and are summarised in Table 1. C_1 was the main component of the hydrate-bound hydrocarbons, whereas C_2 was the second component. C_2 proportions in the hydrocarbons were 2.1–15.6 vol% at Kedr-1 and 4.2–14.1 vol% at Kedr-2. C_3 proportions were smaller than C_2 and distributed over a wider range at Kedr-1 (0.0003–0.3039 vol%) and Kedr-2 (0.0035–0.0349 vol%). Therefore, $C_1/(C_2 + C_3)$ in Fig. 4a is mainly influenced by C_1 and C_2 . $C_1/(C_2 + C_3)$ and $C_1 \delta^{13}C$ of the Kedr MV ranged from 5 to 47 and from -47.8% to -44.0% , respectively. $C_1/(C_2 + C_3)$ of the Kukuy K-2 MV in the central Baikal basin are clearly separated into two distinct groups responding to the gas hydrate structures: sI (21–80) and sII (~ 6)²³. Conversely, there was no clear separation in $C_1/(C_2 + C_3)$ values for Kedr-1, suggesting that some gas hydrate samples contained both sI and sII. The relationship between $\delta^{13}C$ and δ^2H of C_1 is shown in Fig. 4b. $C_1 \delta^2H$ of the Kedr MV was




Figure 3. Powder X-ray diffraction profiles of the gas hydrate samples. 2015St1GC2 (25–35 cmlbf), 2015St1GC8 (168–174 cmlbf), 2016St19GC1 (163–168 cmlbf) and 2017St13GC3 (230 cmlbf) belong to structure II, whereas 2017St13GC3 (260 cmlbf) belongs to structure I. cmlbf, centimetres below lake floor.

between -280.5‰ and -272.8‰ , and these values were $\sim 30\text{‰}$ higher than those from other gas hydrate sites. Figure 4c shows an L-shaped distribution between $C_1 \delta^{13}C$ and $C_2 \delta^{13}C$ in the hydrate-bound gas at Lake Baikal. $C_2 \delta^{13}C$ of the Kedr MV ranged from -27.8‰ to -25.3‰ and the values were almost the same as Kukuy K-0, Kukuy K-2 and Goloustnoe. Iso-butane (*i*- C_4 , 2-methylpropane), n-butane (*n*- C_4), neopentane (*neo*- C_5 , 2,2-dimethylpropane) and *i*- C_5 were detected in the hydrate-bound hydrocarbons (Table 1); however, the concentration of n-pentane (*n*- C_5) was under the detection limit of our gas chromatograph. The compositions of C_3 , *i*- C_4 and *neo*- C_5 detected in the hydrate-bound gases of the sI were smaller than those of sII.

Molecular and isotopic compositions of sediment gases. Figure 5 shows the selected depth profiles of sediment gas in the hydrate-bearing cores obtained using headspace gas method (all data are shown in the Supplementary Information). C_1 concentrations of the all sediment cores increased with depth and reached 1–10 mM at a depth of around 50 cm below lake floor (cmlbf). The CO_2 concentrations of all cores increased slightly with depth. The value of $C_1/(C_2 + C_3)$ generally decreased with depth. Because C_3 concentrations in the sediment gases were three orders of magnitude smaller than C_2 concentrations (average concentrations of C_2 and C_3 for all sediment gases are 146 μM and 0.13 μM , respectively), these results indicated that the Kedr MV area with gas hydrates is characterised by a high C_2 concentration.

$C_1 \delta^{13}C$ of all cores increased with depth; however, it also increased beneath the lake floor (10–20 cmlbf) in 2016St18GC3 and 2017St12GC3 cores, where C_1 concentrations were low, suggesting oxidation of C_1 and




Figure 4. Diagrams of hydrate-bound gases. (a) $C_1/(C_2 + C_3)$ plotted against $C_1 \delta^{13}C$, based on the classification of Milkov and Etiope²⁵; (b) $\delta^{13}C$ of C_1 plotted against δ^2H , based on the classification of Milkov and Etiope²⁵; and (c) $\delta^{13}C$ of C_2 plotted against $\delta^{13}C$ of C_1 , based on the classification of Milkov²⁸. Malenky, Bolshoy, Malyutka, Peschanka P-2, Kukuy K-0, Kukuy K-2 and Goloustnoe data are from Hachikubo et al.²³.

Structure	Kedr-1			Kedr-2		
	sI	sI + sII	sII	sI	sI + sII	sII
Number	39	9	36	2	1	9
Molecular composition [vol%]						
CH ₄	96.5 ± 0.7	91.8 ± 2.6	85.7 ± 0.7	95.4 ± 0.6	94.2	86.7 ± 0.4
C ₂ H ₆	3.5 ± 0.7	8.1 ± 2.6	14.2 ± 0.7	4.6 ± 0.6	5.7	13.3 ± 0.4
C ₃ H ₈	0.0041	0.0397	0.0701	0.0073	0.0297	0.0158
<i>i</i> -C ₄ H ₁₀	0.0003	0.0034	0.0057	0.0005	0.0039	0.0012
<i>n</i> -C ₄ H ₁₀	0.0001	0.0006	0.0007	0.0002	0.0002	0.0002
<i>neo</i> -C ₅ H ₁₂	0.0012	0.0095	0.0175	0.0013	0.0119	0.0079
<i>i</i> -C ₅ H ₁₂	0.0000	0.0001	0.0001	0.0001	0.0000	0.0000
C ₁ /(C ₂ + C ₃)	28.5 ± 6.8	12.3 ± 4.0	6.0 ± 0.4	20.8 ± 2.7	16.4	6.5 ± 0.2
Isotopic composition [$\delta^{13}\text{C}$: V-PDB, $\delta^2\text{H}$: V-SMOW]						
C ₁ $\delta^{13}\text{C}$	-46.6 ± 0.8	-45.8 ± 0.5	-46.5 ± 0.5	-44.5 ± 0.2	-44.5	-44.4 ± 0.3
C ₂ $\delta^{13}\text{C}$	-26.5 ± 0.7	-26.4 ± 0.6	-26.4 ± 0.6	-27.5 ± 0.2	-27.5	-27.2 ± 0.4
C ₃ $\delta^{13}\text{C}$	-10.8 ± 0.9	-9.9 ± 0.9	-10.5 ± 0.8	n.d	-11.6	n.d
C ₁ $\delta^2\text{H}$	-275.0 ± 1.3	-274.9 ± 1.2	-276.9 ± 1.6	-275.7 ± 1.1	-277.4	-276.6 ± 0.9
C ₂ $\delta^2\text{H}$	-212.5 ± 5.7	-218.9 ± 7.0	-216.3 ± 6.2	-214.9 ± 8.6	-220.2	-211.0 ± 2.5

Table 1. Molecular and isotopic compositions of hydrate-bound hydrocarbons at the Kedr mud volcano (MV). These data list mean values and standard deviations, sorted into three groups: sI, where C₂ composition was < 5%; sI + sII, where C₂ composition was 5%–13%; and sII, where C₂ composition was > 13%. n.d., not determined.

consumption of light C₁. C₁ $\delta^{13}\text{C}$ of Kedr-1 and Kedr-2 was around -45‰ and -42‰, respectively. C₁ $\delta^2\text{H}$ of hydrate-bearing cores was between -280‰ and -270‰ although that of 2015St1GC15 was between -300‰ and -270‰. C₂ $\delta^{13}\text{C}$ of all cores was almost constant with depth and averaged at around -26‰, suggesting thermogenic C₂. C₂ $\delta^2\text{H}$ of hydrate-bearing cores was at around -210‰; however, that of some sediment cores (2015St1GC14 and 2015St1GC15) was between -240‰ and -230‰ at their base. CO₂ $\delta^{13}\text{C}$ of all cores generally increased with depth, reaching +20‰ (Kedr-1) and +30‰ (Kedr-2).

Discussion

Origin of hydrate-bound hydrocarbons. A relationship between C₁/(C₂ + C₃) and C₁ $\delta^{13}\text{C}$ has been applied to identify the sources of hydrocarbons in submarine seeps²⁴. Recently, this diagram was revised based on a large dataset²⁵. As shown in Fig. 4a, hydrate-bound hydrocarbons at the Kedr MV have thermogenic and/or secondary microbial origins, whereas those of other gas hydrate sites (Malenky, Bolshoy, Malyutka, Peschanka P-2, Kukuy K-0, Kukuy K-2 and Goloustnoe; Fig. 1) in Lake Baikal demonstrate microbial or early mature thermogenic origins. The hydrate-bound C₁ from all locations except those at the Kedr MV were interpreted to be of microbial origin via methyl-type fermentation²³ according to Whiticar's old diagram²⁶; however, the revised diagram²⁵ suggests early mature thermogenic gases (Fig. 4b). Those of the Kedr MV plot at the boundary of the thermogenic and secondary microbial origin zones. Low C₁ and C₂ $\delta^{13}\text{C}$ at the Peschanka P-2 MV indicated that C₁ and C₂ are of microbial origin^{27,28}, whereas Kedr MV shows high C₁ and C₂ $\delta^{13}\text{C}$ indicating their thermogenic origin (Fig. 4c). At other sites, C₁ and C₂ $\delta^{13}\text{C}$ suggested that gases are mainly of microbial origin (in terms of C₁) with some thermogenic component (¹³C rich and higher concentration in C₂).

Stable isotopes in hydrate-bound C₁ at the Kedr-1 and Kedr-2 areas suggested its thermogenic origin. However, it is close to the field of secondary microbial C₁ in Fig. 4b, and the data are plotted in the overlap between the fields of thermogenic and secondary microbial in Fig. 4a. Milkov²⁹ mentioned that secondary microbial C₁ is characterised by C₁-rich dry gas, large C₁ $\delta^{13}\text{C}$ (between -55‰ and -35‰) and large CO₂ $\delta^{13}\text{C}$ (more than +2‰). Although hydrate-bound and sediment gases in the Kedr MV were not C₁ rich and contained 3%–15% of C₂, C₁ $\delta^{13}\text{C}$ was around -45‰, which agrees with the secondary microbial C₁. Because some data of secondary microbial gas are plotted outside the field on the original graph²⁵, we could include the gas data in the category of secondary microbial C₁ in Fig. 4b.

Figure 6 shows the relationship between C₁ $\delta^{13}\text{C}$ and CO₂ $\delta^{13}\text{C}$ in the sediment gas obtained using headspace gas method. According to the genetic diagram²⁵, gas hydrate cores are plotted at the zones of the thermogenic and secondary microbial origins, whereas the cores at the peripheral area are primary microbial. The headspace gas data of the hydrate-bearing cores in Fig. 6 seem to be plotted in the field of thermogenic gas (low CO₂ $\delta^{13}\text{C}$), but the effect of light CO₂ produced by methane oxidation in the subsurface layer also decreased CO₂ $\delta^{13}\text{C}$ as shown in Fig. 5. These results suggested that secondary microbial C₁ mixes into thermogenic gas. Coal-bearing sediments exist around the Kedr area^{21,22}, and secondary microbial C₁ can also form from coal beds³⁰. Hydrate-bound C₁ of secondary microbial origin has been only reported at the Alaska North Slope³¹. This study is another case for it.




Figure 5. Depth profiles of C₁ and CO₂ concentrations, C₁/(C₂+C₃) values, C₁ δ¹³C, C₁ δ²H, C₂ δ¹³C, C₂ δ²H and CO₂ δ¹³C in the headspace gas. cmblf, centimetres below lake floor.




Figure 6. A diagram of headspace gases. CO₂ δ¹³C plotted against C₁ δ¹³C, based on the classification of Milkov and Etiope²⁵.




Figure 7. Depth profiles of $C_2 \delta^2H$ of hydrate-bound and sediment gases. cmlbf, centimetres below lake floor.

Formation process of the sII gas hydrates. As stated before, the crystallographic structure of gas hydrates at the Kedr MV is mainly due to the composition of thermogenic C_2 in the volatile hydrocarbons. The concentration of C_3 , which is one of the sII-forming components, was two to three orders of magnitude smaller than that of C_2 , because biodegradation occurs and this preferentially reduces C_{3-5} of n -alkanes^{19,32,33}. The concentration of n - C_4 was smaller than that of i - C_4 , whereas that of n - C_5 was not detected (Table 1). $C_3 \delta^{13}C$ was around -10‰ , suggesting that light C_3 is consumed by microbial activity. Assuming that sediment gas C_{3+} can be ignored, sediment gas ratio C_1/C_2 at the study area was 30 ± 17 (mean and standard deviation), and the concentration of C_2 was $\sim 3\%$. Such a composition of thermogenic gas is, therefore, considered to be supplied from a deep sediment layer, forming sI gas hydrates composed of mainly C_1 and C_2 ^{11,12} in the lake floor sediment.

In the cases where sI gas hydrates plug and block migration pathways, upward fluid flow becomes more focused in other areas¹⁶. Once gas supply stops locally, gas hydrates begin to decompose, with the gas dissolving into gas-poor sediment pore water. In the system of C_1 and C_2 , C_2 is prone to be engaged in gas hydrate and decreases the equilibrium pressure of mixed-gas hydrate. Therefore, C_2 -rich gas hydrate forms in parallel with the decomposition of sI gas hydrate. The Colorado School of Mines Hydrate (CSMHYD) program³⁴ showed that C_2 -rich sII gas hydrate (C_2 concentration 17%) forms from mixed gas composed of C_1 and C_2 (C_2 concentration 3%). The C_2 concentration of hydrate-bound gas at the Kedr MV was $\sim 14\%$, agreeing fairly well with the results of the CSMHYD program. Such secondary generation of gas hydrates can produce compositions and crystallographic structures that are different from the original crystals. A calorimetric study of synthetic C_1 and C_2 mixed-gas hydrate revealed that double peaks of heat flow correspond to the dissociation process of C_1 and C_2 mixed-gas hydrate, suggesting that C_2 -rich gas hydrate forms simultaneously from dissociated gas and showed that the second heat flow peak correspond to the dissociation of C_2 -rich gas hydrate¹⁸. The PXRD and solid-state ^{13}C nuclear magnetic resonance techniques demonstrated that C_2 -rich sI gas hydrate forms in the dissociation process of $C_1 + C_2$ sI gas hydrate³⁵.

Among twenty hydrate-bound cores in the Kedr area, four cores contained sI only, seven cores had sII only, and seven cores showed sII at the upper layer and sI at the lower layer, as observed at the Kukuy K-2 MV^{13,16,17}. Furthermore, in the cores 2015St1GC15 and 2016St18GC2, gas hydrate structure had sI at the upper and lower layer, and sII at the middle layer. These results suggested that complex gas hydrate layers are composed of sI and sII in subsurface sediments as shown in the schematic illustration in Poort et al.¹⁶.

Depth profiles of $C_2 \delta^2H$ of gas hydrate cores from the Kedr MV are shown in Fig. 7. $C_2 \delta^2H$ of hydrate-bound gases varied between -227‰ and -206‰ , with a grouping around -210‰ . $C_2 \delta^2H$ of sediment gases was also around -210‰ , indicating that $C_2 \delta^2H$ of the original thermogenic gas is -210‰ . As stated above, $C_2 \delta^2H$ of some cores showed low values at their base. Based on the isotopic fractionation of hydrogen in C_2 during the formation of sI C_2 hydrate³⁶, δ^2H of hydrate-bound C_2 was 1.1‰ lower than that of residual C_2 . However, this is




Figure 8. Concentration of C_{3-5} against C_2 concentration in the hydrate-bound gases.

too small to explain the wide distribution in C_2 δ^2H shown in Fig. 7. On the other hand, Matsuda et al.³⁷ reported that isotopic fractionation of hydrogen in C_2 is dependent on the crystallographic structure: 1‰–2‰ for sI and ~10‰ for sII. Gas hydrates plotting around –220‰ in C_2 δ^2H can be explained as a secondary generation of sII from dissociated gas hydrates, of which C_2 δ^2H was around –210‰. However, some sII samples showed high C_2 δ^2H (around –210‰), whereas some sI samples showed low C_2 δ^2H (around –220‰). These results indicated that formation and dissociation processes of gas hydrates produce complicated isotopic profiles in C_2 δ^2H under non-equilibrium conditions.

Characteristics of hydrate-bound gases in sII. C_3 , $i-C_4$, $n-C_4$ and $neo-C_5$ can be engaged in the larger hexadecahedral cages of sII¹. $n-C_4$ and $neo-C_5$ can be engaged using a help gas (e.g. C_1) to fill in the smaller dodecahedral cages of sII, because they cannot form pure $n-C_4$ and $neo-C_5$ hydrates, respectively. Figure 8 shows the concentration of C_3 , $i-C_4$, $n-C_4$, $neo-C_5$ and $i-C_5$ plotted against C_2 concentration. The figure illustrates a clear division between sI (3–4%) and sII (14%) C_2 concentrations. Data points between C_2 concentrations of 5% and 13% were considered to have a mixture of sI and sII. Concentrations of C_3 , $i-C_4$, $n-C_4$ and $neo-C_5$ had a positive correlation with the concentration of C_2 , and these concentrations in sII were 1 or 2 orders of magnitude larger than those in sI, suggesting that C_3 , $i-C_4$, $n-C_4$ and $neo-C_5$ are engaged with C_2 in the sII formation process.

C_3 values of 0.001%–0.01%, ~0.0001% of $n-C_4$, and 0.0001%–0.01% of $neo-C_5$ were also detected in sI hydrate-bound gas (Fig. 8), despite these hydrocarbons being unable to be engaged in sI. This can be explained by gases being adsorbed with sediments and gas hydrate crystals, which are then trapped in the grain boundary of polycrystalline gas hydrate crystals, and the gases are engaged if a small amount of sII crystals are present. For example, Uchida et al.³⁸ examined natural gas hydrate retrieved at the Mackenzie Delta (onshore Canada) and detected C_3 engaged in sII using Raman spectroscopy, although PXRD results suggested that the sample was sI and the major component of hydrate-bound gas was C_1 (more than 99%).

$neo-C_5$ is considered to form from the decomposition of gem-dimethylcycloalkanes derived from the terpenes of terrestrial organic matter³⁹. It is easily enriched by preferential diffusion due to the nearly spherical molecules and its diffusion coefficient, which is higher than that of less branched isomers⁴⁰. The sII hydrates retrieved at the Kukuy K-2 MV (central Baikal basin) contained 0.026–0.064% of $neo-C_5$ in the volatile hydrocarbons^{13,14}, and those at the Kedr MV had a maximum value of 0.054% of $neo-C_5$ (Supplementary Information Table S1). On the contrary, in the case of natural gas hydrates retrieved at the Joetsu Basin (Japan Sea), $neo-C_5$ was excluded and remained in sediment during the formation of sI gas hydrates from C_1 -rich gas⁴¹. The molecular size of $i-C_5$ is

considerably large to be encaged in the large cages of sII. Maximum concentration of *i*-C₅ in the hydrate-bound gases was in several parts per million in both the fields of sI and sII (Fig. 8), indicating that *i*-C₅ is not a hydrate-bound hydrocarbon and adsorbed with gas hydrate crystals and/or trapped in their grain boundary.

Conclusion

We reported the molecular and stable isotope compositions of hydrate-bound and sediment gases at the Kedr MV in the southern Baikal basin. The empirical classifications of the molecular and isotopic compositions of hydrate-bound hydrocarbons showed that the gas source is mainly thermogenic, one of the end members of hydrate-bound gases in Lake Baikal. Large CO₂ δ¹³C in the sediment gases suggested that secondary microbial C₁ mixes with thermogenic gas. Double-structure gas hydrates composed of sI and sII were observed, likely created by sI crystals partly dissociating and C₂-rich sII crystals forming. The C₂ δ²H values of hydrate-bound gas revealed that light C₂ is preferentially encaged into sII crystals; however, some exceptions indicated that more complicated processes of gas exchange might exist between sI, sII and the dissolved gas in pore water. Because C₂ is preferentially concentrated into the gas hydrate phase, high concentration of thermogenic C₂ produce sII crystals with C₁, and C₂ is encaged into the large cages of sII with C₃, *i*-C₄, *n*-C₄, and *neo*-C₅ in the re-crystallisation process.

Methods

Gas hydrate crystals were collected onboard R/V *G. Yu. Vereshchagin* and stored in liquid nitrogen. Powder X-ray diffraction (PXRD) measurements were performed to check the crystallographic structure. Samples for PXRD were finely ground at a temperature of 77 K and then kept at 173 K. The X-ray diffraction patterns were recorded at 2θ = 5–35° using Cu Kα radiation (λ = 1.5418 Å) and a Bruker D8 Advance diffractometer equipped with a TTK 450 Anton Paar temperature controlling device. The positions of diffraction peaks corresponding to sI and sII hydrates and hexagonal ice (Ih) were calculated with the use of reference data on space group and unit cell parameters of the respective compounds⁴².

Hydrate-bound gases were collected using water displacement method and stored in 5 mL glass vials sealed with butyl septum stoppers. To avoid microbial alteration, 0.3 mL of preservative (50 wt% aqueous solution of benzalkonium chloride) was introduced into the vials. Gas sampling was conducted for each layer of gas hydrate in the hydrate-bearing cores. Several vials of samples were taken from a hydrate nodule. Sediment gases were collected using headspace gas method to calculate the depth profiles of each gas component in the sediment cores. To create a 5 mL headspace, 10 mL of sediment and 10 mL of saturated aqueous solution of NaCl were introduced into 25 mL-glass vials. The headspace was flushed with helium, the carrier gas used in the gas chromatography, to reduce air contamination⁴³. The headspace gases were then placed into the 5 mL glass vials to separate them from sediment particles or water and to prevent any microbial activity during storage.

The molecular compositions of the hydrocarbons (from C₁ to C₅) were determined using a gas chromatograph (GC-2014, Shimadzu, Kyoto, Japan) equipped with a packed column (Shimadzu Sunpak-S; length 2 m, inner diameter [ID] 3 mm), along with a thermal conductivity detector and flame ionisation detector for detecting high and low concentrations of hydrocarbons respectively. The two detectors were connected in series. The detection limit was 0.5 ppmv (C₁–C₃) and 5 ppmv (C₄–C₅). The analytical error estimated by multiple injections of standard gases was < 1.2% for each gas component. Stable carbon and hydrogen isotopic ratios of the hydrocarbons and CO₂ were measured using a continuous-flow isotope-ratio mass spectrometer (CF-IRMS, DELTA V, Thermo Fisher Scientific, Waltham, MA, USA) coupled with a gas chromatograph (TRACE GC Ultra, Thermo Fisher Scientific). The gas chromatograph was equipped with a Carboxen-1006 PLOT capillary column (length 30 m, ID 0.32 mm, film thickness 15 μm, Sigma-Aldrich, St. Louis, MO, USA). In the case of samples with low C₁ concentration, a Sigma-Aldrich Carboxen-1010 PLOT capillary column (length 30 m, ID 0.32 mm, film thickness 15 μm) was also used to separate air components from C₁. Stable isotope compositions were reported as δ values (in per-mille):

$$\delta[\text{‰}] = \left(\frac{R_{\text{sample}} - R_{\text{standard}}}{R_{\text{standard}}} \right) \times 1000 \quad (1)$$

where R denotes the ¹³C/¹²C or ²H/¹H ratio. δ¹³C and δ²H are given with reference to the V-PDB and V-SMOW standards, respectively, determined using NIST RM8544 (NBS19) for δ¹³C and NIST RM8561 (NGS3) for δ²H. The analytical precision was 0.3‰ for hydrocarbon (C₁–C₃) δ¹³C and 1‰ for δ²H.

Data availability

All the gas data are reported in the Supplementary Information.

Received: 16 April 2020; Accepted: 12 August 2020

Published online: 08 September 2020

References

- Sloan, E. D. & Koh, C. A. *Clathrate hydrates of natural gases* (CRC Press, Boca Raton, FL, 2008).
- Makogon, Y. F., Holditch, S. A. & Makogon, T. Y. Natural gas-hydrates—a potential energy source for the 21st Century. *J. Pet. Sci. Eng.* **56**, 14–31. <https://doi.org/10.1016/j.petrol.2005.10.009> (2007).
- Boswell, R. & Collett, T. S. Current perspectives on gas hydrate resources. *Energy Environ. Sci.* **4**, 1206–1215. <https://doi.org/10.1039/C0EE00203H> (2011).
- Kennedy, M., Mrofka, D. & von der Borch, C. Snowball Earth termination by destabilization of equatorial permafrost methane clathrate. *Nature* **453**, 642–645. <https://doi.org/10.1038/nature06961> (2008).

5. Milkov, A. V. Global estimates of hydrate-bound gas in marine sediments: how much is really out there? *Earth Sci. Rev.* **66**, 183–197. <https://doi.org/10.1016/j.earscirev.2003.11.002> (2004).
6. Milkov, A. V. & Sassen, R. Two-dimensional modeling of gas hydrate decomposition in the northwestern Gulf of Mexico: significance to global change assessment. *Glob. Planet Change* **36**, 31–46. [https://doi.org/10.1016/S0921-8181\(02\)00162-5](https://doi.org/10.1016/S0921-8181(02)00162-5) (2003).
7. Milkov, A. V. *et al.* Ethane enrichment and propane depletion in subsurface gases indicate gas hydrate occurrence in marine sediments at southern Hydrate Ridge offshore Oregon. *Org. Geochem.* **35**, 1067–1080. <https://doi.org/10.1016/j.orggeochem.2004.04.003> (2004).
8. Sassen, R., Sweet, S. T., DeFreitas, D. A. & Milkov, A. V. Exclusion of 2-methylbutane (isopentane) during crystallization of structure II gas hydrate in sea-floor sediment Gulf of Mexico. *Org. Geochem.* **31**, 1257–1262. [https://doi.org/10.1016/S0146-6380\(00\)00144-3](https://doi.org/10.1016/S0146-6380(00)00144-3) (2000).
9. Khlystov, O. *et al.* Gas hydrate of lake baikal: discovery and varieties. *J. Asian Earth Sci.* **62**, 162–166. <https://doi.org/10.1016/j.jseaes.2012.03.009> (2013).
10. Khlystov, O. M., Khabuev, A. V., Minami, H., Hachikubo, A. & Krylov, A. A. Gas hydrates in Lake Baikal. *Limnol. Freshwater Biol.* **1**, 66–70 <https://doi.org/10.31951/2658-3518-2018-A-1-66> (2018).
11. Subramanian, S., Kini, R. A., Dec, S. F. & Sloan, E. D. Jr. Evidence of structure II hydrate formation from methane + ethane mixtures. *Chem. Eng. Sci.* **55**, 1981–1999. [https://doi.org/10.1016/S0009-2509\(99\)00389-9](https://doi.org/10.1016/S0009-2509(99)00389-9) (2000).
12. Subramanian, S., Ballard, A. L., Kini, R. A., Dec, S. F. & Sloan, E. D. Jr. Structural transitions in methane + ethane gas hydrates — part I: upper transition point and applications. *Chem. Eng. Sci.* **55**, 5763–5771. [https://doi.org/10.1016/S0009-2509\(00\)00162-7](https://doi.org/10.1016/S0009-2509(00)00162-7) (2000).
13. Kida, M. *et al.* Coexistence of structure I and II gas hydrates in Lake Baikal suggesting gas sources from microbial and thermogenic origin. *Geophys. Res. Lett.* **33**, L24603. <https://doi.org/10.1029/2006GL028296> (2006).
14. Kida, M. *et al.* Natural gas hydrates with locally different cage occupancies and hydration numbers in Lake Baikal. *Geochem. Geophys. Geosyst.* **10**, Q05003. <https://doi.org/10.1029/2009GC002473> (2009).
15. Hachikubo, A. *et al.* Model of formation of double structure gas hydrates in Lake Baikal based on isotopic data. *Geophys. Res. Lett.* **36**, L18504. <https://doi.org/10.1029/2009GL039805> (2009).
16. Poort, J. *et al.* Thermal anomalies associated with shallow gas hydrates in the K-2 mud volcano Lake Baikal. *Geo-Mar. Lett.* **32**, 407–417. <https://doi.org/10.1007/s00367-012-0292-0> (2012).
17. Manakov, A. Yu., Khlystov, O. M., Hachikubo, A. & Ogienko, A. G. A physicochemical model for the formation of gas hydrates of different structural types in K-2 mud volcano (Kukui Canyon, Lake Baikal). *Rus. Geol. Geophys.* **54**, 475–482. <https://doi.org/10.1016/j.rgg.2013.03.009> (2013).
18. Hachikubo, A. *et al.* Dissociation heat of mixed-gas hydrate composed of methane and ethane. In *Proc. 6th Int. Conf. on Gas Hydrates*, 6–10 July, 2008, Vancouver, Canada (2008). <https://hdl.handle.net/2429/2694>
19. Kalmychkov, G. V., Pokrovsky, B. G., Hachikubo, A. & Khlystov, O. M. Geochemical characteristics of methane from sediments of the underwater high Posolskaya Bank (Lake Baikal). *Lithol. Min. Resour.* **52**, 102–110. <https://doi.org/10.1134/S0024490217020055> (2017).
20. Minami, H. *et al.* Hydrogen and oxygen isotopic anomalies in pore waters suggesting clay mineral dehydration at gas hydrate-bearing Kedr mud volcano, southern Lake Baikal Russia. *Geo-Mar. Lett.* **38**, 403–415. <https://doi.org/10.1007/s00367-018-0542-x> (2018).
21. Rasskazov, S. V. *et al.* Sediments in the Tertiary Tankhoi field, south Baikal basin: stratigraphy, correlation and structural transformations in the Baikal region (in Russian). *Geodyn. Tectonophys.* **5**, 993–1032. <https://doi.org/10.5800/GT-2014-5-4-0165> (2014).
22. Khlystov, O. M. *et al.* New evidence on the relief of the southern underwater slope in the south Baikal basin. *Geogr. Nat. Resour.* **39**, 33–38. <https://doi.org/10.1134/S1875372818010055> (2018).
23. Hachikubo, A. *et al.* Molecular and isotopic characteristics of gas hydrate-bound hydrocarbons in southern and central Lake Baikal. *Geo-Mar. Lett.* **30**, 321–329. <https://doi.org/10.1007/s00367-010-0203-1> (2010).
24. Bernard, B. B., Brooks, J. M. & Sackett, W. M. Natural gas seepage in the Gulf of Mexico. *Earth Planet. Sci. Lett.* **31**, 48–54. [https://doi.org/10.1016/0012-821X\(76\)90095-9](https://doi.org/10.1016/0012-821X(76)90095-9) (1976).
25. Milkov, A. V. & Etiope, G. Revised genetic diagrams for natural gases based on a global dataset of >20,000 samples. *Org. Geochem.* **125**, 109–120. <https://doi.org/10.1016/j.orggeochem.2018.09.002> (2018).
26. Whitticar, M. J. Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane. *Chem. Geol.* **161**, 291–314. [https://doi.org/10.1016/S0009-2541\(99\)00092-3](https://doi.org/10.1016/S0009-2541(99)00092-3) (1999).
27. Taylor, S. W., SherwoodLollar, B. & Wassenaar, L. I. Bacteriogenic ethane in near-surface aquifers: Implications for leaking hydrocarbon well bores. *Environ. Sci. Technol.* **34**, 4727–4732. <https://doi.org/10.1021/es001066x> (2000).
28. Milkov, A. V. Molecular and stable isotope compositions of natural gas hydrates: a revised global dataset and basic interpretations in the context of geological settings. *Org. Geochem.* **36**, 681–702. <https://doi.org/10.1016/j.orggeochem.2005.01.010> (2005).
29. Milkov, A. V. Worldwide distribution and significance of secondary microbial methane formed during petroleum biodegradation in conventional reservoirs. *Org. Geochem.* **42**, 184–207. <https://doi.org/10.1016/j.orggeochem.2010.12.003> (2011).
30. Scott, A. R., Kaiser, W. R. & Ayers, W. B. Jr. Thermogenic and secondary biogenic gases, San Juan Basin, Colorado and New Mexico – implications for coalbed gas producibility. *Am. Assoc. Pet. Geol. Bull.* **78**, 1186–1209. <https://doi.org/10.1306/A25FEAA9-171B-11D7-8645000102C1865D> (1994).
31. Lorenson, T. D., Collett, T. S. & Hunter, R. B. Gas geochemistry of the Mount Elbert Gas Hydrate Stratigraphic Test Well, Alaska North Slope: Implications for gas hydrate exploration in the Arctic. *Mar. Petrol. Geol.* **28**, 343–360. <https://doi.org/10.1016/j.marpetgeo.2010.02.007> (2011).
32. James, A. T. & Burns, B. J. Microbial alteration of subsurface natural gas accumulations. *AAPG Bull.* **68**, 957–960 (1984).
33. Katz, B. J. Microbial processes and natural gas accumulations. *Open Geol. J.* **5**, 75–83. <https://doi.org/10.2174/1874262901105010075> (2011).
34. Sloan, E. D. Jr. *Clathrate hydrates of natural gases* (Marcel Dekker, NY, 1998).
35. Kida, M., Jin, Y., Takahashi, N., Nagao, J. & Narita, H. Dissociation behavior of methane-ethane mixed gas hydrate coexisting structures I and II. *J. Phys. Chem. A* **114**, 9456–9461. <https://doi.org/10.1021/jp1055667> (2010).
36. Hachikubo, A. *et al.* Isotopic fractionation of methane and ethane hydrates between gas and hydrate phases. *Geophys. Res. Lett.* **34**, L21502. <https://doi.org/10.1029/2007GL030557> (2007).
37. Matsuda, J., Hachikubo, A., Ozeki, T. & Takeya, S. Effect of crystallographic structure on hydrogen isotope fractionation of ethane in the system of methane and ethane mixed-gas hydrate (in Japanese). *Annu. Rep. on Snow and Ice Studies in Hokkaido* **37**, 27–30 (2018).
38. Uchida, T. *et al.* Physical properties of natural gas hydrate and associated gas-hydrate-bearing sediments in the JAPEX/JNOC/GSC *et al.* Mallik 5L-38 gas hydrate production research well in Scientific Results from the Mallik 2002 Gas Hydrate Production Research Well Program, Mackenzie Delta, Northwest Territories, Canada (eds. Dallimore, S. R. & Collett, T. S.) (Geological Survey of Canada, Bulletin 585, 2005).
39. Hunt, J. M. & Whelan, J. K. Dissolved gases in Black Sea sediments. *DSDP Initial Rep.* **42**, 661–665. <https://doi.org/10.2973/dsdp.proc.42-2.125.1978> (1978).
40. Schaefer, R. G. & Leythaeuser, D. C₂–C₈ hydrocarbons in sediments from Deep Sea Drilling Project Leg 75, holes 530A, Angola Basin, and 532 Walvis Ridge. *DSDP Initial Rep.* **75**, 1055–1067. <https://doi.org/10.2973/dsdp.proc.75.137.1984> (1984).

41. Hachikubo, A., Yanagawa, K., Tomaru, H., Lu, H. & Matsumoto, R. Molecular and isotopic composition of volatiles in gas hydrates and in pore water from Joetsu Basin, eastern margin of Japan Sea. *Energies* **8**, 4647–4666. <https://doi.org/10.3390/en8064647> (2015).
42. Manakov, A. Y., Kosyakov, V. I. & Solodovnikov, S. F. Structural chemistry of clathrate hydrates and related compounds in *Comprehensive Supramolecular Chemistry II* (ed. Atwood, J. L.) 161–206 (Oxford, Elsevier, 2017).
43. Sakagami, H. *et al.* Molecular and isotopic composition of hydrate-bound and sediment gases in the southern basin of Lake Baikal, based on an improved headspace gas method. *Geo-Mar. Lett.* **32**, 465–472. <https://doi.org/10.1007/s00367-012-0294-y> (2012).

Acknowledgements

We greatly appreciate the assistance of the shipboard crews of RV *G. Yu. Vereshchagin* during the Lake Baikal expeditions. We are grateful to Prof. A. Milkov, Dr. G. G. Akhmanov and an anonymous reviewer for their constructive comments. This work was supported by funding agencies in Japan (Japan Society for the Promotion of Science KAKENHI 26303021, 16H05760 and 17H03300 and the presidential grant of the Kitami Institute of Technology). The work was also supported by the project of LIN SB RAS (AAAA-A16-116122110064-7), IGC SB RAS # IX.127.1.2. (0350-2019-0004), RSF-19-17-00226 and 01-FPK-19 of the INRTU. X-ray diffraction studies were supported by paragraph V.44.4.9. ‘Development of the scientific foundations of the physical chemistry of clathrate (gas), semi-clathrate and ionic clathrate hydrates’ of the program of basic scientific research ‘Development of the scientific basis of the directed synthesis of new inorganic and coordination compounds and functional materials based on them’.

Author contributions

A.H. designed the study, performed gas analysis and drafted the manuscript; H.M., S.Y., A.K. and A.K. conducted the fieldwork and prepared the data; G.K. helped with gas analysis; J.P. conceived the model of gas hydrate formation; M.D. revised and edited the manuscript; A.C. conducted geophysical survey; A.M. did the PXRD analysis and O.K. designed the framework of the cruises. All authors contributed to the drafts and gave final approval for publication.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-71410-2>.

Correspondence and requests for materials should be addressed to A.H.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020