



# Crystal structure of 2-amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-*a*]quinoxalin-11-ium bromide monohydrate

Md. Serajul Haque Faizi<sup>a</sup> and Yuliia Parashchenko<sup>b\*</sup>

Received 25 August 2015

Accepted 29 September 2015

Edited by G. Smith, Queensland University of Technology, Australia

**Keywords:** crystal structure; hydrogen bonding; quinoxaline derivatives; pyrido[1,2-*a*]quinoxalines

**CCDC reference:** 1428593

**Supporting information:** this article has supporting information at journals.iucr.org/e


<sup>a</sup>Department of Chemistry, College of Science, Sultan Qaboos University, PO Box 36 Al-Khod 123, Muscat, Sultanate of Oman, and <sup>b</sup>National Taras Shevchenko University, Department of Chemistry, Volodymyrska str. 64, 01601 Kiev, Ukraine. \*Correspondence e-mail: lyulya200288@mail.ru

In the title hydrated salt,  $C_{12}H_8Br_2N_3O^+ \cdot Br^- \cdot H_2O$ , which was synthesized by the reaction of the pyridine derivative Schiff base *N*<sup>1</sup>,*N*<sup>4</sup>-bis(pyridine-2-ylmethylene)benzene-1,4-diamine with bromine, the asymmetric unit contains a 2-amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-*a*]quinoxalin-11-ium cation, with a protonated pyridine moiety, a bromide anion and a water molecule of solvation. The cation is non-planar with the dibromo-substituted benzene ring, forming dihedral angles of 24.3 (4) and 11.5 (4)° with the fused pyridine and pyrazine ring moieties, respectively. In the crystal, the cations are linked through a centrosymmetric hydrogen-bonded cyclic  $R_4^2(8)$   $Br_2(H_2O)_2$  unit by  $N-H \cdots Br$ ,  $N-H \cdots O$  and  $O-H \cdots Br$  hydrogen bonds, forming one-dimensional ribbons extending along *b*, with the planes of the cations lying parallel to (100).


## 1. Chemical context

Quinoxaline and its derivatives are an important class of benzo-heterocycles (Kurasawa *et al.*, 1988; Cheeseman & Werstiuk, 1978), displaying a broad spectrum of biological activities (Seitz *et al.*, 2002; Toshima *et al.*, 2002) which have made them important structures in combinatorial drug-discovery literature (Wu & Ede, 2001; Lee *et al.*, 1997). These compounds have also found applications as dyes (Zaragoza *et al.*, 1999; Sonawane & Rangnekar, 2002) and building blocks in the synthesis of organic semiconductors (Katoh *et al.*, 2000; Dailey *et al.*, 2001) and they also serve as useful rigid subunits in macrocyclic receptors for molecular recognition (Mizuno *et al.*, 2002) and chemically controllable switches (Elwahi, 2000). The present work is a part of an ongoing structural study of Schiff bases and their utilization in the synthesis of new organic and polynuclear coordination compounds (Faizi & Sen, 2014; Moroz *et al.*, 2012). We report here the synthesis and crystal structure of 2-amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-*a*]quinoxalin-11-ium bromide monohydrate (refcode ADOQBM). Previously, we have reported new methods for the preparation of substituted quinoxaline derivatives together with their crystallographic characterization. However, there are very few reported structures of compounds similar to the title compound, one being the doubly protonated dibromide salt 2-azaniumyl-3-bromo-6-oxo-5,6-dihydropyrido[1,2-*a*]quinoxalin-11-ium dibromide (Faizi *et al.*, 2015).

The title singly protonated monobromide monohydrate salt,  $C_{12}H_8Br_2N_3O^+ \cdot Br^- \cdot H_2O$ , was synthesized from the reaction the pyridine derivative Schiff base *N*1,*N*4-bis(pyridine-2-ylmethylene)benzene-1,4-diamine (BPYBD) with molecular bromine. The cyclization occurs by oxidation of BPYBD,




reduction of molecular bromine and finally hydrolysis of the imine bond which creates the charge at the pyridine nitrogen atom in the quinoxaline ring system. The structure is reported herein.



## 2. Structural commentary

The asymmetric unit of the title compound contains a discrete 2-amino-1,3-dibromo-6-oxo-5,6-dihydro-1,2-aquinoxalin-11-ium cation with a protonated pyridine moiety, and a bromide counter-anion and a water molecule of solvation (Fig. 1). The cation is non-planar compared to the previously reported structure (Faizi *et al.*, 2015). The mean plane of the



**Figure 1**  
The molecular conformation and atom-numbering scheme for the title compound, with non-H atoms drawn as 40% probability displacement ellipsoids.

**Table 1**  
Hydrogen-bond geometry (Å, °).


$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N2-H5\cdots Br3^i$	0.86	2.49	3.332 (6)	166
$N3-H3B\cdots Br1$	0.86	2.60	3.048 (7)	113
$N3-H3B\cdots Br3^{iii}$	0.86	2.84	3.581 (7)	145
$N3-H3A\cdots O1^{iii}$	0.86	2.17	2.977 (9)	155
$N3-H3A\cdots Br2$	0.86	2.56	3.006 (7)	113
$O2-H11\cdots Br3^{iv}$	0.89	2.50	3.383 (6)	180
$O2-H12\cdots Br1^v$	0.88	2.61	3.309 (7)	137

Symmetry codes: (i)  $x, y+1, z-1$ ; (ii)  $x, y, z-1$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+1, -y, -z+1$ ; (v)  $x, y, z+1$ .


pyridine ring forms a dihedral angle of  $24.2(4)^\circ$  with the benzene ring and  $14.6(4)^\circ$  with the pyrazine ring of the fused system while the dihedral angle between the pyrazine and the benzene ring is  $11.5(4)^\circ$ . A shorter  $C10-N3$  distance of  $1.367(9)$  Å, compared to the usual aromatic  $C-N_{amine}$  single bond distance of  $1.43(3)$  Å, might be due to the electron-withdrawing effect of the positively charged pyridine N atom, and the *ortho*-substituted bromine atom which decreases the  $C-N_{amine}$  bond order. Other  $C-C$  and  $C-N$  bond distances are well within the limits expected for aromatic rings (Koner & Ray, 2008; Kandal *et al.*, 2005; Fritsky *et al.*, 2006). Present also in the cations are intramolecular  $N3-H\cdots Br1$  and  $N3-H\cdots Br2$  interactions [ $3.048(7)$ ,  $3.006(7)$  Å, respectively, Table 1].

## 3. Supramolecular features

In the crystal, the cations are linked through a centrosymmetric hydrogen-bonded cyclic  $R_4^2(8)$   $Br_2(H_2O)_2$  unit and  $N-H\cdots Br$ ,  $N-H\cdots O$  and  $O-H\cdots Br$  hydrogen bonds (Table 1), forming broad one-dimensional ribbons extending along  $b$  (Fig. 2). The planes of the cations lie parallel to (100). Fig. 3 shows the packing in the unit cell, viewed along the  $b$  axis, in which layers of quinoxalinium cations are embedded between



**Figure 2**  
The one-dimensional hydrogen-bonded ribbon structure, viewed along the  $a$ -axis direction. Inter-species interactions are shown as dashed lines.



**Figure 3**  
The layering of the ribbon structures, viewed along the *b* axis.

ionic layers of anions and *vice versa*, forming an alternating hydrocarbon–ionic layer structure. No intermolecular  $\pi$ – $\pi$  interactions are evident in the hydrocarbon layer in the structure.

#### 4. Database survey

There are very few examples of similar compounds in the literature, a search of the Cambridge Structural Database (Version 5.35, May 2014; Groom & Allen, 2014) revealing the structure of 2-azaniumyl-3-bromo-6-oxo-5,6-dihydropyrido-[1,2-*a*]quinoxalin-11-ium dibromide (Faizi *et al.*, 2015), in which the 2-amino-1,2-dibromide ring in the title compound is replaced by a 2-azaniumyl-3-bromo ring. Other similar structures have been reported (Faizi & Sen, 2014; Koner *et al.*, 2008).

#### 5. Synthesis and crystallization

Molecular bromine (440 mg, 144.0  $\mu$ L, 2.80 mmol) was added to a methanolic solution (10 mL) of Schiff base, *N*1,*N*4-bis(pyridine-2-ylmethylene)benzene-1,4-diamine (BPYBD) (197 mg, 0.70 mmol). The color of the solution immediately changed from yellow to orange. The reaction mixture was stirred for 4 h at room temperature under a fume hood. The resulting yellow precipitate was recovered by filtration, washed several times with small portions of acetone and then with diethyl ether to give 200 mg (yield: 64%) of 2-amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-*a*]quinoxalin-11-ium bromide monohydrate (ADOQBM). The crystal of the title compound suitable for X-ray analysis was obtained within three days by slow evaporation of a solution of the compound in methanol.

#### 6. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. All N-bound H atoms were located

**Table 2**  
Experimental details.

Crystal data	
Chemical formula	$C_{12}H_8Br_2N_3O^+ \cdot Br^- \cdot H_2O$
$M_r$	467.93
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	100
$a, b, c$ (Å)	7.5069 (7), 9.7435 (10), 10.782 (1)
$\alpha, \beta, \gamma$ (°)	88.490 (7), 73.798 (7), 71.981 (7)
$V$ (Å <sup>3</sup> )	718.61 (12)
$Z$	2
Radiation type	Mo $K\alpha$
$\mu$ (mm <sup>-1</sup> )	8.42
Crystal size (mm)	0.20 × 0.15 × 0.11
Data collection	
Diffractometer	Bruker SMART APEX CCD
Absorption correction	Multi-scan ( <i>SADABS</i> ; Bruker, 2003)
$T_{min}$ , $T_{max}$	0.259, 0.365
No. of measured, independent and observed [ $I > 2\sigma(I)$ ] reflections	8077, 2187, 1681
$R_{int}$	0.163
$\theta_{max}$ (°)	23.8
$(\sin \theta/\lambda)_{max}$ (Å <sup>-1</sup> )	0.568
Refinement	
$R[F^2 > 2\sigma(F^2)]$ , $wR(F^2)$ , $S$	0.059, 0.155, 1.00
No. of reflections	2187
No. of parameters	181
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}$ , $\Delta\rho_{min}$ (e Å <sup>-3</sup> )	1.18, -1.16

Computer programs: *SMART* and *SAINT* (Bruker, 2003), *SIR97* (Altomare *et al.*, 1999), *SHELXL97* (Sheldrick, 2008) and *DIAMOND* (Brandenburg & Putz, 2006).

in difference-Fourier maps and their positions were then held fixed. The isotropic displacement parameters were refined for these atoms. Aromatic H atoms were placed in calculated positions and treated as riding on their parent C atoms [ $C-H = 0.93$  Å and  $U_{iso}(H) = 1.2$  or  $1.5U_{eq}(C)$ ].

#### Acknowledgements

The authors are grateful to the Department of Chemistry, Indian Institute of Technology Kanpur, Kanpur, UP-208016, India, for X-ray data collection.

#### References

- Altomare, A., Burla, M. C., Camalli, M., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Brandenburg, K. & Putz, H. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2003). *SMART*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cheeseman, G. W. H. & Werstiuk, E. S. G. (1978). *Adv. Heterocycl. Chem.* **22**, 367–431.
- Dailey, S., Feast, J. W., Peace, R. J., Sage, I. C., Till, S. & Wood, E. L. (2001). *J. Mater. Chem.* **11**, 2238–2243.
- Elwahi, A. H. M. (2000). *Tetrahedron*, **56**, 897–907.
- Faizi, M. S. H. & Sen, P. (2014). *Acta Cryst.* **E70**, m206–m207.
- Faizi, M. S. H., Sharkina, N. O. & Iskenderov, T. S. (2015). *Acta Cryst.* **E71**, o17–o18.
- Fritsky, I. O., Kozłowski, H., Kandal, O. M., Haukka, M., Świątek-Kozłowska, J., Gumienna-Kontecka, E. & Meyer, F. (2006). *Chem. Commun.* pp. 4125–4127.

- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Kanderal, O. M., Kozłowski, H., Dobosz, A., Świątek-Kozłowska, J., Meyer, F. & Fritsky, I. O. (2005). *Dalton Trans.* pp. 1428.
- Katoh, A., Yoshida, T. & Ohkanda, J. (2000). *Heterocycles*, **52**, 911–920.
- Koner, R. R. & Ray, M. (2008). *Inorg. Chem.* **47**, 9122–9124.
- Kurasawa, Y., Sakata, G. & Makino, K. (1988). *Heterocycles*, **27**, 2481–2515.
- Lee, J., Murray, W. V. & Rivero, R. A. (1997). *J. Org. Chem.* **62**, 3874–3879.
- Mizuno, T., Wei, W.-H., Eller, L. R. & Sessler, J. L. (2002). *J. Am. Chem. Soc.* **124**, 1134–1135.
- Moroz, Y. S., Demeshko, S., Haukka, M., Mokhir, A., Mitra, U., Stocker, M., Müller, P., Meyer, F. & Fritsky, I. O. (2012). *Inorg. Chem.* **51**, 7445–7447.
- Seitz, L. E., Suling, W. J. & Reynolds, R. C. (2002). *J. Med. Chem.* **45**, 5604–5606.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sonawane, N. D. & Rangnekar, D. W. (2002). *J. Heterocycl. Chem.* **39**, 303–308.
- Toshima, K., Takano, R., Ozawa, T. & Matsumura, S. (2002). *Chem. Commun.* pp. 212–213.
- Wu, Z. & Ede, N. J. (2001). *Tetrahedron Lett.* **42**, 8115–8118.
- Zaragoza, F. & Stephensen, H. (1999). *J. Org. Chem.* **64**, 2555–2557.

## supporting information

*Acta Cryst.* (2015). E71, 1332-1335 [doi:10.1107/S2056989015018253]

## Crystal structure of 2-amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-a]quinoxalin-11-ium bromide monohydrate

Md. Serajul Haque Faizi and Yuliia Parashchenko

### Computing details

Data collection: *SMART* (Bruker, 2003); cell refinement: *SAINTE* (Bruker, 2003); data reduction: *SAINTE* (Bruker, 2003); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg & Putz, 2006); software used to prepare material for publication: *DIAMOND* (Brandenburg & Putz, 2006).

### 2-Amino-1,3-dibromo-6-oxo-5,6-dihydropyrido[1,2-a]quinoxalin-11-ium bromide monohydrate

#### Crystal data

$C_{12}H_8Br_2N_3O^+ \cdot Br^- \cdot H_2O$

$M_r = 467.93$

Triclinic,  $P\bar{1}$

$a = 7.5069$  (7) Å

$b = 9.7435$  (10) Å

$c = 10.782$  (1) Å

$\alpha = 88.490$  (7)°

$\beta = 73.798$  (7)°

$\gamma = 71.981$  (7)°

$V = 718.61$  (12) Å<sup>3</sup>

$Z = 2$

$F(000) = 448$

$D_x = 2.163$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1023 reflections

$\theta = 1.5$ – $23.5$ °

$\mu = 8.42$  mm<sup>-1</sup>

$T = 100$  K

Block, yellow

$0.20 \times 0.15 \times 0.11$  mm

#### Data collection

Bruker SMART APEX CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

/w-scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2003)

$T_{\min} = 0.259$ ,  $T_{\max} = 0.365$

8077 measured reflections

2187 independent reflections

1681 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.163$

$\theta_{\max} = 23.8$ °,  $\theta_{\min} = 2.0$ °

$h = -8 \rightarrow 8$

$k = -11 \rightarrow 10$

$l = -12 \rightarrow 12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.059$

$wR(F^2) = 0.155$

$S = 1.00$

2187 reflections

181 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0902P)^2]$$

where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 1.18 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -1.16 \text{ e } \text{\AA}^{-3}$$

### Special details

**Experimental.** The OH H-atom was located in difference Fourier map and refined with with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{O})$ . The N- and C-bound H-atoms were positioned geometrically and refined using a riding model: N—H = 0.86 Å and C—H = 0.93 Å with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent atom})$ .

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br3	0.26076 (12)	0.03614 (8)	0.73364 (8)	0.0441 (3)
Br2	0.27561 (13)	0.36785 (8)	0.22264 (8)	0.0446 (3)
Br1	0.30008 (13)	0.45076 (9)	-0.29950 (8)	0.0480 (3)
C3	0.1893 (12)	0.8452 (10)	0.4590 (8)	0.048 (2)
H3	0.2041	0.8791	0.5342	0.058*
C1	0.1158 (12)	0.6740 (8)	0.3479 (7)	0.0388 (19)
H1	0.0617	0.5998	0.3495	0.047*
N1	0.1941 (9)	0.7217 (6)	0.2309 (6)	0.0316 (14)
C2	0.1146 (14)	0.7316 (9)	0.4612 (8)	0.048 (2)
H2	0.0644	0.6953	0.5393	0.057*
C4	0.2417 (12)	0.9077 (9)	0.3430 (8)	0.046 (2)
H4	0.2805	0.9898	0.3414	0.055*
C12	0.2134 (10)	0.6506 (7)	0.1102 (7)	0.0306 (17)
N2	0.2314 (10)	0.8738 (6)	0.0112 (6)	0.0377 (16)
H5	0.2190	0.9269	-0.0528	0.045*
C7	0.2241 (11)	0.7334 (7)	0.0019 (7)	0.0327 (18)
C11	0.2384 (11)	0.5009 (7)	0.0921 (7)	0.0318 (18)
C9	0.2558 (11)	0.5314 (8)	-0.1296 (8)	0.0362 (19)
C10	0.2586 (10)	0.4390 (7)	-0.0288 (7)	0.0324 (18)
C8	0.2379 (11)	0.6744 (7)	-0.1175 (8)	0.0349 (18)
H6	0.2351	0.7313	-0.1879	0.042*
C5	0.2364 (11)	0.8482 (8)	0.2307 (8)	0.0353 (18)
C6	0.2565 (12)	0.9308 (8)	0.1142 (8)	0.042 (2)
O1	0.2841 (12)	1.0476 (6)	0.1172 (7)	0.071 (2)
N3	0.2950 (11)	0.2940 (7)	-0.0508 (7)	0.0457 (18)
H3A	0.3064	0.2371	0.0109	0.055*
H3B	0.3066	0.2597	-0.1263	0.055*
O2	0.3545 (11)	0.2190 (7)	0.4635 (6)	0.072 (2)
H12	0.3415	0.2338	0.5459	0.108*

H11                    0.4550                    0.1520                    0.4120                    0.108\*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br3	0.0499 (5)	0.0360 (5)	0.0489 (5)	-0.0143 (4)	-0.0173 (4)	0.0057 (4)
Br2	0.0516 (6)	0.0273 (5)	0.0573 (6)	-0.0140 (4)	-0.0184 (4)	0.0127 (4)
Br1	0.0565 (6)	0.0389 (5)	0.0510 (6)	-0.0130 (4)	-0.0206 (4)	-0.0041 (4)
C3	0.044 (5)	0.051 (6)	0.051 (5)	-0.013 (4)	-0.017 (4)	-0.003 (4)
C1	0.046 (5)	0.029 (4)	0.043 (5)	-0.014 (4)	-0.012 (4)	0.004 (4)
N1	0.033 (3)	0.022 (3)	0.045 (4)	-0.012 (3)	-0.016 (3)	0.007 (3)
C2	0.061 (6)	0.042 (5)	0.040 (5)	-0.016 (5)	-0.017 (4)	0.012 (4)
C4	0.052 (5)	0.033 (4)	0.060 (6)	-0.018 (4)	-0.021 (4)	-0.006 (4)
C12	0.032 (4)	0.022 (4)	0.040 (4)	-0.011 (3)	-0.010 (3)	0.000 (3)
N2	0.052 (4)	0.018 (3)	0.044 (4)	-0.010 (3)	-0.016 (3)	0.006 (3)
C7	0.036 (4)	0.016 (4)	0.047 (5)	-0.007 (3)	-0.015 (4)	0.006 (3)
C11	0.033 (4)	0.017 (4)	0.051 (5)	-0.014 (3)	-0.014 (3)	0.009 (3)
C9	0.037 (4)	0.030 (4)	0.048 (5)	-0.013 (4)	-0.018 (4)	-0.004 (4)
C10	0.029 (4)	0.016 (4)	0.053 (5)	-0.005 (3)	-0.017 (4)	0.004 (4)
C8	0.038 (4)	0.020 (4)	0.045 (5)	-0.008 (3)	-0.009 (4)	0.002 (3)
C5	0.036 (4)	0.018 (4)	0.049 (5)	-0.002 (3)	-0.014 (4)	-0.004 (3)
C6	0.053 (5)	0.017 (4)	0.057 (5)	-0.013 (4)	-0.016 (4)	0.002 (4)
O1	0.115 (6)	0.030 (3)	0.086 (5)	-0.041 (4)	-0.038 (4)	0.007 (3)
N3	0.069 (5)	0.022 (3)	0.052 (4)	-0.019 (3)	-0.022 (4)	0.003 (3)
O2	0.077 (5)	0.063 (4)	0.064 (4)	-0.005 (4)	-0.020 (4)	0.015 (4)

*Geometric parameters (Å, °)*

Br2—C11	1.907 (7)	N2—C6	1.338 (10)
Br1—C9	1.912 (8)	N2—C7	1.393 (9)
C3—C2	1.384 (12)	N2—H5	0.8600
C3—C4	1.387 (12)	C7—C8	1.388 (11)
C3—H3	0.9300	C11—C10	1.400 (11)
C1—C2	1.355 (11)	C9—C8	1.364 (10)
C1—N1	1.370 (10)	C9—C10	1.394 (11)
C1—H1	0.9300	C10—N3	1.367 (9)
N1—C5	1.364 (9)	C8—H6	0.9300
N1—C12	1.440 (9)	C5—C6	1.471 (11)
C2—H2	0.9300	C6—O1	1.222 (9)
C4—C5	1.373 (11)	N3—H3A	0.8600
C4—H4	0.9300	N3—H3B	0.8600
C12—C7	1.399 (10)	O2—H12	0.8769
C12—C11	1.423 (9)	O2—H11	0.8900
C2—C3—C4	119.0 (8)	C12—C7—N2	120.3 (7)
C2—C3—H3	120.5	C10—C11—C12	121.3 (7)
C4—C3—H3	120.5	C10—C11—Br2	115.3 (5)
C2—C1—N1	122.1 (8)	C12—C11—Br2	123.0 (6)

C2—C1—H1	119.0	C8—C9—C10	124.3 (7)
N1—C1—H1	119.0	C8—C9—Br1	117.2 (6)
C5—N1—C1	118.1 (6)	C10—C9—Br1	118.3 (5)
C5—N1—C12	119.7 (6)	N3—C10—C11	122.5 (7)
C1—N1—C12	121.9 (6)	N3—C10—C9	121.0 (7)
C1—C2—C3	118.9 (8)	C11—C10—C9	116.3 (6)
C1—C2—H2	120.5	C9—C8—C7	118.8 (7)
C3—C2—H2	120.5	C9—C8—H6	120.6
C5—C4—C3	119.9 (8)	C7—C8—H6	120.6
C5—C4—H4	120.1	N1—C5—C4	120.2 (7)
C3—C4—H4	120.1	N1—C5—C6	120.8 (7)
C7—C12—C11	118.4 (7)	C4—C5—C6	118.7 (7)
C7—C12—N1	116.8 (6)	O1—C6—N2	123.8 (8)
C11—C12—N1	124.6 (6)	O1—C6—C5	120.2 (8)
C6—N2—C7	123.7 (6)	N2—C6—C5	115.9 (7)
C6—N2—H5	118.2	C10—N3—H3A	120.0
C7—N2—H5	118.2	C10—N3—H3B	120.0
C8—C7—C12	120.5 (7)	H3A—N3—H3B	120.0
C8—C7—N2	119.1 (7)	H12—O2—H11	123.0
C2—C1—N1—C5	-13.0 (11)	Br2—C11—C10—C9	-172.4 (6)
C2—C1—N1—C12	173.4 (7)	C8—C9—C10—N3	-173.9 (8)
N1—C1—C2—C3	2.1 (12)	Br1—C9—C10—N3	1.0 (10)
C4—C3—C2—C1	7.5 (12)	C8—C9—C10—C11	1.0 (12)
C2—C3—C4—C5	-6.1 (13)	Br1—C9—C10—C11	175.9 (5)
C5—N1—C12—C7	-16.6 (10)	C10—C9—C8—C7	0.9 (12)
C1—N1—C12—C7	156.8 (7)	Br1—C9—C8—C7	-174.0 (6)
C5—N1—C12—C11	157.9 (7)	C12—C7—C8—C9	-4.9 (11)
C1—N1—C12—C11	-28.7 (11)	N2—C7—C8—C9	171.8 (7)
C11—C12—C7—C8	6.7 (11)	C1—N1—C5—C4	14.3 (10)
N1—C12—C7—C8	-178.4 (7)	C12—N1—C5—C4	-172.0 (7)
C11—C12—C7—N2	-169.9 (7)	C1—N1—C5—C6	-159.2 (7)
N1—C12—C7—N2	5.0 (11)	C12—N1—C5—C6	14.4 (10)
C6—N2—C7—C8	-166.8 (7)	C3—C4—C5—N1	-5.0 (12)
C6—N2—C7—C12	9.8 (12)	C3—C4—C5—C6	168.7 (8)
C7—C12—C11—C10	-4.7 (11)	C7—N2—C6—O1	171.7 (9)
N1—C12—C11—C10	-179.2 (7)	C7—N2—C6—C5	-12.2 (11)
C7—C12—C11—Br2	168.1 (6)	N1—C5—C6—O1	176.1 (8)
N1—C12—C11—Br2	-6.4 (11)	C4—C5—C6—O1	2.5 (12)
C12—C11—C10—N3	175.8 (7)	N1—C5—C6—N2	-0.1 (11)
Br2—C11—C10—N3	2.4 (10)	C4—C5—C6—N2	-173.8 (7)
C12—C11—C10—C9	0.9 (11)		

## Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H5...Br3 <sup>i</sup>	0.86	2.49	3.332 (6)	166
N3—H3B...Br1	0.86	2.60	3.048 (7)	113



---

N3—H3B···Br3 <sup>ii</sup>	0.86	2.84	3.581 (7)	145
N3—H3A···O1 <sup>iii</sup>	0.86	2.17	2.977 (9)	155
N3—H3A···Br2	0.86	2.56	3.006 (7)	113
O2—H11···Br3 <sup>iv</sup>	0.89	2.50	3.383 (6)	180
O2—H12···Br1 <sup>v</sup>	0.88	2.61	3.309 (7)	137
O2—H12···Br3	0.88	2.83	3.393 (6)	123

---

Symmetry codes: (i)  $x, y+1, z-1$ ; (ii)  $x, y, z-1$ ; (iii)  $x, y-1, z$ ; (iv)  $-x+1, -y, -z+1$ ; (v)  $x, y, z+1$ .