














ORIGINAL ARTICLE

Fatty acid and amino acid profiles of cheese, butter, and ghee made from buffalo milk

Abu Hena Md. Asif^{1†} , Md. Abid Hasan Sarker^{1†} , Gautam Kumar Deb² , Md. Rezwatul Habib¹ , Sumaiya Arefin³ , Md. Sadakatul Bari¹ , Md. Zakirul Islam¹ , Md. Harun-ur- Rashid¹ , Mohammad Shohel Rana Siddiki¹ , Umma Fatema Shahjadee⁴ , Sharmin Akter Lisa⁴ , Salma Ahmed⁴ , Mohammad Ashiqul Islam¹ 

¹Department of Dairy Science, Bangladesh Agricultural University, Mymensingh, Bangladesh

²Animal Biotechnology Division, Bangladesh Livestock Research Institute, Savar, Bangladesh

³Department of Livestock Services, Ministry of Fisheries and Livestock, Dhaka, Bangladesh

⁴Institute of Food Science Technology IFST, Bangladesh Council of Scientific and Industrial Research BCSIR, Dhaka, Bangladesh

[†]These authors contributed equally to this work.

ABSTRACT

Objective: The objective was to assess the chemical composition, cholesterol, fatty acid (FAs), and amino acid (AAs) profiles of buffalo cheese, butter, and ghee.

Materials and Methods: Buffalo milk (raw) was collected from the Bangladesh Agricultural University (BAU) Dairy Farm, BAU, Mymensingh-2202, Bangladesh. Cheese, butter, and ghee were prepared at the Dairy Chemistry and Technology Laboratory, Department of Dairy Science, BAU, Mymensingh, Bangladesh, and subjected to subsequent analyses. The gross nutritional composition and AAs profile of milk were analyzed prior to the manufacture of cheese, butter, and ghee. The gross nutritional composition of milk and dairy products was analyzed by applying an automated milk analyzer and the Association of Agricultural Chemists techniques, respectively. The cholesterol, FAs, and AAs contents of cheese, butter, and ghee were determined by the Bangladesh Council for Scientific and Industrial Research, Dhaka, Bangladesh. Furthermore, atherogenic and thrombogenic indices were also calculated using reference equations.

Results: The results indicated that the buffalo milk is a good source of first-rate nutrients (dry matter: 16.50%, fat: 7.50%, protein: 3.75%). Findings indicated that the butter was significantly rich with ($p < 0.05$) total solids and fat where higher ($p > 0.05$) protein, carbohydrate, and minerals were found in cheese. The saponification, Reichert-Meissl, Polenski, and Kirschner values of buffalo ghee were found to be 225, 30, 1.2, and 25, respectively. A significant ($p < 0.05$) variation was found in the cholesterol content of buffalo cheese, butter, and ghee. Butter and ghee had 40.14 and 39.57 mg more cholesterol, respectively, than cheese. The results revealed identical FA profiles except for C24:0 among the three dairy products where the major FA compositions were C4:0, C14:0, C16:0, and C18:0 and C18:1 cis-9. The atherogenicity index and thrombogenicity index of cheese, butter, and ghee were statistically similar ($p > 0.05$). Butter was found with the most conducive anti-atherogenic and anti-thrombogenic characteristics due to lower saturated and higher polyunsaturated FAs. However, all the AAs concentrations were statistically higher ($p < 0.05$) in cheese than in butter and ghee.

Conclusion: To conclude, buffalo cheese is superior to butter and ghee as regards nutrient density, but consumers can choose other foods based on their choice.

ARTICLE HISTORY

Received August 13, 2021

Revised December 12, 2021

Accepted December 28, 2021

Published March 13, 2022

KEYWORDS

Buffalo milk; fatty acids; amino acids; cholesterol; atherogenic index; thrombogenic index



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by/4.0>)

Introduction

Milk and dairy products are nutrient-dense foods that contribute significantly to human consumption [1] and

provide energy and substances crucial for the development of humans, particularly infants. Buffalo milk is considered a superior raw material for western and traditional dairy

Correspondence Mohammad Ashiqul Islam ✉ m.a.islam@bau.edu.bd 📧 Department of Dairy Science, Bangladesh Agricultural University, Mymensingh, Bangladesh.

How to cite: Asif AHM, Sarker MAH, Deb GK, Habib MR, Arefin S, Bari MS, et al. Fatty acid and amino acid profiles of cheese, butter, and ghee made from buffalo milk. *J Adv Vet Anim Res* 2022; 9(1):144–154.

products. Dairy products produced with buffalo milk are excellent in the body and firmer in texture owing to the higher content of total solids (TS), protein, and fat. Buffalo milk contains a high concentration of linolenic and conjugated linolenic acids (CLA) [2], as well as fat, lactose, caseins, AAs, and calcium [3]. These nutrients offer excellent opportunities to process and develop different dairy products from buffalo milk [2]. Importantly, buffalo milk and milk products help to prevent metabolic syndrome, chronic diseases, type-2 diabetes, hypertension, and cardiovascular disease due to higher amounts of fatty acids (FAs) and amino acids (AAs) [4].

Cheese is a good source of nutrition and a delicious milk product prepared from the coagulation of casein (with entrapped fat and water) by proteolytic enzyme action. It is expedient, versatile, and offers various flavors, textures, and forms. Cheese can be classified into many groups based on appearance, preparation, ripening, and chemical composition [5]. Cheese is manufactured from different types of milk, and its quality relies on milk composition, microbial quality, preparation, and consistency [6]. Some FAs are found in ruminant cheeses like vaccenic (C18:1 ν 11) and conjugated linoleic acids (CLA; C18:2 c 9, t 11) that are beneficial to human health [7].

Buffalo milk is rich in fat (an average of 8.3%), implying the feasibility of using buffalo milk for manufacturing fat-based dairy products. Butter is a fat-rich dairy product where the liquid phase is dissolved in the oily phase, forming a water-in-oil emulsion. It signifies the concentration of fat as the only one of the three main constituents, like protein, fat, and carbohydrates, of milk. Protein and carbohydrates are present in minor proportions, brought into butter by buttermilk residues after churning [8].

Ghee is an anhydrous dairy product containing no less than 99.5% milk-fat and is treated as the most popular fat-rich dairy product in the Indian subcontinent. Ghee is prepared from cream or churned butter from fresh or ripened cream by heating, where ripening is carried through the fermentation of milk with bacteria inherent to milk or preferred starter cultures. Ghee is composed of mixed glycerides, FAs, phospholipids, sterols, fat-soluble vitamins, hydrocarbons, small amounts of charred casein, and trace amounts of calcium, phosphorus, iron, etc. [8]. According to Ullah et al. [9], 62% of the monounsaturated fats in ghee are rich in CLA. However, buffalo milk fat is proportionally rich in rumenic acid (the principal CLA). The CLA isomers of buffalo milk are considered anti-carcinogenic, anti-atherogenic, anti-obesity, and anti-diabetic components [2].

The buffalo is vastly important in Southeast Asia, and consumers are gladly consuming buffalo milk products all over the Indian sub-continent. The type of milk is majorly responsible for the sound characteristics of cheese, butter,

and ghee with proximate quality, FAs, and AAs. Nutritional composition focuses on consumer preference, and country-specific data on the nutritional composition of milk or products need to be set out [10].

Several articles have been reported on cheese based on biochemical, microbial, and sensory evaluations prepared from the milk of different species [11,12], emphasizing nutritional sources through feeding [13], different types of coagulant [14,15], and types of culture [16]. But no work has been located with detailed FA and AA profiles. Although there is a wealth of research on the physical and processing properties, texture, nutritional value, and FAs composition of butter from cow milk [17,18], the domain is deficient in the case of butter from buffalo milk, with a focus on FAs and AAs profiles. Most of the research work has been focused on the physicochemical and nutritional aspects of ghee from different species [19], the development of the process of arjuna herbal ghee [20], and the effect of cholesterol removal on the composition of cow ghee [21]. However, not much information is available on the detailed composition of ghee with FAs and AAs profiles, whereas, consumers are becoming more apprehensive about food compositions emphasizing FAs and AAs, which may greatly affect the maintenance of human health.

Most importantly, there is insufficient research data on the chemical characteristics highlighting FAs and AAs of cheese, butter, and ghee from indigenous buffalo in the Indian subcontinent. Therefore, the objective of this study was to assess the chemical composition, cholesterol, FAs, and AAs profiles of buffalo cheese, butter, and ghee prepared from buffalo milk. This study will help people decide what to eat based on these products' composition and nutritional differences and how they are made.

Materials and Methods

Ethical approval

Animal management procedures, methods of milking, and collection of milk samples were approved by the Animal Welfare and Experimentation Ethics Committee, Bangladesh Agricultural University (BAU), Mymensingh-2202, Bangladesh [AWEEC/BAU/2020 (21)].

Collection and analysis of raw milk

Morning buffalo milk ($n = 3$, non-descriptive) was collected and pooled separately from the BAU Dairy Farm (24°43'46.5"N, 90°25'22.8"E), Mymensingh-2202, Bangladesh. After that, 5 l of milk was taken from mixed samples and sent to the laboratory for product manufacturing and following analyses. Buffaloes were fed individually on German grass (*Echinochloa crus-galli*) in a cut and carry system supplemented with a concentrated mixture [22].

Buffaloes had free access to clean, fresh drinking water for 24 h. Gross milk compositions were analyzed through an automated milk analyzer (Lactoscan, SLP, MILKOTONIC Ltd., Nova Zagora-8900, Bulgaria) at the Dairy Chemistry and Technology Laboratory, Department of Dairy Science, BAU, Mymensingh-2202. The cholesterol content, FAs, and AAs were assayed at the Bangladesh Council for Scientific and Industrial Research, Dhaka, Bangladesh. Table 1 shows raw buffalo milk's chemical composition, cholesterol level, and AA composition. The FA composition of buffalo milk was previously defined by Islam et al. [4].

Manufacturing of cheese, butter, and ghee

The products were manufactured at the Dairy Chemistry and Technology Laboratory, Department of Dairy Science, BAU, Mymensingh-2202. The brief details are presented below.

Cheese

The cheese-making procedure is illustrated in Figure 1. Fresh buffalo milk was heated to 80°C for 30 min before

Table 1. Proximate composition, cholesterol content, and AA profile (mean ± standard deviation) of buffalo milk.

Parameters	Buffalo milk (n = 3)
pH	6.75 ± 0.15
Acidity (%)	0.13 ± 0.01
TS (%)	16.50 ± 0.30
Fat (%)	7.50 ± 0.26
Carbohydrates (%)	4.70 ± 0.12
Protein (%)	3.57 ± 0.11
Ash (%)	0.72 ± 0.01
Cholesterol (mg/100 g)	21.93 ± 9.36
AAs (gm 100 gm ⁻¹)	
Aspartic acid	0.31 ± 0.01
Threonine	0.18 ± 0.00
Serine	0.22 ± 0.01
Glutamic acid	0.78 ± 0.02
Glycine	0.14 ± 0.00
Alanine	0.19 ± 0.01
Valine	0.37 ± 0.01
Methionine	0.17 ± 0.01
Isoleucine	0.25 ± 0.01
Leucine	0.29 ± 0.01
Tyrosine	0.21 ± 0.00
Histidine	0.12 ± 0.00
Lysine	0.32 ± 0.01
Arginine	0.110 ± 0.005

being slowly cooled to 40°C. For acidification, lactic starter culture (10 gm/l) was added by direct inoculation to milk for desirable acidity. After that, 0.07 gm/l of commercial rennet (CHY-max®, Denmark) was added to make the milk curd. Stirring continued for 1 min to ensure uniform incorporation of the added enzyme into the milk, followed by discontinuation to allow the curd formation. When the curd formed a clean break, it was cut into cubes and salted in brine (10 gm of NaCl/kg). The curds were placed in a perforated container and stored at 4°C for 24 h.

Butter

The manufacturing process of butter from buffalo milk is depicted in Figure 2. First, the cream was separated from raw buffalo milk by applying a cream separator, and its fat content was standardized to 35% (w/w). The cream was kept overnight at 4°C and churned using a 1-l capacity laboratory churn. The butter kernels were washed with cold water (12°C) in the churn. After draining butter whey, the working of butter was done to a continuous fat phase with a finely dispersed water phase to attain a homogenous mixture of butter granules, water, and salt. Approximately, 100 gm of salt (w/w) was added during the work to improve the shelf-life and flavor. The finished butter was then wrapped in cube-sized plastic and stored at 4°C for further analysis.

Ghee

Manufacturing protocols for ghee from buffalo milk are shown in Figure 3. Ghee was made using the direct cream method, in which 35% (w/w) cream was melted in a metal pan and clarified at 115°C–116°C with continuous stirring until almost all of the moisture was removed. The heating was discontinued when the curd particles attained the desired golden yellow or brown color. After heating, the contents were sedimented, decanted, filtered, and packed for analysis.

Proximate component analysis

The acidity of the cheese, butter, and ghee was measured using the titration method against 0.1 M NaOH solution and phenolphthalein indicator, and pH was measured by a pH meter-215 (Corning Diagnostic Ltd., Sudbury, Suffolk, England). The TS content was determined by evaporating at 105°C for 24 h in an oven [J.P. Selecta; S.A. ctra. Nil km: 585.1, Abrera (Barcelona) Spain]. After drying, the samples were ignited at 600°C for 6 h in a muffle furnace (Vulcan A-550, Ney®, USA) to estimate ash. Protein was estimated by the Kjeldahl method [23], and fat content was determined by the Babcock fat test method described by Aggarwala and Sharma [24].




Figure 1. Manufacturing process of fresh cheese from buffalo milk.




Figure 2. Manufacturing process of butter from buffalo milk.

Determination of milk fat constants

Saponification value

The saponification value was determined using the method Cd 3-25 [23]. First, 5 gm of filtered ghee was mixed with 50 ml of 4% alcoholic KOH, and the mixture was mildly boiled until complete saponification. After cooling, the sample was titrated against 0.5 M HCl (43.01 ml/l) in the presence of a phenolphthalein indicator (1 gm/100 ml) to neutralize excess KOH.

Reichert-Meissl, Polenske, and Kirschner value

Reichert-Meissl (chiefly butyric and caproic), Polenske (mostly caprylic, capric, and lauric), and Kirschner (butyric

acid) values were determined according to the method of Cd 5-40 [25]. With 5 gm of ghee sample, 20 ml of glycerol soda solution (20 ml of 500 gm/l NaOH added with 180 ml of glycerol) was added in a round-bottom flask and subsequently heated until complete saponification.

After cooling, the prepared soap solution was diluted with distilled water (135 ml) and acidified with 6 ml of sulphuric acid (H₂SO₄). The flask was connected to the condenser and heated to distillate volatile FAs. The collected volatile FAs were filtrated to separate soluble FAs from insoluble ones. The aqueous solution of the soluble volatile FAs and the ethanolic solution of the insoluble volatile FAs were titrated against 4 gm/l of NaOH solution in the presence of a phenolphthalein indicator (1 gm/100




Figure 3. Manufacturing process of ghee from buffalo milk cream.

ml). Then 0.5 gm of silver sulfate powder was added to the solution obtained from the RM procedure to determine the Kirschner value.

Cholesterol analysis

Extraction of lipid

In the case of milk samples, 20 ml of H_2SO_4 was added to 10 ml of milk and the mixture was kept in a water bath at 100°C for 30 min. To separate the fat layer, after adding 70 ml of diethyl ether, the content was shaken 4–5 times. The content was kept in a water bath at 100°C for 10 min, followed by oven drying at 102°C for 4–5 h to dry the fat layer. To keep them moisture-free, cheese, butter, and ghee samples were dried in an oven at 102°C for 4–5 h. Then further lipid extraction was done from cheese, butter, and ghee as described by the Association of Agricultural Chemists [23].

Analysis of cholesterol

Cholesterol analysis from extracted lipid samples of milk, cheese, butter, and ghee was done using the method Huang et al. [26] with a few modifications. A standard curve of pure cholesterol and glacial acetic acid was made (0.05 gm of pure cholesterol was dissolved into 100 ml of glacial acetic acid), and the sample measurement was done at 625 nm on a Ultraviolet-Spectrophotometer (SPECURD®/250 Plus, Analytikjena Co., Germany).

Fatty acid analysis

Preparation of fatty acid methyl esters (FAMES)

The Rose-Gottlieb method was applied to extract lipids from cheese, butter, and ghee samples [23]. The FA

compositions of cheese, butter, and ghee were determined from their corresponding FAMES. Ten milliliters of each extracted lipid sample were kept in a 15 ml test tube. Then, 3 ml of 0.5 M CH_3ONa solution was added to the tube and digested with continuous stirring in a heating water bath for about 15 min. Subsequently, the test tube was cooled at room temperature, and 1 ml of petroleum ether with a boiling point of 40°C–60°C was added, followed by 10 ml of de-ionized. Following gentle mixing, the tubes were allowed to settle (5–6 min) the contents until the top layer of methyl ester became clear. Then, the top layer of the FAMES was separated with care into the capped Gas chromatography (GC) vial for analysis. Two hundred milligrams of different FAs standard with corresponding methyl esters were liquefied distinctly in 10 ml petroleum ether (boiling point 40°C–60°C) in a series of screw-capped test tubes. Aliquots of 1.0 μ l FAMES were injected, and the peaks of FAs were recorded against their respective column retention times and areas with the help of the data processor unit of GC.

Chromatography

FAMES of cheese, butter, and ghee were quantified by a GC (GC; Shimadzu GC-14B, Japan) with flame ionization detector with a fused silica capillary column (FAMEWAX, Crossbond® polyethylene glycol, 15 m \times 0.25 mm \times 0.25 μ m film thickness, Restek; PA). The temperature program was kept as: the temperature at initial was kept for 5 min at 150°C after injection then programmed to increase at 190°C (at 8°C/min), kept for 10 min. Then the temperature was raised to 200°C (at 2°C/min) and held for 10 min. Both the injection and detector temperatures were 250°C, and part of a 1.0 μ l volume of the sample was injected. The

splitless injection technique was used with nitrogen as a carrier gas at a constant flow rate (20 ml/min). FAMES were identified by applying a standard FAME mix (Food Industry 37 FAME mix, 35077 Restek Co, Bellefonte, PA), and areas of the peak were calculated in the chromatogram and then expressed as percentages (gm/100 g of total FA) by the automated GC software (Class GC-10, version-2.00). The atherogenicity index (AI) and thrombogenicity index (TI) was calculated following the formula recommended by Ulbricht and Southgate [27]:

$$AI = [(12:0 + 4 (14:0) + 16:0) / [(n6 + n3) PUFA + 18:1 + \Sigma MUFA]]$$

$$TI = (14:0 + 16:0 + 18:0) / [(0.5 \times 18:1) + 0.5 (\Sigma MUFA) + 0.5 (n6 PUFA) + 3 (n3 PUFA) + (n3 PUFA/n6 PUFA)].$$

Amino acids analysis

Fat from milk, cheese, butter, and ghee was extracted using n-hexane (Merck, Darmstadt, Germany). Fat-free samples were then dried in an oven at 50°C for 3–5 h. For the estimation of AAs (aspartic acid, threonine, serine, glutamic acid, glycine, alanine, valine, methionine, isoleucine, leucine, tyrosine, histidine, lysine, and arginine), a dried sample of 0.5 gm was taken as described in the Shimadzu manual [28]. This AA analyzer [high-performance liquid chromatography (HPLC)] used a column (Amino-Na) to separate a highly acidic cation exchange resin. The AA adducts have been efficiently separated by using a binary gradient elution technique and detected by using a post-column derivatization detection method and a fluorescence detector at high sensitivity and pressure (88,267.1 mmHg). The HPLC system was composed of several components such as a controller (SCL-10A), liquid chromatography (LC-10AD), pump, auto-injector (SIL-10AD), column oven (CTO-10 AD), column (Amino-Na), and a fluorescence detector. Mobile A contained 0.20 M sodium citrate (adjusted pH 3.2 with perchloric acid) and 7.0% ethanol, and mobile phase B contained 0.60 M sodium citrate and 0.20 M boric acid (adjusted to pH 10.0 with 4 M sodium hydroxide). Washing solution C was composed of 0.20 M sodium hydroxide. In the fluorescence method for AAs, o-phthalaldehyde reagent, RB (o-phthalaldehyde, ethanol, polyoxyethylene lauryl ether (Brij-35), and N-acetyl-L-cysteine) were used to react with primary amines of AAs, peptides, and proteins to enable detection and quantitation. The flow rate of both RA and RB was 20–21 ml/min. To prepare the samples for AAs analysis, 0.5 gm of the oven-dried fat-free sample was crushed by the mortar pestle and mixed with 50 ml of HCl (6 M) to make a fine paste. After that, the fine paste was moved into a 250 ml round-bottom flask via filtration, and the flask containing the filtrate was placed on the heating mantle at 110°C for 22 h for hydrolysis. Thereafter, HCl was eliminated from the filtrate by adding 3–4 times distilled water and evaporating it at 90°C–95°C

in a laboratory water bath. The obtained solution was filtered in a 25 ml volumetric flask using a Whatman filter paper of 41 and columned with a stock solution of 0.1 M HCl. Furthermore, that stock solution was filtered using a 0.45 µm syringe filter (Shimadzu, Japan). Then, stock and internal standard (IS) solutions (Mixed standard, Sigma-Aldrich, St. Louis, MO) were run through the AA analyzer, which displayed the standard curve for the standard solution and the sample. Finally, by comparing both the areas of two curves against stock and sample, AA was calculated in gm/100-gm as follows:

$$AA = (IS \text{ concentration} \times \text{Sample peak area}) \div IS \text{ peak area.}$$

Statistical analysis

The independent sample *t*-test was done to determine the statistical differences in gross nutritional quality between the buffalo cheese and butter. The one-way analysis of variance was carried out to determine any significant differences among buffalo cheese, butter, and ghee in cholesterol, FAs, and AA profile. Minitab was used as statistical software (Minitab-18, Minitab Inc.®, State College, PA).

Results and Discussion

Gross nutritional quality of cheese and butter

In Table 2, the gross nutritional substances of cheese and butter are presented. The findings indicated that all the chemical constituents of cheese and butter were

Table 2. Gross nutritional quality of buffalo milk cheese and butter (mean ± standard deviation).

Parameters	Buffalo milk		p-value
	Cheese	Butter	
pH	5.90 ± 0.09	6.90 ± 0.04	<0.05
Acidity (%)	0.15 ± 0.04	0.12 ± 0.05	>0.05
TS (%)	59.28 ± 5.1	88.86 ± 2.8	<0.05
Fat (%)	29.75 ± 5.88	86.00 ± 2.2	<0.05
Carbohydrates (%)	2.60 ± 0.68	0.62 ± 0.42	>0.05
Protein (%)	22.43 ± 1.6	1.46 ± 0.17	<0.05
Minerals (%)	4.50 ± 0.2	0.78 ± 0.04	<0.05

Table 3. Milk fat constants of buffalo milk ghee (mean ± standard deviation).

Fat constants	Buffalo milk ghee
Saponification value	225 ± 5.66
RM value	30.00 ± 4.24
Polensky value	1.20 ± 0.14
Kirschner value	25.00 ± 2.83

significantly ($p < 0.05$) different, except for acidity and carbohydrates. Butter was found rich in TS and fat content at 29.5% and 56%, respectively. Higher TS and fat content in butter may result from variations in manufacturing technology and the percentage of chemical compositions of milk. Butter is a fat-rich milk product where fat is the major component, forming a water and oil emulsion with little carbohydrate, protein, and minerals. Due to this compositional variation, the fat content of butter is considerably higher than that of cheese. The TS and fat content of butter discovered are consistent with the findings of Hurtaud et al. [29]. However, being a protein-based product, cheese also has a greater portion of milk fat entrapped, and the fat content of buffalo cheese found in this study agrees with another study [30]. In terms of protein and mineral content, there were 21% and 3.7% higher in cheese than in butter. The higher protein and minerals in cheese are imparted due to the cheese-making process where the rennet enzymes precipitate casein. A coagulum contains casein, whey proteins, lactose, fat, and minerals. This finding agrees with Rana et al. [25], who reported 21.87% protein and 3.19% minerals in buffalo cheese. However, both cheese and butter composition conforms to the Bangladesh Standard and Testing Institute [31].

Milk fat constants of ghee

In Table 3, milk fat constants signify FAs present in different kinds of fats. These are also used to detect fat adulteration qualitatively and, in some cases, quantitatively. In this study, the saponification value was 225 in ghee, implying the molecular weight of FAs. The average saponification value varies from 225.00 to 230.00 [32]. The remarkable measure of short-chain FAs implies the remarkable measure of butyric (C4) and caproic (C6) acid. The typical RM value of ruminant milk fat generally ranges from 17–35. Here, we found RM value 30 in buffalo ghee. The result is consistent with Gandhi et al. [33], who reported a range of 31.46 to 34.98 RM value in buffalo ghee. Polenske value, which typically ranges from 1.2 to 2.4 [34], denotes milk fat's volatile water-insoluble FAs. This study found a Polenske value in buffalo ghee of 1.2, which is in the reported range. In another study, Bhatia et al. [35] found a Polenske value of 1.77 in cow ghee, suggesting more volatile water-insoluble FAs in cow milk. In this study, the Kirschner value was 25, which is also in the normal range.

Cholesterol and FAs composition of cheese, butter, and ghee

Table 4 summarizes the cholesterol content and FAs of buffalo cheese, butter, and ghee. A significant ($p < 0.05$) variation was found in cholesterol content among cheese, butter, and ghee, where butter appeared with the highest value (45.38 mg), followed by ghee (44.81 mg) as fat-rich

milk products. On the other hand, cheese had significantly lower cholesterol (5.24 mg) due to manufacturing technology variations. Higher fat content, buffalo milk contains significantly lower amounts of cholesterol than cow milk [36]. Buffalo milk is more popular among health-conscious people due to its lower cholesterol content. The sum of saturated FAs was found to be statistically similar ($p >$), butter (60%), and ghee (68%), as presented in Table 4. This higher percentage of saturated FAs in all three products might be attributed to the higher saturated FAs content in raw buffalo milk [37]. Amid all the saturated FAs, only C24:0 showed significant ($p < 0.05$) variation among the products. The major saturated FAs of cheese, butter, and ghee were found to be C4:0, C14:0, C16:0, and C18:0. The FAs C4:0 (5.28%), C12:0 (2.51%), C14:0 (12.66%), and C16:0 (38.18%) were found to be high in cheese and C18:0 (13.73%) in butter. Here, it was also notable that the lower content of C16:0 had been compensated for by increased C18:1, *cis*-9, C18:2, *cis*-9, and 12 and C18:3, *cis*-6, 9, and 12 in buffalo butter. It is assumed that the types of FAs, *viz.*, saturated, mono- and polyunsaturated, branched, and conjugated, are thought to be either positive or negative for the consumer's health [38]. The sum of unsaturated FAs was statistically ($p > 0.05$) alike in buffalo cheese, butter, and ghee, as presented in Table 4. Oleic acid (C18:1 *cis*-9) content was found to be the second-highest in cheese (19.65%), butter (31.36%), and ghee (24.83%). This FA is very effective against cardiovascular disease and atherosclerosis [39]. In a study, buffalo milk was reported to have a higher C18:1 *cis*-9 [4]. Polyunsaturated FAs such as C18:2, *cis*-9, 12, and C18:3, *cis*-6, 9, 12, were also exponentially higher in buffalo butter. Generally, the CLA percentage of milk products largely depends on the composition of raw milk. In a study, Islam et al. [4] reported C18:2 (0.90%) and C18:3 (0.05%) among all FAs in buffalo milk. Conjugated linoleic acid isomers are thought to be anti-carcinogenic, anti-atherogenic, and anti-obesogenic [2]. The AI describes the ratio between atherogenic and anti-atherogenic FAs. Again, the TI considers the major saturated FAs as pro-thrombogenic while the unsaturated ones are considered anti-thrombogenic FAs. The differences among the atherogenic and thrombogenic indices of cheese, butter, and ghee from buffalo milk were statistically non-significant ($p > 0.05$), suggesting a positive health effect on human health. However, the AI values of cheese, butter, and ghee were 3.72, 1.65, and 2.41, respectively, where butter was recorded with the lower value. Here, atherogenic saturated FAs 12:0, 14:0, and 16:0 were found to be lower in buffalo butter with increased C18:1, *cis*-9. In this study, the AI of butter and ghee was lower than the findings of Bobe et al. [40], who reported that the high AI value in dairy products is 2.71. It is suggested that milk products with minimum AI and TI values are less likely to be negative for

Table 4. Cholesterol content (mg 100 gm⁻¹) and FA profile (% of FAMES) of buffalo milk cheese, butter and ghee (mean ± standard deviation).

Parameters	Buffalo milk			p-value
	Cheese	Butter	Ghee	
Cholesterol	5.24 ^c ± 0.29	45.38 ^a ± 0.25	44.81 ^{ab} ± 9.70	<0.05
ε Saturated FAs	73.18 ± 10.37	59.90 ± 3.38	67.40 ± 3.62	>0.05
ε Unsaturated FAs	25.91 ± 8.57	38.01 ± 2.34	31.12 ± 3.06	>0.05
Saturated FAs				
Butyric acid (C4:0)	5.28 ± 1.19	3.62 ± 4.01	3.38 ± 3.28	>0.05
Valeric acid (C5:0)	0.55 ± 0.15	0.17 ± 0.04	1.13 ± 0.82	>0.05
Caproic acid (C6:0)	0.30 ± 0.15	0.19 ± 0.03	0.64 ± 0.29	>0.05
Caprylic acid (C8:0)	0.89 ± 0.17	0.53 ± 0.08	1.36 ± 0.43	>0.05
Capric acid (C10:0)	0.11	nd	0.09	-
Lauric acid (C12:0)	2.51 ± 0.84	0.90 ± 0.09	1.95 ± 0.45	>0.05
Tridecanoic acid (C13:0)	0.57 ± 0.37	0.07	0.10 ± 0.02	>0.05
Myristic acid (C14:0)	12.66 ± 2.73	8.17 ± 2.24	10.43 ± 1.17	>0.05
Pentadecanoic acid (C15:0)	1.69 ± 0.08	1.57 ± 0.14	1.41 ± 0.02	>0.05
Palmitic acid (C16:0)	38.18 ± 7.97	29.01 ± 0.20	34.42 ± 0.95	>0.05
Margaric acid (C17:0)	0.75 ± 0.06	1.29 ± 0.47	0.74 ± 0.14	>0.05
Stearic acid (C18:0)	9.59 ± 1.34	13.73 ± 2.45	11.35 ± 1.52	>0.05
Arachidic acid (C20:0)	0.10	0.30 ± 0.10	0.19 ± 0.08	>0.05
Heneicosylic acid (C21:0)	Nd	0.08	0.09	-
Behenic acid (C22:0)	nd	0.14	0.06 ± 0.07	>0.05
Lignoceric acid (C24:0)	nd	0.13 ± 0.02	0.06 ± 0.01	<0.05
Unsaturated FAs				
Tridecenoic acid (C13:1)	0.47 ± 0.14	0.26 ± 0.07	0.22 ± 0.02	>0.05
Myristoleic acid (C14:1, <i>cis</i> -9)	1.77 ± 0.39	2.05 ± 0.74	1.73 ± 0.14	>0.05
Pentadecanoic acid (C15:1, <i>cis</i> -10)	0.41 ± 0.13	0.34 ± 0.03	0.29 ± 0.03	>0.05
Palmitoleic acid (C16:1, <i>cis</i> -9)	2.25 ± 0.98	1.72 ± 0.92	2.58 ± 0.26	>0.05
Heptadecatrienoic acid (C17:1, <i>cis</i> -10)	0.29 ± 0.06	0.18 ± 0.15	0.27 ± 0.07	>0.05
Oleic acid (C18:1, <i>cis</i> -9)	19.65 ± 7.28	31.36 ± 2.77	24.83 ± 2.51	>0.05
Linoleic acid (C18:2, <i>cis</i> -9,12)	0.75 ± 1.00	1.48 ± 0.18	0.86 ± 0.40	>0.05
Linolenic acid (C18:3, <i>cis</i> -6,9,12)	0.21 ± 0.26	0.43 ± 0.19	0.22 ± 0.11	>0.05
Eicosenoic acid (C20:1, <i>cis</i> -11)	0.11	0.19 ± 0.08	0.12 ± 0.08	>0.05
ε Others	1.01 ± 1.32	2.19 ± 0.35	1.53 ± 0.42	>0.05
AI	3.72 ± 0.65	1.65 ± 0.02	2.41 ± 0.1	>0.05
TI	4.71 ± 0.98	2.53 ± 0.15	3.32 ± 0.21	>0.05

FAMES, fatty acid methyl esters; nd, not detected.

^{abc} Mean values in a row with uncommon superscript letters differ significantly.

humans [41]. To reduce the risk of cardiovascular disease, the American Heart Association and American College of Cardiology recommended that saturated fat intake be limited to 5% to 6% of total daily calorie intake [42]. The scientific basis for reducing dietary saturated FAs has been focused on the well-established effects of rising low-density lipoprotein cholesterol, along with a decrease in

non-low-density lipoprotein cholesterol, in contributing to atherosclerosis.

AA composition of cheese, butter, and ghee

The quantification of AAs has a significant role in assessing the nutritional quality of foods. The AA contents of cheese, butter, and ghee are given in Table 5. Buffalo

Table 5. AAs profile (gm 100 gm⁻¹) of cheese, butter and ghee prepared from buffalo (mean ± standard deviation)

AAs	Buffalo milk			p-value
	Cheese	Butter	Ghee	
Aspartic	1.53 ^a ± 0.03	0.09 ^b ± 0.01	0.05 ^{bc} ± 0.001	<0.05
Threonine	0.82 ^a ± 0.03	0.05 ^b ± 0.005	0.02 ^{bc} ± 0.001	<0.05
Serine	0.88 ^a ± 0.06	0.05 ^b ± 0.005	0.03 ^{bc} ± 0.001	<0.05
Glutamic	3.28 ^a ± 0.13	0.15 ^b ± 0.01	0.08 ^{bc} ± 0.004	<0.05
Glycin	0.67 ^a ± 0.03	0.04 ^b ± 0.003	0.02 ^{bc} ± 0.001	<0.05
Alanine	0.73 ^a ± 0.04	0.04 ^b ± 0.003	0.02 ^{bc} ± 0.001	<0.05
Valine	1.25 ^a ± 0.04	0.08 ^b ± 0.01	0.04 ^{bc} ± 0.001	<0.05
Methionine	0.63 ^a ± 0.06	0.03 ^b ± 0.002	0.02 ^{bc} ± 0.002	<0.05
Isoleucine	0.84 ^a ± 0.03	0.05 ^b ± 0.004	0.03 ^{bc} ± 0.001	<0.05
Leucine	0.93 ^a ± 0.03	0.07 ^b ± 0.004	0.06 ^{bc} ± 0.002	<0.05
Tyrosine	0.86 ^a ± 0.03	0.05 ^b ± 0.005	0.03 ^{bc} ± 0.001	<0.05
Histidine	0.59 ^a ± 0.03	0.04 ^b ± 0.003	0.02 ^{bc} ± 0.002	<0.05
Lysine	1.08 ^a ± 0.05	0.12 ^b ± 0.006	0.07 ^{bc} ± 0.005	<0.05
Arginine	0.60 ^a ± 0.06	0.07 ^b ± 0.01	0.03 ^{bc} ± 0.002	<0.05

cheese provides a reliable source of AAs, and it is significantly different ($p < 0.05$) compared with butter and ghee. This higher concentration of AAs in cheese might be due to the higher protein content in cheese (22.43%) [43]. In particular, buffalo cheese is rich in essential AAs such as valine (1.25%), lysine (1.08%), and leucine (0.93%). This large amount of branched AAs confers an enormous health benefit. Moreover, the primary non-essential AAs were glutamic (3.28%), aspartic (1.53%), and serine (0.88%). On the other hand, butter and ghee are fat-rich products and contain lower amounts of AAs. Compared to ghee, butter is composed of a significantly higher ($p < 0.05$) content of AAs. It is important that glutamic acid was noticeably higher among the total AAs pool in cheese (3.28 gm), butter (0.15 gm), and ghee (0.08 gm). Glutamic acid has numerous taste perceptions, intermediate metabolism, and neurotransmission functions. Glutamic acid is considered to be a specific forerunner of arginine and proline, along with, more importantly, γ -aminobutyric acid (g amino butyric acid) and glutathione, which are anti-hypertension [44] and anti-diabetic [45].

Conclusion

The manufacturing process and milk types are primarily responsible for the good qualities of cheese, butter, and ghee in their nutritional profiles. The nutritional profile focuses on consumer preference, which is why buffalo milk products' detailed compositional properties need to be set out. The chemical composition results showed that the butter was rich in proximate composition, except for

protein. They found milk-fat constants were within the normal range, and these fat constants would be extremely useful in detecting adulteration of milk fat. The results also revealed that butter represents an essential source of FAs, mainly unsaturated FAs, while cheese was rich in AAs, both essential and non-essential. According to the analysis, the nutritional values of milk products differed significantly. The least atherogenicity and thrombogenicity indices were found in butter rather than cheese and ghee. Even though cheese is a protein-based milk product and butter and ghee are fat-based milk products, consumers would benefit from choosing the appropriate one based on nutritional characteristics. These findings may perhaps assist dairy farmers, milk processors, and product manufacturers with better clarification regarding the types of dairy products.

List of Abbreviations

L, Litre; H₂SO₄, Sulphuric acid; μ l, Microlitler; FAMES, Fatty acid methyl esters; GC, Gas chromatography; HPLC, High-performance liquid chromatography; IS, Internal standard; LC, Liquid chromatography.

Acknowledgment

The funding of this research work was supported by the Buffalo Development Project (Component B), Bangladesh Livestock Research Institutes, Savar, Dhaka, Bangladesh. The authors are also grateful to Professor Shankar Majumder, Department of Agricultural and Applied Statistics, Bangladesh Agricultural University,

Mymensingh-2202, Bangladesh, for his enormous efforts on the statistical part of this manuscript.

Conflicts of interest

The authors declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.

Author contributions

Conceptualization: Islam MA, Deb GK, Rashid MH. Data curation: Asif AHM, Shahjadee UF, Lisa SA, Ahmed S, Sarker MAH. Formal analysis: Islam MA, Asif AHM, Arefin S, Bari MS, Islam MZ, Sarker MAH. Validation: Shahjadee UF, Lisa SA, Ahmed S, Asif AHM, Sarker MAH. Methodology: Shahjadee UF, Lisa SA, Ahmed S, Asif AHM. Investigation: Asif AHM, Sarker MAH, Deb GK, Siddiki MRS, Islam MA. Writing -original draft: Sarker MAH, Habib MR, Arefin S, Bari MS, Islam MZ. Writing -review & editing: Habib MR, Siddiki MRS, Arefin S, Bari MS, Islam MZ, Rashid MH, Islam MA.

References

- [1] Drewnowski A. The contribution of milk and milk products to micronutrient density and affordability of the US diet. *J Am Coll Nutr* 2011; 30:422–8; <https://doi.org/10.1080/07315724.2011.10719986>
- [2] Ménard O, Ahmad S, Rousseau F, Briard-Bion V, Gaucheron F, Lopez C. Buffalo vs. cow milk fat globules: size distribution, zeta-potential, compositions in total fatty acids and in polar lipids from the milk fat globule membrane. *Food Chem* 2010; 120(2):544–51; <https://doi.org/10.1016/j.foodchem.2009.10.053>
- [3] Vieira RG, FalcãoFilho RS, Duarte TF, Pessoa TRB, Queiroga RC, Moreira RT. Acceptability and sensory preference of cheese curds elaborated with female buffalo, goat and cow milk. *Rev Inst Laticín Când Tost* 2013; 36(3):12–6.
- [4] Islam MA, Alam MK, Islam MN, Khan MA, Ekeberg D, Rukke EO. Principal milk components in buffalo, holstein cross, indigenous cattle and red chittagong cattle from Bangladesh. *Asian-Australas J Anim Sci* 2014; 27(6):886–97; <https://doi.org/10.5713/ajas.2013.13586>
- [5] Walstra P, Wouters JTM, Geurts TJ. Principles of cheese making. In: Walstra P, Wouters JTM, Geurts TJ (eds.). *Dairy science and technology*. 2nd edition, CRC Press, Taylor and Francis Group, Boca Raton, FL, pp 577–82, 2006; <https://doi.org/10.1201/9781420028010>
- [6] Fox PF, Guinee TP, Cogan TM, Mc Sweeney PLH. Factors that affect cheese quality. In: Fox PF, Guinee TP, Cogan TM, Mc Sweeney PLH (eds.). *Fundamentals of cheese science*. Springer, Boston, MA, pp 533–42, 2017; https://doi.org/10.1007/978-1-4899-7681-9_15
- [7] Aldai N, de Renobales M, Barron L, Kramer J. What are the trans-fatty acids issues in foods after discontinuation of industrially produced trans fats? Ruminant products, vegetable oils, and synthetic supplements. *Eur J Lipid Sci* 2013; 115(12):1378–401; <https://doi.org/10.1002/ejlt.201300072>
- [8] Mehta BM. Butter, butter oil, and ghee. *Gourmet and health-promoting specialty oils*. AOCS Press, Elsevier Inc, Gujarat, India. pp 527–59, 2009; <https://doi.org/10.1016/B978-1-893997-97-4.50027-9>
- [9] Ullah R, Khan S, Ali H, Bilal M, Saleem M. Identification of cow and buffalo milk based on Beta carotene and vitamin-A concentration using fluorescence spectroscopy. *PLoS One* 2017; 12(5):e0178055; <https://doi.org/10.1371/journal.pone.0178055>
- [10] Schönfeldt H, Hall N, Smit L. The need for country specific composition data on milk. *Food Res Int* 2012; 47(2):207–9; <https://doi.org/10.1016/j.foodres.2011.05.018>
- [11] Tadjine D, Boudalia S, Bousbia A, Khelifa R, Mebirouk L, Tadjine BA, et al. Pasteurization effects on yield and physicochemical parameters of cheese in cow and goat milk. *Food Sci Technol* 2020; 40(3):580–7; <https://doi.org/10.1590/fst.13119>
- [12] Sant'Ana A, Bezerril F, Madruga M. Nutritional and sensory characteristics of Minas fresh cheese made with goat milk, cow milk, or a mixture of both. *J Dairy Sci* 2013; 96(12):7442–53; <https://doi.org/10.3168/jds.2013-6915>
- [13] Bennato F, Ianni A, Innosa D, Martino C, Grotta L, Pomilio F, et al. Influence of licorice root feeding on chemical-nutritional quality of cow milk and straciatina cheese, an Italian Traditional fresh dairy product. *Animals* 2019; 9(12):1153; <https://doi.org/10.3390/ani9121153>
- [14] Jesmin MM, Sarker MAH, Islam MA, Rashid MH, Siddiki MSR. Use of papaya plant (*Carica papaya*) latex for making cottage cheese from cow milk. *J Bangl Agric Univ* 2021; 19(1):85–90; <https://doi.org/10.5455/JBAU.13262>
- [15] Bas D, Kendirci P, Salum P, Govce G, Erbay Z. Production of enzyme-modified cheese (EMC) with ripened white cheese flavour: I-effects of proteolytic enzymes and determination of their appropriate combination. *Food Bioprod Process* 2019; 117:287–301; <https://doi.org/10.1016/j.fbp.2019.07.016>
- [16] Murtaza MA, Rehman SU, Anjum FM, Huma N. Descriptive sensory profile of cow and buffalo milk cheddar cheese prepared using indigenous cultures. *J Dairy Sci* 2013; 96:1380–6; <http://dx.doi.org/10.3168/jds.2012-5992>
- [17] Lopes FCF, Ribeiro CGS, Rodriguez NM, da Gama MAS, Morenz MJF, Antoniassi R, et al. Butter fatty acid composition as a function of soybean oil supplementation and time of milking, and performance of Holstein x Gyr cows fed with chopped elephant grass-based diets. *Semin Ciênc Agrár* 2019; 40(5):2027–44; <http://dx.doi.org/10.5433/1679-0359.2019v40n5p2025>
- [18] Rashid S. Nutritional evaluation of butter prepared from fresh adulterants free cow milk. *Pure Appl Biol* 2017; 6(4):1366–71; <https://doi.org/10.19045/bspab.2017.600147>
- [19] Parmar NB, Mehta MB, Amaranth KD. Composition of ghee prepared from camel, cow and buffalo milk. *J Camel Pract Res* 2018; 25(3):321–6. <https://doi.org/10.5958/2277-8934.2018.00046.2>
- [20] Parmar P, Khamrui K. Development of process for the production of arjuna herbal ghee from buffalo milk. *Indian J Anim Sci* 2017; 87(2):203–7.
- [21] Bhatia P, Sharma V, Arora S, Rao P. Effect of cholesterol removal on compositional and the physicochemical characteristics of anhydrous cow milk fat (cow ghee). *Int J Food Prop* 2019; 22(1):1–8; <https://doi.org/10.1080/10942912.2018.1564762>
- [22] Habib MR, Islam MZ, Bari MS, Sarker MAH, Rashid MH, Islam MA. Effect of concentrate supplementation during transition period on production and reproduction of indigenous buffalo. *Bang J Anim Sci* 2020; 49:83–90; <https://doi.org/10.3329/bjas.v49i1.49385>
- [23] AOAC. Official methods of analysis of AOAC International. 17th edition, AOAC International, Gaithersburg, MD, 2000.
- [24] Aggarwala AC, Sharma RC. A laboratory manual of milk inspection. 4th edition, Asia Publishing House, Kolkata, India, 1961.
- [25] Rana M, Hoque M, Deb G, Nahar T, Habib R, Siddiki M. Preparation of cottage cheese from buffalo milk with different levels of papaya latex. *Bang J Anim Sci* 2017; 46(2):128–33; <https://doi.org/10.3329/bjas.v46i2.34443>
- [26] Huang T, Chen C, Wefler V, Raftery A. A Stable reagent for the Liebermann-burchard reaction. application to rapid serum cholesterol determination. *Anal Chem* 1961; 33(10):1405–7; <https://doi.org/10.1021/ac60178a040>

- [27] Ulbricht TLV, Southgate DAT. Coronary heart disease: seven dietary factors. *Lancet* 1991; 338(8773):985–99; [https://doi.org/10.1016/0140-6736\(91\)91846-m](https://doi.org/10.1016/0140-6736(91)91846-m)
- [28] Shimadzu. Method of Shimadzu HPLC amino acid analysis system. In: Amino acid analysis system instruction manual, Chapter 6, Shimadzu Corporation Publisher, Kyoto, Japan, 1993.
- [29] Hurtaud C, Faucon F, Couvreur S, Peyraud J. Linear relationship between increasing amounts of extruded linseed in dairy cow diet and milk fatty acid composition and butter properties. *J Dairy Sci* 2010; 93(4):1429–43; <https://doi.org/10.3168/jds.2009-2839>
- [30] Bergamo P. Fat-soluble vitamin contents and fatty acid composition in organic and conventional Italian dairy products. *Food Chem* 2003; 82(4):625–31; [https://doi.org/10.1016/s0308-8146\(03\)00036-0](https://doi.org/10.1016/s0308-8146(03)00036-0)
- [31] BSTI. Bangladesh standard specification for dairy products, Bangladesh Standards and Testing Institution, Dhaka, Bangladesh, 2019.
- [32] Sbihi H, Nehdi I, Tan C, Al-Resayes S. Characteristics and fatty acid composition of milk fat from Saudi Aradi goat. *Gras Aceit* 2015; 66(4):e101; <https://doi.org/10.3989/gya.0233151>
- [33] Gandhi K, Upadhyay N, Aghav AD, Sharma V, Lal D. Detection of adulteration of ghee (clarified milk fat) with palmolein and sheep body fat using Reichert-Meissl (RM) value coupled with solvent fractionation technique. *Ind J Dairy Sci* 2014; 67:387–93.
- [34] Ayari F, Mirzaee- Ghaleh E, Rabbani H, Heidarbeigi K. Detection of the adulteration in pure cow ghee by electronic nose method (case study: sunflower oil and cow body fat). *Int J Food Prop* 2018; 21(1):1670–9; <https://doi.org/10.1080/10942912.2018.1505755>
- [35] Bhatia P, Sharma V, Arora S, Rao P. Effect of cholesterol removal on compositional and the physicochemical characteristics of anhydrous cow milk fat (cow ghee). *Int J Food Prop* 2019; 22(1):1–8; <https://doi.org/10.1080/10942912.2018.1564762>
- [36] Ahmad S, Anjum FM, Huma N, Sameen A, Zahoor T. Composition and physico-chemical characteristics of buffalo milk with particular emphasis on lipids, proteins, minerals, enzymes and vitamins. *J Anim Plant Sci* 2013; 23:62–74.
- [37] Cinar MU, Özsoy T, Beyzi SB, Kaliber M, Konca Y. Milk and fatty acid composition of Anatolian water buffalo (*Bubalus bubalis*) from different provinces. *Buff Bull* 2019; 38:107–18.
- [38] Meremäe K, Roasto M, Kuusik S, Ots M, Henno M. Trans fatty acid contents in selected dietary fats in the Estonian market. *J Food Sci* 2012; 77:163–8; <https://doi.org/10.1111/j.1750-3841.2012.02829.x>
- [39] Perdomo L, Beneit N, Otero Y. Protective role of oleic acid against cardiovascular insulin resistance and in the early and late cellular atherosclerotic process. *Cardiovasc Diabetol* 2015; 14:Article number 75; <https://doi.org/10.1186/s12933-015-0237-9>
- [40] Bobe G, Hammond E, Freeman A, Lindberg G, Beitz D. Texture of butter from cows with different milk fatty acid compositions. *J Dairy Sci* 2003; 86(10):3122–7; [https://doi.org/10.3168/jds.s0022-0302\(03\)73913-7](https://doi.org/10.3168/jds.s0022-0302(03)73913-7)
- [41] Allred S, Dhiman T, Brennand C, Khanal R, McMahon D, Luchini N. Milk and cheese from cows fed calcium salts of palm and fish oil alone or in combination with soybean products. *J Dairy Sci* 2006; 89(1):234–48; [https://doi.org/10.3168/jds.s0022-0302\(06\)72088-4](https://doi.org/10.3168/jds.s0022-0302(06)72088-4)
- [42] Eckel RH, Jakicic JM, Ard JD, De Jesus JM, Hubbard VS, Lee IM, et al. AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American college of cardiology/American heart association task force on practice guidelines. *Circulation* 2013; 129:129–75; <https://doi.org/10.1161/01.cir.0000437740.48606.d1>
- [43] Salum P, Govce G, Kendirci P, Bas D, Erbay Z. Composition, proteolysis, lipolysis, volatile compound profile and sensory characteristics of ripened white cheeses manufactured in different geographical regions of Turkey. *Int Dairy J* 2018; 87:26–36; <https://doi.org/10.1016/j.idairyj.2018.07.011>
- [44] Inoue K, Shirai T, Ochiai H, Kasao M, Hayakawa K, Kimura M, et al. Blood pressure lowering effect of a novel fermented milk containing g amino butyric acid (GABA) in mild hypertensives. *Eur J Clin Nutr* 2003; 57:490–5; <https://doi.org/10.1038/sj.ejcn.1601555>
- [45] Hagiwara H, Seki T, Ariga T. The effect of pre-germinated brown rice intake on blood glucose and PAI-1 levels in streptozotocin-induced diabetic rats. *Biosci Biotechnol Biochem* 2004; 68:444–7; <https://doi.org/10.1271/bbb.68.444>