

The Role of Focal Adhesion Kinase in the TGF- β -Induced Myofibroblast Transdifferentiation of Human Tenon's Fibroblasts

Samin Hong, Jong Bok Lee, Yoko Iizuka, Yoo Kyung Song, Gong Je Seong, Sueng-Han Han

Institute of Vision Research, Department of Ophthalmology, Yonsei University College of Medicine, Seoul, Korea

Purpose: To investigate the role of focal adhesion kinase (FAK) in transforming growth factor (TGF)- β -induced myofibroblast transdifferentiation of human Tenon's fibroblasts.

Methods: Primary cultured human Tenon's fibroblasts were exposed to TGF- β 1 for up to 48 hours. The mRNA levels of FAK, α smooth muscle actin (α SMA), and β -actin were determined by quantitative real time reverse transcription polymerase chain reaction. The protein levels of collagen type I, FAK, phospho-FAK, α SMA, and β -actin were determined by Western immunoblots. After the small interfering RNA targeting FAK (siRNA_{FAK}) molecules were delivered into the cells, the expressions of α SMA proteins were determined by Western immunoblots.

Results: In human Tenon's fibroblasts, TGF- β 1 significantly increased the mRNA and protein expressions of α SMA. However, when the action of FAK was inhibited using siRNA_{FAK}, the TGF- β 1-induced expression of α SMA was attenuated.

Conclusions: Our data suggest that FAK may be associated with the TGF- β 1-induced transdifferentiation of human Tenon's fibroblasts to myofibroblasts, which is the essential step of subconjunctival fibrosis.

Key Words: Fibroblast, Focal adhesion protein-tyrosine kinases, Myofibroblast, Transforming growth factors

Myofibroblast transdifferentiation of activated fibroblasts is an essential step of the fibrotic process in most living tissues. Myofibroblasts have characteristics that are taken from both smooth muscle cells and fibroblasts; they express the contractile phenotype of α smooth muscle actin (α SMA) and synthesize extracellular matrix proteins including collagens [1-3]. In subconjunctival fibrosis, the myofibroblast transdifferentiation of Tenon's fibroblasts is known to be mainly caused by transforming growth factor (TGF)- β [4-6]. Many ophthalmic researchers have sought to identify an anti-fibrotic mechanism by inhibiting TGF- β signaling [7-10].

Focal adhesion kinase (FAK), a cytosolic protein tyrosine kinase, provides scaffolding functions at sites of integrin adhesion and promotes cell migration [11,12].

The kinase modulates several basic processes in normal development and cancer metastasis. In addition, it also plays a crucial role in the transdifferentiation of fibroblasts to myofibroblasts induced by TGF- β [13-15]. However, because the precise action of the kinase is somewhat variable based on the tissue [14,15], the role of FAK in myofibroblast transdifferentiation in human Tenon's fibroblasts must be confirmed.

Using primary cultured human Tenon's fibroblasts, the role of FAK in TGF- β -induced myofibroblast transdifferentiation of human Tenon's fibroblasts was investigated in the present study.

Materials and Methods

Cell culture and exposure to transforming growth factor- β 1

After the Institutional Review Board approved our protocol, selected patients received comprehensive information and provided written consent for inclusion. In compliance with the tenets of the Declaration of Helsinki, small human Tenon's capsule specimens were obtained during

Received: November 17, 2010 Accepted: January 1, 2011

Corresponding Author: Sueng-Han Han, MD, PhD. Department of Ophthalmology, Yonsei University College of Medicine, #211 Eonju-ro, Gangnam-gu, Seoul 135-720, Korea. Tel: 82-2-2019-3440, Fax: 82-2-3463-1049, E-mail: shhan222@yuhs.ac

© 2012 The Korean Ophthalmological Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1. Primer sequences for reverse transcription polymerase chain reaction

Gene name	Type	Sequence
FAK	Forward	5'-GAA GCA TTG GGT CGG GAA CTA-3'
	Reverse	5'-CTC AAT GCA GTT TGG AGG TGC-3'
α SMA	Forward	5'-GTG TTA TGT AGC TCT GGA CTT TGA AAA-3'
	Reverse	5'-GGC AGC GGA AAC GTT CAT T-3'
β -actin	Forward	5'-GCG GGA AAT CGT GCG TGA CAT T-3'
	Reverse	5'-GAT GGA GTT GAA GGT AGT TTC GTG-3'

FAK = focal adhesion kinase; α SMA = α smooth muscle actin.

strabismus surgery, and primary human Tenon's fibroblasts were cultured from the explants. Cells were incubated in Dulbecco's modified Eagle's medium (DMEM; Life Technologies, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (Life Technologies), 100 units/mL penicillin, and 100 μ g/mL streptomycin (Life Technologies) at 37°C in 5% CO₂.

After 12 hours of incubation in serum-free media, the cells were treated with 5 ng/mL of recombinant human TGF- β 1 (R&D Systems Inc., Minneapolis, MN, USA) for up to 48 hours. For the control group, the same volume of DMEM was added to the media instead of TGF- β 1.

Small interfering RNAs targeting focal adhesion kinase

Small interfering RNA molecules targeting FAK mRNA (siRNA_{FAK}) were purchased from Santa Cruz Biotechnology Inc. (Santa Cruz, CA, USA) and were delivered into cells according to the manufacturer's instructions. Briefly, the fibroblasts were incubated in media containing 50 nM of siRNA_{FAK} and 2 μ g/mL of Lipofectamine 2000 (Life Technologies) for 16 hours at 37°C in a CO₂ incubator.

Real time reverse transcription polymerase chain reaction

Total RNA was extracted using an RNeasy Mini Kit (Qiagen, Venlo, Netherlands), and cDNAs were synthesized using the SuperScript III First-Strand Synthesis System (Life Technologies). Real-time polymerase chain reaction (PCR) was performed with 50 ng cDNA per reaction using 25 μ L of iQ SYBR Green Supermix (Bio-Rad Laboratories, Hercules, CA, USA) containing 500 nM of specific primers (Table 1) in the iCycler iQ Real-Time PCR Detection System (Bio-Rad Laboratories). The SYBR green data were analyzed with a relative standard curve of β -actin.




Fig. 1. Quantitative reverse transcription polymerase chain reaction for focal adhesion kinase (FAK) and α smooth muscle actin (α SMA) after exposure to 5 ng/mL of transforming growth factor- β 1 for up to 16 hours. The level of target mRNA was calculated using a relative standard curve of β -actin and expressed as the mean \pm SEM. * $p < 0.05$.

Western immunoblots

Whole cellular proteins were extracted from the cells, and equal amounts of protein (10 μ g) were resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis. The proteins were transferred to polyvinylidene fluoride membranes and probed overnight with primary antibodies against human collagen type I, FAK, phospho-FAK (pFAK), α SMA, and β -actin (Santa Cruz Biotechnology). Immunoreactive bands were detected with horseradish peroxidase-conjugated secondary antibodies (Santa Cruz Biotechnology) and visualized with an enhanced chemiluminescent system.

Statistical analysis

Quantitative reverse transcription (RT)-PCR data is expressed as mean \pm SEM and compared with the Kruskal-Wallis one-way analysis of variance using the MedCalc ver. 11.4.2 (MedCalc Software bvba, Mariakerke, Belgium). Image analysis of the immunobands was performed using the ImageJ ver. 1.43u (National Institutes of Health, Bethesda, MD, USA). A p -value less than 0.05 was consid-




Fig. 2. Representative Western immunoblots and densitometric data for collagen type I (190 kDa), focal adhesion kinase (FAK, 125 kDa), phospho-FAK (pFAK, 125 kDa), and β -actin (43 kDa) in human Tenon's fibroblasts stimulated by 5 ng/mL of transforming growth factor- β 1 for up to 48 hours.




Fig. 3. Representative Western immunoblots and densitometric data for α smooth muscle actin (43 kDa) in human Tenon's fibroblasts stimulated by 5 ng/mL of transforming growth factor (TGF)- β 1 for up to 48 hours with or without the treatment of siRNA targeting FAK (siRNA_{FAK}) molecules.

ered statistically significant.

Results

The quantitative data of RT-PCR for FAK and α SMA are presented in Fig. 1. In the human Tenon's fibroblasts, the 16-hour treatment with TGF- β 1 significantly increased the mRNA levels of α SMA (5.11 ± 1.37 times to control, $p = 0.023$) but not of FAK (1.16 ± 0.06 times to control, $p = 0.204$).

On the Western immunoblots, the TGF- β 1 treatment increased the protein expression of FAK as well as that of collagen type I. Expression of pFAK, an active form of FAK, was also increased. These changes were observed after 12 hours. Representative immunobands for collagen type I, FAK, pFAK, and β -actin are shown in Fig. 2.

Regarding α SMA, though serum starvation itself slightly induced the expression of α SMA, the exposure to TGF- β 1 for 48 hours stimulated α SMA expression in human Tenon's fibroblasts. When the siRNA_{FAK} molecules were introduced into the cells, the TGF- β 1-induced α SMA expression was attenuated. Representative immunobands for α SMA in the

control group, TGF- β 1 only treatment group, and siRNA_{FAK}/TGF- β 1 treatment group are shown in Fig. 3.

Discussion

FAK, also known as protein tyrosine kinase 2, participates in the focal adhesion between the cytoskeleton and extracellular matrix and elicits intracellular signal transductions associated with cell migration and survival [11,12,16-18]. It is activated by autophosphorylation via Smad- and p38 MAPK-dependent mechanisms [19-22] and plays a key role in cancer metastasis as well as normal development [11,12]. Even though FAK is also linked to the myofibroblast transdifferentiation of fibroblasts in response to TGF- β 1 [13-15], the precise role of the kinase has not been well studied in the fibrotic process in the subconjunctival space. Excessive subconjunctival fibrosis causes ocular morbidity in not only patients with oculocutaneous disorders, but also patients who have undergone strabismus surgery and/or glaucoma-filtering surgery. A better understanding and modulation of FAK signaling in this process may result in a novel therapeutic strategy for those patients.

Using primary cultured human Tenon's fibroblasts, we investigated the role of FAK in TGF- β -induced myofibroblast transdifferentiation in the present study and found that silencing of FAK action using siRNA_{FAK} duplex significantly attenuated the α SMA expression induced by TGF- β 1 in human Tenon's fibroblasts. Our data implies that FAK may participate in the myofibroblast transdifferentiation of those cells.

Myofibroblasts contain characteristics found in smooth muscle cells and fibroblasts in that they possess a contractile phenotype of α SMA and synthesize extracellular matrix proteins [1-3]. Since they play a crucial role in most fibrotic responses, myofibroblast transdifferentiation from activated fibroblasts is an essential step in the fibrotic process. In the subconjunctival space, TGF- β typically initiates this myofibroblast transdifferentiation of fibroblasts [4-6].

On quantitative RT-PCR, the mRNA expression of α SMA was dramatically increased by TGF- β 1, but FAK was not significantly altered. However, on Western immunoblots, the protein levels of both FAK and pFAK were minimally stimulated by TGF- β 1 treatment. The increase in FAK seems to result from an increase in translation rather than transcription.

Currently, many researchers have tried to develop anti-TGF- β neutralizing antibodies to modulate postoperative scarring in patients undergoing ocular surface surgery [23-25]. FAK, as a participant in TGF- β -associated intracellular signaling of myofibroblast transdifferentiation from fibroblasts, might also be valuable as a novel therapeutic

strategy for anti-fibrosis.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

Acknowledgements

This work was supported by a Faculty Research Grant from the Yonsei University College of Medicine, Seoul, Republic of Korea (6-2008-0274).

References

1. Eyden B. The myofibroblast: an assessment of controversial issues and a definition useful in diagnosis and research. *Ultrastruct Pathol* 2001;25:39-50.
2. Eyden B. Electron microscopy in the study of myofibroblastic lesions. *Semin Diagn Pathol* 2003;20:13-24.
3. Desmouliere A, Guyot C, Gabbiani G. The stroma reaction myofibroblast: a key player in the control of tumor cell behavior. *Int J Dev Biol* 2004;48:509-17.
4. Meyer-Ter-Vehn T, Grehn F, Schlunck G. Localization of TGF-beta type II receptor and ED-A fibronectin in normal conjunctiva and failed filtering blebs. *Mol Vis* 2008;14:136-41.
5. Saika S, Yamanaka O, Okada Y, et al. TGF beta in fibroproliferative diseases in the eye. *Front Biosci (Schol Ed)* 2009;1:376-90.
6. Seong GJ, Hong S, Jung SA, et al. TGF-beta-induced interleukin-6 participates in transdifferentiation of human Tenon's fibroblasts to myofibroblasts. *Mol Vis* 2009;15:2123-8.
7. Nakamura H, Siddiqui SS, Shen X, et al. RNA interference targeting transforming growth factor-beta type II receptor suppresses ocular inflammation and fibrosis. *Mol Vis* 2004;10:703-11.
8. Meyer-Ter-Vehn T, Gebhardt S, Sebald W, et al. p38 inhibitors prevent TGF-beta-induced myofibroblast transdifferentiation in human tenon fibroblasts. *Invest Ophthalmol Vis Sci* 2006;47:1500-9.
9. Yamanaka O, Saika S, Ikeda K, et al. Interleukin-7 modulates extracellular matrix production and TGF-beta signaling in cultured human subconjunctival fibroblasts. *Curr Eye Res* 2006;31:491-9.
10. Meyer-Ter-Vehn T, Katzenberger B, Han H, et al. Lovastatin inhibits TGF-beta-induced myofibroblast transdifferentiation in human tenon fibroblasts. *Invest Ophthalmol Vis Sci* 2008;49:3955-60.
11. Chatzizacharias NA, Kouraklis GP, Theocharis SE. The role of focal adhesion kinase in early development. *Histol Histopathol* 2010;25:1039-55.
12. Schwock J, Dhani N, Hedley DW. Targeting focal adhesion kinase signaling in tumor growth and metastasis. *Expert Opin Ther Targets* 2010;14:77-94.
13. Thannickal VJ, Lee DY, White ES, et al. Myofibroblast differentiation by transforming growth factor-beta1 is dependent on cell adhesion and integrin signaling via focal adhesion kinase. *J Biol Chem* 2003;278:12384-9.
14. Mimura Y, Ihn H, Jinnin M, et al. Constitutive phosphorylation of focal adhesion kinase is involved in the myofibroblast differentiation of scleroderma fibroblasts. *J Invest Dermatol* 2005;124:886-92.
15. Brenmoehl J, Miller SN, Hofmann C, et al. Transforming growth factor-beta 1 induces intestinal myofibroblast differentiation and modulates their migration. *World J Gastroenterol* 2009;15:1431-42.
16. McLean GW, Carragher NO, Avizienyte E, et al. The role of focal-adhesion kinase in cancer: a new therapeutic opportunity. *Nat Rev Cancer* 2005;5:505-15.
17. Cox BD, Natarajan M, Stettner MR, Gladson CL. New concepts regarding focal adhesion kinase promotion of cell migration and proliferation. *J Cell Biochem* 2006;99:35-52.
18. Liu S, Xu SW, Kennedy L, et al. FAK is required for TGF-beta-induced JNK phosphorylation in fibroblasts: implications for acquisition of a matrix-remodeling phenotype. *Mol Biol Cell* 2007;18:2169-78.
19. Horowitz JC, Rogers DS, Sharma V, et al. Combinatorial activation of FAK and AKT by transforming growth factor-beta1 confers an anoikis-resistant phenotype to myofibroblasts. *Cell Signal* 2007;19:761-71.
20. Parsons CJ, Takashima M, Rippe RA. Molecular mechanisms of hepatic fibrogenesis. *J Gastroenterol Hepatol* 2007;22 Suppl 1:S79-84.
21. Walsh MF, Ampasala DR, Hatfield J, et al. Transforming growth factor-beta stimulates intestinal epithelial focal adhesion kinase synthesis via Smad- and p38-dependent mechanisms. *Am J Pathol* 2008;173:385-99.
22. Walsh MF, Ampasala DR, Rishi AK, Basson MD. TGF-beta1 modulates focal adhesion kinase expression in rat intestinal epithelial IEC-6 cells via stimulatory and inhibitory Smad binding elements. *Biochim Biophys Acta* 2009;1789:88-98.
23. Cordeiro MF. Technology evaluation: lerdelimumab, Cambridge Antibody Technology. *Curr Opin Mol Ther* 2003;5:199-203.
24. Mead AL, Wong TT, Cordeiro MF, et al. Evaluation of anti-TGF-beta2 antibody as a new postoperative anti-scarring agent in glaucoma surgery. *Invest Ophthalmol Vis Sci* 2003;44:3394-401.
25. CAT-152 0102 Trabeculectomy Study Group, Khaw P, Grehn F, et al. A phase III study of subconjunctival human anti-transforming growth factor beta(2) monoclonal antibody (CAT-152) to prevent scarring after first-time trabeculectomy. *Ophthalmology* 2007;114:1822-30.